The Prognostic and Predictive Value of *ESR1* Fusion Gene Transcripts in Primary Breast Cancer

Silvia R. Vitale

University of Catania, Department of Clinical and Experimental Medicine and Center of Exper. Onco. and Hem., A.O.U. Policlinico "G. Rodolico- San Marco", Catania

Kirsten Ruigrok-Ritstier

Erasmus MC Cancer Institute, Erasmus University Medical Center, Department of Medical Oncology. Rotterdam

A. Mieke Timmermans

Erasmus MC Cancer Institute, Erasmus University Medical Center, Department of Medical Oncology. Rotterdam

Renée Foekens

Erasmus MC Cancer Institute, Erasmus University Medical Center, Department of Medical Oncology. Rotterdam

Anita M.A.C. Trapman-Jansen

Erasmus MC Cancer Institute, Erasmus University Medical Center, Department of Medical Oncology. Rotterdam

Corine M. Beaufort

Erasmus MC Cancer Institute, Erasmus University Medical Center, Department of Medical Oncology. Rotterdam

Paolo Vigneri

University of Catania, Department of Clinical and Experimental Medicine and Center of Exper. Onco. and Hem., A.O.U. Policlinico "G. Rodolico- San Marco", Catania

Stefan Sleijfer

Erasmus MC Cancer Institute, Erasmus University Medical Center, Department of Medical Oncology. Rotterdam

John W.M. Martens

Erasmus MC Cancer Institute, Erasmus University Medical Center, Department of Medical Oncology and Cancer Genomics Netherlands. Rotterdam

Anieta M. Sieuwerts

Erasmus MC Cancer Institute, Erasmus University Medical Center, Department of Medical Oncology and Cancer Genomics Netherlands. Rotterdam

Maurice P.H.M. Jansen (✉ m.p.h.m.jansen@erasmusmc.nl)

Erasmus MC Cancer Institute, Erasmus University Medical Center, Department of Medical Oncology. Rotterdam

Research Article

Keywords: fusion genes, ESR1, CCDC170, breast cancer, prognosis, RT-qPCR

Posted Date: December 1st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1110533/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at BMC Cancer on February 12th, 2022. See the published version at <https://doi.org/10.1186/s12885-022-09265-1>.

Abstract

Background: In breast cancer (BC), recurrent fusion genes of estrogen receptor alpha (*ESR1*) and *AKAP12*, *ARMT1* and *CCDC170* have been reported. In these gene fusions the ligand binding domain of *ESR1* has been replaced by the transactivation domain of the fusion partner constitutively activating the receptor. As a result, these gene fusions can drive tumor growth hormone independently as been shown in preclinical models, but the clinical value of these fusions have not been reported. Here, we studied the prognostic and predictive value of different frequently reported *ESR1* fusion transcripts in primary BC.

Methods: We evaluated 732 patients with primary BC (131 *ESR1*-negative and 601 *ESR1*-positive cases), including two ER-positive BC patient cohorts: one cohort of 322 patients with advanced disease who received first-line endocrine therapy (ET) (predictive cohort), and a second cohort of 279 patients with lymph node negative disease (LNN) who received no adjuvant systemic treatment (prognostic cohort). Fusion gene transcript levels were measured by reverse transcriptase quantitative PCR. The presence of the different fusion transcripts was associated, in uni- and multivariable Cox regression analysis taking along current clinic-pathological characteristics, to progression free survival (PFS) during first-line endocrine therapy in the predictive cohort, and disease-free survival (DFS) and overall survival (OS) in the prognostic cohort.

Results: The *ESR1-CCDC170* fusion transcript was present in 27.6% of the *ESR1*-positive BC subjects and in 2.3% of the *ESR1*-negative cases. In the predictive cohort, none of the fusion transcripts were associated with response to first-line ET. In the prognostic cohort, the median DFS and OS were respectively 37 and 93 months for patients with an *ESR1-CCDC170* exon 8 gene fusion transcript and respectively 91 and 212 months for patients without this fusion transcript. In a multivariable analysis, this *ESR1-CCDC170* fusion transcript was an independent prognostic factor for DFS (HR) (95% confidence interval (CI): 1.8 (1.2–2.8), $P=0.005$) and OS (HR (95% CI: 1.7 (1.1–2.7), $P=0.023$).

Conclusions: Our study shows that in primary BC only *ESR1-CCDC170* exon 8 gene fusion transcript carries prognostic value. None of the *ESR1* fusion transcripts, which are considered to have constitutive ER activity, was predictive for outcome in BC with advanced disease treated with endocrine treatment.

Background

The estrogen receptor (ER) plays a key role in cellular growth and tumor development in a large fraction of breast cancers. As a result, endocrine therapy has been and still is a successful treatment in patients with *ESR1*-positive (*ESR1+*) breast cancers (BC) [1]. However, in the metastatic setting, nearly half of the patients are *de novo* resistant to endocrine therapy while the remaining cases acquire resistance over time [2, 3]. One of the primary characterized mechanisms of acquired resistance to endocrine therapy is the acquisition of mutations within the ligand-binding domain (LBD) of the estrogen receptor alpha gene (*ESR1*) activating the receptor constitutively thereby rendering tumor cells less dependent on estrogen [4–7]. Another mechanism that lead to less estrogen dependency of BC cells is the occurrence of *ESR1* fusion proteins. Through analysis of RNA-sequencing data in breast cancer, recurrent intragenic fusions of 5' end of *ESR1* and the 3' ends of *AKAP12*, *ARMT1* or *CCDC170* have been identified [8–13]. Gene fusions were preferentially detected in high-grade disease and/or endocrine-resistant forms of *ESR1+* BC [10, 13]. Despite the diversity among these fusions, they share a common structure retaining the hormone-independent transactivation domain as well as the DNA-binding domain whereas their ligand-binding domain is lost and replaced with a functional (transactivating) domain of the fusion partner, suggesting a pathological impact in *ESR1+* BC [13]. However, the clinical significance of these fusions has not yet been properly addressed in uniform and well annotated cohorts.

In this study, we explored the occurrence of fusion transcripts of three of the most commonly reported fusion partners of *ESR1* (i.e. *CCDC170*, *AKAP12* and *ARMT1*) and determined the associations of their presence with clinical outcome in a cohort of 732 breast cancer patients allowing us to investigate their predictive value for endocrine treatment failure as well as their prognostic value.

Methods

Study Cohorts

The protocol to study biological markers associated with disease outcome was approved by the medical ethics committee of the Erasmus Medical Centre Rotterdam, The Netherlands (MEC 02.953) and was performed in accordance with the Code of Conduct of the Federation of Medical Scientific Societies in The Netherlands (<https://www.federa.org/codes-conduct>). The use of coded left-over material for scientific purposes and, therefore, for the greater good, does not require informed consent according to Dutch law and the new European general data protection regulation (GDPR).

In this retrospective study (see Fig. 1A for the consort diagram of the study), female patients were included, who underwent surgery for invasive primary breast cancer between 1980 and 2000 in the Netherlands. A further selection criterion was no previously diagnosed cancers with the exception of basal cell carcinoma or stage Ia/Ib cervical cancer. Within this study, only data from sections of primary tumors with at least 30% invasive tumor cells were included. The details of tissue processing, RNA isolation, cDNA synthesis and QC of this cohort have been described previously [14, 15]. Tumor grade was assessed according to standard procedures at the time of inclusion. For the classification of patients' RNA samples regarding expression of the estrogen and progesterone receptors, as well as the human epidermal growth factor receptor 2 (HER2) amplification status, reverse transcriptase quantitative PCR (RT-qPCR) was used with cut-offs previously described by us [14, 15].

The total cohort consisted of 732 patients with primary breast cancer (131 *ESR1*-negative and 601 *ESR1*-positive cases) (Fig. 1B). The clinical relevance of the gene fusion transcripts was evaluated in a predictive and a prognostic cohort of *ESR1+* BC patients.

The predictive cohort consisted of 322 breast cancer patients with *ESR1+* primary tumors of which 235 patients received tamoxifen (40 mg daily) and 87 patients an aromatase inhibitor (AI: anastrozole, letrozole, exemestane [16]) as a 1st-line treatment for recurrent disease. Clinical response to tamoxifen

therapy was defined as previously described [14, 17]. The prognostic cohort included primary tumors from 279 lymph node negative (LNN) *ESR1*+ BC patients who had not received any systemic (neo) adjuvant therapy. Of note, 122 of these LNN *ESR1*+ patients were also included in the predictive cohort. Clinicopathological characteristics of each of these 2 cohorts are described in Table 1. Association of *ESR1* fusions with clinical parameters of patients enrolled in the predictive cohort and in the prognostic cohort are reported in Table 2 and Table 3, respectively.

Table 1
Clinicopathological characteristics of ESR1-positive breast cancer patient cohorts

	Predictive Endocrine Therapy Cohorts		Prognostic Cohort
	Tamoxifen	Aromatase inhibitors	Lymph node negative (LNN)
Total	235	87	279
Median age (range)	61 (29-90)	66 (35-86)	55 (26-85)
Menopausal Status:			
Premenopausal	60	4	120
Postmenopausal	175	82	159
Surgery:			
Lumpectomy	87	8	178
Ablation	147	22	101
Adjuvant hormonal therapy:			
no	235	17	279
yes	0	69	0
Adjuvant chemotherapy:			
no	198	69	279
yes	37	18	0
Lymph node status:			
negative	102	20	279
positive	81	49	0
not applicable (M1)	42	17	0
Distant metastasis:			
yes	235	87	165
no	0	0	114
Disease -Free Interval:			
<1 year	59	13	20
1-3 year	108	29	71
>3 year	68	45	188
Median Follow-up time (in months):			
after surgery	62 (3-272)	103 (7-295)	93 (5-337)
after start therapy	30 (1-208)	45 (2-108)	
PR status*:			
Positive	186	72	217
Negative	48	15	62
HER2 status*:			
Amplified	31	10	43
Not amplified	202	77	233
CCDC170 status*:			
Positive	206	81	252
Negative	28	3	26
*as measured by RT-qPCR			

PR: Progesterone receptor; ERBB2: Erb-B2 Receptor Tyrosine Kinase 2; CCDC170: coiled-coil domain containing 170; qRT-PCR: Real-Time quantitative reverse transcription PCR.

Table 2
Association of *ESR1* fusions with clinical parameters in the predictive cohort

Predictive Endocrine Therapy Cohorts													
Parameters	n	at least one ESR1- CCDC170 (exon 2 to 8) fusion		P- Value	ESR1- CCDC170 (exon 2) fusion		PValue	ESR1- CCDC170 (exon 8) fusion		P- Value	ESR1- AKAP12		P- Value
		n	%		n	%		n	%		n	%	
All patients	322	89	27.6%		50	15.5%		51	15.8%		13	4.0%	
Age at start 1st line treatment (years)													
≤ 50	63	19	30.2%	0.63	12	19.0%	0.62	8	12.7%	0.029	1	1.6%	0.36
>50≤ 70	161	37	23.0%		23	14.3%		24	14.9%		7	4.3%	
>70	98	33	33.7%		15	15.3%		19	19.4%		5	5.1%	
Menopausal status at start of 1st line treatment													
Premenopausal	64	17	26.6%	0.82	10	15.6%	0.99	8	12.5%	0.41	1	1.6%	0.26
Postmenopausal	257	72	28.0%		40	15.6%		43	16.7%		12	4.7%	
Surgery type													
Lumpectomy	95	25	26.3%	0.79	14	14.7%	0.90	15	15.8%	0.83	2	2.1%	0.89
Ablation	169	42	24.9%		24	14.2%		25	14.8%		4	2.4%	
Radiotherapy													
No	105	30	28.6%	0.33	20	19.0%	0.08	16	15.2%	0.98	2	1.9%	0.74
Yes	159	37	23.3%		18	11.3%		24	15.1%		4	2.5%	
Nodal status													
No lymph nodes	122	33	27.0%	0.88	19	15.6%	0.99	20	16.4%	0.95	4	3.3%	0.2
Positive lymph nodes	130	38	29.2%		21	16.2%		22	16.9%		9	6.9%	
Tumor outside lymph nodes	53	15	28.3%		8	15.1%		7	13.2%		0	0.0%	
Not applicable (M1)	16	3	18.8%		2	12.5%		2	12.5%		0	0.0%	
Pathological Tumor classification													
pT1	85	22	25.9%	0.60	13	15.3%	0.21	14	16.5%	0.90	2	2.4%	0.36
pT2 + unknown	186	50	26.9%		25	13.4%		30	16.1%		10	5.4%	
pT3+pT4	51	17	33.3%		12	23.5%		7	13.7%		1	2.0%	
Tumor grade													
Poor	160	45	28.1%	0.36	27	16.9%	0.60	27	16.9%	0.60	7	4.4%	0.078
Unknown	81	18	22.2%		10	12.3%		10	12.3%		0	0.0%	
Moderate/Good	74	24	32.4%		13	17.6%		13	17.6%		5	6.8%	
Tumor cell content													
30-49%	27	7	25.9%	0.96	4	14.8%	0.99	2	7.4%	0.25	2	7.4%	0.63
50-70%	98	28	28.6%		15	15.3%		13	13.3%		4	4.1%	
>70%	197	54	27.4%		31	15.7%		36	18.3%		7	3.6%	
Hormone/ growth factor status (RT-qPCR)*													
ESR1-negative	0	0			0			0			0		
ESR1-positive	322	89	27.6%		50	15.5%		51	15.8%		13	4.0%	
PR-negative	63	18	28.6%	0.87	11	17.5%	0.65	11	17.5%	0.70	6	9.5%	0.014
PR-positive	258	71	27.5%		39	15.1%		40	15.5%		7	2.7%	

Predictive Endocrine Therapy Cohorts														
HER2 non-amplified	279	77	27.6%	0.63	44	15.8%	0.85	45	16.1%	0.81	13	4.7%	0.16	
HER2 amplified	41	12	29.3%		6	14.6%		6	14.6%		0	0.0%		
CCDC170 negative	31	5	16.1%	0.13	2	6.5%	0.15	4	12.9%	0.62	0	0.0%	0.23	
CCDC170 positive	287	83	28.9%		47	16.4%		47	16.4%		13	4.5%		
Adjuvant endocrine therapy														
No	252	66	26.2%	0.24	38	15.1%	0.64	36	14.3%	0.13	7	2.8%	0.030	
Yes (AI cohort only)	69	23	33.3%		12	17.4%		15	21.7%		6	8.7%		
Adjuvant chemotherapy														
No	267	76	28.5%	0.47	40	15.0%	0.55	45	16.9%	0.27	12	4.5%	0.36	
Yes	55	13	23.6%		10	18.2%		6	10.9%		1	1.8%		
Disease-free interval														
≤ 1 year disease-free	72	23	31.9%	0.47	14	19.4%	0.62	12	16.7%	0.99	2	2.8%	0.45	
1-3 years disease-free	137	37	27.0%		20	14.6%		20	14.6%		8	5.8%		
>3 years disease-free	113	29	25.7%		16	14.2%		19	16.8%		3	2.7%		
Dominant site of metastasis														
Local regional	29	10	34.5%	0.51	7	24.1%	0.32	4	13.8%	0.36	0	0.0%	0.40	
Bone	159	40	25.2%		25	15.7%		21	13.2%		6	3.8%		
Other distant metastasis	130	38	29.2%		17	13.1%		25	19.2%		7	5.4%		
Response type														
Complete response	11	3	27.3%	0.87	2	18.2%	0.73	1	9.1%	0.29	0	0.0%	0.46	
Partial response	39	9	23.1%		3	7.7%		6	15.4%		2	5.1%		
Stable disease over 6 months (SD > 6m)	115	32	27.8%		16	13.9%		23	20.0%		1	0.9%		
Stable disease for 6 months or less (SD ≤ 6m)	13	2	15.4%		2	15.4%		1	7.7%		0	0.0%		
Progressive disease (PD)	83	20	24.1%		14	16.9%		8	9.6%		3	3.6%		
Response type														
No response	96	22	22.9%	0.50	16	16.7%	0.38	9	9.4%	0.05	3	3.1%	0.50	
Response	165	44	26.7%		21	12.7%		30	18.2%		3	1.8%		

ESR1: Estrogen Receptor 1 gene; ERBB2: Erb-B2 Receptor Tyrosine Kinase 2; CCDC170: coiled-coil domain containing 170. Statistically significant differences are indicated in bold.

Table 3
Associations of *ESR1* fusions with clinical parameters in prognostic clinical cohort

Associations of ESR1 fusions with clinical parameters in prognostic clinical cohort													
LNN ESR+ Prognostic cohort													
Parameters	n	at least one ESR1-CCDC170 (exon 2 to 8) fusion		P- value	ESR1-CCDC170 (exon 2) fusion		P- value	ESR1-CCDC170 (exon 8) fusion		P- value	ESR1- AKAP12		P- value
		n	%		n	%		n	%		n	%	
All patients	279	70	25.1%		33	11.8%		39	14.0%		5	1.8%	
Age at primary surgery													
≤ 40 years	29	6	20.7%	0.001	4	13.8%	0.38	4	13.8%	0.26	1	3.4%	0.27
41-50 years	81	11	13.6%		5	6.2%		5	6.2%		0	0.0%	
51-70 years	125	36	28.8%		16	12.8%		21	16.8%		3	2.4%	
>70 years	44	17	38.6%		8	18.2%		9	20.5%		1	2.3%	
Menopausal status													
Premenopausal	120	19	15.8%	0.002	10	8.3%	0.12	11	9.2%	0.044	1	0.8%	0.29
Postmenopausal	159	51	32.1%		23	14.5%		28	17.6%		4	2.5%	
Surgery type													
Lumpectomy	178	44	24.7%	0.85	19	10.7%	0.43	25	14.0%	0.97	4	2.2%	0.45
Ablation	101	26	25.7%		14	13.9%		14	13.9%		1	1.0%	
Radiotherapy													
No	84	24	28.6%	0.38	14	16.7%	0.10	12	14.3%	0.92	1	1.2%	0.62
Yes	195	46	23.6%		19	9.7%		27	13.8%		4	2.1%	
Nodal status													
No lymph nodes	279	70	25.1%		33	11.8%		39	14.0%		5	1.8%	
Positive lymph nodes	0	0			0			0			0		
Tumor outside lymph nodes	0	0			0			0			0		
Pathological Tumor classification													
pT1	151	34	22.5%	0.28	17	11.3%	0.61	16	10.6%	0.08	2	1.3%	0.1
pT2 + unknown	119	32	26.9%		14	11.8%		20	16.8%		2	1.7%	
pT3+pT4	9	4	44.4%		2	22.2%		3	33.3%		1	11.1%	
Tumor grade													
Poor	131	36	27.5%	0.60	21	16.0%	0.06	21	16.0%	0.56	3	2.3%	0.84
Unknown	81	20	24.7%		9	11.1%		11	13.6%		1	1.2%	
Moderate/Good	67	14	20.9%		3	4.5%		7	10.4%		1	1.5%	
Tumor cell content													
30-49%	31	9	29.0%	0.82	6	19.4%	0.38	4	12.9%	0.86	1	3.2%	0.81
50-70%	69	16	23.2%		7	10.1%		11	15.9%		1	1.4%	
>70%	179	45	25.1%		20	11.2%		24	13.4%		3	1.7%	
Hormone/ growth factor status (RT-qPCR)*													
ESR1 negative	0	0			0			0			0		
ESR1 positive	279	70	25.1%		33	11.8%		39	14.0%		5	1.8%	
PR negative	62	16	25.8%	0.88	9	14.5%	0.46	8	12.9%	0.78	2	3.2%	0.93
PR positive	217	54	24.9%		24	11.1%		31	14.3%		3	1.4%	

LNN ESR+ Prognostic cohort													
<i>HER2</i> non-amplified	233	62	26.6%	<i>0.15</i>	29	12.4%	<i>0.30</i>	34	14.6%	<i>0.61</i>	4	1.7%	<i>0.78</i>
<i>HER2</i> amplified	43	7	16.3%		3	7.0%		5	11.6%		1	2.3%	
<i>CCDC170</i> negative	26	4	15.4%	<i>0.23</i>	2	7.7%	<i>0.49</i>	3	11.5%	<i>0.70</i>	0	0.0%	<i>0.47</i>
<i>CCDC170</i> positive	252	66	26.2%		31	12.3%		36	14.3%		5	2.0%	
Disease-free interval													
≤ 1 year disease-free	20	7	35.0%	0.011	2	10.0%	<i>0.08</i>	4	20.0%	0.006	0	0.0%	<i>0.57</i>
1-3 years disease-free	71	18	25.4%		10	14.1%		14	19.7%		2	2.8%	
>3 years disease-free	188	45	23.9%		21	11.2%		21	11.2%		3	1.6%	
Adjuvant endocrine therapy													
No	279	66	23.7%		33	11.8%		39	14.0%		5	1.8%	
Yes	0	0			0			0			0		
Adjuvant chemotherapy													
No	279	66	23.7%		33	11.8%		39	14.0%		5	1.8%	
Yes	0	0			0			0			0		

ESR1: Estrogen Receptor 1 gene; ERBB2: Erb-B2 Receptor Tyrosine Kinase 2; CCDC170: coiled-coil domain containing 170. Statistically significant differences are indicated in bold.

RNA isolation and RT-qPCR

Total RNA isolation from human breast cancer tissue, breast cancer cell line models and quality control were performed as previously described [14]. Next, cDNA was generated by a cycle at 48°C for 30 minutes with RevertAid H-minus (Applied Biosystems, Carlsbad, CA), according to the manufacturer's instructions. The cDNA was then pre-amplified for specific genes as previously described [14]. Briefly, 2 µL of cDNA (0.1 to 1 ng/ µL) was subject to a pre-amplification of 15 cycles using a multiple loci target-specific amplification for *ESR1* fusions with *AKAP12*, *ARMT1* and *CCDC170* and two reference genes, the Epithelial Cell Adhesion Molecule (*EPCAM*) and the Hypoxanthine Phosphoribosyltransferase 1 (*HPRT1*), with TaqMan PreAmp Master Mix (Applied Biosystems), as recommended by the manufacturer. Pre-amplified products were then diluted 12-fold in LoTE buffer (3 mM Tris-HCl/0.2 mM EDTA, pH 8.0) prior to downstream analysis. Next, 5 µL diluted pre-amplified samples were subjected to a TaqMan probe based real-time quantitative PCR (qPCR) for each gene combination, according to the manufacturer's instructions, in a MX3000P Real-Time PCR System (Agilent, Santa Clara, CA). The average expression of *HPRT1* and the epithelial marker *EPCAM* was used as reference to control RNA quality and calculate the expression levels of target genes, as previously described [14]. Only those samples with a $\Delta Cq > 25$ relative to the two reference genes were used for further evaluation of gene fusions, as previously described [18–20]. **Additional File 1** describes the primer sets used in the pre-amplification combination, as well as the Taqman qPCR used to quantify the fusions and reference genes. For *ESR1-CCDC170* fusion transcripts, the variants in which exon 2 of *ESR1* is fused to the coding region (exon 2 to 11) of *CCDC170* were examined (E2-E2, E2-E3, E2-E4, E2-E5, E2-E6, E2-E7, E2-E8, E2-E10 and E2-E11). Samples with a $\Delta Cq > 25$ relative to the reference genes were afterwards validated by MultiNA analysis (Shimadzu Europe, Duisburg, Germany). Only those samples with a MultiNA fusion product of the expected size were considered positive for the fusion transcripts (**Additional File 2**). The detection of *ESR1-CCDC170* fusion transcripts with RT-qPCR and MultiNA analysis was verified and confirmed in a set of fusion-positive reported breast cancer cell lines (**Additional File 3 - 4 - 5**).

Statistical analysis

All data were entered in SPSS version 24 (IBM Corp., Armonk, NY, USA) to generate the tables and perform the statistical analyses. For contingency tables, the Pearson Chi-Square Test was used. All *P*-values are 2-sided and *P*<0.05 was considered statistically significant.

Results

Association of *ESR1* with its *CCDC170*, *AKAP12* and *ARMT1* fusion partner

The presence of the *ESR1* fusions with *AKAP12*, *ARMT1* and *CCDC170* (exon 2 to exon 11) was evaluated in breast cancer tissue samples from 732 breast cancer patients. Fusion transcripts were predominantly detected in the *ESR1*+ population, with *CCDC170*, *AKAP12* or *ARMT1* fusion transcripts observed in 27.6%, 4.04% and 1.4% of the ER-positive cases respectively, and seen in 2.3%, 0.8% and 0% of the *ESR1*- cases respectively (*P*<0.001, Fisher's exact test two tailed. Table 4 and **Additional File 6**). In ER-positive tumors, full length *ESR1* and *CCDC170* mRNA levels were strongly correlated ($R^2=0.31$, *P*<0.0001)

(Additional File 7A) and transcript levels of both were significantly higher in the group of samples with an *ESR1-CCDC170* fusion transcript when compared to the group without [Student T-Test $P = 0.0316$ and 0.0001 , respectively (Additional File 7B)].

Table 4
Prevalence of *ESR1* fusions in the different analyzed cohorts.

	Total Count		At least one <i>ESR1-CCDC170</i> (exon 2 to 8) fusion				<i>ESR1-CCDC170</i> exon 2			<i>ESR1-CCDC170</i> exon 8			<i>ESR1_AKAP12</i>			
			no	yes	%	% of total count	no	yes	%	no	yes	%	no	yes	%	% of total count
All samples studied	788	ESR1 negative	128	3	2.3%	22.0%	128	3	2.29%	130	1	0.76%	130	1	0.76%	2.7%
		ESR1 positive	487	170	25.9%		565	92	14.00%	556	101	15.37%	637	20	3.04%	
1st line Tamoxifen	235	ESR1 negative	0	0	0%	24.7%	0	0	0%	0	0	0%	0	0	0%	2.6%
		ESR1 positive	177	58	24.7%		204	31	13.19%	201	34	14.47%	229	6	2.55%	
1st line AI	87	ESR1 negative	0	0	0%	35.6%	0	0	0%	0	0	0%	0	0	0%	8.0%
		ESR1 positive	56	31	35.6%		68	19	21.84%	70	17	19.54%	80	7	8.05%	
1st line endocrine cohort	322	ESR1 negative	0	0	0%	27.6%	0	0	0%	0	0	0%	0	0	0%	4.0%
		ESR1 positive	233	89	27.6%		272	50	15.53%	271	51	15.84%	309	13	4.04%	
Primary cohort	566	ESR1 negative	113	3	2.6%	17.8%	113	3	2.59%	115	1	0.86%	115	1	0.86%	1.9%
		ESR1 positive	352	98	21.8%		403	47	10.44%	392	58	12.89%	440	10	2.22%	
Primary LNP cohort	192	ESR1 negative	26	0	0.0%	15.6%	26	0	0.00%	26	0	0.00%	26	0	0.00%	2.6%
		ESR1 positive	136	30	18.1%		152	14	8.43%	148	18	10.84%	161	5	3.01%	
Primary LNN cohort	369	ESR1 negative	87	3	3.3%	18.7%	87	3	3.33%	89	1	1.11%	89	1	1.11%	1.6%
		ESR1 positive	213	66	23.7%		246	33	11.83%	240	39	13.98%	274	5	1.79%	
Normal breast tissue of breast cancer patients	36	ESR1 negative	0	0	0%	66.7%	0	0	0%	0	0	0%	0	0		0.0%
		ESR1 positive	12	24	66.7%		18	18	50.0%	23	13	36.1%	36	0	0.0%	
Tissue of breast fibroadenoma's	16	ESR1 negative	0	0	0%	25.0%	0	0	0%	0	0	0%	0	0	0%	0.0%
		ESR1 positive	12	4	20.0%		16	0	0.0%	16	4	20.0%	16	0	0.0%	
Tissue of breast DCIS	13	ESR1 negative	0	0	0%	7.7%	0	0	0%	0	0	0%	0	0	0%	0.0%
		ESR1 positive	12	1	7.7%		13	0	0.0%	13	0	0.0%	13	0	0.0%	
Normal breast tissue of healthy women	10	ESR1 negative	0	0	0%	10.0%	0	0	0%	0	0	0%	0	0	0%	0.0%
		ESR1 positive	9	1	10.0%		9	1	10.0%	10	0	0.0%	10	0	0.0%	

ESR1: Estrogen Receptor 1 gene; AI: Aromatase Inhibitor; LNP: Lymph node positive; LNN: Lymph node negative; DCIS: ductal carcinoma in situ. Statistically significant differences are indicated in bold.

Prevalence of *ESR1* fusion genes in normal mammary tissue, benign lesions and carcinoma in situ of the breast

While *AKAP12* and *ARMT1* fusion transcripts were not found in 36 non-malignant breast tissues taken at a distance of the primary tumor, *ESR1-CCDC170* fusion transcripts were detected in 67% of these normal breast tissues of patients with diagnosed breast cancer (Table 4). Note that *CCDC170*, but not *ESR1*, mRNA levels were significantly higher in these normal (adjacent to tumor) tissues than in cancer tissue (Kruskal Wallis Test $P < 0.0001$, (Fig. 2). To investigate this unexpectedly high incidence in more detail, we analyzed normal breast tissues of ten women without diagnosed breast cancer, 16 benign fibroadenomas and 13 ductal carcinomas in situ (DCIS) tissues, all of them *ESR1*-positive. In addition, we measured the fusion transcripts in three sets of patient-matched normal breast and primary tumor carcinomas and four patient-matched sets of primary breast tumors and metastatic lymph nodes, also all *ESR1*-positive. In none of these cases did we detect an *ESR1* fusion transcripts with *AKAP12* or *ARMT1*. However, one of the breast tissues of women without breast cancer diagnosis (10%) showed *ESR1-CCDC170* exon 2 (E2-E2) fusion transcripts, one of the DCIS cases (7.7%) had *ESR1-CCDC170* exon 6 (E2-E6) fusion transcripts, and four patients with fibroadenoma (25%) had *ESR1-CCDC170* exon 8 (E2-E8) fusion transcripts (Table 4 and **Additional File 6**). For one out of the three matched normal-tumor cases we found an *ESR1-CCDC170* exon 8 fusion in both the primary tumor and the normal breast tissue taken at a distance from the primary tumor. Finally, for two out of the four patients of which we had a matched primary tumor and lymph node metastasis, an *ESR1-CCDC170* exon 2 fusion was present in both the primary tumor and the lymph node metastasis.

Prevalence of *ESR1* fusion genes in breast tumor tissues

Since fusion transcripts were predominantly detected in the *ESR1+* population, we decided to investigate the clinical relevance of these transcripts in primary tumors. To this end, we stratified *ESR1+* patients in two distinct cohort: a predictive cohort of advanced BC patients treated with first-line endocrine therapy and a prognostic cohort of primary BC patients with lymph node negative disease (LNN) who did not receive any adjuvant systemic treatment.

In these two *ESR1+* cohorts, *ESR1-ARMT1* fusion transcripts were detected in four patients of the predictive cohort (1.2%) and in three patients of the prognostic cohort (1.1%). Due to the low incidence of this *ESR1-ARMT1* fusion transcript, it was not further pursued. *ESR1-AKAP12* fusion transcripts were more common, and observed in 13 patients of the predictive cohort (4.0%) and in five patients of the prognostic cohort (1.8%). The *ESR1-CCDC170* fusion transcripts, however, were the most prevalent and detected in the predictive cohort in 89 patients (27.6%) and in the prognostic cohort in 70 patients (25.1%). Interestingly, all patients harboring an *ESR1-ARMT1* or an *ESR1-AKAP12* fusion were also positive for an *ESR1-CCDC170* rearranged transcript. Moreover, we noticed the coexistence of the three fusions in two subjects. Of all the breast tissue samples studied, the most prominent *ESR1-CCDC170* fusion transcripts found involved exon 2 of *ESR1* fused with exon 2 (14%) and exon 8 (15.37%) of *CCDC170* (Table 4).

Association of *ESR1* fusion genes with DFS and OS in the prognostic cohort

The presence of *ESR1-CCDC170* fusion transcripts in the primary tumor of our *ESR1+* LNN patients predicted a shorter disease-free survival in a Cox proportional hazards regression survival analysis (HR $\pm$ 95% CI: 1.44 (1.01 – 2.05), $P = 0.044$) (Table 5). We decided to investigate the two frequently present *ESR1-CCDC170* fusion transcripts (E2-E2 and E2-E8). Analyzing the *ESR1-CCDC170* exon 2 and exon 8 separately, showed that the fusion with exon 8 of *CCDC170* on its own associated with a short disease free survival (DFS; HR $\pm$ 95% CI: 1.95 (1.30 – 2.93), $P = 0.001$). No association with disease free survival was seen for *ESR1-AKAP12* fusion transcripts (HR $\pm$ 95% CI: 1.23 (0.39 – 3.87), $P = 0.72$). Concerning overall survival, only the presence of an *ESR1-CCDC170* exon 8 fusion predicted a shorter overall survival time (HR $\pm$ 95% CI: 1.85 (1.18 – 2.90), $P = 0.007$) The DFS and OS Kaplan Meier curves as a function of *ESR1-CCDC170* exon 8 fusion transcripts are shown in Fig. 3A and Fig. 3B, respectively. A multivariate analysis was performed in which age at primary surgery, pathological tumor classification, tumor grade, progesterone receptor and HER2 status were included. The analysis revealed HER2 status as a significant prognostic factor for overall survival, but not for DFS ($P = 0.36$) (Table 5). In this analyses, the presence of *ESR1-CCDC170* exon 8 fusion transcripts was an independent prognostic factor for both DFS (HR $\pm$ 95% CI: 1.82 (1.20 – 2.75), $P = 0.005$) and OS (HR $\pm$ 95% CI: 1.71 (1.08 – 2.72), $P = 0.001$).

Table 5
Uni- and multivariate Cox proportional hazards regression survival analysis

Univariate model DFS						Multivariate model DFS				Univariate model OS				Multivariate model OS			
Parameters	n	HR	(95% CI)		P	HR	(95% CI)		P	HR	(95% CI)		P	HR	(95% CI)		P
	279																
Age at primary surgery					0.25				0.31				0.19				0.18
≤ 40 years	29	1				1				1				1			
41-50 years	81	0.59	0.35	1.00	0.049	0.60	0.35	1.02	0.06	0.53	0.30	0.96	0.036	0.51	0.28	0.94	0.032
51-70 years	125	0.73	0.44	1.19	0.20	0.72	0.44	1.18	0.19	0.75	0.44	1.28	0.30	0.72	0.42	1.26	0.25
>70 years	44	0.78	0.43	1.40	0.41	0.71	0.39	1.28	0.25	0.73	0.37	1.43	0.35	0.73	0.37	1.47	0.38
Menopausal status																	
Premenopausal	120	1								1							
Postmenopausal	159	1.01	0.73	1.38	0.96					1.06	0.74	1.53	0.73				
Pathological Tumor classification					0.009				0.037				0.017				0.019
pT1	151	1				1				1				1			
pT2 + unknown	119	1.54	1.12	2.11	0.007	1.35	0.98	1.88	0.069	1.30	0.90	1.87	0.165	1.19	0.81	1.74	0.375
pT3+pT4	9	2.31	1.00	5.32	0.049	2.47	1.07	5.75	0.035	3.26	1.39	7.62	0.006	3.45	1.45	8.19	0.005
Grade					< 0.001				0.001				0.033				0.082
poor	131	1				1				1				1			
unknown	81	1.36	0.97	1.91	0.076	1.40	0.98	1.99	0.064	0.89	0.59	1.34	0.577	0.97	0.64	1.48	0.894
moderate and good	67	0.52	0.33	0.82	0.004	0.57	0.36	0.89	0.014	0.51	0.31	0.85	0.009	0.57	0.34	0.94	0.029
ER	279	1.11	0.98	1.25	0.10					0.99	0.86	1.14	0.92				
PR																	
negative	62	1				1				1				1			
positive	217	0.66	0.46	0.93	0.019	0.68	0.47	0.98	0.037	0.49	0.33	0.73	< 0.001	0.56	0.37	0.85	0.007
HER2 status*																	
not amplified	233	1								1				1			
amplified	43	1.21	0.80	1.84	0.36					1.82	1.17	2.84	0.008	1.72	1.08	2.73	0.022
Univariate model PFS						Univariate model post-relapse survival											
1st line Tamoxifen	235																
at least one ESR1-CCDC170 (exon 2 to 8) fusion		0.96	0.71	1.30	0.81					1.16	0.85	1.60	0.35				
ESR1-AKAP12		1.37	0.61	3.10	0.44					1.92	0.84	4.35	0.12				
1st line AI	87																
at least one ESR1-CCDC170 (exon 2 to 8) fusion		0.85	0.53	1.37	0.50												
ESR1-AKAP12		1.62	0.73	3.60	0.24												

Univariate model DFS						Multivariate model DFS				Univariate model OS				Multivariate model OS					
						Separately added to the base model													
						Univariate model DFS						Multivariate model DFS				Univariate model OS			
at least one ESR1-CCDC170 (exon 2 to 8) fusion																			
negative	213	1				1				1				1					
positive	66	1.44	1.01	2.05	0.044	1.33	0.92	1.92	0.13	1.67	1.13	2.47	0.010	1.54	1.02	2.33	0.042		
ESR1-CCDC170 (exon 2) fusion																			
negative	246	1								1				1					
positive	33	1.40	0.89	2.21	0.14					1.75	1.07	2.87	0.026	1.38	0.82	2.33	0.22		
ESR1-CCDC170 (exon 8) fusion																			
negative	240	1				1				1				1					
positive	39	1.95	1.30	2.93	0.001	1.82	1.20	2.75	0.005	1.85	1.18	2.90	0.007	1.71	1.08	2.72	0.023		
ESR1-AKAP12																			
negative	274	1								1									
positive	5	1.23	0.39	3.87	0.72					2.45	0.90	6.65	0.08						

DFS: Disease free survival; OS: Overall survival; ER: Estrogen Receptor; PR: Progesterone receptor; HER2: human epidermal growth factor receptor 2; PFS: Progression free survival. *Due to the unknown data, numbers do not add up to 279. Differences statistically significant are indicated in bold.

Association of *ESR1* fusion genes with clinical characteristics, PFS and post-relapse overall survival in advanced BC patients

The fusion transcripts were related with traditional clinical parameters, with response to first-line endocrine therapy in the predictive cohort (n=322; tamoxifen (n=235), aromatase inhibitors (n=87)) (Table 2). In the predictive cohort *ESR1-CCDC170* fusion transcripts showed an association with age at start of first-line treatment, whereas *ESR1-AKAP12* fusion transcripts were enriched in patients with progesterone-negative primary tumors at time of surgery and in AI-treated patients who received adjuvant tamoxifen. No relation with PFS after first-line tamoxifen (n=235) was found in our Cox proportional hazards regression survival analysis for the *ESR1-CCDC170* fusion transcripts (HR ± 95% CI: 0.96 (0.71 – 1.30), P = 0.81) nor for the *ESR1-AKAP12* fusion transcripts (HR ± 95% CI: 1.37 (0.61 – 3.10), P = 0.44) (Table 5). In addition, the presence of these fusion transcripts did not affect the time from relapse to death (post-relapse survival, HR ± 95% CI: 1.16 (0.85 – 1.60), P = 0.35 and 1.92 (0.84 – 4.35), P = 0.12, for *ESR1* fusions with *CCDC170* and *AKAP12*, respectively) (Table 5). Similarly, also no association with PFS for first-line aromatase inhibitors (n=87) was found for *ESR1-CCDC170* fusion transcripts (HR ± 95% CI: 0.85 (0.53 – 1.37), P = 0.50) nor for the *ESR1-AKAP12* fusion transcripts (HR ± 95% CI: 1.62 (0.73 – 3.60), P = 0.24). With data available for only 27 patients post-relapse, we did not analyze post-relapse survival for aromatase inhibitors. Moreover, no-significant associations with PFS were seen when the *ESR1-CCDC170* exon 2 and exon 8 fusion transcripts were analyzed separately (Table 5).

Discussion

The genetic landscape contributing to *de novo* or acquired resistance to endocrine therapy in breast cancer patients is not completely understood yet. In this study, we investigated the occurrence of recurrent fusion transcripts between *ESR1* and three different loci adjacent to *ESR1* (*CCDC170*, *AKAP12* and *ARMT1*) and correlated their presence with clinical outcome. All of the fusion transcripts analyzed are recurrent and most frequently present in ER-positive disease and among them *ESR1-CCDC170* fusion transcripts were the most predominant. As proposed by others [10, 13], the presumption was that these fusion transcripts, which are considered to cause constitutive ER signaling, might signify resistance to endocrine therapy. However, in patients with advanced breast cancer, we did not find that the presence of any of these fusion transcripts is associated with outcome to endocrine therapy whether it concerned first line tamoxifen or an aromatase inhibitor. Importantly, smaller size effects from these the variants may be undetected due to the relatively small sample size of the study cohort, 87 patients treated with aromatase inhibitors and 235 subjects with tamoxifen. In contrast, in patients with primary BC and not receiving adjuvant systemic hormone treatments, we found that fusion between *ESR1* and *CCDC170* in general, and between exon 2 of *ESR1* and exon 8 of *CCDC170* in particular, predicted in uni- and multivariable analyses shorter disease free survival as well as shorter overall survival. Thus, *ESR1* and *CCDC170* fusion transcript pinpoint cancers with an adverse outcome.

Understanding the molecular mechanisms that underlay the origin of fusion transcripts could help to comprehend the role of these fusions in carcinogenesis as well as improve the diagnosis of cancer patients [10, 13]. Although the progress in DNA sequencing enhanced detection of recurrent and pathological breast cancer fusions, the complexity of underlying genomic rearrangement patterns makes their characterization at the DNA level often difficult. The fusion between *ESR1* and its neighboring gene *CCDC170* are potentially generated by tandem duplication [9, 13, 21, 22], which is also causing other genetic rearrangements in cancer [9, 23, 24]. Kim *et al.* found a region within the *ESR1* genomic locus most vulnerable to DNA strand breakage, which often included

intron 6 region of its neighboring gene *CCDC170*, resulting in oncogenic mRNA *ESR1-CCDC170* fusion transcript of exons 2 of *ESR1* connected to exon 2-11 of *CCDC170*, i.e. the C-terminal domain of *CCDC170* [25]. Irrespective of mechanisms causing the gene fusions, they occur in a patient-specific manner, which makes their identification at the DNA level less suitable for routine diagnostics. Our method to analyze fusion transcripts is much less dependent on exact position of the underlying gene fusion at the DNA level and is therefore better suited to evaluate as a general biomarker in large patient cohorts. However, an important caveat for detecting gene fusions at the transcript level is the fact that it cannot distinguish between fusion transcripts arising from actual genetic rearrangements and those that arise from transcription reading from one gene into the next without a genetic cause. Finally, the generation of artefactual fusion sequences, which are randomly ligated during the sequencing procedure, might happen, as previously reported by *Veeraraghavan et al.* [13]. Overall, we performed RT-qPCR analysis and investigated RNA not DNA, therefore we cannot tell whether fusion transcripts are the results of (DNA) rearrangements.

Another interesting question regards the biological significance of clinically relevant fusion transcripts. Gene fusions and their products (RNAs and proteins) are assumed to be exclusive to cancer. However, RNA-sequencing analyses from normal appearing margins of cancerous specimens showed fusion transcripts also in normal tissues [26]. In fact, oncogenic rearrangements, such as the *EML4-ALK* [27], *NPM-ALK* [28], *JAZF1-JJAZ1* [29] and *BCR-ABL1* [30] fusions are also expressed at a low level in histologically non-neoplastic tissues [9]. In our study, expression of *ESR1* fused to exons 2 and exon 8 of *CCDC170* was found in mammary epithelial tissues derived from women without diagnosis of breast cancer, and in cases with (benign) fibroadenomas, respectively. Also in early stages of breast cancer, like DCIS, we detected fusion transcripts. Moreover, *ESR1-CCDC170* fusion transcripts were also detected in normal breast tissues of patients with diagnosed breast cancer. This argues that a percentage may be transcript read-through instead of fusion transcripts arising from gene fusions.

According to our results, the expression of *ESR1-CCDC170* exon 2 and exon 8 fusion transcripts were linked to a less favorable disease in BC patients who not received adjuvant systemic treatment. Overall, our results are in agreement with those reported by *Veeraraghavan et al.* which showed that *ESR1-CCDC170* fusions, when introduced into ER-positive breast cancer cells, leads to a markedly increase of cell motility and colony-forming ability, increase in S-G2/M phase cells and a decrease in G0/G1 phase cells. Although several functional studies [9, 31] demonstrated a role of *ESR1-CCDC170* fusions in endocrine therapy resistance, no relationship between fusion transcripts and treatment outcome was observed in our predictive cohort. Overall, since *ESR1-CCDC170* fusions in our study demonstrated no predictive value for endocrine therapy resistance, their prognostic value might be explained by the recurrent incidence of read-through events during cell cycle progression. This latter has been exemplified with the abundance of *CTSD-IFITM10* readthrough fusions during breast cancer cell proliferation [32].

Conclusions

The most important conclusion from our work is that among the fusion transcripts evaluated measuring *ESR1-CCDC170* exon 8 fusion transcripts in primary breast cancers has diagnostic potential as it identifies a more aggressive subset of ER-positive breast cancer patients. Furthermore, with our study we demonstrated that *ESR1-CCDC170* fusion transcript does not predict endocrine therapy resistance in our setting.

Abbreviations

AI: aromatase inhibitor; ARMT1: Acidic Residue Methyltransferase 1; AKAP12: A-Kinase Anchoring Protein 12; BC: breast cancer; CCDC170: Coiled-Coil Domain Containing 170; DCIS: ductal carcinomas in situ; DFS: disease-free survival; ER: estrogen receptor; ERBB2: Erb-B2 Receptor Tyrosine Kinase 2; ESR1: estrogen receptor alpha; ET: endocrine therapy; HPRT1: Hypoxanthine Phosphoribosyltransferase 1; HR: Hazard ratio; IHC: immunohistochemical staining; LBD: ligand-binding domain; LNN: lymph node negative; LNP: Lymph node positive; OS: Overall survival; PFS: progression free survival; PR: Progesterone receptor; RT-qPCR: reverse transcriptase quantitative PCR; TAM: Tamoxifen.

Declarations

Ethics approval and consent to participate: The protocol of the study was approved by the medical ethics committee of the Erasmus Medical Centre Rotterdam, The Netherlands (MEC 02.953) and was performed in accordance with the Code of Conduct of the Federation of Medical Scientific Societies in The Netherlands (<https://www.federa.org/codes-conduct>). The use of coded left-over material for scientific purposes does not require informed consent according to Dutch law and the new European general data protection regulation (GDPR).

Consent for publication: Not applicable

Availability of data and materials: All data generated or analysed during this study are included in this published article and its supplementary information files.

Disclosures: None declared.

Grants and acknowledgements: AMS and JWM were supported by the Cancer Genomics Netherlands (CGC.nl)/Netherlands Organization for Scientific Research (NWO). SRV was supported by an ERC Advanced grant (No 322737). The authors are especially grateful to all patients, to Steven van Laere and Luc Dirix - Department Oncology, Antwerp University, Belgium and to Sabine C. Linn – Department of Medical Oncology, Netherlands Cancer Institute, Amsterdam, the Netherlands for their contribution of tissues from patients treated with 1st line aromatase inhibitor.

Authors' contributions: SRV and AMS designed the study; SRV, KR-R, CB, AMT, RF, AMT-J and AMS performed the laboratory experiments (RT-qPCR, western blotting analyses and immunohistochemical stainings). AMS and MPHMJ analyzed the data and compiled statistics; JWM and MPHMJ supplied the patient

materials (tissues and clinical information); PV, SS and JWM supervised the study and provided the financial budget; SRV, AMS and MPHMJ wrote the manuscript, which was reviewed, edited and approved by all authors.

References

1. Pritchard KI: **Endocrine therapy: is the first generation of targeted drugs the last?** *J Intern Med* 2013, **274**:144-152.
2. Riggins RB, Schrecengost RS, Guerrero MS, Bouton AH: **Pathways to tamoxifen resistance.** *Cancer Lett* 2007, **256**:1-24.
3. Osborne CK, Schiff R: **Mechanisms of endocrine resistance in breast cancer.** *Annu Rev Med* 2011, **62**:233-247.
4. Schiavon G, Hrebien S, Garcia-Murillas I, Cutts RJ, Pearson A, Tarazona N, Fenwick K, Kozarewa I, Lopez-Knowles E, Ribas R, et al: **Analysis of ESR1 mutation in circulating tumor DNA demonstrates evolution during therapy for metastatic breast cancer.** *Sci Transl Med* 2015, **7**:313ra182.
5. Pejerrey SM, Dustin D, Kim JA, Gu G, Rechoum Y, Fuqua SAW: **The Impact of ESR1 Mutations on the Treatment of Metastatic Breast Cancer.** *Horm Cancer* 2018, **9**:215-228.
6. Beijer N, Sieuwerts AM, Kraan J, Van NM, Onstenk W, Vitale SR, van der Vlugt-Daane M, Dirix LY, Brouwer A, Hamberg P, et al: **Estrogen receptor mutations and splice variants determined in liquid biopsies from metastatic breast cancer patients.** *Mol Oncol* 2018, **12**:48-57.
7. Vitale SR, Sieuwerts AM, Beijer N, Kraan J, Angus L, Mostert B, Reijm EA, Van NM, van Marion R, Dirix LY, et al: **An Optimized Workflow to Evaluate Estrogen Receptor Gene Mutations in Small Amounts of Cell-Free DNA.** *J Mol Diagn* 2019, **21**:123-137.
8. Robinson DR, Kalyana-Sundaram S, Wu YM, Shankar S, Cao X, Ateeq B, Asangani IA, Iyer M, Maher CA, Grasso CS, et al: **Functionally recurrent rearrangements of the MAST kinase and Notch gene families in breast cancer.** *Nat Med* 2011, **17**:1646-1651.
9. Veeraraghavan J, Tan Y, Cao XX, Kim JA, Wang X, Chamness GC, Maiti SN, Cooper LJ, Edwards DP, Contreras A, et al: **Recurrent ESR1-CCDC170 rearrangements in an aggressive subset of oestrogen receptor-positive breast cancers.** *Nat Commun* 2014, **5**:4577.
10. Giltane JM, Hutchinson KE, Stricker TP, Formisano L, Young CD, Estrada MV, Nixon MJ, Du L, Sanchez V, Ericsson PG, et al: **Genomic profiling of ER(+)** breast cancers after short-term estrogen suppression reveals alterations associated with endocrine resistance. *Sci Transl Med* 2017, **9**.
11. Li S, Shen D, Shao J, Crowder R, Liu W, Prat A, He X, Liu S, Hoog J, Lu C, et al: **Endocrine-therapy-resistant ESR1 variants revealed by genomic characterization of breast-cancer-derived xenografts.** *Cell Rep* 2013, **4**:1116-1130.
12. Yoshihara K, Wang Q, Torres-Garcia W, Zheng S, Vegesna R, Kim H, Verhaak RG: **The landscape and therapeutic relevance of cancer-associated transcript fusions.** *Oncogene* 2015, **34**:4845-4854.
13. Veeraraghavan J, Ma J, Hu Y, Wang XS: **Recurrent and pathological gene fusions in breast cancer: current advances in genomic discovery and clinical implications.** *Breast Cancer Res Treat* 2016, **158**:219-232.
14. Sieuwerts AM, Meijer-van Gelder ME, Timmermans M, Trapman AM, Garcia RR, Arnold M, Goedheer AJ, Portengen H, Klijn JG, Foekens JA: **How ADAM-9 and ADAM-11 differentially from estrogen receptor predict response to tamoxifen treatment in patients with recurrent breast cancer: a retrospective study.** *Clin Cancer Res* 2005, **11**:7311-7321.
15. Sieuwerts AM, Usher PA, Meijer-van Gelder ME, Timmermans M, Martens JW, Brunner N, Klijn JG, Offenberg H, Foekens JA: **Concentrations of TIMP1 mRNA splice variants and TIMP-1 protein are differentially associated with prognosis in primary breast cancer.** *Clin Chem* 2007, **53**:1280-1288.
16. Ramirez-Ardila DE, Helmijs JC, Look MP, Lurkin I, Ruigrok-Ritstier K, van Laere S, Dirix L, Sweep FC, Span PN, Linn SC, et al: **Hotspot mutations in PIK3CA associate with first-line treatment outcome for aromatase inhibitors but not for tamoxifen.** *Breast Cancer Res Treat* 2013, **139**:39-49.
17. Hayward JL, Rubens RD, Carbone PP, Heuson JC, Kumaoka S, Segaloff A: **Assessment of response to therapy in advanced breast cancer. A project of the programme on clinical oncology of the International Union against Cancer, Geneva, Switzerland.** *Eur J Cancer* 1978, **14**:1291-1292.
18. Onstenk W, Sieuwerts AM, Kraan J, Van M, Nieuweboer AJ, Mathijssen RH, Hamberg P, Meulenbeld HJ, De Laere B, Dirix LY, et al: **Efficacy of Cabazitaxel in Castration-resistant Prostate Cancer Is Independent of the Presence of AR-V7 in Circulating Tumor Cells.** *Eur Urol* 2015, **68**:939-945.
19. Sieuwerts AM, Kraan J, Bolt-de Vries J, van der Spoel P, Mostert B, Martens JW, Gratama JW, Sleijfer S, Foekens JA: **Molecular characterization of circulating tumor cells in large quantities of contaminating leukocytes by a multiplex real-time PCR.** *Breast Cancer Res Treat* 2009, **118**:455-468.
20. Sieuwerts AM, Mostert B, Bolt-de Vries J, Peeters D, de Jongh FE, Stouthard JM, Dirix LY, van Dam PA, Van Galen A, de Weerd V, et al: **mRNA and microRNA expression profiles in circulating tumor cells and primary tumors of metastatic breast cancer patients.** *Clin Cancer Res* 2011, **17**:3600-3618.
21. Lei JT, Gou X, Ellis MJ: **ESR1 fusions drive endocrine therapy resistance and metastasis in breast cancer.** *Mol Cell Oncol* 2018, **5**:e1526005.
22. Paratala BS, Dolfi SC, Khiabani H, Rodriguez-Rodriguez L, Ganesan S, Hirshfield KM: **Emerging Role of Genomic Rearrangements in Breast Cancer: Applying Knowledge from Other Cancers.** *Biomark Cancer* 2016, **8**:1-14.
23. Jones DT, Kocalkowski S, Liu L, Pearson DM, Backlund LM, Ichimura K, Collins VP: **Tandem duplication producing a novel oncogenic BRAF fusion gene defines the majority of pilocytic astrocytomas.** *Cancer Res* 2008, **68**:8673-8677.
24. Lipson D, Capelletti M, Yelensky R, Otto G, Parker A, Jarosz M, Curran JA, Balasubramanian S, Bloom T, Brennan KW, et al: **Identification of new ALK and RET gene fusions from colorectal and lung cancer biopsies.** *Nat Med* 2012, **18**:382-384.
25. Kim RN, Moon HG, Han W, Noh DY: **Perspective Insight into Future Potential Fusion Gene Transcript Biomarker Candidates in Breast Cancer.** *Int J Mol Sci* 2018, **19**.
26. Babiceanu M, Qin F, Xie Z, Jia Y, Lopez K, Janus N, Facemire L, Kumar S, Pang Y, Qi Y, et al: **Recurrent chimeric fusion RNAs in non-cancer tissues and cells.** *Nucleic Acids Res* 2016, **44**:2859-2872.
27. Martelli MP, Sozzi G, Hernandez L, Pettirossi V, Navarro A, Conte D, Gasparini P, Perrone F, Modena P, Pastorino U, et al: **EML4-ALK rearrangement in non-small cell lung cancer and non-tumor lung tissues.** *Am J Pathol* 2009, **174**:661-670.

28. Maes B, Vanhentenrijk V, Wlodarska I, Cools J, Peeters B, Marynen P, de Wolf-Peeters C: **The NPM-ALK and the ATIC-ALK fusion genes can be detected in non-neoplastic cells.** *Am J Pathol* 2001, **158**:2185-2193.

29. Li H, Wang J, Mor G, Sklar J: **A neoplastic gene fusion mimics trans-splicing of RNAs in normal human cells.** *Science* 2008, **321**:1357-1361.

30. Biernaix C, Loos M, Sels A, Huez G, Stryckmans P: **Detection of major bcr-abl gene expression at a very low level in blood cells of some healthy individuals.** *Blood* 1995, **86**:3118-3122.

31. Li Li JV, Yiheng Hu, Xian Wang, Ying Tan, Rachel Schiff and Xiaosong Wang: **Abstract 376: Therapeutic role of ESR1-CCDC170 gene fusion in breast cancer endocrine resistance.** *American Association for Cancer Research* 2019, **79**:Supplement, pp. 376.

32. Varley KE, Gertz J, Roberts BS, Davis NS, Bowling KM, Kirby MK, Nesmith AS, Oliver PG, Grizzle WE, Forero A, et al: **Recurrent read-through fusion transcripts in breast cancer.** *Breast Cancer Res Treat* 2014, **146**:287-297.

Figures

Figure 1

Figure 1

Overview of the study and selection of available patients. A) Flow diagram of the study; B) Workflow of processing samples: fusion gene mRNA levels were measured in 322 ER-positive primary tumors (predictive cohort) by quantitative reverse transcriptase PCR (RT-qPCR). All patients in this cohort were hormone-naïve and all experienced a disease recurrence and subsequently received 1st line endocrine therapy. The association of the presence of ESR1 fusion genes in the primary tumor progression-free survival (PFS) after start with 1st line tamoxifen (n=235) or aromatase inhibitors (n=87), were evaluated. Similarly, disease free interval (DFS) and overall survival (OS) were investigated in 279 lymph node negative ER-positive breast cancer patients (prognostic cohort) who had not received any (neo)adjuvant systemic therapy. ESR1: Estrogen Receptor 1 gene; AI: Aromatase Inhibitor; LNN: Lymph node negative; ER: Estrogen Receptor; RT-qPCR: Quantitative reverse transcriptase PCR.

Figure 2

Figure 2

Expression of ESR1 and CCDC170 genes in breast tissues. CCDC170 (blue box) and ESR1 (green box) mRNA levels are showed in the y-axis and were measured by RT-qPCR in normal (adjacent to tumor), benign (DCIS) and carcinoma (LNN and LNP) breast tissues. The box plots show interquartile ranges (IQR) together with the median (black horizontal line) of the ESR1 and CCDC170 mRNA levels for the different conditions. DCIS: ductal carcinomas in situ; LNP: Lymph node positive; LNN: Lymph node negative.

Figure 3A

Figure 3B

Figure 3

Figure 3

Disease free survival (DFS) and overall survival (OS) in the prognostic cohort. The DFS and OS Kaplan Meier curves in positive LNN patients. A) DFS of patients with or without ESR1-CCDC170 exon 8 fusion gene; (B) OS of patients with or without ESR1-CCDC170 exon 8 fusion gene. The reported P-value is from a log-rank test and the test statistics from Cox regression analyses.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AdditionalFile3BMCCancer.jpg](#)
- [AdditionalFile4BMCCancer.jpg](#)
- [AdditionalFile7BMCCancer.jpg](#)
- [AdditionalFileFusionpaperBMCCancer.xlsx](#)