Mosquito Host-Seeking Diel Rhythm and Chemosensory Gene Expression is Affected by Age and Plasmodium Stages

Melika Hajkazemian

Stockholm University

Sharon R. Hill

Swedish University of Agricultural Sciences

Raimondas Mozūraitis

Stockholm University

Lisa Ranford-Cartwright

University of Glasgow

S. Noushin Emami (✉ noushin.emami@su.se)

Stockholm University

Rickard Ignell

Swedish University of Agricultural Sciences

Research Article

Keywords: Plasmodium falciparum, parasite stages, Anopheles gambiae, vector, locomotion, olfaction, antennae, transcriptomics

Posted Date: December 9th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1114857/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Malaria parasites can affect vector-related behaviours, increasing transmission success. Using *Anopheles gambiae* and *Plasmodium falciparum*, we consider the effect of interaction between infection stage and vector age on diel locomotion in response to human odour and the expression of antennal chemosensory genes. We demonstrate age-dependent behavioural diel compartmentalisation by uninfected females. Infection disrupts overall and diel activity patterns compared with age-matched controls. Mosquitoes carrying transmissible sporozoites are more active, shifting activity periods to coincide with human host availability, in response to human odour. Older, uninfected females reduce activity during their peak host-seeking period in response to human odour. Age- and infection stage-specific changes in odour-mediated locomotion coincide with altered transcript abundance of select chemosensory genes providing a possible molecular mechanism regulating the behaviour. Our results support the hypothesis that vector-related behaviours of female mosquitoes are altered by infection stage and further modulated by the age of the vector, and have important implications for malaria transmission and disease dynamics.

Summary

The presence of different life-cycle stages of the malaria parasite, *Plasmodium falciparum*, together with the age of the vector, affect the host-seeking daily rhythm and antennal chemosensory gene expression of *Anopheles gambiae* to indirectly increase the odds of successful disease transmission.

Introduction

Malaria parasites depend on mosquito vectors for transmission from one host to another. Increasing empirical evidence demonstrates that insect-borne pathogens may manipulate both their human host and their insect vector to enhance transmission success^{1–4}. This manipulation is not static, as vector-related behaviours can vary significantly between uninfected and infected vectors over different time scales relevant to the disease transmission dynamics^{3–5}. Pathogens are known to affect vector behaviours, including locomotion, flight, host seeking, probing and feeding, which can manifest in increased avidity to locate and obtain a blood meal or in alterations in host preference, resulting in an enhanced transmission through the increased contact between vector and host^{3,5,6}. The molecular mechanisms by which these behavioural traits may be modulated have only recently received attention, identifying pathways in the neuroendocrine circuitry as key targets^{2,5,7,8}.

Mosquitoes acquire *Plasmodium* parasites, the causative agent of malaria, through a blood meal from an infected vertebrate host. The pathogen undergoes several developmental stages in the mosquito, from midgut invasion and oocyst formation (infected mosquitoes, 7 days post-infection; dpi) through to the invasion of the salivary glands, when a mosquito can transmit the malaria sporozoites (infectious mosquitoes, 14 dpi)⁹. The avidity of mosquitoes to engage in vector-related behaviours, including host seeking and blood feeding, depends on the developmental stage of the pathogen in the mosquito, rather

than the infection status of the mosquito alone³. Available data for *Plasmodium* parasites that cause avian, rodent and human malaria emphasise that vector-related behaviours are suppressed when mosquitoes are infected, but not able to transmit the pathogen, whereas infectious mosquitoes are more active^{10–12}. This modulation appears to be conserved across vectors and other pathogens, *e.g.*, *Aedes aegypti* that transmit dengue and Zika virus^{5,13}. The observed changes in the odour-mediated host-seeking behaviour, demonstrated for both infected and infectious mosquitoes, are reflected in the selective sensitivity of the peripheral olfactory system to key host-related volatile organic compounds^{5,10,14}. Such modulation may be regulated through alterations in neural signalling and chemosensory gene expression, as demonstrated in malaria- and dengue-infected mosquitoes^{2,5}.

In depth analysis of the molecular basis of olfaction in mosquitoes, has allowed for the identification and functional characterisation of chemosensory gene families, including odorant receptors (*Ors*), gustatory receptors (*Grs*), ionotropic receptors (*Irs*), odorant binding proteins (*Obps*) and chemosensory proteins (*Csps*)^{15–19}. The expression of select members of these gene families in olfactory-related tissues change in response to adult maturation, age and physiological state^{20–25}, including infection status⁵. These state-dependent changes in gene abundance coincide with concerted alterations in the sensitivity of olfactory sensory neurones and the behaviour of the mosquitoes^{20,21,25}.

This study analysed the stage-dependent effect of *P. falciparum* infection on the locomotion activity of female *Anopheles gambiae* in the presence and absence of human odour, demonstrating that infected and infectious mosquitoes display differential locomotor activity correlating with age, infection status and time of day. The potential molecular mechanisms underlying the age- and infection-dependent odour-mediated locomotor activity were assessed by transcriptome analysis, identifying differentially abundant chemosensory-related genes in the antennae coinciding with the demonstrated differences in mosquito behaviour. A better understanding of the effect of infection on locomotor activity and odour-mediated host seeking throughout the day has the potential to more accurately direct vector measures. Identifying the molecular mechanisms by which these behavioural traits are modulated by *P. falciparum* may lead to a better understanding of vectorial capacity and transmission dynamics potentially inspiring the development of innovative control strategies.

Results

Mosquito age and the stage of *Plasmodium falciparum* infection affect locomotor activity

Locomotor activity of uninfected female *An. gambiae* in the absence of human odour significantly reduced with age when assessed in *Drosophila* activity monitors ($\chi^2_1 = 5.86$, $p = 0.02$; Fig. 1a-b). This age effect was not observed following infection with *P. falciparum* ($\chi^2_1 = 0.83$, $p = 0.35$; Fig. 1b). In the presence of human odour, the age-dependent reduction in locomotor activity was observed in both uninfected ($\chi^2_1 = 21.12$, $p < 0.001$) and *P. falciparum* carrying mosquitoes ($\chi^2_1 = 5.73$, $p = 0.02$; the GLMM:lmer model was constrained by age and the random effect of experimental replication; Fig. 1b).

Infection status significantly affected the activity of mosquitoes when comparing the overall locomotion profiles of 7 dpi and 14 dpi individuals with age-matched uninfected females in the presence and absence of human odour (β -estimated = weighted mean value based on all variables plus the random effect of the replication $\pm$ SE; presence of human odour: $\chi^2_1 = 18.06$, $p < 0.001$; absence of human odour: $\chi^2_1 = 5.73$, $p < 0.01$). In the absence of human odour, 7 dpi females were less active compared to age-matched controls ($\chi^2_1 = 8.21$, $p = 0.004$), whereas 14 dpi females were as active as their age-matched controls ($\chi^2_1 = 2.27$, $p = 0.13$; Fig. 1c). The presence of human odour differentially altered the locomotor activity. While there were no significant differences between 7 dpi and age-matched females in the presence of human odour ($\chi^2_1 = 0.12$, $p = 0.73$), the infectious 14 dpi mosquitoes were more active than age-matched uninfected females (14 days control; 14 dc $\chi^2_1 = 8.80$, $p = 0.002$; Fig. 1c).

Mosquito locomotor activity varies temporally in response to host odour and *P. falciparum* infection

Diurnal cycles in *An. gambiae* locomotor and host-seeking activities were evident, with distinct crepuscular activity peaks culminating in the highest activity at the onset of photophase (ZT 23-24), and low to moderate activity during scotophase (ZT 12-23) and low activity during photophase (ZT 0-11)²⁶. To assess the diurnal effect on the locomotor activity of mosquitoes with *P. falciparum* infection and their respective controls, a mixed model analysis was used (lmer model: constrained by two main explanatory variables, infection status and period of time, as well as the random effect of replication, Supplementary Table 1-2). In the absence of human odour, the diurnal locomotor activity profile of uninfected mosquitoes confirmed the high activity at dawn (ZT 23-24), with the younger cohort being more active during scotophase with a peak at dawn, whereas the older control females displayed an increased activity at dusk that was maintained until dawn (ZT 0-11: $\chi^2_1 = 27.43$, $p < 0.001$; ZT 11-12: $\chi^2_1 = 4.58$, $p = 0.049$; ZT 12-23: $\chi^2_1 = 4.33$, $p = 0.03$; ZT 23-24: $\chi^2_1 = 32.32$, $p < 0.001$, Fig. 1d, Supplementary Table 2.). In the presence of human odour, the locomotor activity of the younger uninfected cohort increased at dawn, and remained at this activity level until dusk, while the older females only displayed an increase in activity at dusk (ZT 0-11: $\chi^2_1 = 17.54$, $p < 0.001$; ZT 11-12: $\chi^2_1 = 7.67$, $p = 0.005$; ZT 12-23: $\chi^2_1 = 15.21$, $p < 0.001$; ZT 23-24: $\chi^2_1 = 18.25$, $p < 0.001$, Fig. 1d, Supplementary Table 2.).

The time-of-day, and the stage of *P. falciparum* sporogony, differentially affected the locomotor activity of the vector (Fig. 1d). In mosquitoes with parasites undergoing sporogony, there was no effect of age on locomotion throughout the 24 h period in the absence of human odour (ZT 11-12: $\chi^2_1 = 1.71$, $p = 0.18$; ZT 12-23: $\chi^2_1 = 0.31$, $p = 0.57$; ZT 23-24: $\chi^2_1 = 3.49$, $p = 0.06$, Supplementary Table 2.), except during photophase, in which infectious (14 dpi) mosquitoes demonstrated lower activity than infected (7 dpi) individuals (ZT 0-11: $\chi^2_1 = 21.97$, $p < 0.001$). However, in the presence of human odour, the infectious cohort (14 dpi) was more active at dawn (ZT 23-24: $\chi^2_1 = 15.40$, $p < 0.001$), while being less active than the younger mosquitoes throughout the rest of the diurnal period (ZT 0-11: $\chi^2_1 = 9.35$, $p = 0.002$; ZT 11-12: $\chi^2_1 = 20.02$, $p < 0.001$; ZT 12-23: $\chi^2_1 = 9.73$, $p = 0.001$, Supplementary Table 2.).

Individual locomotor profiles of mosquitoes at 7 dpi and 14 dpi differed significantly from that of age-matched uninfected controls (Fig. 1d). Infection significantly reduced the locomotor activity in 7 dpi females in the absence of human odour when compared with their uninfected counterparts during most time periods (ZT 0-11: $\chi^2_1 = 8.49$, $p = 0.03$; ZT 11-12: $\chi^2_1 = 1.008$, $p = 0.31$; ZT 12-23: $\chi^2_1 = 7.84$, $p = 0.005$; ZT 23-24: $\chi^2_1 = 13.78$, $p < 0.001$, Supplementary Table 1-2). Mosquitoes carrying the transmissible stage of the parasite (14 dpi) displayed a lower locomotor activity than the uninfected counterparts throughout photophase and scotophase (ZT 0-11: $\chi^2_1 = 5.92$, $p = 0.01$; ZT 12-23: $\chi^2_1 = 11.87$, $p < 0.001$, Supplementary Table 1-2), but not at dusk and dawn (ZT 11-12: $\chi^2_1 = 2.68$, $p = 0.10$; ; ZT 23-24: $\chi^2_1 = 2.20$, $p = 0.10$). In the presence of human odour, the infected (7 dpi) females showed lower activity than age-matched controls at dawn (ZT 23-24: $\chi^2_1 = 7.29$, $p = 0.006$), while the inverse activity pattern was observed during photophase (ZT 0-11: $\chi^2_1 = 12.28$, $p = 0.001$). The infectious mosquitoes (14 dpi) demonstrated a significantly increased locomotor activity at dawn (ZT 23-24: $\chi^2_1 = 23.78$, $p < 0.001$) and during scotophase (ZT 12-23: $\chi^2_1 = 25.42$, $p = 0.001$), while there was no difference in activity of the infectious and uninfected females during photophase (ZT 0-11: $\chi^2_1 = 2.26$, $p = 0.13$), with a significant reduction in dusk (ZT 11-12: $\chi^2_1 = 16.06$, $p < 0.001$, Supplementary Table 1- 2).

Plasmodium falciparum **modulates antennal transcript abundance**

Paired-end sequencing of each of the libraries constructed from antennal RNA, with a total of 2 400 antennae, generated an average mapping of 26 399 740 million cleaned reads per library. Out of the 13 832 coding genes annotated in the genome of *An. gambiae* (Agam4.10), a total of 10 115 transcripts were reliably detected above 1 transcript per million (TPM) mapped reads in the antennae, among all experimental and control groups, demonstrating an adequate level of coverage.

A principal component analysis (PCA) of the antennal transcripts was conducted to demonstrate the overall variation among the antennal transcriptomes (infected, infectious, and age-matched uninfected conditions; 4 replicates each; Fig. 2a). The PCA identified that 49.8% of the variation among the libraries was based on the relative infection status in each age group (PC1), while 15.3% of the variance was dependent on age and infection status relative to the controls (PC2; Fig. 2a). All of the biological replicates of the same age and infection status clustered tightly together in the principal component space, except for the infectious samples (14 dpi), demonstrating that variation in the libraries due to handling and processing was successfully minimised. The separation of the four libraries of the 14 dpi samples into two clusters correlates with demonstrated differences in parasite load of the mosquitoes (Fig. 2a; Supplementary Fig. 1).

Antennal transcripts were significantly differentially regulated between the two age-matched cohorts (6 187), of which 3 465 were differentially abundant only between 14 dpi and 14 dc, whereas in 7 dpi and 7 dc, 850 were differentially abundant (Fig. 2b). In total, 4 807 transcripts were significantly differentially regulated between the two age-matched control groups and the two groups carrying *P. falciparum* parasites, of which 1 071 transcripts were shared between them (Fig. 2c). Within the age-matched control

groups, 2 614 transcripts were uniquely regulated, whereas 1 122 were uniquely regulated within the two groups carrying *P. falciparum* parasites (Fig. 2c; Supplementary Fig. 2). Overall, differentially abundant transcripts were not condition-dependently regulated, except those regulated post-infection during both the infected and infectious stages (1 872), of which more than 80% of the transcripts were down-regulated post-infection (Fig. 2b).

To characterise the functional ontology of the differentially abundant genes in the antennae of infected and infectious mosquitoes with their age-matched controls, a gene ontology (GO) analysis of molecular function (level three) was conducted (Fig. 3). Of the 3 685 differentially abundant transcripts between the two uninfected control groups, the majority (>75%) were functionally classified as structural constituent of cuticle (GO: 0042302) and enzyme inhibitor activity (GO: 0004857; Fig. 3a). Both of these classes were more abundant in older compared to the younger individuals. None of the age-dependent GO terms identified in the uninfected cohort comparison (Fig. 3a) were detected in the pairwise comparisons of the antennal transcriptomes among infected, infectious and their age-matched controls. The three most represented functional classes in the pairwise comparisons between the infected and infectious groups with their age-matched controls, as well as between the infected and infectious antennal transcriptomes, were heterocyclic compound binding (GO: 1901363), organic cyclic compound binding (GO: 0097159) and ion binding (GO: 0043167; Fig. 3b). Two functional classes were regulated differently in the antenna of mosquitoes with a *P. falciparum* infection, odorant binding (GO: 0005549) and carbohydrate derivative binding (GO:0097367). The number of genes in the functional class odorant binding were regulated in both infected and infectious samples, while those in the carbohydrate derivative binding class were differentially regulated only in infectious samples (Fig. 3b).

A detailed analysis of the major chemosensory gene families associated with the odorant binding functional class, *Ors*, *Irs*, *Grs*, *Csps* and *Obps*, was conducted. As the mosquitoes aged, 34% of the chemosensory genes were significantly regulated in the antennae of uninfected females, with chemosensory receptors demonstrating higher abundance in the older females, while the binding proteins were both up- and down-regulated (Fig. 4). Following *P. falciparum* infection, the abundance of 18 *Obps* reduced with age, while only one chemosensory gene, *Ir41a*, demonstrated an increased abundance in the antennae of older females (Fig. 4). The differential abundance of these 18 *Obps*, along with 3 others, appear to be a result of an increased abundance at 7 dpi compared to the age-matched controls. The other chemosensory gene that was regulated at this age, *Ir7u*, was down-regulated upon infection.

The age-dependent regulation of chemosensory genes was affected following *P. falciparum* infection, as 73% of those genes that were up-regulated with age (Fig. 4 column 1), were shown not to increase in abundance post-infection (Fig. 4 columns 2 and 3), resulting in a higher abundance of these transcripts in the controls compared with those at 14 dpi (Fig. 4 column 4). The only exception to this was *Ir7u*, which was down-regulated at 7 dpi, exacerbating the decreased abundance observed at 14 dpi. Interestingly, when the 7 dpi mosquitoes were compared to 14 dpi, the abundance of the above-mentioned *Obps* were reversed, returning to pre-infection levels (Fig. 4 column 2).

Discussion

The stage of infection of *P. falciparum* in *An. gambiae* has a significant impact on vector-related behaviour, which affects malaria transmission and disease dynamics^{2,5,27,28}. This study confirms that infected mosquitoes are less active than infectious females, and furthermore identifies this change in behaviour to be a result of an interaction between infection stage and age, which can be modulated by the presence of human odour and diel rhythms. Transcriptomic profiling of antennal tissue of *P. falciparum*-infected and -infectious *An. gambiae* identifies members of several chemosensory gene families as changing with age and infection status, suggesting a molecular mechanism, which may regulate the sensitivity of the peripheral olfactory system of the vector to human odour during the sporogonic cycle. As age is intrinsic to the infection status of the female, controlling for age provides a more accurate indication of which behaviours and molecular correlates are regulated by *P. falciparum* infection.

Host seeking in mosquitoes is dependent on age^{20,21,23,29,30} and physiological state, including infection³⁻⁵. Our results demonstrated that the overall locomotor activity of uninfected females significantly reduced with age, in line with previous studies on mosquitoes^{21,23}, further emphasising the role of age in regulating host seeking. As mosquitoes age, energy reserves decrease, resulting in older females reducing their search activities and becoming more selective to reduce their energy requirements^{31,32}. The presence of human odour did not alter the overall locomotor activity, similar to that described by Cator et al.^{10,28}, suggesting that the difference in the level of activity with age is independent of the presence or absence of human odour. While previous studies on the effects of *Plasmodium* infection on mosquito host seeking have shown a reduced activity in infected compared with infectious mosquitoes, these have not controlled for the age of the insects^{10,11}. Our results indicate an interaction between age and infection in infected females that display a similar locomotor activity as an age-matched cohort, which is not seen in infectious individuals. When controlling for age and observing the effect of infection, infected females reduced their overall locomotor activity compared to age-matched controls in the absence of human odour, while in the presence of human odour there was no discernible difference in activity. In contrast, infectious mosquitoes increased their overall locomotor activity in the presence of human odour, but not in its absence, compared to age-matched controls. These results provide a rationale to explain why previous studies have presented opposing findings concerning the activity of oocyst-infected and sporozoite-infectious females^{10,33}. The mechanisms underlying the changes in host-seeking and blood-feeding behaviours are still unclear, however several hypotheses have been advanced, chief among these are that the response in infected and infectious females is a result of a general immune response, and/or a direct manipulation of the vector behaviour by the pathogen^{3,34-36}. The increased response to human odour by infectious mosquitoes coincides with increased feeding frequency, probing rate and meal size^{28,35,37,38}, and has substantial effects on malaria transmission ecology and overall disease dynamics³⁶.

Diel locomotor activity in mosquitoes is regulated by the age and infection status of female malaria vectors^{26,39,40}. In the presence and absence of human odour, younger uninfected mosquitoes (13 days post-emergence, dpe) were more active than the older mosquitoes (19 dpe) during photophase, scotophase and peaking at dawn⁴¹. The peak of activity for older females, on the other hand, occurred at dusk. The age-dependent change in diel locomotor activity was not modulated by the presence of human odour, which is in line with the diel patterns of the peripheral olfactory system responsiveness^{42–44}. The shift in peak activity of older females co-occurs with an increased propensity of older females to visit nectar sources at dusk⁴⁵, indicating a behavioural compartmentalisation due to an age-related reduction in energy reserves^{31,46}. Infection with *P. falciparum* altered the diel locomotor activity in both the presence and absence of human odour, generally decreasing activity in infected and infectious females in the absence of human odour, similar to that described in *An. stephensi* infected with *Plasmodium yoelii*⁴⁷. In the presence of human odour, infected and infectious females demonstrated a disrupted activity pattern associated with host seeking. In particular, infectious females shifted their odour-associated activity to scotophase and dawn, which is in line with observations of field-caught mosquitoes carrying sporozoites^{39,48}. By selectively increasing their activity during times of low-risk of predation and high access to human hosts, the infectious females increase the probability of a successful blood meal and pathogen transmission^{49–53}. While the infectivity of the bite is not affected by diel rhythms⁵⁴, the altered host-related activity pattern in infectious females may have explicit implications on the most effective deployment of vector control tools and thus our ability to affect malaria transmission dynamics efficiently³⁶. Whether this is a result of an active manipulation by the pathogen or a general immune response remains to be clarified^{3, 34–36,55,56}.

Age and *P. falciparum* infection significantly affected overall antennal transcript abundance, with infection systemically down-regulating gene abundance, and differentially affecting the functional ontological class odorant binding (GO:0005549). Among the differentially abundant transcripts, antennal abundance generally increased with age across all of the major olfactory-related gene families, whereas only *Obps*, the most abundantly expressed transcripts of the chemosensory gene families, exhibited higher abundance in infected compared with infectious females. Odorant binding proteins are involved in binding and transporting odorants to the receptors and in the overall sensitivity of the peripheral olfactory system by providing gain control^{22,57,58}. The *Obps* appear to play a major role in regulating the sensitivity and selectivity of the peripheral olfactory system of *An. gambiae* in response to aging and infection status, as 67% of the *Obp* transcripts reliably expressed in the antennae were age- and infection-dependently regulated. As the females age, older individuals exhibited a higher transcript abundance in two thirds of the differentially expressed classic, and all of the plus-C, classified *Obps*^{17,57}. The other third of the differentially regulated classic *Obps*, along with all 13 of the two-domain *Obps*^{17,57}, were expressed abundantly in 7 dpi antennae compared with not being reliably expressed in their age-matched controls and infectious females. Bimolecular protein interaction models indicate that the most energetically favourable ligands of the reliably expressed classic *Obps*, including those that were differentially abundant, are distinct sets of carboxylic acids, aromatics and terpenes¹⁷, known to regulate host and

floral seeking^{45,46}. While no clear pattern of ligand binding was apparent associated with aging, those classic *Obps* increasing in abundance 7 dpi generally did not include carboxylic acids among their modelled ligands, likely resulting in a reduced sensitivity of the peripheral olfactory system to these salient human odorants by infected females^{59–61}. This corresponds to the reduction in human odour-associated activity exhibited by infected females.

The functionally characterised differentially abundant olfactory receptors, *Or11*, *Or39* and *Ir41a*, all respond to human odour constituents^{16,21,62}. The receptor *Or39* has been postulated to modulate the onset of host seeking in *Anopheles coluzzii* through the concerted downregulation of the receptor and the sensitivity to sulcatone of its cognate olfactory sensory neuron²¹. Whether the lower abundance of *Or39*, in conjunction with *Or11*, observed in the antennae of infectious compared with uninfected females underlies the observed increased locomotor activity of infectious females in the presence of human odour, remains to be investigated. The only receptor to have a higher abundance in the antennae of infectious compared to infected females, *Ir41a*, responds to the human spermoscyclic amine pyrrolidine^{16,63}, suggesting that this receptor is involved in the increased attraction to human odour 14 dpi. The differential regulation of transcript abundance of genes modulating the chemosensory pathway in response to parasite development within the insect vector provides a mechanism by which the physiological¹⁴ and behavioural sensitivity^{4,10,11,33,35} to human odour can be modulated during the sporogonic cycle of *P. falciparum*.

Behavioural activity of *An. gambiae* and its expression of chemosensory-related genes are altered in an age- and *P. falciparum* stage-dependent manner. Irrespective of the mechanism regulating these changes, be it active manipulation by the parasite and/or general response to immune challenge, the differential behavioural response to human odour of females as they age or become infected significantly affects vectorial capacity and malaria transmission. Our ability to increase the resolution concerning which behavioural change is affected by age and/or infection, allows us to fine tune the epidemiological models for malaria transmission³³, and integrate both factors and their interactions into the development and implementation of control measures.

Methods

Ethics

Human blood (type O) was provided in citrate-phosphate-dextrose-adenine anti-coagulant/preservative, and serum (type AB) was obtained from the blood transfusion service (blood bank) at the Karolinska University Hospital (Solna, Sweden), in accordance with the Declaration of Helsinki, and approved by the Ethical Review Board in Stockholm (2011/850-32). The human materials were obtained anonymously from the blood bank and did not require an informed consent statement. The Swedish Work Environment Authority (Stockholm, Sweden, SU FV-2.10.2-2905-13/31-01-2022) approved the class 3 biological agent laboratory and its practices, including insectary design and equipment to work with *P. falciparum*-infected

mosquitoes. The regulations are mainly based on the EU directive 2000/54/EC on the protection of workers from risks related to exposure to biological agents at work.

Mosquito rearing and blood feeding

Anopheles gambiae (Keele) were reared and maintained at 27 ± 1 °C, $80 \pm 1\%$ relative humidity and a photoperiod of 12 h light: 12 h dark cycle, 11 h full light of ~ 300 lx and 11 h darkness, separated by 1 h dawn and dusk transitions, respectively. The Keele colony was established at Keele University as a result of balanced interbreeding of four *An. gambiae* s.l. strains: G3 from MacCarthy Island, the Gambia; and three Tanzanian strains, ZAN U from Zanzibar, Ifakara strain from Njage, and KIL from Marangu⁶⁴, and set up at Stockholm University in 2014. For experiments, pupae (ca. 350) were collected and transferred to cages (custom made: 30 cm $\times$ 20 cm $\times$ 20 cm) to emerge as adults, which were fed *ad libitum* on 5% glucose and 0.05% para-amino benzoic acid (PABA, Sigma-Aldrich, Steinheim, Germany). Females (5 dpe) were provided with a meal of cultured red blood cells (RBCs) through glass membrane-feeders connected to a custom-built heating system kept at 37 °C. The RBCs were washed with Roswell Park Memorial Institute (RPMI-1640) medium (Gibco™, ThermoFisher Scientific, Drieich, Germany), and stored in RPMI-1640 at 50% haematocrit at 4 °C prior to use.

Parasite culture and mosquito infection

The strain of *P. falciparum* used in this study, NF₅₄-SU, was initially isolated in the Netherlands and donated by Klavs Berzins. *In vivo* culturing of *P. falciparum* NF₅₄-SU followed the standard protocol⁶⁵. In short, *P. falciparum* NF₅₄-SU was cultured using washed RBCs (5% haematocrit) from type-O blood donors, and 10% serum mixed with RPMI-1640 medium (Gibco™, ThermoFisher Scientific), supplemented with 5.94 g l⁻¹ HEPES buffer (Sigma Aldrich), 2.1 g l⁻¹ sodium hydrogen carbonate (Sigma Aldrich), and 0.05 g l⁻¹ hypoxanthine (Sigma Aldrich). The cultures were kept in controlled conditions under gas (1% O₂, 3% CO₂ and 96% N₂) at 37 °C. *Plasmodium falciparum* gametocyte cultures were set up at 0.5-0.7% parasitaemia, 6% haematocrit in RPMI-1640 medium. On the day of experimental infection, uninfected RBCs were supplemented with *P. falciparum* gametocytes to a final gametocytaemia of ca. 0.7%, which generates a high infection prevalence (>50%) in *An. gambiae* s.s.⁶⁵. For each infectious meal, mixtures of gametocytes from 14- and 17-day gametocyte cultures were used. Twenty-eight groups of ca. 100 mosquitoes were fed on mature gametocyte-infected RBCs or on uninfected washed RBCs using a membrane feeder. All age-matched control females (7 and 14 days control ; 7 and 14 dc) were handled in the same manner as the infected and infectious cohorts, including access to mates, blood meals and oviposition substrates.

Mosquito odour-mediated locomotor activity

For recording the behavioural activity of the mosquitoes, DAM2 *Drosophila* activity monitors (TriKinetics Inc., Waltham, MA, USA) were used. Two hours prior to the experiments, mosquitoes were placed individually in glass tubes (65 mm $\times$ 7 mm), using an aspirator, for pre-conditioning. Cotton plugs

saturated with 5% glucose with 0.05% PABA (as above) were placed at one end of the glass tubes to provide *ad libitum* access to sustenance. Human odour was supplied by a previously worn cotton sock (ca. 24 h) placed inside a 200 ml glass wash bottle (Sigma Aldrich). The odour was introduced, via Teflon tubing, into the glass tubes through a MAN2 gas distribution manifold (TriKinetics Inc.). A clean unworn sock was similarly used as a control. Each experiment was replicated three times with infected and infectious mosquitoes assessed in parallel with age-matched controls, for a total of 384 mosquitoes assayed (96 per condition). Behavioural experiments were conducted in the class 3 biological agent laboratory (BSL3) within a climate chamber kept at 27 ± 1 °C, 70% relative humidity with a photoperiod of 11 h light: 11 h dark with 60 min of both dawn and dusk lighting conditions. Both assays were conducted over 24 h, with activity data recorded every 30 s.

The influence of age, the presence and absence of human odour, and the effect of the sporogonic stages of *P. falciparum* on the *An. gambiae* (7 and 14 days post infection; 7 and 14 dpi), diel locomotor activity was analysed using general linear mixed models (GLMM), in which the effect of replication (experimental blocks) and the weighting of multiple replications (random variables) were taken into account. For all results, the significance of the best maximal model containing all explanatory effects was evaluated by using likelihood ratio test. All analyses were performed using R (R Core team⁶⁶: R v.3.2.3 and RStudio 1.1.463⁶⁷). The *b*-estimated values of locomotor activity were estimated in all final significant models (GLMM: lmer function; in Fig 1b: The mixed model was build based on the effect of age as a main fixed variable combined with the random effect of the other variables; Fig 1c: The model was demonstrated the effect of the main fixed variable of infection including the rest of the variables as random variables; Fig 1d: The mixed model was estimated the effect of all four main fixed variables including age, infection, time, human odour, as well as the random effect of the experimental replication combined with the interactions among all these variables). In all analyses, all of the data met the assumptions of the test for normality and error homogeneity. Backward elimination was used for sequential removal of non-significant variables, to obtain the minimal statistically significant model. All graphical visualisations and statistical analyses were preformed using GraphPad Prism⁶⁸ and R (R Core team⁶⁶: R v.3.2.3 and RStudio 1.1.463⁶⁷). The precise levels of significance are presented in the results section, except the universally accepted $p < 0.001$.

Transcriptome analyses

Pairs of antennae from infected, infectious and age-matched control mosquitoes were collected from individual females following flash freezing on dry ice for <30 s, and placed into 24-well culture plates (ThermoFisher Scientific) in RNAlater (500 µl; Thermo Fisher Scientific). Individual carcasses (bodies missing antennae) of infected and infectious mosquitoes were also collected for subsequent qPCR analysis, to determine the parasite abundance of individual females (see below). Following the quantification of the parasite load, the antennae of successfully infected individuals were pooled into tubes of 150 pairs of antennae, according to the level of infection, and stored at -80 °C. Four replicates of each condition were collected. RNA libraries were constructed using a TruSeq RNA Library Prep Kit (Illumina, Berlin, Germany) from total RNA from the pooled antennae following extraction and DNase I

digestion (RNeasy Mini kit, Qiagen, Hilden, Germany). Quality and quantity control of total RNA aliquots was performed using a Nanodrop spectrophotometer (Thermo Fisher Scientific), an Agilent 2100 Bioanalyser (Santa Clara, CA, USA) and a Qubit 2.0 Fluorometer (Thermo Fisher Scientific), before sending to BGI Genomics (MGI Tech Co., Ltd., China) for Illumina paired-end indexed sequencing, using the Illumina HiSeq2000.

RNA-Seq and differential expression analyses

High-quality reads were determined using the following criteria: low quality reads were clipped from the start and end of each read using a sliding window and reads shorter than 40 nt were removed. The cleaned reads were mapped using CLC Genomics Workbench version 11 (<https://digitalinsights.qiagen.com>) to the *An. gambiae* genome Agam4 with reference to the Agam4.10 gene set (www.vectorbase.org). On average 94.11% of all reads mapped to the genome. Transcript abundance was reported as transcript per million (TPM) with a threshold level of abundance above 1 TPM. Differential transcript abundance was determined using the β -binomial general linear model algorithms in CLC Genomics Workbench 11. To control for false discovery rate (FDR), the Benjamini-Hochberg correction was applied⁶⁹. This analysis generated weighted fold changes (FC) and FDR-corrected *p*-values that were used to detect differential expression. Significantly differential gene abundance was determined based on a $FC \geq 2$ and a FDR-corrected *p*-value < 0.05.

Quantitative real-time polymerase chain reaction analysis

Quantitative real-time polymerase chain reaction (qPCR) was used to estimate the abundance of *P. falciparum* in individual mosquitoes subjected to an infectious meal at 7 and 14 dpi, and used for both behavioural and transcriptomic analyses, allowing quantification of the numbers of parasite genomes. In all analyses, a single body of an uninfected female mosquito was used as a negative control. Reactions were performed on a Roche Light Cycler using SYBR Green (Roche, Mannheim, Germany). DNA standards containing known numbers of *P. falciparum* parasites were produced from asexual cultures, and used to generate standard curves of *P. falciparum*. A similar statistical approach, as described for the behavioural analyses, was used to test for variation in the number of parasites (oocyst load and total number of sporozoites per whole mosquito body)⁷⁰. Given the highly over-dispersed nature of parasite abundance data, a negative binomial distribution was assumed in these analyses (GLMM: glmmADMB, nlme package; R Core team⁶⁶: R v.3.2.3 and RStudio 1.1.463⁶⁶). A backwards elimination approach was used to test for the significance of all fixed effects and interactions, while controlling for random variation due to replicate, as for the analyses of oocyst and sporozoite prevalence.

Declarations

Data availability

All data supporting the findings of this study are available within the article and its Supplementary Information files. The sequencing data are available at the GEO database (BioProject ID PRJNA756244),

and are available from the corresponding authors upon request.

Acknowledgements

We thank the Swedish Research Council for funding to RI (VR/2014-3331) and SNE (VR/2017-01229; VR/2017-05543 UFNW). We extend our thanks to Jeansson's Stiftelser (SJ-2018) for the valuable support for setting up the essential equipment of the BSL3 laboratory. We highly appreciate the donation of the NF₅₄-SU strain of *P. falciparum* by Prof. Klavs Berzins.

Author Contributions

RI and SNE conceived the study and designed the experiments. SNE and MH cultured the parasites, performed the mosquito infections as well as conducting the behavioural analyses. All statistical and modelling analyses for behavioural assays were carried out by MH and SNE for the deep analysing of the mixed model data. LRC, RM and SNE dissected the antenna and prepared RNA for Illumina sequencing. RI, SRH and SNE designed the transcriptome study, and MH and SRH performed the analyses in consultation SNE. MH, RI, SRH and SNE wrote the manuscript. All authors read and approved the manuscript.

Competing interests

The authors declare no competing interests. S. Noushin Emami is a co-founder of Molecular Attraction AB and there are no commercial aspects or financial gain to this company resulting from this study.

Materials and correspondence

Correspondence and materials requests should be directed to S. Noushin Emami, noushin.emami@su.se; and Rickard Ignell, rickard.ignell@slu.se

References

1. De Moraes, C. M. *et al.* Malaria-induced changes in host odors enhance mosquito attraction. *Proc Natl Acad Sci U S A* **111**, 11079-11084, doi:10.1073/pnas.1405617111 (2014).
2. Emami, S. N. *et al.* A key malaria metabolite modulates vector blood seeking, feeding, and susceptibility to infection. *Science* **10**, doi:10.1126/science.aah4563 (2017).
3. Hurd, H. Manipulation of medically important insect vectors by their parasites. *Annu Rev Entomol* **48**, 141-161, doi:10.1146/annurev.ento.48.091801.112722 (2003).
4. Lefevre, T. & Thomas, F. Behind the scene, something else is pulling the strings: emphasizing parasitic manipulation in vector-borne diseases. *Infect Genet Evol* **8**, 504-519, doi:10.1016/j.meegid.2007.05.008 (2008).
5. Tallon, A. K. *et al.* Dengue infection modulates locomotion and host seeking in *Aedes aegypti*. *PLoS Negl Trop Dis* **14**, e0008531, doi:10.1371/journal.pntd.0008531 (2020).

6. Botto-Mahan, C., Cattan, P. E. & Medel, R. Chagas disease parasite induces behavioural changes in the kissing bug *Mepraia spinolai*. *Acta Tropica* **98**, 219-223, doi:10.1016/j.actatropica.2006.05.005 (2006).
7. Lefevre, T. *et al.* Malaria *Plasmodium* agent induces alteration in the head proteome of their *Anopheles* mosquito host. *Proteomics* **7**, 1908-1915, doi:10.1002/pmic.200601021 (2007).
8. Biron, D. G. *et al.* Behavioural manipulation in a grasshopper harbouring hairworm: a proteomics approach. *Proceedings of the Royal Society B-Biological Sciences* **272**, 2117-2126, doi: 10.1098/rspb.2005.3213 (2005).
9. Beier, J. C. Malaria parasite development in mosquitoes. *Annual Review of Entomology* **43**, 519-543, doi: 10.1146/annurev.ento.43.1.519 (1998).
10. Cator, L. J. *et al.* 'Manipulation' without the parasite: altered feeding behaviour of mosquitoes is not dependent on infection with malaria parasites. *Proc Biol Sci* **280**, 20130711, doi:10.1098/rspb.2013.0711 (2013).
11. Smallegange, R. C. *et al.* Malaria infected mosquitoes express enhanced attraction to human odor. *Plos One* **8**, e63602, doi:10.1371/journal.pone.0063602 (2013).
12. Cornet, S., Nicot, A., Rivero, A. & Gandon, S. Avian malaria alters the dynamics of blood feeding in *Culex pipiens* mosquitoes. *Malar J* **18**, 82, doi:10.1186/s12936-019-2690-5 (2019).
13. Padilha, K. P. *et al.* Zika infection decreases *Aedes aegypti* locomotor activity but does not influence egg production or viability. *Mem Inst Oswaldo Cruz* **113**, e180290, doi:10.1590/0074-02760180290 (2018).
14. Stanczyk, N. M. *et al.* Species-specific alterations in *Anopheles* mosquito olfactory responses caused by *Plasmodium* infection. *Sci Rep* **9**, 3396, doi:10.1038/s41598-019-40074-y (2019).
15. Lu, T. *et al.* Odor coding in the maxillary palp of the malaria vector mosquito *Anopheles gambiae*. *Curr Biol* **17**, 1533-1544, doi:10.1016/j.cub.2007.07.062 (2007).
16. Pitts, R. J., Derryberry, S. L., Zhang, Z. & Zwiebel, L. J. Variant ionotropic receptors in the malaria vector mosquito *Anopheles gambiae* tuned to amines and carboxylic acids. *Sci Rep* **7**, 40297, doi:10.1038/srep40297 (2017).
17. Manoharan, M. *et al.* Comparative genomics of odorant binding proteins in *Anopheles gambiae*, *Aedes aegypti*, and *Culex quinquefasciatus*. *Genome Biol Evol* **5**, 163-180, doi:10.1093/gbe/evs131 (2013).
18. Bohbot, J. D. *et al.* Conservation of indole responsive odorant receptors in mosquitoes reveals an ancient olfactory trait. *Chem Senses* **36**, 149-160, doi:10.1093/chemse/bjq105 (2011).
19. Erdelyan, C. N., Mahood, T. H., Bader, T. S. & Whyard, S. Functional validation of the carbon dioxide receptor genes in *Aedes aegypti* mosquitoes using RNA interference. *Insect Mol Biol* **21**, 119-127, doi:10.1111/j.1365-2583.2011.01120.x (2012).
20. Omondi, B. A., Majeed, S. & Ignell, R. Functional development of carbon dioxide detection in the maxillary palp of *Anopheles gambiae*. *J Exp Biol* **218**, 2482-2488, doi:10.1242/jeb.116798 (2015).

21. Omondi, A. B., Ghaninia, M., Dawit, M., Svensson, T. & Ignell, R. Age-dependent regulation of host seeking in *Anopheles coluzzii*. *Sci Rep* **9**, 9699, doi:10.1038/s41598-019-46220-w (2019).
22. Rinker, D. C. *et al.* Blood meal-induced changes to antennal transcriptome profiles reveal shifts in odor sensitivities in *Anopheles gambiae*. *Proc Natl Acad Sci U S A* **110**, 8260-8265, doi:10.1073/pnas.1302562110 (2013).
23. Tallon, A. K., Hill, S. R. & Ignell, R. Sex and age modulate antennal chemosensory-related genes linked to the onset of host seeking in the yellow-fever mosquito, *Aedes aegypti*. *Sci Rep* **9**, 43, doi:10.1038/s41598-018-36550-6 (2019).
24. Hill, S. R. & Ignell, R. Modulation of odour-guided behaviour in mosquitoes. *Cell and Tissue Research* **383**, 195-206, doi:10.1007/s00441-020-03368-6 (2021).
25. Hill, S. R., Ghaninia, M. & Ignell, R. Blood meal induced regulation of gene expression in the maxillary palps, a chemosensory organ of the mosquito *Aedes aegypti*. *Front Ecol Evol* **7**, doi:ARTN 33610.3389/fevo.2019.00336 (2019).
26. Jones, M. D. R. & Gubbins, S. J. Changes in the circadian flight activity of the mosquito *Anopheles gambiae* in relation to insemination, feeding and oviposition. *Physiol Entomol* **3**, 213-220, doi: 10.1111/j.1365-3032.1978.tb00151.x (1978).
27. Koella, J. C., Sorensen, F. L. & Anderson, R. A. The malaria parasite, *Plasmodium falciparum*, increases the frequency of multiple feeding of its mosquito vector, *Anopheles gambiae*. *Proc Biol Sci* **265**, 763-768, doi:10.1098/rspb.1998.0358 (1998).
28. Anderson, R. A., Koella, J. C. & Hurd, H. The effect of *Plasmodium yoelii* nigeriensis infection on the feeding persistence of *Anopheles stephensi* Liston throughout the sporogonic cycle. *Proceedings of the Royal Society B-Biological Sciences* **266**, 1729-1733, doi:10.1098/rspb.1999.0839 (1999).
29. Bohbot, J. D., Durand, N. F., Vinyard, B. T. & Dickens, J. C. Functional development of the octenol response in *Aedes aegypti*. *Front Physiol* **4**, 39, doi:10.3389/fphys.2013.00039 (2013).
30. Foster, W. A. & Takken, W. Nectar-related vs. human-related volatiles: behavioural response and choice by female and male *Anopheles gambiae* (Diptera: Culicidae) between emergence and first feeding. *Bull Entomol Res* **94**, 145-157, doi:10.1079/ber2003288 (2004).
31. Rowley, W. A. & Graham, C. L. The effect of age on the flight performance of female *aedes aegypti* mosquitos. *J Insect Physiol* **14**, 719-728, doi:10.1016/0022-1910(68)90230-8 (1968).
32. Nayar, J. K. & Sauerman, D. M., Jr. A comparative study of flight performance and fuel utilization as a function of age in females of Florida mosquitoes. *J Insect Physiol* **19**, 1977-1988, doi:10.1016/0022-1910(73)90192-3 (1973).
33. Vantaux, A. *et al.* Host-seeking behaviors of mosquitoes experimentally infected with sympatric field isolates of the human malaria parasite *Plasmodium falciparum*: no evidence for host manipulation. *Front Ecol Evol* **3**, doi:10.3389/fevo.2015.00086 (2015).
34. Cator, L. J. *et al.* Immune response and insulin signalling alter mosquito feeding behaviour to enhance malaria transmission potential. *Sci Rep* **5**, 11947, doi:10.1038/srep11947 (2015).

35. Koella, J. C., Rieu, L. & Paul, R. E. L. Stage-specific manipulation of a mosquito's host-seeking behavior by the malaria parasite *Plasmodium gallinaceum*. *Behavioral Ecology* **13**, 816-820, doi:10.1093/beheco/13.6.816 (2002).
36. Cator, L. J., Lynch, P. A., Read, A. F. & Thomas, M. B. Do malaria parasites manipulate mosquitoes? *Trends in Parasitology* **28**, 466-470, doi:10.1016/J.Pt.2012.08.004 (2012).
37. Rossignol, P. A., Ribeiro, J. M. & Spielman, A. Increased intradermal probing time in sporozoite-infected mosquitoes. *Am J Trop Med Hyg* **33**, 17-20, doi: 10.4269/ajtmh.1984.33.17 (1984).
38. Rossignol, P. A., Ribeiro, J. M. & Spielman, A. Increased biting rate and reduced fertility in sporozoite-infected mosquitoes. *Am J Trop Med Hyg* **35**, 277-279, doi:10.4269/ajtmh.1986.35.277 (1986).
39. Gillies, M. T. Age-groups and the biting cycle in *Anopheles gambiae*. A preliminary investigation *B Entomol Res* **48**, 553-559, doi:10.1017/S0007485300002728 (1957).
40. Bockarie, M. J. & Dagoro, H. Are insecticide-treated bednets more protective against *Plasmodium falciparum* than *Plasmodium vivax*-infected mosquitoes? *Malar J* **5**, 15, doi:10.1186/1475-2875-5-15 (2006).
41. Lindsay, S. W., Shenton, F. C., Snow, R. W. & Greenwood, B. M. Responses of *Anopheles gambiae* complex mosquitoes to the use of untreated bednets in The Gambia. *Med Vet Entomol* **3**, 253-262, doi:10.1111/j.1365-2915.1989.tb00225.x (1989).
42. Rund, S. S. *et al.* Daily rhythms in antennal protein and olfactory sensitivity in the malaria mosquito *Anopheles gambiae*. *Sci Rep* **3**, 2494, doi:10.1038/srep02494 (2013).
43. Eilerts, D. F., VanderGiessen, M., Bose, E. A., Broxton, K. & Vinauger, C. Odor-specific daily rhythms in the olfactory sensitivity and behavior of *Aedes aegypti* Mosquitoes. *Insects* **9**, doi:10.3390/insects9040147 (2018).
44. Tanoue, S., Krishnan, P., Krishnan, B., Dryer, S. E. & Hardin, P. E. Circadian clocks in antennal neurons are necessary and sufficient for olfaction rhythms in *Drosophila*. *Curr Biol* **14**, 638-649, doi:10.1016/j.cub.2004.04.009 (2004).
45. Gary, R. E., Jr. & Foster, W. A. Diel timing and frequency of sugar feeding in the mosquito *Anopheles gambiae*, depending on sex, gonotrophic state and resource availability. *Med Vet Entomol* **20**, 308-316, doi:10.1111/j.1365-2915.2006.00638.x (2006).
46. Foster, W. A. Mosquito sugar feeding and reproductive energetics. *Annu Rev Entomol* **40**, 443-474, doi:10.1146/annurev.en.40.010195.002303 (1995).
47. Rowland, M. Changes in the circadian flight activity of the mosquito *Anopheles-Stephensi* Associated with Insemination, blood-feeding, oviposition and nocturnal Light-Intensity. *Physiol Entomol* **14**, 77-84, doi:DOI 10.1111/j.1365-3032.1989.tb00939.x (1989).
48. Djenontin, A., Bouraima, A., Soares, C., Egbinola, S. & Cottrell, G. Human biting rhythm of *Anopheles gambiae* Giles, 1902 (Diptera: *Culicidae*) and sleeping behaviour of pregnant women in a lagoon area in Southern Benin. *BMC Res Notes* **14**, 200, doi:10.1186/s13104-021-05615-7 (2021).
49. O'Donnell, A. J., Rund, S. S. C. & Reece, S. E. Time-of-day of blood-feeding: effects on mosquito life history and malaria transmission. *Parasit Vectors* **12**, 301, doi:10.1186/s13071-019-3513-9 (2019).

50. Takken, W., van Loon, J. J. & Adam, W. Inhibition of host-seeking response and olfactory responsiveness in *Anopheles gambiae* following blood feeding. *J Insect Physiol* **47**, 303-310, (2001).
51. Reece, S. E., Prior, K. F. & Mideo, N. The Life and Times of Parasites: Rhythms in Strategies for Within-host Survival and Between-host Transmission. *J Biol Rhythms* **32**, 516-533, doi:10.1177/0748730417718904 (2017).
52. Martinez-Bakker, M. & Helm, B. The influence of biological rhythms on host-parasite interactions. *Trends Ecol Evol* **30**, 314-326, doi:10.1016/j.tree.2015.03.012 (2015).
53. Mideo, N., Reece, S. E., Smith, A. L. & Metcalf, C. J. The Cinderella syndrome: why do malaria-infected cells burst at midnight? *Trends Parasitol* **29**, 10-16, doi:10.1016/j.pt.2012.10.006 (2013).
54. Githeko, A. K. *et al.* Confirmation that *Plasmodium falciparum* has aperiodic infectivity to *Anopheles gambiae*. *Med Vet Entomol* **7**, 373-376, doi: 10.1111/j.1365-2915.1993.tb00708.x (1993).
55. Poulin, R. The evolution of parasite manipulation of host behavior - a theoretical-analysis. *Parasitology* **109**, S109-S118, doi: 10.1017/S0031182000085127 (1994).
56. Emami, S. N., Hajkazemian, M. & Mozuraitis, R. Can *Plasmodium*'s tricks for enhancing its transmission be turned against the parasite? New hopes for vector control. *Pathog Glob Health* **113**, 325-335, doi:10.1080/20477724.2019.1703398 (2019).
57. Li, Z. X., Pickett, J. A., Field, L. M. & Zhou, J. J. Identification and expression of odorant-binding proteins of the malaria-carrying mosquitoes *Anopheles gambiae* and *Anopheles arabiensis*. *Arch Insect Biochem Physiol* **58**, 175-189, doi:10.1002/arch.20047 (2005).
58. Larter, N. K., Sun, J. S. & Carlson, J. R. Organization and function of *Drosophila* odorant binding proteins. *Elife* **5**, doi:10.7554/eLife.20242 (2016).
59. Acree, F., Jr., Turner, R. B., Gouck, H. K., Beroza, M. & Smith, N. L-Lactic acid: a mosquito attractant isolated from humans. *Science* **161**, 1346-1347, doi:10.1126/science.161.3848.1346 (1968).
60. Raji, J. I. *et al.* *Aedes aegypti* mosquitoes detect acidic volatiles found in human odor using the IR8a Pathway. *Curr Biol* **29**, 1253-1262 e1257, doi:10.1016/j.cub.2019.02.045 (2019).
61. Cork, A. & Park, K. C. Identification of electrophysiologically-active compounds for the malaria mosquito, *Anopheles gambiae*, in human sweat extracts. *Med Vet Entomol* **10**, 269-276, doi:10.1111/j.1365-2915.1996.tb00742.x (1996).
62. Carey, A. F., Wang, G., Su, C. Y., Zwiebel, L. J. & Carlson, J. R. Odorant reception in the malaria mosquito *Anopheles gambiae*. *Nature* **464**, 66-71, doi:10.1038/nature08834 (2010).
63. Amoore, J. E., Forrester, L. J. & Buttery, R. G. Specific Anosmia to 1-Pyrroline: The Spermos Primary Odor. *J Chem Ecol* **1**, 299-310, doi:10.1007/Bf00988831 (1975).
64. Hurd, H., Taylor, P. J., Adams, D., Underhill, A. & Eggleston, P. Evaluating the costs of mosquito resistance to malaria parasites. *Evolution* **59**, 2560-2572, doi: 10.1554/05-211.1 (2005).
65. Carter, R., Ranford-Cartwright, L. & Alano, P. The culture and preparation of gametocytes of *Plasmodium falciparum* for immunochemical, molecular, and mosquito infectivity studies. *Methods Mol Biol* **21**, 67-88, doi:10.1385/0-89603-239-6:67 (1993).

66. R Core Team. *R: a language and environment for statistical computing*, <<https://www.R-project.org/>> (2020).

67. RStudio-Team. RStudio: Integrated Development for R. . *RStudio*, doi: <http://www.rstudio.com/> (2021).

68. GraphPad Prism. *GraphPad Software version 6.04*, <<http://www.graphpad.com/>> (2020).

69. Trapnell, C. *et al.* Differential analysis of gene regulation at transcript resolution with RNA-seq. *Nat Biotechnol* **31**, 46-53, doi:10.1038/nbt.2450 (2013).

70. Emami, S. N., Ranford-Cartwright, L. C. & Ferguson, H. M. The transmission potential of malaria-infected mosquitoes (*An.gambiae*-Keele, *An.arabiensis*-Ifakara) is altered by the vertebrate blood type they consume during parasite development. *Sci Rep* **7**, 40520, doi:10.1038/srep40520 (2017).

Figures

Figure 1

Plasmodium falciparum infection stage differentially affects mosquito locomotor activity. Schematic design of the locomotor assay (a) used to assess the activity of female *Anopheles gambiae*, 7 days post-infection (dpi) and 14 dpi with *P. falciparum* compared to age-matched uninfected females (days control, dc), in the presence (with human odour) or absence of human odour (w/o human odour). Individual locomotor profiles (n = 384), assessed for variation using a redundancy analysis, demonstrated a differential effect of age (The mixed model was build based on the effect of age as a main fixed variable combined with the random effect of the other variables; β -lmer $\pm$ SE= age + experimental replication|1; b) and infection (The model was demonstrated the effect of the main fixed variable of infection including the rest of the variables as random variables ; β -lmer $\pm$ SE= infection + experimental replication|1; c) on locomotor activity in the presence and absence of human odour. (d) Diel locomotor activity profiles of 7 and 14 dpi females, along with age-matched controls, demonstrated effects of both age and infection in the presence and absence of human odour during informative activity periods (photophase, Zeitgeber Time (ZT) 0-11; dusk, ZT 11-12; scotophase, ZT 12-23; and dawn, ZT 23-34) over a 24 hour period (The mixed model was estimated the effect of all four main fixed variables including age, infection, time, human odour, as well as the random effect of the experimental replication combined with the interactions among all these variables; β -lmer $\pm$ SE= age + infection + time + human odour + experimental replication|1). Significant effects were determined using GLMM models (lmer). Bars represented by β -estimation generated by the mixed model $\pm$ SE (β -lmer $\pm$ SE); asterisks denote significant differences (*p < 0.05; **p < 0.01; ***p < 0.001; ns, non-significant). Letters indicate the pairwise comparisons between infected and uninfected groups; a = p < 0.001, b = p < 0.01, c = p < 0.05, d = non-significant).

Figure 2

Plasmodium falciparum infection stage affects antennal transcript abundance in *Anopheles gambiae*. Principal component analysis (a) of the overall transcript abundance in the antennae of infected (7 days post-infection, dpi) and infectious (14 dpi) mosquitoes compared to age-matched uninfected mosquitoes (7 days control, dc and 14 dc). *Plasmodium falciparum* infection load is indicated (see Supplementary Figure 1). Proportional Venn diagrams (b-c) depicting pairwise comparisons among the antennal

transcriptomes of infected, infectious, and age-matched uninfected females. Overlapping regions represent the subsets of transcripts that are shared between the different conditions. Significant differences in transcript abundance were determined as a fold change of greater than 1.5, and an FDR-corrected p-value < 0.05.

Figure 3

Plasmodium falciparum infection stage affects the functional ontology of differentially abundant antennal transcripts. A level 3 gene ontology analysis of molecular function of Anopheles gambiae antennal transcriptomes comparing those of 13 and 19 days post-emergence females (7 days control, dc and 14 dc, respectively) (a), as well as those of 7 and 14 days post-infection (dpi) females along with age-matched controls (b). Gene ontology classes that were included comprised greater than 1% of the total number of differentially abundant genes identified. Those with less than 1 % are grouped together in “other”.

Figure 4

Antennal chemosensory transcript abundance in female *Anopheles gambiae* is affected by *Plasmodium falciparum* infection stage. Chemosensory transcripts demonstrating reliable expression (>1 TPM) in the antennae of infected (7 days post-infection; dpi) and infectious (14 dpi) female mosquitoes, along with their age-matched controls (7 days control; 7dc and 14 dc) were analysed in pairwise comparisons displayed as log₂ fold change in abundance. Chemosensory genes are arranged into gene families, odorant binding proteins (OBPs), chemosensory proteins (CSPs), odorant receptors (Ors), ionotropic receptors (Irs) and gustatory receptors (Grs), in descending order of abundance (grey gradient bar, log₁₀ TPM). Significant differences in abundance (fold change > 2 ; FDR-corrected $p < 0.05$) are indicated by black dots.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementaryinformation101121.pdf](#)