Cytokine and Chemokine mRNA Expressions After Mycobacterium Tuberculosis-Specific Antigen Stimulation in Whole Blood From Hemodialysis Patients with Latent Tuberculosis Infection

Ji Young Park

Department of Internal Medicine, Park Clinic, Busan

Sung-Bae Park

Department of Clinical Laboratory Science, College of Health Sciences, Catholic University of Pusan, Busan

Heechul Park

Department of Clinical Laboratory Science, College of Health Sciences, Catholic University of Pusan, Busan

Jungho Kim

Department of Clinical Laboratory Science, College of Health Sciences, Catholic University of Pusan, Busan

Ye-Na Kim

Department of Internal Medicine, Kosin University Gospel Hospital, Busan

Sunghyun Kim (✉ shkim0423@cup.ac.kr)

Department of Clinical Laboratory Science, College of Health Sciences, Catholic University of Pusan, Busan

Research Article

Keywords: Latent tuberculosis infection, cytokine, chemokine, mRNA expression, hemodialysis

Posted Date: December 2nd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-111540/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

There have been few reports on the kinetics of hemodialyzed (HD) patients' immune responses in latent tuberculosis infection (LTBI). Nine cytokines and chemokines were screened through relative mRNA expression levels in whole blood samples after stimulation with *Mycobacterium tuberculosis* (MTB)-specific antigens from HD patients with LTBI (HD/LTBI), HD patients without LTBI, and healthy individuals and results were compared with QuantiFERON-TB Gold In-Tube (QFT-GIT) test. Only the CCL11 mRNA expression level of the HD/LTBI group was significantly higher than the other two groups ($P = 0.028$). CCL11 mRNA expression level of the > 0.7 IU/mL group in QFT-GIT test was significantly higher than the < 0.2 IU/mL group in QFT-GIT test and the 0.2-0.7 IU/mL group in QFT-GIT test ($P = 0.0043$). The present study reveals that the relative mRNA expression of CCL11 was statistically different in LTBI based on the current cut-off value (i.e., ≥ 0.35 IU/mL) and in the > 0.7 IU/mL group. These results suggest that CCL11 might be an alternative biomarker for LTBI diagnosis in HD patients.

Introduction

There have been a few reports on the accurate prevalence of tuberculosis (TB) in high-risk groups, such as the elderly, patients with diabetes or chronic kidney disease, and etc. According to the Korean Society of Nephrology end-stage renal disease (ESRD) registry data in 2018, TB incidence was 0.5% in the hemodialyzed (HD) patients¹. Patients with ESRD receiving dialysis is a risk factor of active TB, which is 25.3 times more prevalent in this group than in individuals with normal renal function². This relationship is likely associated with alterations in the immune system due to various factors including uremia.

Although interferon gamma (IFN- γ) release assays (IGRAs) are currently used to identify LTBI cases and are useful even in immunocompromised hosts, it should be interpreted carefully because of low sensitivity and specificity in a high-burden setting³. Moreover, 7.7% of IGRA display discordant results in a duplicated test, and most are located within a range near the cut-off value^{4,5}. Additionally, there is a high proportion of IGRA reversion in serial follow-up studies, and lower positive IGRA response is associated with reversion^{6,7}. Thus, some investigators have suggested that use of alternative borderline zone instead of the cut-off value in commercial IGRA test to avoid over-diagnosing LTBI⁸. However, little is known about the impact of impaired cellular immunity in dialysis patients in terms of IGRA results.

In addition to plasma IFN- γ level after *Mycobacterium tuberculosis* (MTB)-specific antigen stimulation, several cytokines and chemokines have been investigated as potential biomarkers for MTB infection and disease. However, sparse research on cytokines and chemokines in LTBI or different forms of TB in immunocompromised patients such as ESRD patients receiving dialysis has been conducted. The kinetics of immune responses in HD patients with LTBI has not been fully investigated.

In the present study, mRNA expression levels of nine immune markers including IFN- γ , tumor necrosis factor- α (TNF- α), interleukin-2 receptor (IL-2R), C-X-C motif ligand 9 (CXCL9), CXCL11, IFN- γ -induced protein 10 (IP-10), C-C motif chemokine 11 (CCL11), granulocyte-macrophage-colony stimulating factor

(GM-CSF), and TNF receptor (TNFR) were analyzed using quantitative reverse transcription polymerase chain reaction (qRT-PCR) TaqMan probe assay with whole blood samples after MTB-specific antigen stimulation from HD patients and the healthy control groups to identify biomarker candidates in the discovery stage toward the development of a diagnostic test to distinguish HD patients with LTBI from healthy controls (HC).

Materials And Methods

Study design and population

All enrolled individuals were more than 19 years old with ESRD and were undergoing more than three months hemodialysis treatment or were HCs. Those with HIV infection, liver cirrhosis of Child-Pugh class C, cancer or autoimmune disease and those who had received chemotherapy within the last three months and who had any history of active TB treatment were excluded. HD patients without repeated QFT-GIT test (Qiagen, Hilden, Germany) within six months were excluded in the examination of the reversion and conversion in the HD group. LTBI was diagnosed based on positive results of QFT-GIT test, with no symptoms or signs of active TB and no history of previous TB. HCs all displayed no symptoms or signs of active TB and no signs of active TB on chest radiographs. Whole blood bioresources of HD patients were collected from Kosin University Gospel Hospital Biobank. All the blood samples of HD patients were the remaining samples after IGRA test, and the test was performed before hemodialysis.

Study procedures

Demographic and clinical data were collected with an electronic medical chart review that included the following: sex, age, TB contact history, prior TB, BCG scar or vaccination history, co-morbidity diabetes mellitus (DM) and ischemic heart disease (IHD), smoking history, body mass index (BMI), chest X-ray (CXR), and LTBI treatment history if available. Prior TB contact was defined as a contact history of any kind with active lifelong TB patients. Prior TB was defined as a previous treatment history that included treatment for active TB. Findings suggestive of abnormal CXR were fibrotic infiltrates with pleural thickening or calcified nodules over the upper lung fields or other fibrotic lesions documented from previous TB.

Low and normal BMI were defined based on the categories used by the National Heart, Lung, and Blood Institute and the World Health Organization (WHO) (low BMI < 18.5 kg/m² and normal BMI between 18.5 and 24.9 kg/m²).

QuantiFERON-TB Gold In-Tube (QFT-GIT) assay

A peripheral whole blood samples were collected in three QFT-GIT collection tubes (nil, MTB-specific antigens, and T-cell mitogen, which contain lithium heparin). The QFT-GIT collection tubes were processed within two hour and incubated for 24 hours at 37°C. Plasma was collected by centrifugation at 2,000 x g for 5 min and stored at -80 °C until it was assayed.

After incubation, 50 µL of 1x conjugate was dispensed into an ELISA plate well before dispensing 50 µL of plasma samples into the prepared plate wells. Plasma samples, standards and conjugate were mixed well using a microwell plate shaker for 1 min. Each plate was covered with a lid and incubated at room temperature (RT) for 120 min. After incubation, 400 µL of 1x wash buffer (WB) was applied to each well and washing was performed at least six times. Extra WB was removed by tapping the plate into a towel. Then 100 µL of enzyme substrate solution was dispensed into each well and mixed in a microwell plate shaker. The plate was covered with lid and incubated at RT for 30 min after which 50 µL of enzyme-stopping solution was added to each well. The reaction was stopped and the optical density value of each well was measured within 5 min using a microwell plate reader equipped with a 450 nm filter and a 620-650 nm reference filter. T-cell mitogen stimulation served as an intrinsic control for blood sample quality. An IFN-γ concentration of ≥ 0.35 IU/mL (after subtraction of nil control IFN-γ), following exposure to MTB-specific antigens was, considered positive for QFT-GIT test; a concentration of < 0.35 IU/mL was considered negative. If the IFN-γ response to mitogen was < 0.5 IU/mL higher than that for the nil control, or > 8 IU/mL higher than that for the nil control, the result was deemed indeterminate⁹.

Total RNA isolation and reverse transcription

In order to analyze relative mRNA expressions of IFN-γ, TNF-α, IL-2R, CXCL9, IP-10, CXCL11, CCL11, GM-CSF, and TNFR after MTB-specific antigen stimulation, a blood cell pellets were treated using 500 µL of RNA/DNA stabilization reagent for Blood/Bone marrow (Roche Diagnostics, Mannheim, Germany). And then, MagNA Pure LC RNA Isolation Kit-High Performance kit (Roche Diagnostics) was used for RNA isolation¹⁰. The sample was centrifuged at 8,000 xg for 15 sec, and after centrifugation, the flow through in the collection tube was discarded. 50 µL of DNase $\times$ was then added, and the sample was incubated for 15 min at RT. Next, 500 µL of wash buffer $\times$ was added to the upper reservoir of the filter tube assembly and the sample was again centrifuged under the same conditions as above. Flow through was discarded before again washing with wash buffer $\times$ under the same conditions. After two washes, 200 µL of wash buffer was again added and the sample was centrifuged for 2 min at 13,000 xg to remove any residual wash buffer. Then the collection tube was discarded and the filter tube was inserted into a clean, sterile 1.5 mL microcentrifuge tube. After the transfer, 50 µL of elution buffer was added to the upper reservoir of the filter tube and it was centrifuged at 8,000 xg for 1 min. Finally, the concentration and purity of extracted total RNA was measured with a NanoDrop 2,000 spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA), and the samples were stored at -80°C until further use.

Next, complementary DNA (cDNA) was synthesized using a Moloney murine leukemia virus reverse transcriptase (M-MLV) reverse transcriptase kit (Invitrogen, Carlsbad, CA, USA). 10 µL of total RNA was added to a master mix containing 4 µL of 2.5 mM dNTP MIX AND 1 µL of random hexamer in a PCR tube, and the reaction mixture was incubated at 65 °C for 5 min before being quickly chilled on ice. Next, a mixture of 4 µL of 5X first-strand synthesis buffer, 2 µL of 0.1 M dithiothreitol (DTT), and 1 µL of M-MLV-RT were added to the previous reaction mixture in PCR tubes and then incubated at 25 °C for 1min, 50 °C for 50 min, and 70 °C for 15 min. All reactions were performed using a SimpliAmp (Life Technologies, Carlsbad, CA, USA) thermal cycler.

qRT-PCR TaqMan probe assay targeting multiple immune marker mRNAs

Oligonucleotide primer sets and TaqMan probe were designed using Primer3 software (http://biotools.umassmed.edu/bioapps/primer3_www.cgi) for performing the qRT-PCR TaqMan probe assay targeting multiple immune markers include IFN- γ , TNF- α , IL-2R, CXCL9, IP-10, CXCL11, CCL11, GM-CSF, and TNFR. The oligonucleotide sequences of the PCR primer pairs and TaqMan probes were based on the National Center for Biotechnology Information (NCBI) reference mRNA or cDNA sequences of human target genes. Quantitative PCR was carried out with THUNDERBIRD™ Probe qPCR Master Mix (TOYOBO, Osaka, Japan) using 3 μ L of cDNA as the template in a total volume of 20 μ L. The thermal cycling conditions were 10 min at 95°C, followed by 40 cycles of 10 sec at 95°C and 30 sec at 60°C. All reactions were performed using an ABI 7500 FAST Real-time PCR system (Applied Biosystems, Waltham, MA, USA) and QuantSTUDIO™ 7 Flex Real-time PCR system (Thermo Fisher Scientific, Waltham, MA, USA). Relative quantification was performed by the 2-ddCT method; relative expression was calculated as the ratio between the mean threshold cycle (CT) values of the target genes and reference gene (GAPDH) in each stimulated sample in relation to a reference sample (not stimulated)¹¹.

Statistical analysis

Geometric means and medians were used for measurements of central tendency. Data were expressed as mean $\pm$ SD for continuous variables and percentages for categorical variables. Paired numeric data were compared using Student *t*-test or one-way analysis of variance (ANOVA) tests with Dunnett's multiple comparisons. Chi-square analysis or Fisher's exact test were used to evaluate differences in proportions. The kappa coefficient was calculated to check the correlation between two QFT-GITs. Statistical analyses were performed using SPSS version 24.0 (SPSS Inc., Chicago, IL, USA) and Graphpad Prism 8.2.1 software (Graphpad Software 8.2.1, San Diego, CA, USA). All tests of significance were two sided; *P* values < 0.05 were considered statistically significant.

Results

Characteristics of the study population

The present study included 28 HD patients with LTBI (HD/LTBI), 47 HD patients without LTBI (HD/normal) and 48 HCs (see Table 1). The average age was 62.0 ± 11.18 years for the HD/LTBI group, 60.70 ± 12.46 years for the HD/normal group, and 23.94 ± 2.72 years for the HC group. HD groups were older than the HC group; however, there were no difference between HD/LTBI and HD/normal groups. The male proportion of HD/LTBI group (18/28) was significantly higher than the HD/normal (16/47) and HC groups (21/48) (*P* = 0.038).

Previous TB contact and abnormal CXR lesion of the HD/LTBI groups was more frequent than the other two groups (*P* = 0.002 and *P* = 0.005, respectively); however, BCG vaccination or scar of HD/LTBI groups was less frequent than other two groups (*P* < 0.001). There was no significant difference in terms of

underlying diseases like DM and IHD between HD/LTBI and HD/normal groups. Low BMI in HD/LTBI groups was significantly more frequent than in the HD/normal group ($P = 0.012$).

The amount of IFN- γ released after T-cell mitogen stimulation in QFT-GIT was analyzed in HD and HC groups (Table 1). There was statistically significant difference among the HD/LTBI, HD/normal, and HC groups ($P = 0.005$). In addition, the amounts of IFN- γ in the HD group were lower than in the HC group ($P = 0.001$).

QFT-GIT test results and previous QFT-GIT test results within 6 months

Most of the HD patients had repeated QFT-GIT test within 4 to 6 months. QFT-GIT test responses according to positive (> 0.7 IU/mL), borderline ($0.2-0.7$ IU/mL), and negative (< 0.2 IU/mL) criteria were as follows: Negative results were 97.9% in the HD/normal group, and positive results were 75% in the HD/LTBI group (see Table 2). Borderline results were 25% in the HD/LTBI group, 2.1% in the HD/normal group, and 4.2% in the HC group.

Previous QFT-GIT test response in HD patients was as follows: Positive results were 92.9% in the HD/LTBI group and 21.3% in the HD/normal group. Negative results were 3.6% in the HD/LTBI group and 51.5% in the HD/normal group. Borderline results were 3.6% in the HD/LTBI group and 27.7% in HD/normal group.

Based on the cut-off value of 0.35 IU/mL, conversion rate was 3.1% (1/32), and reversion rate was 37.2% (16/43). In addition, the reversion rate of the 0.35-0.7 IU/mL response group (85.7%, 6/7) was statistically higher than the > 0.7 IU/mL response group (27.8%, 10/36) ($P = 0.0162$). According to the three QFT-GIT test response groups, 92.7% (13/14) of previous 0.35-0.7 IU/mL response cases and 25% (9/36) of previous > 0.7 IU/mL response cases changed to negative results following QFT-GIT test (see Fig 1, $P < 0.001$).

Comparison of target gene mRNA expression levels after stimulation of MTB-specific Ags for 24 hours according to LTBI diagnosis

There were no statistically significant differences for mRNA expression levels of IFN- γ , TNF- α , IL-2R, CXCL9, CXCL11, IP-10, GM-CSF, and TNFR (Figure 1-a, b, c, d, e, f, h, i) among the three groups. However, there were statistically significant difference in only CCL11 mRNA expression levels ($P = 0.028$): mRNA expression level was higher in the HD/LTBI group (126.36 ± 406.25) than in the other two groups (HD/normal, 6.12 ± 17.45 vs HC, 19.58 ± 55.86 , Figure 1-g). Within HD groups, the CCL11 of the HD/LTBI group was significantly higher than the HD/normal group ($P = 0.0208$).

Comparison of target gene mRNA expression levels after stimulation of MTB-specific antigens for 24 hours according to QFT-GIT response groups

There was no significant difference among three QFT-GIT response groups in terms of IFN- γ , TNF- α , IL-2R, CXCL9, CXCL11, and IP-10 (figure 2-a, b, c, d, e, f). However, CCL11 mRNA expression levels in the > 0.7 IU/mL response group (168 ± 464) was significantly higher than in the < 0.2 IU/mL response group

(13.1 ± 42.5) and 0.2 to 0.7 IU/mL response group (2.55 ± 6.43 , $P = 0.0043$, Figure 2-g). Compared to CCL11 mRNA expression level, GM-CSF mRNA expression levels in the 0.2 to 0.7 IU/mL response group ($1,512,106 \pm 4,781,492$) were significantly higher than the < 0.2 IU/mL response group (626 ± 4129) and > 0.7 IU/mL response group (15 ± 53.9 , $P = 0.0029$, Figure 2-h). TNFR mRNA expression levels in the < 0.2 IU/mL response group ($227,937 \pm 720,687$) were significantly higher than the QFT-GIT-1 group (2.31 ± 4.44) and > 0.7 IU/mL response group (4.01 ± 8.81 , $P=0.0029$, Figure 2-i). Although there was no significant difference ($P = 0.0981$), mRNA expression level of IL-2R in the > 0.7 IU/mL response group was higher than in the less than 0.7 IU/mL groups (Figure 2-c).

Comparison of Target gene mRNA expression levels after stimulation of MTB-specific antigens for 24 hours according to QFT-GIT response groups in HD

There was statistically significant difference in mRNA expression of CCL11 among QFT-GIT test response groups in HD patients ($P = 0.041$), for which the expression levels in the > 0.7 IU/mL response group was higher than that of the other two groups (Figure 3-g).

The mRNA expressions of GM-CSF and TNFR differed significantly ($P = 0.0132$), and the expression levels of the 0.2 to 0.7 IU/mL response group was higher than the other two groups compared to CCL11 (Figure 3- h, Figure 3-i). The mean value of TNFR mRNA expression in the 0.2 to 0.7 IU/mL response group was higher than the other two groups; however, the median value of TNFR mRNA expression in the > 0.7 IU/mL response group was higher than in the other two groups. If one case with extreme values of GM-CSF and TNFR mRNA expression among the 0.2 to 0.7 IU/mL response group was excluded, there were no statistical differences for GM-CSF and TNFR mRNA expression among the three QFT-GIT test response groups. However, there were significant differences of GM-CSF and TNFR mRNA expression between the < 0.2 IU/mL response group and 0.2 to 0.7 IU/mL response group ($P = 0.0083$). There was no significant difference among three QFT-GIT response groups in terms of IFN- γ , TNF- α , IL-2R, CXCL9, CXCL11, and IP-10 (Figure 3-a, b, c, d, e, f).

Comparison of clinical features, and mRNA expression of cytokine and chemokine between IGRA reversion and persistent group in HD group

There were no statistically significant differences between the reversion group ($n = 16$) and the persistent group ($n = 58$) in terms of sex, age, DM, IHD, plasma IFN- γ protein level after T-cell mitogen stimulation and LTBI treatment. Also, there was no statistically significant differences in terms of cytokines and chemokines.

Discussion

Although an accurate detection of LTBI among HD patients is important, little is known about the kinetics of HD immune response in LTBI. Our study reveals several features concerning the differences in MTB-specific antigen-stimulated mRNA expression of cytokines and chemokines between HD and HC groups.

The IFN- γ mRNA expression level of MTB-specific antigen-stimulated whole blood sample was not significantly different among the three groups according to LTBI diagnosis and QFT-GIT test response, although the mRNA expression levels of HC groups were higher than HD groups ($P = 0.3609$). However, for comparison within HD groups, the mRNA expression level of IFN- γ for the > 0.7 IU/mL response group was higher than the < 0.7 IU/mL response group ($P = 0.0988$). The IFN- γ mRNA expression decreased in HD patients, which may cause false negative result for QFT-GIT test, although there was no statistical significance.

TNF- α and IFN- γ signaling have gained particular attention because their functions in host resistance to TB have been well documented in both mouse models and infected humans. Consistent with the previous study in patients with immune-mediated inflammatory disease, this study revealed that the amount of IFN- γ after mitogen stimulation was lower in the HD group than in the HC group¹².

Establishment of the TB granuloma is controlled by the synchronized expression of various chemokines. This synchronization has recently been shown to be critical in the control of TB disease along with CD4⁺ T-cell recruitment into the lung parenchyma¹³⁻¹⁵.

IP-10 has been suggested as an alternative biomarker to IFN- γ in the QFT-plus format, as on study reported that IP-10 in response to TB1 was increased in subjects with LTBI compared to those with active TB¹⁶. However, this study discovered no significant difference between LTBI groups and normal groups, including HD and HC groups. IP-10 would not be useful in the diagnosis of LTBI, especially in immunocompromised patients.

CCL11 is produced by IFN- γ -stimulated endothelial cells and TNF-activated monocytes. It has been reported that CCL11, CCL24, and CCL26, which are produced by Th2 cells and other cells that induce Th2 development, are increased in TB patients compared to controls. It appears that TB suppresses Th1 and subsequently the classic function of macrophages by inducing chemokine expression¹⁷. This study showed that CCL11 mRNA expression level was significant higher in the HD/LTBI group and > 0.7 IU/mL response group with no significance in terms of the receiver operating characteristic (ROC) analysis. Although CCL11 is little known in LTBI immune response, Yang et al.(2014) reported that the level of CCL11 continued to increase at the end of 12 months of follow-up after anti-TB treatment¹⁸. However, although this study lacked sufficient subjects to evaluate the effect of anti-TB treatment, CCL11 level was not related to anti-TB treatment in the present study. On the other hand, CCL11 was not significantly different among HC, LTBI, and ATB groups ($P = 0.4592$)¹⁹. This study suggests the need to investigate the kinetics of CCL11 response in LTBI.

In the present study, the mRNA expression levels of GM-CSF and TNFR were significantly higher in the 0.2 to 0.7 IU/mL response group than in the < 0.2 IU/mL response group and the > 0.7 IU/mL response group. However, there were no significant differences among the HD/LTBI, HD/normal, and HC groups. GM-CSF is involved in supporting the productive phase of inflammatory responses^{20,21}. T cell production of GM-CSF is required for control of TB. It has been reported that GM-CSF level was significantly different

between MTB-infected and uninfected populations and suggested GM-CSF could be the promising alternative biomarker to detect MTB¹⁹.

However, there were insufficient subjects in the 0.2 to 0.7 IU/mL response group and the existence of extreme values. These reasons could explain the different significance of mean and median values among the three QFT-GIT response groups in the HD and HC populations and the only HD population. Based on our results, mRNA expression levels of CCL11, GM-CSF, and TNFR require further investigation to distinguish LTBI and noninfected TB among the immunocompromised patients in the future.

Although the reversion rate observed in the present study was not as high as the 72% observed in the United States (US), the reversion rate within six months observed in the present study was 34% (16/43) regardless of the LTBI treatment²². In addition, the reversion rate of the 0.35 to 0.8 IU/mL response group was 74% (14/19), which was higher than that of the > 0.81 IU/mL response group (24%, 6/25) in the previous study²³. Similarly, in the present study, the reversion rate of the 0.35-0.7 IU/mL response group was 85.7% (6/7), which was significantly higher than the > 0.7 IU/mL response group (27.8%, 10/36) ($P = 0.0162$).

The reasons for IGRA variability are for the most part unexplained. Shu et al. (2013) suggested that the reversion occurs as an exponential decay, and the half-life of IGRA positivity was approximately two months and 18 months, respectively²⁴. This phenomenon may be due to the attenuated cellular immunity in dialysis patients. However, in the present study, there was no significant difference in terms of demographic and clinical factors, and mRNA expression levels of cytokines and chemokines between the reversion group and the persistent positive group. We could not match and analyze the differences of cytokines and chemokines using the previous samples of QFT-GIT test within the past six months^{25,26}.

In conclusion, the present study revealed that the mRNA expression of CCL11 was statistically different in LTBI patients compared to HD patients without LTBI and HCs based on the current cut-off value (≥ 0.35 IU/mL). Additionally, there was a significant difference of CCL11 mRNA expression level among the different QFT-GIT responses. Our study suggests that CCL11 may be a complementary or alternative biomarker for LTBI diagnosis, especially in immunocompromised patients.

The present study had certain limitations. The present study did not include an active TB group with HD and healthy donors. In addition, no peritoneal dialyzed patients with LTBI were included. The comparison of the mRNA expression level of cytokines and chemokines among the three different QFT-GIT test response groups was not efficient because of the insufficient number of subjects in the 0.2 to 0.7 IU/mL response group. Therefore, future large-scale studies and cohort designs are needed to evaluate the diagnostic performance of the cytokines and chemokines selected in the present study.

Declarations

Ethical approval

The present study was approved by the Institutional Review Board of the Kosin University Gospel Hospital (KUGH 2017-11-042) and the Catholic University of Pusan (CUPIRB-2017-036). Remaining blood samples of the HD group were used after Kosin University Gospel Hospital Biobank approval and whole blood samples of the HC group were used and collected from participants who provided their written informed consent. All experiments were performed in accordance with relevant guidelines and regulations.

Funding

This study was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (NRF-2016R1C1B108888 and 2020R1C1007169) and Brain Busan 21 Plus project.

Compliance with ethical standards

Conflict of interest the authors declare that they have no conflict of interest

Informed consent all subject provided informed consent for this study

References

1. ESRD Registry Committee, Korean Society of Nephrology. Current renal replacement therapy in Korea- Insan memorial dialysis registry 2018. Available from: <http://www.ksn.or.kr>. (assessed 21 May 2020)
2. Christopoulos AI, Diamantopoulos AA, Dimopoulos PA, Goumenos DS, Barbaliass GA. Risk factors for tuberculosis in dialysis patients: a prospective multi-center clinical trial. *BMC Nephrol* 2009;10:36. doi:10.1186/1471-2369-10-36.
3. Kang YA, Lee HW, Hwang SS, et al. Usefulness of whole-blood interferon-gamma assay and interferon-gamma enzyme-linked immunospot assay in the diagnosis of active pulmonary tuberculosis. *Chest* 2007;132(3):959-65. doi:10.1378/chest.06-2805.
4. Metcalfe JZ, Cattamanchi A, McCulloch CE, Lew JD, Ha NP, Graviss EA. Test variability of the QuantiFERON-TB gold in-tube assay in clinical practice. *Am J Respir Crit Care Med* 2013;187(2):206-11. doi:10.1164/rccm.201203-0430OC .
5. Southern J, Sridhar S, Tsou CY, Hopkins S, Collier S, Nikolayevskyy V, et al. Discordance in latent tuberculosis (TB) test results in patients with end-stage renal disease. *Public Health* 2019;166:34-39. doi:10.1016/j.puhe.2018.09.023.
6. Ringshausen FC, Nienhaus A, Torres Costa J, Knoop H, Schlösser S, Schultze-Werninghaus G, et al. Within-subject variability of Mycobacterium tuberculosis-specific gamma interferon responses in German health care workers. *Clin Vaccine Immunol* 2011;18:1176-82. doi: 10.1128/CI.05058-11.
7. Schablon A, Nienhaus A, Ringshausen FC, Preisser AM, Peters C. Occupational screening for tuberculosis and the use of a borderline zone for interpretation of the IGRA in German healthcare workers. *PLoS One* 2014;9(12):e115322. doi: 10.1371/journal.pone.0115322.

8. Nemes E, Rozot V, Geldenhuys H, Bilek N, Mabwe S, Abrahams D, et al. Optimization and Interpretation of Serial QuantiFERON Testing to Measure Acquisition of Mycobacterium tuberculosis Infection. *Am J Respir Crit Care Med* 2017;196(5):638-48. doi:10.1164/rccm.201704-0817OC.
9. Mori T, Sakatani M, Yamagishi F, Takashima T, Kawabe Y, Nagao K, et al. Specific detection of tuberculosis infection: an interferon-gamma-based assay using new antigens. *Am J Respir Crit Care Med* 2004;170:59–64. doi: 10.1164/rccm.200402-1790C.
10. Mannhalter C, Koizar D, Mitterbauer G. Evaluation of RNA isolation methods and reference genes for RT-PCR analyses of rare target RNA. *Clin Chem Lab Med* 2000;38(2):171-7. doi:10.1515/CCLM.2000.026.
11. Kubista M, Andrade JM, Bengtsson M, Forootan A, Jonak J, Lind K, et al. The real-time polymerase chain reaction. *Mol Aspects Med* 2006;27:95-125. doi: 10.1016/j.mam.2005.12.007.
12. Latorre I, Mínguez S, Carrascosa JM, Naves J, Hernandez RV, Muriel B, et al. Immune-mediated inflammatory diseases differently affect IGRAs' accuracy for latent tuberculosis infection diagnosis in clinical practice. *PLoS One* 2017;12(12);e0189202. doi:10.1371/journal.pone.0189202.
13. Sakai S, Kauffman KD, Schenkel JM, McBerry CC, Mayer-Barber KD, Masopust D, et al. Cutting edge: control of Mycobacterium tuberculosis infection by a subset of lung parenchyma-homing CD4 T cells. *J Immunol* 2014;192(7):2965-9. doi: 10.4049/jimmunol.1400019.
14. Moguche AO, Shafiani S, Clemons C, Larson RP, Dinh C, Higdon LE, et al. ICOS and Bcl6-dependent pathways maintain a CD4 T cell population with memory-like properties during tuberculosis. *J Exp Med* 2015;212: 715–28. doi: 10.1084/jem.20141518.
15. Torrado E, Fountain JJ, Liao M, Tighe M, Reiley WW, Lai RP, et al. Interleukin 27R regulates CD4+ T cell phenotype and impacts protective immunity during Mycobacterium tuberculosis infection. *J Exp Med* 2015;212:1449–63. doi:10.1084/jem.20141520.
16. Petrone L, Vanini V, Chiacchio T, Petruccioli E, Cuzzi G, Schininà V, et al. Evaluation of IP-10 in Quantiferon-Plus as biomarker for the diagnosis of latent tuberculosis infection. *Tuberculosis* 2018;111: 147-53. doi: 10.1016/j.tube.2018.06.005.
17. Olsson AK, Dimberg A, Kreuger J, Claesson-Welsh L. VEGF receptor signalling - in control of vascular function. *Nat Rev Mol Cell Biol* 2006;7(5):359-71. doi:10.1038/nrm1911.
18. Yang Q, Cai Y, Zhao W, Wu F, Zhang M, Luo K, et al. IP-10 and MIG are compartmentalized at the site of disease during pleural and meningeal tuberculosis and are decreased after antituberculosis treatment. *Clin Vaccine Immunol* 2014;21:1635-44. doi: 10.1128/CI.00499-14.
19. Won EJ, Choi JH, Cho YN, Jin HM, Kee HJ, Park YW, et al. Biomarkers for discrimination between latent tuberculosis infection and active tuberculosis disease. *J Infect* 2017;74:281-93. doi: 10.1016/j.jinf.2016.11.010.
20. Cheknev SB, Apresova MA, Moryakova NA, Efremova IE, Mezdrokhina AS, Piskovskaya LS, et al. Production of the growth factors GM-CSF, G-CSF, and VEGF by human peripheral blood cells induced with metal complexes of human serum γ -globulin formed with copper or zinc ions. *Mediators Inflamm* 2014;2014:518265. doi: 10.1155/2014/518265.

21. Gray J, Reves R, Johnson S, Belknap R. Identification of false-positive QuantiFERON-TB Gold In-Tube assays by repeat testing in HIV-infected patients at low risk for tuberculosis. Clin Infect Dis 2012; 54:e20-3. doi:10.1093/cid/cir792.

22. Zwerling A, van den Hof S, Scholten J, Cobelens F, Menzies D, Pai M. Interferon-gamma release assays for tuberculosis screening of healthcare workers: a systematic review. Thorax 2012;67(1):62-70. doi:10.1136/thx.2010.143180.

23. Schablon A, Nienhaus A, Ringshausen FC, et al. Occupational screening for tuberculosis and the use of a borderline zone for interpretation of the IGRA in German healthcare workers. PLoS One. 2014;9(12):e115322.

24. Shu CC, Wu VC, Yang FJ, et al. Dynamic changes in positive interferon-gamma release assay in a dialysis population: An observational cohort study. J Infect 2013;67(6):529-535.

25. Pai M, Elwood K. Interferon-gamma release assays for screening of health care workers in low tuberculosis incidence settings: dynamic patterns and interpretational challenges . Can Respir J. 2012;19:81–3. doi:10.1155/2012/420392.

26. Griveas I, Visvardis G, Fleva A, Papadopoulou D, Mitsopoulos E, Kyriklidou P, et al. Comparative analysis of immunophenotypic abnormalities in cellular immunity of uremic patients undergoing either hemodialysis or continuous ambulatory peritoneal dialysis. Ren Fail 2005;27:279–82.

Tables

Table 1. Demographics and clinical characteristics and QuantiFERON-TB Gold In-Tube (QFT-GIT) response results.

	Hemodialyzed patients (<i>n</i> = 75)		Healthy control ^c (<i>n</i> = 48)	<i>P</i> -value
	LTBI ^a (<i>n</i> = 28)	Normal ^b (<i>n</i> = 47)		
Sex (Male)	18 (64.3%)	16 (34%)	21 (43.8%)	0.038*
Age	62.00 ± 11.18	60.70 ± 12.46	23.94 ± 2.72	<0.001***
< 65 years old	16 (57.1%)	30 (63.8%)	21 (100.0%)	0.796
65-74 years old	7 (25%)	11 (23.4%)	0 (0.0%)	
≥ 75 years old	5 (17.9%)	6 (12.8%)	0 (0.0%)	
Previous TB contact	7 (25%)	6 (12.8%)	0 (0%)	0.002**
Previous TB treatment	0	0	0	
BCG vaccination or scar	19 (67.9%)	34 (72.3%)	48 (100.0%)	<0.001***
Abnormal Chest X-ray lesion	3 (10.7%)	0 (0.0%)	0 (0.0%)	0.005**
Underlying diseases				
Diabetes mellitus	17 (60.7%)	20 (42.6%)	0 (0.0%)	0.100
Ischemic heart disease	9 (32.1%)	6 (12.8%)	0 (0.0%)	0.043*
BMI (kg/m ²)				0.026*
< 18.5	9 (32.1%)	4 (8.5%)	-	
18.5-22.9	9 (32.1%)	26 (55.3%)	-	
23.0-24.9	6 (21.4%)	5 (10.6%)	-	
25.0-29.9	4 (14.3%)	9 (19.1%)	-	
30.0-34.9	0 (0%)	3 (6.4%)	-	
Low BMI (< 18.5 kg/m ²)	9 (32.1%)	4 (8.5%)	-	0.012*
Smoking	1 (3.6%)	5 (10.6%)	-	0.040*

QFT-GIT results					
Positive		28 (100.0%)	0 (0.0%)	0 (0.0%)	
Indeterminate		0 (0.0%)	0 (0.0%)	0 (0.0%)	
Negative		0 (0.0%)	47 (100.0%)	48 (100.0%)	
Previous QFT-GIT results within 6 months					
Positive		27 (96.4%)	16 (34%)	-	<0.001***
Indeterminate		0 (0.0%)	0 (0.0%)	-	
Negative		1 (3.6%)	31 (66%)	-	
Reversion			16 (37.2%)	-	
Conversion		1 (2.9%)			
LTBI treatment history		6 (22.2%)	6 (37.5%)	-	0.232
QFT-GIT responses groups (IU/mL)					
Negative	<0.2	0 (0.0%)	46 (97.9%)	46 (95.8%)	<0.001***
Borderline	0.2 – 0.34	0 (0.0%)	1 (2.1%)	2 (4.2%)	
	0.35 – 0.7	7 (25%)	0 (0.0%)	0 (0.0%)	
Positive	> .0.7	21 (75.0%)	0 (0.0%)	0 (0.0%)	
Previous QFT-GIT responses group within 6 months (IU/mL)					
Negative	< 0.2	1 (3.6%)	24 (51.1%)	-	<0.001***
Borderline	0.2 – 0.34	0 (0.0%)	7 (14.9%)	-	
	0.35 – 0.7	1 (3.6%)	6 (12.8%)	-	
Positive	> 0.7	26 (92.9%)	10 (21.3%)	-	
Plasma IFN- γ level after T cell mitogen stimulation (Mitogen-Nil, IU/ml)		9.40 $\pm$ 1.68	9.24 $\pm$ 1.34	10 $\pm$ 0	0.005** (a = b < c)

Note: Data were expressed as mean $\pm$ SD for continuous variables and as percentages for categorical variables. *P*-values were calculated using one-way ANOVA tests with Dunnett's multiple comparisons or

Chi-square analysis. * $P < 0.05$, ** 0.01 , *** <0.001 . *Abbreviation* : QFT-GIT, QuantiFERON-TB Gold In-Tube;TB, tuberculosis;BMI, body mass index;LTBI, latent tuberculosis infection;IFN, interferon.

Table 2. Comparison of QuantiFERON-TB Gold In-Tube (QFT-GIT) assay results between the current study and previous QFT-GIT within the past 6 months

Previous QFT-GIT responses within 6 months (IU/mL)	QFT-GIT responses (IU/ml)				Total
	< 0.2	0.20 – 0.34	0.35 – 0.70	> 0.7	
< 0.2	24 (96.0%)	0 (0.0%)	0 (0.0%)	1 (4.0%)	25 (100.0%)
0.20 – 0.34	7 (100.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)	7 (100.0%)
0.35 – 0.7	6 (85.7%)	0 (0.0%)	1 (14.3%)	0 (0.0%)	7 (100.0%)
> 0.7	9 (25.0%)	1 (2.8%)	6 (16.7%)	20 (55.5%)	36 (100.0%)
Total	46	1	7	21	75
Cohen κ coefficient = 0.386, $P < 0.001$					

Note: QFT-GIT, QuantiFERON-TB Gold In-Tube assay

Figures

Figure 1

Target gene mRNA expression levels after stimulation of Mycobacterium tuberculosis (MTB)-specific antigens for 24 h according to latent tuberculosis infection (LTBI) diagnosis. Note : Whole blood of study subjects [hemodialyzed patients (HD) without LTBI (HD/normal) groups, HD with LTBI (HD/LTBI) group, and healthy control (HC) group] were stimulated with MTB-specific antigens (early secretory antigen target 6, culture filtrate protein 10, and TB 7.7) for 24 h and the mRNA expression levels for target genes

were analyzed using real-time RT-PCR. The graphs represented median value (■, HD/normal;●, HD/LTBI;▲, HC) with interquartile range (bar). (a) interferon (IFN)- γ , (b) tumor necrosis factor- α (TNF- α), (c) IL-2R, (d) CXCL9, (e) IFN- γ -induced protein 10 (IP-10), (f) CXCL11, (g) CCL11, (h) granulocyte-macrophage-colony stimulating factor (GM-CSF), and (i) TNF receptor (TNFR). *P < 0.05.

Figure 2

Target gene mRNA expression levels after stimulation of *Mycobacterium tuberculosis* (MTB)–specific antigens for 24 hours according to QuantiFERON-TB Gold In-Tube (QFT-GIT) response groups. Note : Whole blood of study subjects [< 0.2 IU/ml QFT-GIT response group, $0.2–0.7$ IU/ml QFT-GIT response group, and > 0.7 IU/ml QFT-IT response group] were stimulated with MTB-specific antigens (early secretory antigen target 6, culture filtrate protein 10, and TB 7.7) for 24 hours and the mRNA expression levels for target genes were analyzed by real-time RT-PCR. The graphs represented median value (■, HD/normal; ●, HD/LTBI; ▲, HC) with interquartile range (bar). (a) interferon (IFN)- γ , (b) tumor necrosis factor- α (TNF- α), (c) IL-2R, (d) CXCL9, (e) IFN- γ -induced protein 10 (IP-10), (f) CXCL11, (g) CCL11, (h) granulocyte-macrophage-colony stimulating factor (GM-CSF), and (i) TNF receptor (TNFR). $**P < 0.01$.

Figure 3

Target gene mRNA expression levels after stimulation of Mycobacterium tuberculosis (MTB)-specific antigens for 24 hours according to QuantiFERON-TB Gold In-Tube (QFT-GIT) response groups in hemodialyzed patients (HD). Note: Whole blood of study subjects [the < 0.2 IU/ml QFT-GIT response group, 0.2–0.7 IU/ml QFT-GIT response group, and > 0.7 IU/ml QFT-GIT response group] in HD. were stimulated with MTB-specific antigens (early secretory antigen target 6, culture filtrate protein 10, and TB

7.7) for 24 hours, and the mRNA expression levels for target genes were analyzed using real-time RT-PCR. The graphs represented median value (■, HD/normal;●, HD/LTBI;▲, HC) with interquartile range (bar). (a) interferon (IFN)- γ , (b) tumor necrosis factor- α (TNF- α), (c) IL-2R, (d) CXCL9, (e) IFN- γ -induced protein 10 (IP-10), (f) CXCL11, (g) CCL11, (h) granulocyte-macrophage-colony stimulating factor (GM-CSF), and (i) TNF receptor (TNFR). *P < 0.05.