

# Assessment of heavy metals and potential health risks associated with the consumption of vegetables grown on the roadside soils

**Fazal Mabood**

University of Malakand

**Fazal Hadi**

University of Malakand

**Amin Ullah Jan**

Shaheed Benazir Bhutto University Sheringal

**Allah Ditta** (✉ [allah.ditta@sbbu.edu.pk](mailto:allah.ditta@sbbu.edu.pk))

Shaheed Benazir Bhutto University <https://orcid.org/0000-0003-1745-4757>

**Ziaul Islam**

Shaheed Benazir Bhutto University Sheringal

**Manzer H. Siddiqui**

King Saud University

**Hayssam M. Ali**

King Saud University

**Ayman EL Sabagh**

Kafr el-Sheikh University: Kafrelsheikh University

---

## Research Article

**Keywords:** Rumex dentatus, Trachyspermum ammi, Spinacia oleracea, Allium cepa, lead, nickel

**Posted Date:** November 29th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-1115842/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

Vegetables cultivated near roads absorb toxic metals from polluted soil, which enter into the human body through the food chain and cause serious health problems to humans. The present study investigated the contamination level of lead (Pb) and nickel (Ni) in soils and vegetables grown along the roadside of District Swat, Pakistan, and the health risks associated with the consumption of the tested vegetables have been investigated. The plant samples were collected from the cultivated field within a 120-meter range from the roadside. *Spinacia oleracea*, *Allium cepa*, *Rumex dentatus*, and *Trachyspermum ammi* were selected based on their importance as vegetables. In results, Pb concentration was higher in plants located at the distance between 0-10 m away from the roadside than the WHO permissible limit. In such plants, Pb concentration was higher than Ni. *Rumex dentatus* contained the highest concentration of Pb among the tested vegetables while Ni concentration was highest in *Trachyspermum ammi* as compared to other plants. Concentration and accumulation of both the metals decreased in soil and plants with increasing distance from the road. Similarly, target hazard quotient values noted for Pb and Ni were greater than unity, which shows that there is a potential risk associated with the consumption of tested vegetables near the road. Moreover, the values of target cancer risk were greater than 0.0001, which shows that there is a risk of cancer with the consumption of tested vegetables. In conclusion, the consumption of the tested vegetables was very dangerous as it may lead to higher risks of cancer.

## Introduction

Fossil fuel combustion in automobiles releases toxic heavy metals into the air, which are then deposited on the soils nearby roadsides (Suzuki et al. 2008; Bai et al. 2009). The concentration and toxicity vary in various plant species (Mehmood et al. 2021a; Murtaza et al. 2021a). High concentrations of toxic heavy metals in roadside soils have been reported in many regions across the globe (Irshad et al. 2021; Mehmood et al. 2021b; Murtaza et al. 2021b; Naveed et al. 2021a). It has been disclosed that vehicle exhaust is the main cause of the contamination of nearby roadside soils (Bai et al. 2011; Li et al. 2020). Nickel (Ni), lead (Pb), zinc (Zn), and cadmium (Cd) are toxic heavy metals that are frequently found in the soil near roadsides (Akoto et al. 2008; Mehmood et al. 2018).

Lead (Pb) is a common pollutant present in the air, water, and soil due to the widespread use of this metal in industrial processes (Ullah et al. 2020; Kanwal et al. 2021). Combustion of petroleum products, smelting, and sewage wastes are the main sources of Pb pollution (Grover et al. 2010). It has been reported that Australia, the USA, and China produced 400, 516, and 1690 thousand metric tons of Pb in the year 2009, respectively (United State geological survey, 2009). Lead deposits on the surface of bio-particles, organic and clay particles (Sammur et al. 2010). Several biochemical factors control its adsorption in the soil such as level of inorganic and organic ligands (Shahid et al. 2011), cation exchange capacity (Vega et al. 2010), microbial and biological conditions (Arias et al. 2010), redox state (Tabelin and Igarashi, 2009), and competing cations level (Komjarova and Blust, 2009). Plants may uptake Pb through several pathways. High negative potential on the surface of plant roots is required for Pb

absorption. Due to similarity with calcium (Ca), Pb enters into the plant mainly through channels specified for Ca transport (Wang et al. 2007).

No biological role of Pb has been reported until now and thus it is a non-essential element for living organisms. On contrary, it has been considered as the second most hazardous metal after arsenic (Maestri et al 2010; Mehmood et al. 2018, 2019). Besides plants, Pb also adversely affects human health. A decrease in the number of erythrocytes has been reported during Pb exposure because of the inhibition of enzymes (coproporphyrinogen and ferrochelatase, delta-aminolevulinic acid ALAD) needed for the synthesis of hemoglobin and red blood cell (Sipos et al. 2003; Warren et al. 2003). It has been found that Pb exposure increased the excretion of amino acids and glucose in urine due to its toxic effect on the proximal convoluted portion of the nephron (Loghman-Adham, 1997). Long-term Pb exposure causes kidney failure, increased blood uric acid concentration, high blood pressure, and joint infection (Alasia, 2010). The highest amounts of Pb (about 1/3rd of the entire body) were accumulated in the liver (Mudipalli, 2009). As Pb is analogous to Ca, therefore it can cross the blood-brain barrier easily where it can replace and activate calcium-mediated activities, which disturb brain physiology and development (Sanders et al. 2009). People working in the lead storage industry, show disorders of the male gonads (testes), reduced testosterone production, and low sperm count in semen (Queiroz and Waissmann 2006). Threshold limit of Pb that can cause reproductive abnormalities is about  $40 \mu\text{g dL}^{-1}$  (Quintanilla-Vega et al. 2000). Lead affects the menstrual cycle, reduces the duration of pregnancy, and causes abnormal birth (Han et al. 2000).

Ni is the 22nd most abundant element found in the earth's crust, occurring mostly in rocks as a free metal or bonded with iron metal. Ni is a hard, ductile, and silvery-white metal (McIlveen and Negusanti, 1994). Concentrations of Ni in soil and drinking waters are lower than 100 and 0.005 ppm, respectively (Naveed et al. 2020, 2021b). Anthropogenic sources of Ni pollution include vehicular emissions, fossil fuel burning, mining, smelting, municipal and industrial wastes (Alloway, 1995). Ni concentrations may reach up to 26000 ppm in contaminated soils (Alloway, 1995).

Ni is absorbed by plants mainly through the root system via both passive and active transport (Seregin, 2006). The ratio of Ni uptake between active and passive transport changes with plant species, Ni form, and Ni concentration in the soil (Dan et al. 2002; Naveed et al. 2020). Moreover, soluble compounds of Ni might also be absorbed through the Mg ion transport system, due to the similar charge/size ratio of the two metal ions (Oller et al. 1997). However, its concentration in most of the plant species is extremely low i.e. 0.05-10 ppm (Nieminen et al. 2007). Extremely high concentrations of Ni have made some farmland soils unsuitable for growing crops, vegetables, and fruits (Naveed et al. 2021).

In humans, Ni is an abundant metal commonly responsible for skin allergies and is one of the greatest causes of allergic contact dermatitis, as revealed by positive dermal patch tests (Cavani, 2005). Ni is a carcinogenic element in several animal species but the basic mechanisms behind are still unknown (Chang, 1996). Ni can act as a tumor inducer by inhibiting natural killer cell activity (Costa and Klein,

1999). Another possible way by which Ni induces cell death and/or damage is through lipid peroxidation (Misra et al. 1990; Chen et al. 1998; Janicka and Cempel, 2001).

Based on the above discussion, it has been found that both Pb and Ni cause toxic effects on plants as well as humans. The increasing needs of the human population for food have resulted in intensive farming, even near the roadsides. In addition, there is an increasing trend regarding the production of vegetables along the roadside near urban areas. Therefore, it is necessary to evaluate the contamination level of heavy metals especially Pb and Ni in soils as well as plants growing along the roadsides, as it would help in exploring the contamination level of these metals in those soils and plants and to calculate health risks associated with the consumption of these edible plants growing along the roadside. Moreover, it would also help in identifying the safe distance for the production of different crop plants along the roadside. Up to our knowledge, no study has focused on this aspect of the present research work conducted. Based on these hypotheses, the present study was conducted to investigate the contamination level of Pb and Ni in soils and plants that grew along the roadside of Bara Bandai, Ningolai, and Ghalegay, District Swat, Khyber Pakhtunkhwa. Moreover, different parameters related to the health risks associated with the consumption of the tested vegetables have been investigated.

## Materials And Methods

### Study area

The present study was conducted to investigate the contamination level of Pb and Ni in soils and plants that grew along the roadside of Bara Bandai, Ningolai and Ghalegay, District Swat, Khyber Pakhtunkhwa. Plants and soil samples were collected from the cultivated fields near the roadside at Bara Bandai, Ningolai, and Ghalegay.

### Collection of soil and plant samples

Plant samples were collected from the cultivated field within a 120-meter range from the roadside. Collection of plants was made in four groups based on their distance range from road i.e. group-I (0-10 m distance), group-II (10-40 m distance), group-III (40-80 m distance), and group-IV (80-120 m distance). Five replicates were taken for each plant at each range. Similarly, five soils samples were randomly collected from each range. Plants (*Spinacia oleracea*, *Allium cepa*, *Rumex dentatus*, and *Trachyspermum ammi*) were selected for the present investigation based on their importance as vegetables.

### Acid digestion of soil and plant samples

Collected soil samples were ground and sieved (2 mm) to remove large particles. Parameters such as electrical conductivity and pH were calculated using a pH meter (Model CON.3173) and electrical conductivity meter (Model CON 5). For acid digestion, 0.25 g from each soil sample was added in acid solution (5 ml nitric acid and 1 ml sulfuric acid) in digestion tubes and placed overnight in a fume hood. The next day, each soil sample in acid solution was heated until a clear aliquot was obtained. The aliquot

was filtered and the total volume was raised to 50 mL with distilled water. Each solution was stored in plastic bottles until analyzed for heavy metals.

Collected plant samples were cut into parts (roots, stem, and leaves) and then kept in paper envelopes. The samples were dried in the oven at 80 °C for 48 h and then crushed into a powdered form using a commercial blender. Each sample was digested by the method of Allen (1974). A 0.25gm of the sample was taken in the conical flask and then a mixture of acids (5 ml nitric acid and 1 ml sulfuric acid) was added to it and boiled on a hotplate until digested completely. After digestion, each sample was cooled, filtered and then the final volume is raised to 50 mL by adding distilled water. The solutions were stored in plastic bottles for metal analysis. The metal analysis of both soil and plants samples was carried out at Central Resource Laboratory, Peshawar using atomic absorption spectrophotometer (AA2407, USA).

## Health risk assessment

### Bioconcentration factor (BCF)

It is the ratio of heavy metal concentration in the edible part of the plant to heavy metal concentration in a soil sample (Sharma et al. 2018). BCF was calculated using the following formula:

$$BCF = \frac{C_{plant}}{C_{soil}}$$

where  $C_{plant}$  is heavy metal content in the edible part of plant and  $C_{soil}$  is heavy metal content in respective soil. The value of BCF greater than 1 indicates that the plant is a potential accumulator of the metal being considered for analysis.

### Estimated daily intake (EDI)

The estimated daily intake of the metals was determined based on their mean concentration in each plant sample and the estimated daily consumption of the vegetables in grams. The EDI value of each metal of interest was determined by the formula used by Chen et al. (2011) with slight modification as presented in the following equation:

$$EDI = \frac{E_f \times E_d \times F_{IR} \times C_m \times C_f}{B_w \times T_A} \times 0.001$$

where  $E_f$  is exposure frequency (365 day/year),  $E_d$  is the exposure duration (65 years), equivalent to an average lifetime (Woldetsadik et al. 2017),  $F_{IR}$  is the average food (vegetable) consumption (240 g/person/day) which were obtained from the World Health Report (WHO, 2002) for low vegetable intake,  $C_M$  is the metal concentration (mg/kg dry weight),  $C_f$  is the concentration conversion factor for fresh vegetable weight to dry weight i.e. 0.085 (Harmanescu et al. 2011),  $B_w$  is reference body weight for an

adult, which is 70 kg (Woldetsadik et al. 2017),  $T_A$  is the average exposure time (65yrs x 365 days) and 0.001 is the unit conversion factor.

### Target hazard quotient (THQ)

The target hazard quotient (THQ) values were estimated to assess non-carcinogenic human health risk from the consumption of vegetables contaminated by heavy metals. The THQ values were calculated using the following equation as described by Chen et al., (2011).

$$THQ = \frac{EDI}{RfD}$$

Where EDI is the estimated daily metal intake of the population in mg/day/kg body weight and RfD is the oral reference dose (mg/kg/day) values which were 0.0035 for Pb and 0.02 for Ni. If the value of THQ is < 1, it is generally presumed to be safe for the risk of non-carcinogenic effects and if it is > 1, it is supposed that there is a chance of non-carcinogenic effects with an increasing probability as the value upsurges (Chen et al. 2011, Antoine et al. 2017).

### Hazard Index (HI)

It has been documented that the individual health risks of the analyzed heavy metals in the same vegetable are accumulative and that is expressed as hazard index (Chen et al. 2011, Antoine et al. 2017). Accordingly, the HI of target metals considered in this study were calculated using the following equation proposed by Antoine et al. (2017):

$$HI = \sum_{n=1}^i THQ_n, i = 1, 2, 3, \dots, n$$

where HI is the sum of various metals hazards. If the HI value became < 1.0, there is no apparent health impact due to the metals considered. However, an HI value of > 1.0 indicates potential health impact implications. A serious chronic health impact has been suggested for HI > 10.0 (Antoine et al. 2017).

### Target cancer risk (TCR)

The cancer risk posed to human health due to the ingestion of individual possibly carcinogenic metals was estimated using the following equation as described by Sharma et al. (2018). Then, the target cancer risk (TCR) resulting from heavy metals (Pb and Ni) ingestion, which may promote carcinogenic effect depending on the exposure dose, was calculated using the following equation as described by Kamunda et al. (2016).

$$CR = EDI \times CPS_o$$

$$TCR = \sum_{n=1}^i CR, i = 1, 2, 3, \dots, n$$

where CR represents cancer risk over a lifetime by individual heavy metal ingestion, EDI is the estimated daily metal intake of the population in mg/day/kg body weight,  $CPS_o$  is the oral cancer slope factor in (mg/kg/day)<sup>-1</sup> and n is the number of heavy metals considered for cancer risk calculation. The  $CPS_o$  values used for Pb and Ni were 0.0085 and 1.7, respectively. It has been pointed out that the slope factor converts the estimated daily intake of the metal averaged over a lifetime of the exposure directly to the incremental risk of an individual developing cancer (Kamunda et al. 2016).

### Statistical analysis

The mean value of the data was calculated and then it was subjected to analysis of variance (ANOVA) and correlations between different parameters were established using statistical software SPSS 16 and MS Excel 2010.

## Results

### Physicochemical properties of soil

Different physicochemical properties of soils collected at different distances from the roadside are presented in Table 1. Almost all the properties varied with distance from the roadside. A decreasing trend in soil pH while an increasing trend in electrical conductivity (EC) was noted with increasing distance from the roadside. Similarly, the concentrations of lead (Pb) and nickel (Ni) in the soil decreased as the distance from the road increased. Concentrations of Pb and Ni were significantly higher at 0-10 meters from the road compared to the other distances. The highest concentrations of Pb and Ni recorded at 0-10 m from the road were 60.6 and 35.0. The concentration of Pb in the soil was under the normal limit (85 mg kg<sup>-1</sup>) but that of the Ni was at the edge of the permissible limit (35 mg kg<sup>-1</sup>) by WHO, 2001. The soil was loamy in texture, which is suitable for agricultural activities purposes because it has constituents more or less the same amount required for plants growth.

### Lead and nickel concentrations in *Rumex dentatus* L.

As clear from Figure 1, the highest accumulation of both Pb and Ni was in the root portion of the plants followed by stem and leaves. The Pb concentration in all the parts of *Rumex dentatus* L. was above the permissible limits (0.3 mg kg<sup>-1</sup>) set by WHO/FAO, 2001 up to 80 m distance from the road. At 80-120 m distance, the Pb concentration was above the permissible limits only in the root portion of *Rumex dentatus* L. In the case of Ni, the concentration was above the permissible limits in root, leaves, and stem portions of *Rumex dentatus* L. up to 10 m distance from the road. At 10-40 m away from the road, only

the root portion accumulated Ni above the permissible limit. For the rest of the plants after 40-120 m, the concentration of Ni was below the permissible limits as given by WHO, FAO, 2001.

#### **Lead and nickel concentrations in *Trachyspermum ammi* (L.) Sprague ex Turrill**

Nickel (Ni) and lead (Pb) concentration in different parts of *Trachyspermum ammi* (L.) Sprague ex Turrill collected at different distances from the roadside is presented in Figure 2. The red line along the horizontal axis represents the permissible limits given by World Health Organization (WHO) and Food and Agriculture Organization (FAO), 2001 for Pb ( $0.3 \text{ mg kg}^{-1}$ ) and Ni ( $1.5 \text{ mg kg}^{-1}$ ) concentration in plants to be used as food. From the data presented, as the distance from the road was increased, the bioaccumulation of Pb and Ni was decreased and vice versa. The highest accumulation of both Pb and Ni was in the root portion of the plants followed by leaves, stems, and fruit. The Pb concentration in all the parts of *Trachyspermum ammi* (L.) Sprague ex Turrill was above the permissible limits ( $0.3 \text{ mg kg}^{-1}$ ) by WHO/FAO, 2001 up to 40 m distance from the road. At 40-80 m distance, the Pb concentration was above the permissible limits in the root, leaves, and stem portion of the plants while 80-120 m distance, the concentration was above the permissible limits only in the root portion of *Trachyspermum ammi* (L.) Sprague ex Turrill. In the case of Ni, the concentration was above the permissible limits in the root, leaves, and stem portions of *Trachyspermum ammi* (L.) Sprague ex Turrill up to 40 m distance from the road. For the rest of the plants after 40-120 m, the concentration of Ni was below the permissible limits as given by WHO, FAO, 2001.

#### **Lead and nickel concentrations in *Spinacia oleracea* L.**

From the data presented in Figure 3, the bioaccumulation of Pb and Ni in *Spinacia oleracea* L. was decreased as the distance from the road was increased and vice versa. The highest accumulation of both Pb and Ni was in the root portion followed by leaves, and stems. The Pb concentration in all the parts of *Spinacia oleracea* L. was above the permissible limits ( $0.3 \text{ mg kg}^{-1}$ ) by WHO/FAO, 2001 up to 80 m distance from the road. At 80-120 m distance, the Pb concentration was above the permissible limits only in the root of *Spinacia oleracea* L. In the case of Ni, a similar decreasing trend in Ni concentration with increasing distance was recorded in different parts of *Spinacia oleracea* L. The concentration was above the permissible limits in root, leaves, and stem portions of *Spinacia oleracea* L. up to 10 m distance from the road. At 10-40 distance, the Ni concentration was above the permissible limit in only the root portion of the *Spinacia oleracea* L. For the rest of the plants after 40-120 m, the concentration of Ni was below the permissible limits as given by WHO, FAO, 2001.

#### **Lead and nickel concentrations in *Allium cepa* L.**

A decreasing trend with increasing distance from the roadside regarding the concentrations of Ni and Pb in different parts of *Allium cepa* L. was recorded (Figure 4). As clear from the data, the root portion accumulated the maximum amount of Pb, followed by stem and leaves. In the case of Ni, the maximum accumulation was observed in the case of the stem, followed by stem and leaves. Regarding the permissible limit of Pb concentration, only stem and leaves portion at 80-120 away from the roadside



were below the permissible concentration of Pb while all the others were above the permissible limit of Pb i.e.  $0.3 \text{ mg kg}^{-1}$ . In the case of Ni, the concentration was above the permissible limit ( $1.5 \text{ mg kg}^{-1}$ ) in root, stem, and leaves at 0-10 m away from the road while only root and stem portions had Ni concentrations above the permissible limits.

### Lead and nickel bioaccumulation

As clear from the data presented in Table 2, the maximum bioaccumulation of Pb was recorded in *Rumex dentatus* L., followed by *Trachyspermum ammi* (L.) Sprague ex Turrill, *Spinacia oleracea* L., and *Allium cepa* L. The shoot and root portion of *Rumex dentatus* L. bioaccumulated  $46.2$  and  $29.4 \text{ mg kg}^{-1}$  DW.

### Correlation between Ni and Pb concentration in soil and plant samples

A positive strong correlation ( $R^2 = 0.984$ ) was found between Ni and Pb concentration in the soil as given in Figure 5. Similarly, a positive correlation existed between the concentration of Ni and Pb in all the studied plants as shown in Figure 6. The correlation was found to be highly significant in *Trachyspermum ammi* (L.) Sprague ex Turrill ( $R^2 = 0.969$ ), *Spinacia oleracea* ( $R^2 = 0.988$ ), *Allium cepa* ( $R^2 = 0.959$ ) and *Rumex dentatus* ( $R^2 = 0.906$ ).

### Bioconcentration factor

As clear from the data presented in Table 2, the bioconcentration factor (BCF) decreased with increasing distance from the road in all the species studied. The maximum value of BCF ( $0.7629$ ) was observed in *Rumex dentatus* for the Pb, followed by  $0.4949$  in *Trachyspermum ammi* for Ni. The values of BCFs in *Spinacia oleracea*, *Allium cepa*, and *Rumex dentatus* were more for Pb as compared to that observed for Ni in the same species. The minimum BCF value ( $0.0312$ ) was observed in the case of *Trachyspermum ammi* at 80-120 m away from the road for Pb. A decreasing trend in BCF values was observed in all the studied species as the distance from the road was increased.

### Health risk assessment

After calculation of BCF and TF, the data on heavy metals concentration in different plant species studied was analyzed regarding health risks associated. It was found that the values of estimated daily intake (EDI), target hazard quotient (THQ), hazard index (HI), cancer risk (CR), and target cancer risk (TCR) were decreased as the distance of sampling site from the road was increased. In general, the values of EDI and THQ were higher in the case of Pb as compared to those observed for Ni. The maximum EDI ( $0.0135$ ) and THQ ( $3.3682$ ) were recorded in the case of *Rumex dentatus* for Pb at 0-10 m away from the road. Similarly, the values of EDI and THQ in the case of *Spinacia oleracea* L., *Trachyspermum ammi* (L.) Sprague ex Turrill and *Allium cepa* L. were  $0.0086$  and  $2.1507$ ,  $0.0057$  and  $1.4236$ , and  $0.0052$  and  $1.2932$ , respectively. In the case of hazard index, the maximum value ( $3.4445$ ) was observed in the case of *Rumex dentatus*, followed by *Spinacia oleracea* L. ( $2.2466$ ), *Trachyspermum ammi* (L.) Sprague ex Turrill ( $1.6760$ ), and *Allium cepa* L. ( $1.3836$ ). Moreover, the values of cancer risk (CR) regarding Ni were

higher than that of Pb and these values showed a decreasing trend with increasing distance from the road. The value of CR for Ni was maximum (0.8413) in the case of *Trachyspermum ammi* (L.) Sprague ex Turrill at 0-10 m away from the road. A similar trend was observed in the case of target cancer risk (TCR). The maximum value of TCR (0.8413) in the case of *Trachyspermum ammi* (L.) Sprague ex Turrill at 0-10 m away from the road, followed by *Trachyspermum ammi* (L.) Sprague ex Turrill (0.5989) at 10-40 m away from the road, *Spinacia oleracea* L. (0.3197) at 0-10 m away from the road, *Allium cepa* L. (0.3154) at 10-40 m away from the road.

## Discussion

Soil not only provides anchorage and support to the plants but also acts as a reservoir of different nutrients. The physicochemical characteristics of soil are very important and have a crucial role in the mobilization and distribution of toxic heavy metals. A strong correlation existed between concentrations of Ni and Pb in the soil under study, which clearly shows that their sources might be the same on roadsides i.e. traffic emission (Patel and Jain, 2021). Our study showed that Pb and Ni concentration in soil was lower than the threshold limit (0.3 ppm for Pb and 1.50 ppm for Ni) set by World Health Organization. The concentration of Pb and Ni in soils at the road edges was significantly high ( $Pb = 60.60 \pm 5.60$  and  $Ni = 35.00 \pm 3.24$ ) and decreased rapidly with increasing distance from the road which is in line with earlier studies (Huan et al. 2018; Li et al. 2020). Earlier, it has been found that Pb concentration on the surface of the earth naturally occurs below  $50 \text{ mg kg}^{-1}$  (Arias et al 2010). Its concentration in soil mainly changes due to anthropogenic activities. As Pb occurs in the petroleum products and combustion of this fuel in internal combustion engine of cars releases Pb into the atmosphere, which then settles down on the soil near roadsides (Sammut et al 2010). Similar is the case with nickel, which also occurs naturally in petroleum products. Nickel in soils can be found in several different forms: adsorbed or complex with organic cation surfaces or on inorganic cation exchange surfaces, inorganic crystalline minerals, water-soluble, chelated metal complexes, or free ion in soil solution. Lead does not seem to be of big concern outside urban areas but may ultimately become a problem due to decreased pH of soil caused by limited use of soil liming in agriculture and mobilization because of increased acid rain (Bai et al. 2011).

Plants grown in contaminated soils are exposed to the contaminants. Since plants have the natural ability to absorb dissolved substances from the soil solution and in doing so, plants also absorb toxic metals present in the soil. Comparison among lead concentrations in different plant species demonstrated significant variations with distances from roads. Pb concentration in plants occurred in the order of *Rumex dentatus* > *Spinacia oleracea* > *Trachyspermum ammi* > *Allium cepa*. Differences in Pb concentration in plants depend on several factors such as developmental stage, genetic makeup, transpiration coefficient, plant roots system in soil, plant growth rate, and nutrients needed by plants (Cenkci et al. 2010). Roots of plants absorbed the highest Pb concentration as compared to the other plant parts. This is because roots are more exposed to the pollutants in the soil as compared to the other parts of the plant. Many plants have been reported to retain up to 95% of Pb in their root portions and a

little concentration is transported to aerial parts of a plant (Shahid et al., 2011). This restricted transport of Pb to the aerial parts of the plants might be due to the immobilization of Pb on plant cell walls by the negatively charged pectin (Arias et al. 2010). Despite this, some of the Pb moves through intercellular spaces present between plant cells and translocation with water to endodermis (Wang et al. 2007; Shahid et al. 2011). Pb is either stored in a plasma membrane attached with pectin, crystallized in intercellular spaces, or stored in outer root cortical cells (Yang et al. 2007). Plants collected near the roadside were more exposed to lead as compared to the plants away from roadsides. Similar results have been reported by many investigators at different locations of the world (Lone et al., 2006; Bai et al. 2011; Li et al. 2020). Nickel is an essential nutrient for plants required in very low concentrations. Nickel compounds are mainly emitted by the combustion of fossil fuel in automobiles, which results in the contamination of roadside areas with Ni particles.

The most serious problem in toxic metal pollution in agricultural soil is the absorption and accumulation of toxic metals in plants tissues. The food chain is the main route of entry of heavy metals into the human body, which accounts for about 90% of heavy metals source in the human body. Therefore, health risk assessment in the form of estimated daily intake (EDI), target hazard quotient (THQ), and hazard index (HI), cancer risk (CR), and target cancer risk (TCR) was done in the present study. In the present study, EDI values for Pb and Ni were below maximum tolerable daily intake i.e. 0.21 and 0.1-0.3 as reported by Chen et al. (2011) and Woldetsadik et al. (2017), respectively. In the case of THQ, values noted for Pb and Ni were greater than unity, which shows that there is potential risk associated with the consumption of all the tested vegetables up to 0-10 m away from the road. Moreover, the consumption of *Rumex dentatus* was not safe up to 40 m away from the road. All the other species were safe to use after 10 m away from the road except *Rumex dentatus*, which was safe to utilize after 40 m away from the road. A similar trend in HI was observed regarding the usage of different vegetables tested in the present study as observed in the case of THQ. In the case of TCR, all the values were greater than  $1 \times 10^{-4}$ , which shows that there is a risk of cancer with the consumption of all the tested vegetables. Based on these indices, it was found that the consumption of the tested vegetables was very dangerous and may lead to higher risks of cancer.

## Conclusions

The results confirmed the presence of Pb and Ni in selected soils and plant samples. The concentration of Pb in *Rumex dentatus*, *Trachyspermum ammi*, *Spinacia oleracea*, and *Allium cepa* was above the safe limits of WHO within 80 m range from the road. The results demonstrated that plants collected beyond the 80 m range contained Pb concentration below the WHO safe limit. While the plant was found safe in the case of Ni beyond the 10 m range where the concentration of Ni in plant tissues was found below the WHO safe limit. Moreover, the consumption of studied vegetables grown along the roadside was not safe based on the values of different indices. In the case of target hazard quotient (THQ), the values noted for Pb and Ni were greater than unity, which shows that there is potential risk associated with the consumption of all the tested vegetables up to 0-10 m away from the road. In the case of target cancer

risk (TCR), all the values were greater than  $1 \times 10^{-4}$ , which shows that there is a risk of cancer from Ni with the consumption of vegetables to the people in the study area. Based on these results, it is highly recommended to have strict regulatory control on the cultivation of these vegetables along the roadside in the study area.

## Declarations

### Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

### Conflict of interest

The authors declare that they have no competing interests.

### Ethics approval

Not applicable. This manuscript does not involve researching about humans or animals.

### Consent to participate

All of the authors consented to participate in the drafting of this manuscript.

### Consent to publish

All of the authors consented to publish this manuscript.

### Authors' contribution

All authors contributed to the study's conception and design. Fazal Mabood, Fazal Hadi, Amin Ullah Jan, Allah Ditta: Conceptualization, Methodology, Software. Fazal Mabood, Amin Ullah Jan, Allah Ditta, Ziaul Islam.: Data curation, Writing- Original draft preparation. Manzer H. Siddiqui: Visualization, Investigation. Fazal Hadi, Amin Ullah Jan, Allah Ditta: Supervision. Hayssam M. Ali, Ayman EL Sabagh: Software, Validation. Fazal Hadi, Amin Ullah Jan, Allah Ditta: Writing- Reviewing and Editing. All authors read and approved the final manuscript.

### Availability of data and materials

All the data and tools/models used for this work are publicly available.

## References

1. Akoto O, Ephraim JH, Darko G (2008) Heavy metal pollution in surface soils in the vicinity of abundant railway servicing workshop in Kumasi Ghana. *Int J Environ Res* 2(4):359–364.

2. Alasia DD (2010) Lead nephropathy: Revisiting an overlooked cause of kidney disease. *Nephrol Rev* 2:e8. <https://doi.org/10.4081/nr.2010.e8>
3. Allen SE, Grimshaw HM, Parkinson JA, Quarmby C (1974) Chemical analysis of ecological materials. Oxford: Blackwell Scientific.
4. Alloway BJ (1995) Heavy Metal in Soils (Ed: B. J. Alloway), 2<sup>nd</sup> ed., Blackie Academic and Professional, London, pp. 25 – 34.
5. Antoine JMR, Fung LAH, Grant CN (2017) Assessment of the potential health risks associated with the aluminum, arsenic, cadmium, and lead content in selected fruits and vegetables grown in Jamaica. *Toxicol Reports* 4:181–187. <https://doi.org/10.1016/j.toxrep.2017.03.006>
6. Arias JA, Peralta-Videa JR, Ellzey JT, Ren M, Viveros MN, Gardea-Torresdey JL (2010) Effects of *Glomus deserticola* inoculation on Prosopis: enhancing chromium and lead uptake and translocation as confirmed by X-ray mapping, ICP-OES, and TEM techniques. *Environ Exp Bot* 68(2):139–148. <https://doi.org/10.1016/j.envexpbot.2009.08.009>
7. Bai J, Cui B, Wang Q, Gao H, Ding Q (2009) Assessment of heavy metal contamination of roadside soils in Southwest China. *Stoch Environ Res Risk Assess* 23(3):341-347. <https://doi.org/10.1007/s00477-008-0219-5>
8. Bai J, Cui B, Wang Q, Gao H, Ding Q (2009) Assessment of heavy metal contamination of roadside soils in Southwest China. *Stoch Environ Res Risk Assess* 23:341–347. <https://doi.org/10.1007/s00477-008-0219-5>
9. Bai J, Wang Q, Zhang K, Cui B, Liu X, Huang L, Xiao R, Gao H (2011) Trace element contaminations of roadside soils from two cultivated wetlands after abandonment in a typical plateau lakeshore, China. *Stoch Environ Res Risk Assess* 25:91–97. <https://doi.org/10.1007/s00477-010-0406-z>
10. Cenkci S, Cigerci IH, Yildiz M, Özay C, Bozdog A, Terzi H (2010) Lead contamination reduces chlorophyll biosynthesis and genomic template stability in *Brassica rapa* L. *Environ Exp Bot* 67(3):467–473. <https://doi.org/10.1016/j.envexpbot.2009.10.001>
11. Chang LW (1996) Toxicology of Metals, Lewis Publishers: New York, pp 245-246.
12. Chen C, Qian Y, Chen Q, Li C (2011) Assessment of daily intake of toxic elements due to consumption of vegetables, fruits, meat, and seafood by inhabitants of Xiamen, China. *J Food Sci* 76:T181–T188. <https://doi.org/10.1111/j.1750-3841.2011.02341.x>
13. Chen C. H.Y., Huang Y.L., Lin T.H. (1998). Lipid peroxidation in liver of mice administrated with nickel chloride: with special reference to trace elements and antioxidants. *Biol Trace Elem Res* 61(2):193-199. <https://doi.org/10.1007/BF02784030>
14. Costa M, Klein CB (1999) Nickel carcinogenesis, mutation, epigenetics, or selection. *Environ Health Perspect* 107(9):438-439. <https://doi.org/10.1289/ehp.99107a438>
15. Dan TV, Krishnaraj S, Saxena PK (2002) Cadmium and nickel uptake and accumulation in scented Geranium (*Pelargonium* sp.' Frensham'). *Water Air Soil Pollut* 137:355 – 364. <https://doi.org/10.1023/A:1015590007901>

16. Grover P, Rekhadevi P, Danadevi K, Vuyyuri S, Mahboob M, Rahman M (2010) Genotoxicity evaluation in workers occupationally exposed to lead. *Int J Hyg Environ Health* 213(2):99–106.  
<https://doi.org/10.1016/j.ijheh.2010.01.005>
17. Han S, Pfizenmaier DH, Garcia E, Eguez ML, Ling M, Kemp FW, Bogden JD (2000) Effects of lead exposure before pregnancy and dietary calcium during pregnancy on fetal development and lead accumulation. *Environ Health Perspect* 108:527-531. <https://doi.org/10.1289/ehp.00108527>
18. Harmanescu M, Alda L, Bordean D, Gogoasa I, Gergen I (2011) Heavy metals health risk assessment for population via consumption of vegetables grown in old mining area, a case study: Banat County, Romania. *Chem Cent J* 5:64. <https://doi.org/10.1186/1752-153X-5-64>
19. Irshad S, Xie Z, Mehmood S, Nawaz A, Ditta A, Mahmood Q (2021) Insights into conventional and recent technologies for arsenic bioremediation: A systematic review. *Environ Sci Pollut Res* 28(15):18870–18892. <https://doi.org/10.1007/s11356-021-12487-8>
20. Janicka K., Cempel M. (2001). Lipid peroxidation and selected antioxidants in rat liver after oral exposure to nickel(II) chloride. *Bromatol Chem Toksykol* 34:291-295.
21. Kamunda C, Mathuthu M, Madhuku M (2016) Health risk assessment of heavy metals in soils from Witwatersrand gold mining basin, South Africa. *Int J Environ Res Public Health* 13:663.  
<https://doi.org/10.3390/ijerph13070663>
22. Kanwal U, Ibrahim M, Abbas F, Yamin M, Jabeen F, Shahzadi A, Farooque AA, Imtiaz M, Ditta A, Ali S (2021) Phytoextraction of Lead using a Hedge Plant [*Alternanthera bettzickiana* (Regel) G. Nicholson]: Physiological and Bio-chemical Alterations through Bioresource Management. *Sustainability* 13(9):5074. <https://doi.org/10.3390/su13095074>
23. Komjarova I., Blust R. (2009). Effect of Na, Ca and pH on simultaneous uptake of Cd, Cu, Ni, Pb, and Zn in the water flea *Daphnia magna* measured using stable isotopes. *Aquat Toxicol* 94(2):81–86.  
<https://doi.org/10.1016/j.aquatox.2009.05.018>
24. Li Y, Yuan Y, Sun C, Sun T, Liu X, Li J, Fang L, Fan Z (2020) Heavy metals in the soil of an urban industrial zone in a metropolis: risk assessment and source apportionment. *Stoch Environ Res Risk Assess* 34(2):435-446. <https://doi.org/10.1007/s00477-020-01779-z>
25. Loghman-Adham M (1997) Renal Effects of environmental and Occupational Lead Exposure. *Environ Health Perspect* 105(9):928-938. <https://doi.org/10.1289/ehp.97105928>
26. Lone MI, Raza SH, Muhammad S, Naeem MA, Khalid M (2006) Lead content in soil and wheat tissue along roads with different traffic loads in Rawalpindi district. *Pak J Bot* 38(4):1035-1042.
27. Maestri E, Marmiroli M, Visioli G, Marmiroli N (2010) Metal tolerance and hyperaccumulation: costs and trade-offs between traits and environment. *Environ Exp Bot* 68(1):1–13.  
<https://doi.org/10.1016/j.envexpbot.2009.10.011>
28. McIlveen W. D., Negusanti J. J. (1994). Nickel in the terrestrial environment. *Sci Tot Environ* 148:109-138. [https://doi.org/10.1016/0048-9697\(94\)90390-5](https://doi.org/10.1016/0048-9697(94)90390-5)
29. Mehmood S, Ahmed W, Rizwan M, Imtiaz M, Elnahal ASMA, Ditta A, Irshad S, Ikram M, Li W (2021a) Comparative efficacy of raw and HNO<sub>3</sub>-modified biochar derived from rice straw on vanadium

- transformation and its uptake by rice (*Oryza sativa* L.): Insights from photosynthesis, antioxidative response, and gene-expression profile. *Environ Pollut* 289:117916.  
<https://doi.org/10.1016/j.envpol.2021.117916>
30. Mehmood S, Imtiaz M, Bashir S, Rizwan M, Irshad S, Yuvaraja G, Ikram M, Aziz O, Ditta A, Rehman SU, Shakeel Q, Mumtaz MA, Ahmed W, Mahmood S, Chen D, Tu S (2019) Leaching behavior of Pb and Cd and transformation of their speciation in co-contaminated soil receiving different passivators. *Environ Eng Sci* 36(6):749-759. <https://doi.org/10.1089/ees.2018.0503>
  31. Mehmood S, Saeed DA, Rizwan M, Khan MN, Aziz O, Bashir S, Ibrahim M, Ditta A, Akmal M, Mumtaz MA, Ahmed W, Irshad S, Imtiaz M, Tu S, Shaheen A (2018) Impact of different amendments on biochemical responses of sesame (*Sesamum Indicum* L.) plants grown in lead-cadmium contaminated soil. *Plant Physiol Biochem* 132:345-355.  
<https://doi.org/10.1016/j.plaphy.2018.09.019>
  32. Mehmood S, Wang X, Ahmed W, Imtiaz M, Ditta A, Rizwan M, Irshad S, Bashir S, Saeed Q, Mustafa A, Li W (2021b) Removal mechanisms of slag against potentially toxic elements in soil and plants for sustainable agriculture development: A critical review. *Sustainability* 13(9):5255.  
<https://doi.org/10.3390/su13095255>
  33. Misra M, Rodrigez RE, Kasp-Rzak KS (1990) Nickel induced lipid peroxidation in the rat: correlation with nickel effect on antioxidant defense systems. *Toxicology* 64(1):1-17.  
[https://doi.org/10.1016/0300-483X\(90\)90095-X](https://doi.org/10.1016/0300-483X(90)90095-X)
  34. Mudipalli A (2009) Lead hepatotoxicity and potential health effects. Review Article. National Center for Environmental Assessment-RTP Division, Office of Research and Development, U.S. EPA, North Carolina
  35. Murtaza G, Ditta A, Ahmed Z, Usman M, Faheem M, Tariq A (2021a) Co-biosorption potential of *Acacia nilotica* bark in removing Ni and amino azo benzene from contaminated wastewater. *Desalina Water Treat* 233:261-271. <https://doi.org/10.5004/dwt.2021.27514>
  36. Murtaza G, Ditta A, Ullah N, Usman M, Ahmed Z (2021a) Biochar for the management of nutrient impoverished and metal contaminated soils: Preparation, applications, and prospects. *J Soil Sci Plant Nutrit* 21:2191–2213. <https://doi.org/10.1007/s42729-021-00514-z>
  37. Naveed M, Bukhari SS, Mustafa A, Ditta A, Alamri S, El-Esawi MA, Rafique M, Ashraf S, Siddiqui MH (2020) Mitigation of nickel toxicity and growth promotion in sesame through the application of a bacterial endophyte and zeolite in nickel contaminated soil. *Int J Environ Res Public Health* 17(23):8859. <https://doi.org/10.3390/ijerph17238859>
  38. Naveed M, Ditta A, Ahmad M, Mustafa A, Ahmad Z, Conde-Cid M, Tahir S, Shah SAA, Abrar MM, Fahad S (2021b) Processed animal manure improves morpho-physiological and biochemical characteristics of *Brassica napus* L. under nickel and salinity stress. *Environ Sci Pollut Res* 28:45629–45645 <https://doi.org/10.1007/s11356-021-14004-3>
  39. Naveed M, Tanvir B, Xiukang W, Brtnicky M, Ditta A, Kucerik J, Subhani Z, Nazir MZ, Radziemska M, Saeed Q, Mustafa A (2021a) Co-composted biochar enhances growth, physiological and

- phytostabilization efficiency of *Brassica napus* and reduces associated health risks under Cr stress. *Front Plant Sci* <https://doi.org/10.3389/fpls.2021.775785>
40. Nieminen TM, Ukonmaanaho L, Rausch N, Shotyk W (2007) Biogeochemistry of nickel and its release into the environment. In: Sigel A, Sigel H, Sigel RKO (eds) *Metal ions in life sciences*, vol 2. Wiley, Chichester, pp 1–30. <https://doi.org/10.1002/9780470028131.ch1>
  41. Oller AR, Costa M, Oberd G (1997) Carcinogenicity assessment of selected nickel compounds. *Toxicol Appl Pharmacol* 143(1):152-166. <https://doi.org/10.1006/taap.1996.8075>
  42. Patel DK, Jain MK. Contamination and health risk assessment of potentially harmful elements associated with roadside dust in Dhanbad India. *Stoch Environ Res Risk Assess* <https://doi.org/10.1007/s00477-021-02061-6>
  43. Queiroz EK, Waissmann W (2006) Occupational exposure and effects on the male reproductive system. *Cad Saude Publica* 22(3):485-493. <https://doi.org/10.1590/S0102-311X2006000300003>
  44. Quintanilla-Vega B, Hoover DJ, Bal W, Silbergeld EK, Waalkes MP, Anderson LD (2000) Lead interaction with human protamine (HP2) as a mechanism of male reproductive toxicity. *Chem Res Toxicol* 13(7):594-600. <https://doi.org/10.1021/tx000017v>
  45. Sammut M, Noack Y, Rose J, Hazemann J, Proux O, DepouxZiebel M, Fiani E (2010) Speciation of Cd and Pb in dust emitted from sinter plant. *Chemosphere* 78(4):445–450. <https://doi.org/10.1016/j.chemosphere.2009.10.039>
  46. Sanders T, Liu Y, Buchner V, Tchounwou PB (2009) Neurotoxic effects and biomarkers of lead exposure: A Review. *Rev Environ Health* 24(1):15-45. <https://doi.org/10.1515/REVEH.2009.24.1.15>
  47. Seregin IV, Kozhevnikova AD (2006) Physiological role of nickel and its toxic effects on higher plants. *Russ J Plant Physiol* 53:257–277. <https://doi.org/10.1134/S1021443706020178>
  48. Shahid M., Pinelli E., Pourrut B. Silvestre J., Dumat C. (2011). Lead-induced genotoxicity to *Vicia faba* L. roots in relation with metal cell uptake and initial speciation. *Ecotoxicol Environ Saf* 74(1):78–84. <https://doi.org/10.1016/j.ecoenv.2010.08.037>
  49. Sharma S, Nagpal AK, Kaur I (2018) Heavy metal contamination in soil, food crops and associated health risks for residents of Ropar wetland, Punjab, India and its environs. *Food Chem* 255:15–22. <https://doi.org/10.1016/j.foodchem.2018.02.037>
  50. Sipos P, Szentmihályi K, Fehér E, Abaza M, Szilyági M, Blázovics A (2003) Some effects of lead contamination on liver and gall bladder bile. *Acta Biol Szeged* 47(1-4):139-142.
  51. Suzuki K, Yabuki T, Ono Y (2008) Roadside *Rhododendron pulchrum* leaves as bioindicators of heavy metal pollution in traffic areas of Okayama Japan. *Environ Monit Assess* 149:133-141. <https://doi.org/10.1007/s10661-008-0188-7>
  52. Tabelin C, Igarashi T (2009) Mechanisms of arsenic and lead release from hydrothermally altered rock. *J Hazard Mat* 169(1–3):980–990. <https://doi.org/10.1016/j.jhazmat.2009.04.049>
  53. Ullah I, Ditta A, Imtiaz M, Mehmood S, Rizwan M, Rizwan MS, Jan AU, Ahmad I (2020) Assessment of health and ecological risks of heavy metal contamination: A case study of agricultural soils in Thall, Dir-Kohistan. *Environ Monit Assess* 192:786. <https://doi.org/10.1007/s10661-020-08722-3>



54. US Geological Survey (2009) Retrieved on 22 Dec 2012 from <http://minerals.usgs.gov/minerals/pubs/commodity/lead>
55. Vega F, Andrade M, Covelo E (2010) Influence of soil properties on the sorption and retention of cadmium, copper and lead, separately and together, by 20 soil horizons: comparison of linear regression and tree regression analyses. *J Hazard Mat* 174(1–3):522–533. <https://doi.org/10.1016/j.jhazmat.2009.09.083>
56. Wang H, Shan X, Wen B, Owens G, Fang J, Zhang S (2007) Effect of indole-3-acetic acid on lead accumulation in maize (*Zea mays* L.) seedlings and the relevant antioxidant response. *Environ Exp Bot* 61(3):246–253. <https://doi.org/10.1016/j.envexpbot.2007.06.004>
57. Warren MJ, Cooper JB, Wood SP, Shoolingin-Jordan PM (2003) X-ray structure of a putative reaction intermediate of 5-aminolaevulinic acid dehydratase. *Biochem J* 373(3):733–738. <https://doi.org/10.1042/bj20030513>
58. WHO (2002) The world health report 2002: reducing risks, promoting healthy life, World Health Organization, 2002. <https://apps.who.int/iris/handle/10665/42510>
59. Woldetsadik D, Drechsel P, Keraita B, Itanna F, Gebrekidan H (2017) Heavy metal accumulation and health risk assessment in wastewater-irrigated urban vegetable farming sites of Addis Ababa. Ethiopia. *Int J Food Contam* 4:9. <https://doi.org/10.1186/s40550-017-0053-y>
60. Yang X, Li T, Liu D, Jin X, Meng F (2007) Effect of Pb toxicity on root morphology, physiology and ultrastructure in the two ecotypes of *Elsholtzia argyi*. *J Hazard Mat* 147(3), 806–816. <https://doi.org/10.1016/j.jhazmat.2007.01.117>

## Tables

**Table 1:** Physical and chemical properties of soil collected at different distances from roadside.

Distance from road side (m)	pH	EC (dS m <sup>-1</sup> )	Texture	Pb (ppm)	Ni (ppm)
0 – 10	7.01	5.91	Loamy soil	60.60 ± 5.60 <sup>a*</sup>	35.00 ± 3.24 <sup>a</sup>
10 – 40	6.70	5.99	Loamy soil	20.60 ± 4.56 <sup>b</sup>	16.01 ± 2.56 <sup>b</sup>
40 – 80	6.46	6.03	Loamy soil	13.00 ± 3.24 <sup>b</sup>	10.33 ± 1.25 <sup>b</sup>
80 – 120	6.39	6.04	Loamy soil	5.13 ± 2.92 <sup>b</sup>	8.67 ± 1.87 <sup>b</sup>

\*Significant variations between values is represented by different letters.


**Table 2:** Bioaccumulation of Pb and Ni in root and shoot, bioconcentration factor (BCF), and translocation factor (TF) calculated for Pb and Ni concentration in different plant species sampled at different distances from the road

Distance from road	Bioaccumulation of Pb (mg kg <sup>-1</sup> DW)			Bioaccumulation of Ni (mg kg <sup>-1</sup> DW)			Bioconcentration factor (BCF)	
	Root	Shoot	Total	Root	Shoot	Total	Pb	Ni
<b><i>Rumex dentatus</i> L.</b>								
0-10 m	29.4	46.23	75.63	2.83	5.24	8.07	0.7629	0.1497
10-40 m	10.13	14.58	24.71	1.50	2.61	4.11	0.7078	0.1630
40-80 m	0.93	1.00	1.93	0.90	1.42	2.32	0.0769	0.1375
80-120 m	0.30	0.19	0.49	0.72	1.00	1.72	0.0370	0.1153
<b><i>Trachyspermum ammi</i> (L.) Sprague ex Turrill</b>								
0-10 m	24.90	19.54	44.44	10.25	17.32	27.57	0.3224	0.4949
10-40 m	12.12	6.54	18.66	3.17	5.64	8.81	0.3175	0.3523
40-80 m	2.20	1.04	3.24	0.90	1.67	2.57	0.0800	0.1617
80-120 m	0.45	0.16	0.61	0.56	1.11	1.67	0.0312	0.1280
<b><i>Spinacia oleracea</i> L.</b>								
0-10 m	18.00	29.52	47.52	4.24	6.58	10.82	0.4871	0.1880
10-40 m	4.96	5.76	10.72	1.67	2.67	4.34	0.2796	0.1668
40-80 m	1.04	1.05	2.09	1.00	1.58	2.58	0.0808	0.1530
80-120 m	0.62	0.45	1.07	0.64	1.00	1.64	0.0877	0.1153
<b><i>Allium cepa</i> L.</b>								
0-10 m	12.54	17.75	30.29	2.90	6.20	9.1	0.2929	0.1771
10-40 m	6.39	8.66	15.05	1.53	2.97	4.5	0.4204	0.1855
40-80 m	0.79	0.92	1.71	0.72	1.28	2.0	0.0708	0.1239
80-120 m	0.30	0.38	0.68	0.50	0.85	1.35	0.0741	0.0980

**Table 3:** Estimated daily intake (EDI), target hazard quotient (THQ), and hazard index (HI), cancer risk (CR) and target cancer risk (TCR) calculated for Pb and Ni concentration in different plant species sampled at different distances from the road


Distance from road	Estimated daily intake (EDI)		Target Hazard Quotient (THQ)		Hazard index (HI)	Cancer risk (CR)		Target cancer risk (TCR)
	Pb	Ni	Pb	Ni		Pb	Ni	
<i>Rumex dentatus</i> L.								
0-10 m	0.0135	1.53E-03	3.3682	0.0764	3.4445	1.14518E-04	0.2545	0.2546
10-40 m	0.0042	7.61E-04	1.0623	0.0380	1.1003	3.61167E-05	0.2771	0.2772
40-80 m	0.0003	4.14E-04	0.0729	0.0207	0.0935	2.47714E-06	0.2337	0.2337
80-120 m	0.0001	2.91E-04	0.0138	0.0146	0.0284	4.70657E-07	0.1961	0.1961
<i>Trachyspermum ammi</i> (L.) Sprague ex Turrill								
0-10 m	0.0057	5.05E-03	1.4236	0.2524	1.6760	4.84034E-05	0.8413	0.8413
10-40 m	0.0019	1.64E-03	0.4765	0.0822	0.5587	1.62005E-05	0.5989	0.5989
40-80 m	0.0003	4.87E-04	0.0758	0.0243	0.1001	2.57623E-06	0.2748	0.2748
80-120 m	0.0000	3.23E-04	0.0117	0.0162	0.0278	3.96343E-07	0.2176	0.2176
<i>Spinacia oleracea</i> L.								
0-10 m	0.0086	1.92E-03	2.1507	0.0959	2.2466	7.31253E-05	0.3196	0.3197
10-40 m	0.0017	7.78E-04	0.4197	0.0389	0.4586	1.42683E-05	0.2835	0.2835
40-80 m	0.0003	4.60E-04	0.0765	0.0230	0.0995	0.000002601	0.2600	0.2600
80-120 m	0.0001	2.91E-04	0.0328	0.0146	0.0474	1.11471E-06	0.1961	0.1961
<i>Allium cepa</i> L.								
0-10 m	0.0052	1.81E-03	1.2932	0.0903	1.3836	4.39693E-05	0.3011	0.3012
10-40 m	0.0025	8.66E-04	0.6309	0.0433	0.6742	2.14521E-05	0.3154	0.3154
40-80 m	0.0003	3.73E-04	0.0670	0.0187	0.0857	2.27897E-06	0.2106	0.2107

Figures



**Figure 1**

Lead (Pb) and Nickel (Ni) concentration in different part of *Rumex dentatus* L. at different distances from road. Significant variations between the mean values (n = 3) is represented by different letters. The red line along horizontal axis represents the permissible limits given by World Health Organization and Food and Agriculture Organization, 2001 for Pb (0.3 mg kg<sup>-1</sup>) and Ni (1.5 mg kg<sup>-1</sup>) concentration in plants to be used as food



**Figure 2**

Lead (Pb) and Nickel (Ni) concentration in different part of *Trachyspermum ammi* (L.) Sprague ex Turill at different distances from road. Significant variations between the mean values (n = 3) is represented by different letters. The red line along horizontal axis represents the permissible limits given by World Health Organization and Food and Agriculture Organization, 2001 for Pb (0.3 mg kg<sup>-1</sup>) and Ni (1.5 mg kg<sup>-1</sup>) concentration in plants to be used as food




Figure 3

Lead (Pb) and Nickel (Ni) concentration in different part of *Spinacia oleracea* L. at different distances from road. Significant variations between the mean values (n = 3) is represented by different letters. The red line along horizontal axis represents the permissible limits given by World Health Organization and Food and Agriculture Organization, 2001 for Pb (0.3 mg kg<sup>-1</sup>) and Ni (1.5 mg kg<sup>-1</sup>) concentration in plants to be used as food




Figure 4

Lead (Pb) and Nickel (Ni) concentration in different part of *Allium cepa* L. at different distances from road. Significant variations between the mean values (n = 3) is represented by different letters. The red

line along horizontal axis represents the permissible limits given by World Health Organization and Food and Agriculture Organization, 2001 for Pb (0.3 mg kg<sup>-1</sup>) and Ni (1.5 mg kg<sup>-1</sup>) concentration in plants to be used as food




Figure 5

Correlation between Pb and Ni in soil.




Figure 6

Correlation between Pb and Ni accumulation in *Trachyspermum ammi* (A), *Spinacea oleracea* (B), *Allium cepa* (C) and *Rumex dentatus* (D).