

Pregnancy-Associated Listeriosis in a Tertiary Hospital in Ningbo, Zhejiang, China: A Retrospective Study

Yefang Ke

Ningbo Women and Children's Hospital

Lina Ye

Ningbo Women and Children's Hospital

Pan Zhu

Ningbo Women and Children's Hospital

Ying Sun

Ningbo Women and Children's Hospital

Zhe Zhu (✉ zhuzhemcgrady@163.com)

University of Chinese Academy of Sciences

Research Article

Keywords: pregnancy-associated listeriosis, *Listeria monocytogenes*, maternal listeriosis, neonatal listeriosis, outcome.

Posted Date: December 21st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1116269/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background

Pregnancy-associated listeriosis is a severe infectious disease and potentially leading to fetal/neonatal fatal. Limited information on pregnancy-associated listeriosis is available in China.

Methods

We performed a retrospective study on maternal and neonatal patients with pregnancy-associated listeriosis. The clinical characteristics of pregnancy-associated listeriosis were studied, and the outcome determinants of neonatal listeriosis were explored.

Results

14 cases of pregnancy-associated listeriosis were identified. The incidence of pregnancy-associated listeriosis in our hospital was 16.69/100,000 births. All of the 14 maternal patients eventually recovered after delivery shortly with no sequelae. None of the 12 mothers who delivered in this hospital received antepartum first-line empirical treatment. Among the 14 neonatal cases, 1 was late onset listeriosis and 13 were early onset cases; 11 survived and 3 died. Fatality rates were significantly higher in outborn neonates ($P=0.005$). Besides, higher mortality rates were observed in neonates with lower birth weight ($P=0.038$), gestational age < 28 weeks ($P=0.056$), and Apgar score (5th min) < 5 ($P=0.056$), with marginally significant differences.

Conclusions

There is a neglected burden of pregnancy-associated listeriosis in our hospital. Timely and optimal resuscitative measures are considered to be important to reduce mortality of neonatal listeriosis. Listeriosis should be considered in the differential diagnosis of pregnancy-associated infections.

Background

Listeriosis is a rare but severe disease caused by *Listeria monocytogenes* (*L. monocytogenes*), a ubiquitous food-borne bacterial pathogen[1, 2]. People principally get listeriosis by the ingestion of contaminated food[3, 4]. Especially, it has a predilection to infect pregnant women, which is at approximately 10-100 times greater risk for infection than the general population[5-7]. In Europe and North America, it has been observed previously that the proportion of pregnancy-associated listeriosis out of the total listeria infections ranges from 9% to 20%[2, 8-10]. Moreover, recent studies performed in China showed that more than half of listeria cases were pregnancy-associated infections[11, 12].

The high incidence of listeriosis during pregnancy may due to the increased progesterone levels which would affect immune functions [13, 14]. *L. monocytogenes* can actively cross the intestinal barrier, disseminate within the circulation, cross the placental barrier owing to its specific placental tropism, and eventually leading to placental and fetal infections[15]. Although maternal illness is usually mild, neonatal illness is frequently severe and potentially fatal. Pregnancy-associated listeriosis can lead to miscarriage, preterm delivery or stillbirth. It was reported that more than 80% of infected mothers would experience major fetal or neonatal complications[9]. Therefore, pregnancy-associated listeriosis is recognized as one of the worst maternal infections.

It has been reported that the incidence of pregnancy-associated listeriosis varied between different minorities[4, 9, 16]. Recently, a study in Singapore found that the incidence of congenital listeriosis was highest in the infants of Chinese origin[17]. Besides, a study observed that the rate of pregnancy-associated listeriosis in Beijing Obstetrics and Gynecology Hospital was 13.7 per 100,000 births[18], which was higher than that in United States, United Kingdom and New Zealand (3.4, 3.4 and 12.3 per 100,000 births, respectively)[19-21]. Listeriosis has not yet been included in the notifiable disease in China, the surveillance of listeriosis has not been thoroughly implemented. Moreover, owing to the nonspecific obstetrical signs and long incubation period, the diagnosis of pregnancy-associated listeriosis would be challenging[22, 23].

To better understand the maternal and neonatal listeriosis in Ningbo, Zhejiang Province, China and to provide more information for formulating appropriate therapeutic and controlling strategies, we performed a retrospective study to analyze the clinical characteristics and outcomes of pregnancy-associated listeriosis cases.

Methods

Study design and patient selection

This was a retrospective study conducted at Ningbo women and children's hospital, a tertiary maternity and pediatric hospital, where the number of births is about 10,000 every year. This study was approved by the the institutional ethics board (EC2021-028), and the informed consent was waived for this retrospective study.

We conducted a retrospective study to identify all perinatal cases of culture-confirmed *L. monocytogenes* infections at Ningbo women and children's hospital, from August 2013 to September 2021. Data on past medical history, characteristics at admission, underlying medical conditions, laboratory results and treatments were collected from medical record database.

Definitions

A pregnancy-associated listeriosis case was defined based on isolation of *L. monocytogenes* from a normally sterile site of the pregnant women and/or newborns aged ≤ 28 days. Each Mother and her newborn were counted as a single case. Neonatal listeriosis cases were classified as early onset (diagnosed between birth and days 6) listeriosis or late onset (diagnosed between days 7 and 28) listeriosis. We defined “inevitable miscarriage” as fetal loss before 24 weeks of gestation, “stillbirth” as death of the fetus between 24 and 41 weeks of gestation, “maternal leukocytosis” as a white blood cell (WBC) count $> 12 \times 10^9/L$, “neonatal leukocytosis” as a WBC count $> 34 \times 10^9/L$, “neonatal leukopenia” as a WBC count $< 5 \times 10^9/L$, “neonatal thrombocytopenia” as a platelet count $< 150 \times 10^9/L$, “neonatal severe thrombocytopenia” as a platelet count $< 5 \times 10^9/L$, “inborn neonate” as a neonate born in our hospital, and “outborn neonate” as a neonate born in other hospitals and transferred to our hospital after birth.

Microbiological methods

For blood culturing, whole blood for bacterial culture was collected and incubated in an automatic blood culture system (BACT/ALERT 3D) for 5 days or until rated positive. A positive blood culture was incubated on blood agar at $36^\circ C$ for 18-24 h. CSF, cervical secretion, eye secretion, gastric aspirate of newborn and products of conception samples were cultured directly on blood agar at $36^\circ C$ for 18-48 h, then, bacteria with a positive growth were isolated and cultured onto blood agar at $36^\circ C$ for 18-24 h.

The identification of bacteria was performed by using the VITEK 2 COMPACT automatic analysis system (BioMérieux, France). Standard strains (ATCC700323, ATCC25922, ATCC700327, ATCC29213) were used to control the microbiological quality.

Statistical analysis

Clinical, laboratory and meteorology data were recorded in a Microsoft Excel database (Microsoft, Richmond, US). For continuous variables, non-normally distributed variables were expressed as medians and ranges, for categorical variables, data were presented as percentages. The differences between two groups were determined by using the Wilcoxon Manne-Whitney U test for quantitative data and Fisher’s exact test for qualitative data, respectively. A $P < 0.05$ was considered statistically significant. All statistical analyses were performed by using SPSS statistical 22.0 software (IBM, Armonk, NY, USA).

Results

Epidemiological characteristics of pregnancy-associated listeriosis cases

During the study period, 14 cases of pregnancy-associated listeriosis were identified. 12 cases were inborn neonates, the rest (B2 and B3) were outborn neonates. The incidence of pregnancy-associated listeriosis in our hospital was 16.69/100,000 (14/83,875) births during our study period. They were occurred in all seasons, and a half were occurred in summer (50.0%, 7/14).

Characteristics of maternal cases

The characteristics of the 14 maternal listeriosis cases of pregnancy-associated listeriosis were summarized in table 1. Among them, 13 (M1-11, 13, 14) were singleton pregnancies and 1 (M12) was a twin pregnancy. Positive *L. monocytogenes* cultures from blood, amniotic fluid or placental tissue samples were found in eight (57.1%, 8/14) mothers.

Table 1. Characteristics of maternal listeriosis cases

No.	Gestation	Clinical presentation	Culture sites			MSAF	WBC (10 ⁹ /L)	CRP (mg/L)	Mode of delivery	Antibiotic(s) Before Delivery	Admission-delivery interval(hour/s)	C di
			Blood	Amniotic fluid	Placental tissue							
M1	38	None	/	/	/	No	8.5	4.7	Cesarean section	Cefuroxime	24	P
M2	32	NA	NA	NA	NA	NA	NA	NA	Vaginal delivery	NA	NA	N
M3	26	NA	NA	NA	NA	NA	NA	NA	Vaginal delivery	NA	NA	N
M4	30	Fever Tmax 38.2°C, Abd pain, vaginal bleeding	/	/	/	Yes	19.0	25	Vaginal delivery	Cefuroxime	22.5	G
M5	29	Abd pain, vaginal bleeding, RFM	/	/	/	Yes	13.9	54.4	Vaginal delivery	No	4	C
M6	34	RFM	/	/	/	Yes	17.3	30	Cesarean section	Azithromycin	3.5	T cl
M7	30	Abd pain, vaginal bleeding	/	(+)	(+)	Yes	27.6	56.8	Cesarean section	Piperacillin and sulbactam	10	U O
M8	30	Fever Tmax 39.7°C	(+)	(+)	(+)	Yes	17.0	52.9	Cesarean section	Ceftriaxone	1	G cc cl se al G pr
M9	37	Abd pain	(+)	/	/	No	7.1	64	Cesarean section	Cefuroxime	1	C se
M10	28	Abd pain, ROM, flu-like symptoms	/	(+)	(+)	Yes	24.8	62.8	Cesarean section	Ceftriaxone	1.5	C
M11	32	Abd pain, ROM	/	(+)	/	No	23.6	35	Cesarean section	Piperacillin and sulbactam+ Metronidazole	5.5	N
M12	36	Fever Tmax 38.4°C	/	(-)	(+)	No	24.0	85.2	Cesarean section	Cefotaxime	4	C h; pi or s)
M13	18	Fever Tmax 38°C, Abd pain, ROM, vaginal bleeding	(-)	(+)	/	No	17.2	96.9	Inevitable miscarriage	Piperacillin and sulbactam	17.5	C M
M14	35	Fever Tmax 38°C, Abd pain	(-)	(+)	(+)	Yes	23.1	36.3	Cesarean section	Ceftriaxone	3.5	B nr

MSAF meconium-stained amniotic fluid, WBC white blood cell, CRP C-reactive protein, NA not available, Tmax maximal temperature, Abd pain abdominal pain, GDM gestational diabetes mellitus, HSP hereditary spastic paraplegia, RFM reduced fetal movements, URI upper respiratory infection, HBV hepatitis B virus, GBS Group B streptococcus, ROM rupture of membranes.

Two cases were infected with *L. monocytogenes* in the second trimester pregnancy (between 14 and 27 weeks), and others were infected in the trimester pregnancy (between 28 and 41 weeks). Their median gestational age at the time of infection was 31 weeks (range: 18-38 weeks). Among the 14 maternal patients, 12 had premature deliveries, 1 experienced intrauterine fetal death and 1 had term delivery.

One of the 14 maternal patients experienced inevitable miscarriage, 4 had nature labor, and caesarean sections were performed in 9 cases, 8 due to fetal distress or intrauterine infection, 1 due to scarred uterus. Among the 12 mothers who delivered in this hospital, 41.7% (5/12) had prenatal fever at admission, only one patient (1/12, 8.3%) had flu-like symptoms before delivery. No patient with gastrointestinal symptoms before delivery has been observed, and only

one patient had diarrhea after delivery. Various obstetrical symptoms were observed, including abdominal pain (8/12, 66.7%), vaginal bleeding (4/12, 33.3%), rupture of membranes (9/12, 75.0%), and reduced fetal movement (2/12, 16.7%).

83.3% (10/12) maternal listeriosis patients had leukocytosis, and 91.7% (11/12) had elevated CRP levels. 58.3% (7/12) maternal cases had meconium-stained amniotic fluid (MSAF), and 58.3% (7/12) had chorioamnionitis. 91.7% (11/12) mothers were treated with pre-delivery antibiotics, but none received antepartum first-line empirical treatment (ampicillin or in combination with gentamicin). All of them eventually recovered after delivery shortly with no sequelae. However, the perinatal data of two maternal patients who delivered outside this hospital were unacquirable for analysis.

Characteristics of the neonatal cases

The characteristics of neonatal listeriosis cases were summarized in table 2. There were 11 culture-confirmed cases and 3 probable cases.

Table 2. Characteristics of neonatal cases

No.	Sex	GA (wk)	BW (g)	Apgar ^a	Clinical presentation	Culture sites			Intubation	WBC (10 ⁹ /L)	PLT (10 ⁹ /L)	Empirical antibiotic treatment	Complications
						Blood	Gastric fluid	Others					
B1	F	38	3760	9-10	Fever Tmax 38.1°C	(+)	/	/	No	8	171	Amoxicillin and Clavulanate Potassium	Septicemia, foramen ovale
B2	F	32	1650	5-8	RD, apnea, cyanosis, petechiae	(+)	/	Catheter (+)	No	9.6	220	Azlocillin	Abnormal liver function, pneumonia, septicemia, low birth weight
B3	F	26	1000	4-4	RD, apnea, petechiae, rash, cyanosis	(+)	/	/	Yes	6.3	103	Penicillin	Pneumonia, extremely premature in very low birth weight
B4	M	30	1650	5-7	RD, apnea, cyanosis	(+)	(+)	Eye secretions (+)	Yes	3.0	100	Penicillin + Meropenem	Septicemia, purulent meningitis, pneumonia, congenital heart disease, neonatal anaemia, BII neonatal intracranial hemorrhage, premature in low birth weight
B5	F	29	1350	7-8	RD, apnea, hypoglycemia, cyanosis, DOIC 5min	(+)	(+)	Sputum (+)	Yes	18.4	111	Cefotiam+ Penicillin	Pneumonia, septicemia, purulent meningitis, neonatal anaemia, BPD, hypoglycemia, premature in very low birth weight, apnea neonatorum
B6	M	34	2250	1-6	RD, apnea, lethargy, DOIC 5min	(+)	(+)	/	Yes	9.2	177	Penicillin+ Cefazidime	Sepsis, PNEU, PDA, preterm low birth weight
B7	F	30	1400	6-9	RD, apnea, DOIC 1min	(+)	(+)	/	Yes	12.3	186	Penicillin + Meropenem	Pneumonia, septicemia, purulent meningitis, anaemia, RC preterm birth, low birth weight
B8	M	30	1900	5-7	RD, apnea, cyanosis, DOIC 5min	(+)	(+)	/	Yes	9.3	35	Vancomycin+ Meropenem	Sepsis, shock, respiratory failure, cardiac insufficiency, capillary leak syndrome, purulent meningitis, intracranial hemorrhage, seizure, patent ductus arteriosus, coagulation disorders, thrombocytopenia, anaemia, low birth weight, preterm birth, pneumonia, hypoproteinemia, hypernatremia
B9	F	37	2700	9-10	Lethargy, rash	(+)	/	/	Yes	1.5	115	Penicillin+ Cefazidime	Sepsis, septic shock, respiratory failure, pneumonia, acidosis, coagulation disorders,

													thrombocytopenia, hypoproteinemia, seizure, hypocalcemia, purulent meningitis, and renal injury
B10	M	28	1250	8-9	RD	(-)	(+)	/	Yes	13.5	140	Penicillin + Meropenem	Respiratory infection, pneumonia, septicemia, low birth weight, patent ductus arteriosus, hypoproteinemia, anaemia, congenital hypothyroidism
B11	F	32	1600	9-10	RD	(-)	/	/	No	19.6	350	Azlocillin	Pneumonia, preterm birth, low birth weight
B12.1	F	36	2550	9-10	RD, lethargy	(-)	/	/	No	19.6	236	Ceftazidime	Pneumonia, preterm birth
B12.2	M	36	2250	9-10	RD, lethargy, petechiae	(-)	/	/	No	16.2	223	Ceftazidime	Pneumonia, preterm birth, low birth weight
B13	M	35	2570	9-10	Cyanosis	(-)	(+)	/	No	17.5	230	Penicillin+ Ceftazidime	Pneumonia, low birth weight, septicemia, neonatal hypocalcemia, neonatal anaemia, neonatal hypokalemia

^a Apgar score at 1st min - 5th min,

GA gestational age, BW birth weight, PLT platelet, A alive, RD respiratory distress, D dead, BIPI brain injury in premature infants, DOIC delay of initial crying, BPD bronchopulmonary dysplasia, PDA patent ductus arteriosus, ROP retinopathy of prematurity, AKI acute kidney injury, ASD atrial septal defect

Culture-confirmed neonatal listeriosis cases

In the 11 culture-confirmed cases, blood and newborn gastric fluid were the sensitive samples for microbiological diagnosis (positive in 9 [81.8%] of 11 blood samples and 7 [100.0%] of 7 gastric fluid samples), 9 (81.8%) were inborn neonates and 2 (18.2%) were outborn neonates, 6 (54.5%) were female and 5 (45.5%) were male. Most of them were preterm infants (9/11, 81.8%), with a median gestational age (GA) of 30 weeks (range: 26-38 weeks) and a median birth weight (BW) of 1650g (range: 1000-3760g), 8 (72.7%) weighed <2500g and 4 (36.4%) weighed <1500g.

Most of the 11 confirmed neonatal listeriosis cases were early onset (90.9%, 10/11) (B2-B10, B13). Among the 10 early onset cases, 8 cases (80.0%, 8/10) had respiratory distress and 7 case (70.0%, 7/10) received resuscitation at birth. The late onset one was a female who was delivered by caesarean section at GA of 38 weeks, and admitted to hospital with fever for one day at 8 days after birth.

Laboratory analyses were performed immediately on admission. Thrombocytopenia was observed in 54.5% (6/11) of culture-confirmed cases. Cerebrospinal fluid tap was performed in 81.8% (9/11) of neonates, and 4 of them were diagnosed as purulent meningitis, however, their CSF culture were all negative. Cranial imaging revealed intracranial hemorrhage in 2 neonates (B4 and B8), and one of them (B4) was diagnosed as brain injury in premature infants (BIPI).

All of the neonates with confirmed listeriosis were treated empirically, and most of them treated with Penicillin or Cephalosporin or Meropenem. Eight newborns received intubation. Three neonates (B2, B3 and B10) died, and two of them were outborn.

Probable neonatal listeriosis cases

Three newborns (B11, B12.1 and B12.2) were identified as probable neonatal listeriosis cases, including a pair of twins (B12.1 and B12.2). Both of their mothers were confirmed maternal listeriosis, but the culture results for these three neonates were negative. All of them were preterm infants, with GA of 32, 36 and 36 weeks, and BW of 1600g, 2550g and 2250g, respectively. They all had respiratory distress at birth, and were diagnosed as pneumonia subsequently, their laboratory parameters were mild abnormal. All of them were recovered soon after birth.

Risk factors for mortality in neonates with pregnancy-associated listeriosis

We compared the characteristics of the surviving neonates (n=11) with those of the non-surviving neonates (n=3), trying to identify risk factors for mortality in neonates with pregnancy-associated listeriosis (Table 3).

Fatality rates were significantly higher in outborn neonates ($P=0.005$). Besides, higher mortality rates were observed in neonates with lower BW ($P=0.038$), GA <28 weeks ($P=0.056$), and Apgar score (5th min) <5 ($P=0.056$), with marginally significant differences.

Table 3. Comparison between the survival and fatal cases of pregnancy-associated listeriosis

	Survival (n=11)	Fatality (n=3)	P value
BW			
BW (g) ^a	2250 (1600-2570)	1250 (1000-1650)	0.038
VLBW<1500g	2 (18.2%)	2 (66.7%)	0.112
GA			
GA (weeks) ^a	34 (30-36)	28 (26-32)	0.060
GA < 28 weeks	0 (0.0%)	1 (33.3%)	0.056
Apgar score			
Apgar score (1 st min) ^a	9 (5-9)	5 (4-8)	0.225
Apgar score (1 st min) <5	1 (9.1%)	1 (33.3%)	0.305
Apgar score (5 th min) ^a	10 (7-10)	8 (4-9)	0.225
Apgar score (5 th min) <5	0 (0.0%)	1 (33.3%)	0.056
Outborn (n, %)	0 (0.0%)	2 (66.7%)	0.005
Clinical Presentation			
Respiratory distress	8 (72.7%)	3 (100.0%)	0.325
Apnea	5 (45.5%)	2 (66.7%)	0.530
Cyanosis	3 (27.3%)	2 (66.7%)	0.224
Laboratory parameters			
WBC, 10 ⁹ /L ^a	12.3 (8.0-18.4)	9.6 (6.3-13.5)	0.659
PLT, 10 ⁹ /L ^a	177 (111-230)	140 (103-220)	0.659

^a Median(ranges),

BW birth weight, VLBW very low birth weight, GA gestational age, WBC white blood cell, PLT platelet

Discussion

To our knowledge, this is the first detailed study of pregnancy-associated listeriosis infections among children in Ningbo, Zhejiang province, China. The incidence of pregnancy-associated listeriosis in our hospital was 16.69/100,000 deliveries, most of the neonatal listeriosis cases (92.9%, 13/14) were early onset. 3 cases died, all of them were early onset and culture-confirmed neonates, 2 of them were outborn neonates. All of the maternal patients recovered after delivery shortly with no sequelae.

The diagnosis of maternal listeriosis would be challenging, apart from nonspecific obstetrical signs, fever or flu-like symptoms are the only clinical signs in infected mothers[24], but they were not always present. In our study, only one patient had flu-like symptoms before delivery. The difficulty of diagnose was associated with adverse fetal or neonatal outcome. First, we and others all observed that most maternal listeriosis cases were appeared during the second and third trimester period of pregnancy, mainly based on the adverse signs of their infants. While, the occurrence of first trimester maternal listeriosis was likely underestimated as the non-specific nature of disease presentation, thus may causing early fetal losses. Second, non-specific signs in infected pregnant women would influence the antibiotic prophylaxis to prevent neonatal listeriosis[22]. Previous studies have showed that adequate maternal antimicrobial treatment before delivery was associated with a significant decrease of infants' severity[24, 25]. Unfortunately, in our study and others performed in China[18], none of the maternal listeriosis cases received adequate first-line antimicrobial treatment[9]. This may partly account for the adverse outcome of *L. monocytogenes* infected neonates in the present study. In China, there are no national guidelines for the treatment of listeriosis currently. Although rare, listeriosis should be considered in the differential diagnosis of pregnancy-associated infections.

Pregnant women have much higher rates of *L. monocytogenes* infection than the general population[5-7]. Meanwhile, the consequences of listeriosis during pregnancy and in neonates are often severe[9]. Listeriosis neonates are always in critical condition, especially the early onset ones[15]. In the present study, 80.0% (8/10) of early onset confirmed neonatal listeriosis cases had respiratory distress, 70.0% (7/10) received resuscitation at birth, and 30.0% (3/10) died. Identified risk factors for mortality in neonates with pregnancy-associated listeriosis is important to treat and manage the high-risk neonates. It was found that fatality rates were significantly higher in the neonates who were born in other hospitals ($P=0.005$) in our study, these hospitals were lack of neonatal intensive care unit. It has been observed in a previous study that preterm neonates, who were delivered in hospitals with lower levels of neonatal care, suffered from higher rates of adverse outcomes[26]. The capability to provide timely and optimal resuscitative measures is important to reduce mortality of neonatal listeriosis.

Previous studies have also shown that lower GA was associated with a higher mortality[9]. While, later infection, particularly which occurred in the third trimester, is typically associated with more favorable outcomes than earlier infection. In our study, higher mortality rates also were observed in neonates with lower BW ($P=0.038$), GA <28 weeks ($P=0.056$), however, the differences of later one were not statistically significant. We did not show it may due to the low sample size. Further studies are needed to explore their relationship.

Some studies suggested that *L. monocytogenes* was more commonly associated with sporadic episodes and outbreaks rather than being affected by factors such as climate and season[20, 27]. However, seasonal trends of pregnancy-associated listeriosis cases have been observed in the Beijing and Taiwan[18, 28], with peaks reported in summer. Similarly, in our study, more than a half pregnancy-associated listeriosis cases were occurred in summer (50.0%, 7/14). As *L. monocytogenes* was a psychrophilic organism, the ready-to-eat foods and foods stored at refrigeration temperatures are the main sources of *L. monocytogenes*[29]. And people have more opportunity to eat food that is inadequately cooked and improperly stored in the refrigerator during the warmer seasons. Therefore, they were more likely to be *L. monocytogenes* infected in summer.

Unfortunately, the incidence of pregnancy-associated listeriosis in our hospital (16.69/100,000) was higher than that in the United States, United Kingdom and New Zealand (3.4, 3.4 and 12.3 per 100,000 births, respectively)[19-21], where listeriosis was a notifiable disease. The information of listeriosis especially pregnancy-associated cases is limited, and this might due to the lack of a good listeriosis surveillance system in China[30]. Besides, the incidence of pregnancy-associated listeriosis in our hospital was also higher than that in one hospital of Beijing (13.7/100,000), this might because our hospital is the only tertiary maternity and pediatric hospital in this area that admits many critically ill pregnant women and neonates.

There are several limitations in our study. First, because the miscarriage products at the first trimester GA were usually not available for microbiological investigations, and the blood cultures were not routine examinations for pregnant women without severe performance of infections in our hospital, the occurrence of maternal listeriosis was likely to be underestimated. Second, it was a hospital-based study, the sample size was relatively small, and some information were missing (information of two mothers of outborn listeriosis neonates), so our results could not be generalized to whole population. Third, it was a pity that we did not preserve the strains of *L. monocytogenes*, molecular epidemiology data were unavailable. Therefore, a study with a large sample size and relevant experiments is strongly encouraged.

Conclusion

We find that there is a neglected burden of pregnancy-associated listeriosis in our hospital. Listeriosis can have a devastating effect on fetus/neonates. The outborn neonates, who born in hospitals with lower levels of neonatal care, were probably associated with the increased severity. Although rare, listeriosis should be considered in the differential diagnosis of pregnancy-associated infections.

Declarations

Ethics approval and consent to participate

This study was approved by Ningbo Women and Children's Hospital Ethics Committee and our committee's reference number was EC2021-028. The informed consent was waived for this retrospective study by Ningbo Women and Children's Hospital Ethics Committee. All methods were carried out in accordance with relevant guidelines and regulations.

Consent for publication:

Not applicable.

Competing interests

The authors declare that they have no conflict of interest.

Funding

This study was supported by the Ningbo Clinical Research Center for Children's Health and Diseases (No. 2019A21002) and the Ningbo Public Service Technology Foundation, China (No. 2021S155).

Authors' contributions

YK contributed to the study design, data analysis, and manuscript writing and revision. LY, PZ and YS contributed to data collection and management. ZZ contributed to the data analysis and manuscript revision. All authors read and approved the final manuscript.

Acknowledgements

Not applicable.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

References

1. de Noordhout CM, Devleesschauwer B, Angulo FJ, Verbeke G, Haagsma J, Kirk M, et al. The global burden of listeriosis: a systematic review and meta-analysis. *Lancet Infect Dis*. 2014,14(11):1073-82. [https://doi.org/10.1016/S1473-3099\(14\)70870-9](https://doi.org/10.1016/S1473-3099(14)70870-9).
2. Wilking H, Lachmann R, Holzer A, Halbedel S, Flieger A, Stark K. Ongoing High Incidence and Case-Fatality Rates for Invasive Listeriosis, Germany, 2010-2019. *Emerg Infect Dis*. 2021,27(9):2485-8. <https://doi.org/10.3201/eid2709.210068>.
3. Kaptchouang Tchatchouang CD, Fri J, De Santi M, Brandi G, Schiavano GF, Amagliani G, et al. Listeriosis Outbreak in South Africa: A Comparative Analysis with Previously Reported Cases Worldwide. *Microorganisms*. 2020,8(1). <https://doi.org/10.3390/microorganisms8010135>.
4. Jackson KA, Gould LH, Hunter JC, Kucerova Z, Jackson B. Listeriosis Outbreaks Associated with Soft Cheeses, United States, 1998-2014(1). *Emerg Infect Dis*. 2018,24(6):1116-8. <https://doi.org/10.3201/eid2406.171051>.
5. Goulet V, Hebert M, Hedberg C, Laurent E, Vaillant V, De Valk H, et al. Incidence of listeriosis and related mortality among groups at risk of acquiring listeriosis. *Clin Infect Dis*. 2012,54(5):652-60. <https://doi.org/10.1093/cid/cir902>.
6. Pouillot R, Hoelzer K, Jackson KA, Henao OL, Silk BJ. Relative risk of listeriosis in Foodborne Diseases Active Surveillance Network (FoodNet) sites according to age, pregnancy, and ethnicity. *Clin Infect Dis*. 2012,54 Suppl 5:S405-10. <https://doi.org/10.1093/cid/cis269>.
7. Elinav H, Hershko-Klement A, Valinsky L, Jaffe J, Wiseman A, Shimon H, et al. Pregnancy-associated listeriosis: clinical characteristics and geospatial analysis of a 10-year period in Israel. *Clin Infect Dis*. 2014,59(7):953-61. <https://doi.org/10.1093/cid/ciu504>.
8. Pohl AM, Pouillot R, Bazaco MC, Wolpert BJ, Healy JM, Bruce BB, et al. Differences Among Incidence Rates of Invasive Listeriosis in the U.S. FoodNet Population by Age, Sex, Race/Ethnicity, and Pregnancy Status, 2008-2016. *Foodborne Pathog Dis*. 2019,16(4):290-7. <https://doi.org/10.1089/fpd.2018.2548>.
9. Charlier C, Perrodeau E, Leclercq A, Cazenave B, Pilmis B, Henry B, et al. Clinical features and prognostic factors of listeriosis: the MONALISA national prospective cohort study. *Lancet Infect Dis*. 2017,17(5):510-9. [https://doi.org/10.1016/S1473-3099\(16\)30521-7](https://doi.org/10.1016/S1473-3099(16)30521-7).
10. Herrador Z, Gherasim A, Lopez-Velez R, Benito A. Listeriosis in Spain based on hospitalisation records, 1997 to 2015: need for greater awareness. *Euro Surveill*. 2019,24(21). <https://doi.org/10.2807/1560-7917.ES.2019.24.21.1800271>.
11. Li W, Bai L, Ma X, Zhang X, Li X, Yang X, et al. Sentinel Listeriosis Surveillance in Selected Hospitals, China, 2013-2017. *Emerg Infect Dis*. 2019,25(12):2274-7. <https://doi.org/10.3201/eid2512.180892>.
12. Zhang X, Niu Y, Liu Y, Lu Z, Wang D, Cui X, et al. Isolation and Characterization of Clinical *Listeria monocytogenes* in Beijing, China, 2014-2016. *Front Microbiol*. 2019,10:981. <https://doi.org/10.3389/fmicb.2019.00981>.
13. Yang M, Huang G, Qian H, Wang J, Zhao S, Li S, et al. Progesterone decreased indoleamine 2,3-dioxygenase 1 (IDO1) expression in early pregnancy chorionic villi and decidua. *Autoimmunity*. 2021,54(3):156-62. <https://doi.org/10.1080/08916934.2021.1907572>.
14. Piccinni MP, Raghupathy R, Saito S, Szekeres-Bartho J. Cytokines, Hormones and Cellular Regulatory Mechanisms Favoring Successful Reproduction. *Front Immunol*. 2021,12:717808. <https://doi.org/10.3389/fimmu.2021.717808>.
15. Charlier C, Disson O, Lecuit M. Maternal-neonatal listeriosis. *Virulence*. 2020,11(1):391-7. <https://doi.org/10.1080/21505594.2020.1759287>.
16. Jalava K, Amar C, McLauchlin J, Godbole G. Increased incidence of listeriosis among pregnant women belonging to ethnic minorities in England. *J Infect*. 2021,82(6):276-316. <https://doi.org/10.1016/j.jinf.2021.03.024>.
17. Vidal EN, Rajadurai VS, Anand AJ, Chandran S. Listeriosis During Pregnancy and in Newborns: 18 Years of Data From a Large Tertiary Hospital in Singapore. *J Pediatric Infect Dis Soc*. 2020,9(4):498-501. <https://doi.org/10.1093/jpids/piz059>.
18. Li C, Zeng H, Ding X, Chen Y, Liu X, Zhou L, et al. Perinatal listeriosis patients treated at a maternity hospital in Beijing, China, from 2013-2018. *BMC Infect Dis*. 2020,20(1):601. <https://doi.org/10.1186/s12879-020-05327-6>.
19. Sapuan S, Kortsalioudaki C, Anthony M, Chang J, Embleton ND, Geethanath RM, et al. Neonatal listeriosis in the UK 2004-2014. *J Infect*. 2017,74(3):236-42. <https://doi.org/10.1016/j.jinf.2016.11.007>.
20. Jeffs E, Williman J, Brunton C, Gullam J, Walls T. The epidemiology of listeriosis in pregnant women and children in New Zealand from 1997 to 2016: an observational study. *BMC Public Health*. 2020,20(1):116. <https://doi.org/10.1186/s12889-020-8221-z>.
21. Silk BJ, Date KA, Jackson KA, Pouillot R, Holt KG, Graves LM, et al. Invasive listeriosis in the Foodborne Diseases Active Surveillance Network (FoodNet), 2004-2009: further targeted prevention needed for higher-risk groups. *Clin Infect Dis*. 2012,54 Suppl 5:S396-404. <https://doi.org/10.1093/cid/cis268>.
22. Rouse DJ, Keimig TW, Riley LE, Letourneau AR, Platt MY. CASE RECORDS of the MASSACHUSETTS GENERAL HOSPITAL. Case 16-2016. A 31-Year-Old Pregnant Woman with Fever. *N Engl J Med*. 2016,374(21):2076-83. <https://doi.org/10.1056/NEJMcp1516451>.
23. Angelo KM, Jackson KA, Wong KK, Hoekstra RM, Jackson BR. Assessment of the Incubation Period for Invasive Listeriosis. *Clin Infect Dis*. 2016,63(11):1487-9. <https://doi.org/10.1093/cid/ciw569>.
24. Fouks Y, Amit S, Many A, Haham A, Mandel D, Shinar S. Listeriosis in pregnancy: under-diagnosis despite over-treatment. *J Perinatol*. 2018,38(1):26-30. <https://doi.org/10.1038/jp.2017.145>.
25. Charlier C, Goffinet F, Azria E, Leclercq A, Lecuit M. Inadequate management of pregnancy-associated listeriosis: lessons from four case reports. *Clin Microbiol Infect*. 2014,20(3):246-9. <https://doi.org/10.1111/1469-0691.12281>.
26. Fang JL, Mara KC, Weaver AL, Clark RH, Carey WA. Outcomes of outborn extremely preterm neonates admitted to a NICU with respiratory distress. *Arch Dis Child Fetal Neonatal Ed*. 2020,105(1):33-40. <https://doi.org/10.1136/archdischild-2018-316244>.
27. Jemmi T, Stephan R. *Listeria monocytogenes*: food-borne pathogen and hygiene indicator. *Rev Sci Tech*. 2006,25(2):571-80.
28. Tai YL, Chi H, Chiu NC, Lin CY, Cheng JL, Hsu CH, et al. Clinical features of neonatal listeriosis in Taiwan: A hospital-based study. *J Microbiol Immunol Infect*. 2020,53(6):866-74. <https://doi.org/10.1016/j.jmii.2019.08.001>.

29. Salazar JK, Stewart D, Shazer A, Tortorello ML. Short communication: Long-term -20 degrees C survival of *Listeria monocytogenes* in artificially and process-contaminated ice cream involved in an outbreak of listeriosis. *J Dairy Sci.* 2020,103(1):172-5. <https://doi.org/10.3168/jds.2019-16774>.
30. Chen S, Meng F, Sun X, Yao H, Wang Y, Pan Z, et al. Epidemiology of Human Listeriosis in China During 2008-2017. *Foodborne Pathog Dis.* 2020,17(2):119-25. <https://doi.org/10.1089/fpd.2019.2683>.