Laterality of Blood Perfusion in the Lower Extremities after Drinking Saline at Different Temperatures

Shuyong Jia

Institute of Acupuncture and Moxibustion

Qizhen Wang

Institute of Basic Research in Clinical Medicine

Hongyan Li

Institute of Acupuncture and Moxibustion

Xiaoqing Song

Institute of Acupuncture and Moxibustion, China Academy of Chinese Medical Sciences

Shuyou Wang

Institute of Acupuncture and Moxibustion, China Academy of Chinese Medical Sciences

Weibo Zhang

Institute of Acupuncture and Moxibustion

Guangjun Wang (✉ tjuwgj@gmail.com)

Institute of Acupuncture and Moxibustion <https://orcid.org/0000-0001-5344-4594>

Article

Keywords: Laser Doppler blood perfusion, laterality, wavelet analysis, saline, EGG, ECG, dominant frequency, RMSSD

Posted Date: April 20th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-112441/v4>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Skin blood flux (SkBF) changes caused by drinking cold water are generally associated with vagal tone and osmotic factors in the digestive system. However, the distribution and regulation of bilateral blood flow are not symmetrical but exhibit a certain laterality. The aim of this study was to analyze the laterality of skin blood flux after drinking saline (0.9%) at different temperatures by monitoring blood flux in the bilateral lower extremities.

Methods: A total of 60 subjects were recruited for this study, and all subjects completed all measurements. Subjects were randomly divided into a 4 °C group, 10 °C group and 30 °C group. For every subject, skin blood perfusion of bilateral Zusanli acupoints (ST 36) was recorded simultaneously before and after drinking 500 ml of 0.9% saline using a PeriFlux System 5000. The electrogastrogram (EGG) was also monitored, and the dominant frequency of the EGG and heart rate variability were analyzed.

Results: The results indicated that after drinking saline at different temperatures, the laterality index of skin blood flux at the lower extremities was different. Drinking 30 °C saline can increase the laterality index of the lower extremities. The results of wavelet analysis showed that the laterality index changes of skin blood flux were mainly reflected in the frequency interval V (0.4-1.6 Hz). Stimulation at 4 °C and 10 °C resulted in a decrease in the dominant frequency (DF) of the electrogastrogram and an increased root mean square of successive differences (RMSSD) of the RR interval. There was a weak negative correlation between the laterality index of frequency interval I or II and root mean square of successive differences.

Conclusion: There was laterality of blood perfusion in the lower extremities after drinking saline at different temperatures.

Background

Although people prefer to drink cold water[1], few reports have studied the effects of water temperature on healthy humans[2-4]. Especially, studies about the effect of water temperature on blood flux are scarce. Scientists have suggested that the water effect does not depend on gastric distension, but on the water osmolality, which could cause an autonomic cardiovascular response in humans through osmotic sensing nerve fibers in the intestinal or portal circulation[5,6]. On the other hand, the results shown that water intake activates distinct gastrointestinal vagal afferent in a temperature-dependent pattern and might affect cardiac vagal tone[7].

A previous work suggested that oral glucose intake inhibits hypothalamic neuronal activity than intravenous (IV) glucose administration more effectively[8], but oral intake saline results in minimal variations in serum albumin, hematocrit, and hemoglobin compared to IV infusion[9]. These studies indicated that, in addition to osmotic pressure and temperature mechanisms, the different autonomic responses of digestive tract might result in systemic regulation of peripheral blood perfusion. Furthermore, some works have indicated that the cutaneous microvasculature mirrors the vascular

function of other parts of the body[10-12]. Therefore, skin blood flow can be used to explore the regulation of gastrointestinal autonomic function after stimulation. According to previous studies[13,14], the cutaneous blood perfusion of both sides is different.

Therefore, we hypothesized that ingestion of saline at different temperatures influences human cutaneous blood perfusion, which will lead to lateralization of the regulation of blood perfusion. To test these hypotheses, we compared, with a randomized control design, the cutaneous blood perfusion changes in response to different temperatures of saline and additionally evaluated their potentially differential impacts on EGG DF and cardiovascular variables.

Methods

Inclusion and exclusion criteria

Eligible subjects had to be healthy and between the ages of 18 and 60 years. All participants were requested to avoid alcohol, tea and coffee for at least 24 hours prior to the test. None of the subjects had any diseases or were taking any medication affecting cardiovascular or autonomic regulation.

Participants and groups

A total of 60 healthy subjects were enrolled in the study, and all subjects completed measurements and were included in the statistical analysis. All of the subjects lived in Beijing, and all of them were Chinese. The general characters are presented in Table 1. All experiments took place in a quiet, temperature-controlled (24-26 °C) laboratory. On arrival at the laboratory, subjects were asked to empty their bladders. Following a period of cardiovascular stability (40 min), a baseline recording was then made for at least 15 min. Then, over a 5 min period, the test subjects ingested 500 mL of 0.9% saline at different temperatures, and EGG and skin blood flux were monitored. The experimental design, signal recording and analysis process were shown in Figure S1.

Electrogastrogram measurement protocol and analysis

The subjects maintained a supine position. Eight cutaneous electrodes were placed on the abdomen of the participant and connected to DC amplifiers. The EGG recordings were processed with a NeurOne system (NeurOne, MEGA electronics Ltd, Finland). The data were digitized with a sampling rate of 10000 Hz and then downsampled at 1000 Hz. EGG data were analyzed with the FieldTrip toolbox[15], and the analysis code for EGG analysis was provided by Rebollo, I[16].

RMSSD and mean RR interval analysis

The analysis method can be referenced in our previous study[17-19]. The raw data of channel one were exported in EDF format, imported into Kubio HRV software and analyzed [20]. The analysis parameter was default. The root mean square of successive differences (RMSSD) was given by (1).

$$RMSSD = \sqrt{\frac{1}{N-1} \sum_{n=1}^{N-1} (RR_{n+1} - RR_n)^2} \quad (1)$$

Protocol for measurement of blood perfusion

Both legs were exposed, and bilateral Zusanli acupoints (ST 36) were marked by senior acupuncture doctors. The acupuncture point ST 36 is located in the tibialis anterior muscle, 4 fingerbreadths below the kneecap and 1 fingerbreadth lateral from the anterior crest of the tibia[21]. Blood perfusion signals were recorded using a PeriFlux System 5000 (Perimed AB, Stockholm, Sweden) with a 64 Hz sample rate and 0.2 s time constant. An optical fiber probe connected with a Periflux 5000 was used to illuminate and collect the scattered light from the skin tissue. The probe was attached to the surface of interest with two-sided adhesive tape.

Mean blood perfusion analysis and wavelet analysis

The recorded file of each subject was opened in PeriSoft (version 2.5.5, Perimed AB, Stockholm, Sweden) for Windows. The detailed data were exported in txt format and then imported into MATLAB software and analyzed. The laterality index of blood flux every 10 min on both sides of ST36 was calculated as (1).

$$Laterality\ Index = \frac{mean(|Post_L - Post_R|)}{mean(|Pre_L - Pre_R|)} \times 100 \quad (2)$$

Previous studies indicated that blood flux oscillations at frequencies from 0.0095 to 1.6 Hz might reflect different physiological rhythms[22], which can be separated into five frequency bands in the frequency domain[23-26]. In the present study, wavelet analysis was performed on the blood flux signal (10 min) using a Morlet wavelet (The Mathworks Inc., Natick, MA, USA). For every frequency interval, the laterality index was also calculated as (2).

Statistical analysis

Data are presented as the mean±SE. Mixed repeated-measures ANOVA was used to analyze between-subject factors with R software[27] and fdrtool package[28]. All correlation analyses were conducted using MATLAB software (MathWorks, Natick, Massachusetts, USA). All reported *P* values were two-sided, and the level of significance was defined as *P*<0.05.

Results

In this study, a total of 60 subjects were recruited, and all subjects were included in the final statistical analysis. The order of stimulations was randomly generated for each subject. Detailed information on the subjects is summarized in Table 1 (4 °C, N=20; 10 °C, N=20 and 30 °C, N=20).

Table 1. Subject's sex composition and average age, height and weight

Group	n	Sex (female/male)	Age (years, mean±SD)	Height (cm, mean±SD)	Weight (kg, mean±SD)
4 °C	20	17/3	27±3	165.0±7.2	59.6±9.3
10 °C	20	18/2	25±3	163.6±5.6	56.2±10.7
30 °C	20	17/3	26±1	161.9±6.1	54.6±13.6

EGG and ECG results

The results of EGG analysis are shown in Figure 1, the recording electrode position is shown in Figure 1A, and the raw data of 8-channel gastrointestinal electrical signals recorded are shown in Figure 1B. The spectral analysis of each signal is shown in Figure 1C. The statistical results of the main frequency corresponding to the maximum power in each subject's 8-channel gastrointestinal electrical signals are shown in Figure 1D. In the resting state, there was no significant difference in the dominant frequency of the subjects.

Cold stimulation of the digestive tract will not only cause a change in the electrogastrogram but also be accompanied by systemic reactions. The raw data of ECG and RR interval signal are shown in Figure 2A and Figure 2B, respectively. The changes in the RR interval and RMSSD were also analyzed. The results showed that the RR interval (Figure 2C) and RMSSD (Figure 2D) were increased with both 4 °C or 10 °C stimulation compared with 30 °C stimulation.

Skin blood flux

The average responses of blood perfusion to stimulation at different temperatures over time are shown in Figure 3. The recording position of both sides for blood perfusion is shown in Figure 3A, the raw data of bilateral blood perfusion are shown in Figure 3B, the absolute value of bilateral blood flow difference is shown in Figure 3C, and the laterality change in bilateral blood perfusion under different temperature stimulation conditions is shown in Figure 3D. In the first 10 min after stimulation, there was a significant difference between the changes induced by 30 °C stimulation and 4 °C stimulation. At the second and third 10 min after stimulation, there were significant differences between the changes induced by 30 °C stimulation and those caused by 4 °C and 10 °C. However, there were no significant changes in the changes induced by 4 °C and 10 °C.

Because the blood flow signal is composed of many components, bilateral blood flow was transformed by wavelet (Figure 4A), and the laterality index in different frequency intervals was calculated. The results

showed that the difference was mainly reflected in frequency V (0.4-1.6 Hz) (Figure 4F). There was no significant difference between other frequency intervals (Figure 4B-E).

To further analyze the possibility of blood flow laterality, we analyzed the correlation between these parameters. First, we confirmed that there was no correlation between the DF and RMSSD (Figure S2) at different temperature stimulation. There was also no apparent correlation between the laterality index and the DF (Figure S3A-C) or RMSSD (Figure S3D-F). After wavelet transformation, we found that there was a negative correlation between RMSSD and the laterality index of frequencies I at 4 °C stimulation (Figure 5A and D), which was independent of other frequency intervals (Figure 5G, J and M). There was a positive correlation between RMSSD and the laterality index of frequency II (Figure 5F), rather than other frequencies (Figure 5C, I, L and O) at 30 °C stimulation. There was no correlation between the laterality index of each frequency and RMSSD at 10 °C (Figure 5B, E, H, K and N). However, there is no correlation between the laterality index and DF in any frequency interval (Figure S4).

Discussion

As Figure 6 shows, the main significance of current work is that there was response laterality of blood perfusion in the lower extremities after drinking saline at different temperatures, which means that autonomic nerve function is different between left and right in the process of blood flow regulation. The present study demonstrated that drinking saline at different temperatures produced the changes of cutaneous vascular movement, which means that regulation of cutaneous vascular might play an important role in water temperature digestion.

A previous study[5] indicated that the water effect does not depend on gastric distension, but on the water osmolality, which could cause an autonomic cardiovascular response in humans through osmotic sensing nerve fibers in the intestinal or portal circulation[5,6]. However, in the current study, osmotic pressure of saline at different temperatures is same, it is difficult to conclude that the change in sympathetic tone is caused by osmolality.

In peripheral, small changes in cutaneous vascular blood perfusion can shift cardiovascular control. A decrease in the skin blood perfusion can lead to relative changes of sympathetic activation in peripheral[29]. Which can be explained by the activation of temperature sensitive vagal afferent fibers found in the digestive tract[30]. Water intake seems to activate different gastrointestinal vagal afferent fibers in a temperature dependent manner and may affect subsequent cardiac vagal tone[7]. Our results demonstrated that different water temperatures not only decreased the DF of EGG but also were associated with a change in the mean RR interval and RMSSD. However, as shown in Fig 6, there is no direct linear relationship between bilateral blood flow laterality and RMSSD, but after decomposing the blood flow components, it is found that there is a linear relationship between specific components and RMSSD. It is suggested that the regulation of peripheral blood flow induced by stimulation at different temperatures is selective. In the present study, the change trend of SkBF at bilateral ST 36 was different

after drinking saline at different temperatures. As in previous findings[13,31], this phenomenon is defined as laterality and can be described quantitatively by the laterality index.

Laterality tends to focus on the differentiation of left and right handedness and cerebral hemisphere function. However, little attention has been given to the lateralization of blood flow distribution. Our previous studies have found that there are significant differences in the distribution and regulation of blood flow in the same parts of both sides of the body[13,14], and such results have been confirmed by other research groups[32]. A previous study indicated that in young subjects, sympathetic vasoconstrictor activation after drinking water was not accompanied by an increase in arterial blood pressure[33], which suggested that the change in vascular tone in the limbs may be partially compensated by opposing changes in other vascular beds[33]. Therefore, it is acceptable that the distribution of bilateral blood flow and its variation are asymmetric.

In this study, because the age of the recruited subjects is relatively concentrated, it can not represent all age groups. The majority of the subjects recruited are female. Therefore, it is impossible to analyze the changes of lower extremity blood flow laterality index in different age groups and different gender groups, which is the limitation of this study. In addition, the span of stimulation temperature setting is too large. This is also the limitation of this study.

Conclusion

There was laterality of blood perfusion in the lower extremities after ingestion of saline at different temperatures.

List Of Abbreviations

SkBF: skin blood flux;

DF: dominant frequency;

RMSSD: root mean square of successive differences;

ST 36: Zusanli acupoint;

ECG: electrocardiogram;

EKG: electrogastrogram

HRV: heart rate variability;

TCM: Traditional Chinese Medicine;

Fig: figure;

SD: standard deviation;

SE: standard error;

ANOVA: analysis of variance

Declarations

Data availability

The majority of supporting data are presented within this article and its [Supplement](#). Raw data are available from the corresponding author upon reasonable request.

Ethics approval

This study was approved by the Institutional Research Ethics Boards of Acupuncture & Moxibustion, China Academy of Chinese Medical Sciences. In accordance with the Declaration of Helsinki each subject provided informed consent and had an adequate understanding of the procedure and purpose of this study.

Competing interests

The authors declare that they have no competing interests.

References

1. Burdon, C.A.; Johnson, N.A.; Chapman, P.G.; O'Connor, H.T. Influence of beverage temperature on palatability and fluid ingestion during endurance exercise: a systematic review. *Int J Sport Nutr Exerc Metab* **2012**, *22*, 199-211.
2. Boschmann, M.; Steiniger, J.; Hille, U.; Tank, J.; Adams, F.; Sharma, A.M.; Klaus, S.; Luft, F.C.; Jordan, J. Water-induced thermogenesis. *J Clin Endocrinol Metab* **2003**, *88*, 6015-6019, doi:10.1210/jc.2003-030780.
3. Brown, C.M.; Dulloo, A.G.; Montani, J.P. Water-induced thermogenesis reconsidered: the effects of osmolality and water temperature on energy expenditure after drinking. *J Clin Endocrinol Metab* **2006**, *91*, 3598-3602, doi:10.1210/jc.2006-0407.
4. Wang, G.; Jia, S.; Li, H.; Wang, Z.; Tian, Y.; Zhang, W. Response of Blood Perfusion at ST 36 Acupoint after Drinking Cold Glucose or Saline Injection. *Evidence-based complementary and alternative medicine : eCAM* **2017**, *2017*, 4212534, doi:10.1155/2017/4212534.
5. May, M.; Jordan, J. The osmopressor response to water drinking. *Am J Physiol Regul Integr Comp Physiol* **2011**, *300*, R40-46, doi:10.1152/ajpregu.00544.2010.

6. Brown, C.M.; Barberini, L.; Dulloo, A.G.; Montani, J.P. Cardiovascular responses to water drinking: does osmolality play a role? *Am J Physiol Regul Integr Comp Physiol* **2005**, *289*, R1687-1692, doi:10.1152/ajpregu.00205.2005.
7. Girona, M.; Grasser, E.K.; Dulloo, A.G.; Montani, J.P. Cardiovascular and metabolic responses to tap water ingestion in young humans: does the water temperature matter? *Acta physiologica* **2014**, *211*, 358-370, doi:10.1111/apha.12290.
8. Smeets, P.A.; Vidarsdottir, S.; de Graaf, C.; Stafleu, A.; van Osch, M.J.; Viergever, M.A.; Pijl, H.; van der Grond, J. Oral glucose intake inhibits hypothalamic neuronal activity more effectively than glucose infusion. *Am J Physiol Endocrinol Metab* **2007**, *293*, E754-758, doi:10.1152/ajpendo.00231.2007.
9. de Aguilar-Nascimento, J.E.; Valente, A.C.; Oliveira, S.S.; Hartmann, A.; Shlessarenko, N. Changes in body composition, hematologic parameters, and serum biochemistry after rapid intravenous infusion or oral intake of 2 liters of 0.9% saline solution in young healthy volunteers: randomized crossover study. *World J Surg* **2012**, *36*, 2776-2781, doi:10.1007/s00268-012-1756-0.
10. Holowatz, L.A.; Thompson-Torgerson, C.S.; Kenney, W.L. The human cutaneous circulation as a model of generalized microvascular function. *J. Appl. Physiol.* **2008**, *105*, 370-372, doi:10.1152/japplphysiol.00858.2007.
11. Stewart, J.; Kohen, A.; Brouder, D.; Rahim, F.; Adler, S.; Garrick, R.; Goligorsky, M.S. Noninvasive interrogation of microvasculature for signs of endothelial dysfunction in patients with chronic renal failure. *Am. J. Physiol.-Heart Circul. Physiol.* **2004**, *287*, H2687-H2696, doi:10.1152/ajpheart.00287.2004.
12. Rossi, M.; Bradbury, A.; Magagna, A.; Pesce, M.; Taddei, S.; Stefanovska, A. Investigation of skin vasoreactivity and blood flow oscillations in hypertensive patients: effect of short-term antihypertensive treatment. *J. Hypertens.* **2011**, *29*, 1569-1576, doi:10.1097/HJH.0b013e328348b653.
13. Wang, G.; Han, J.; Litscher, G.; Zhang, W. System identification algorithm analysis of acupuncture effect on mean blood flux of contralateral hegu acupoint. *Evidence-based complementary and alternative medicine : eCAM* **2012**, *2012*, 951928, doi:10.1155/2012/951928.
14. Wang, G.; Tian, Y.; Jia, S.; Wang, Z.; Zhang, W. Laterality and coherence analysis of Laser Doppler Flowmetry signals in bilateral Nèi guān(PC6 内关): a potential non-invasive method to assess microcirculatory changes in different ages. *World Journal of Acupuncture-Moxibustion* **2017**, *27*, 47-52.
15. Oostenveld, R.; Fries, P.; Maris, E.; Schoffelen, J.M. FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Comput Intel Neurosc* **2011**, Artn 15686910.1155/2011/156869, doi:Artn 15686910.1155/2011/156869.
16. Wolpert, N.; Rebollo, I.; Tallon-Baudry, C. Electrogastrography for psychophysiological research: Practical considerations, analysis pipeline, and normative data in a large sample. *Psychophysiology* **2020**, *57*, e13599, doi:10.1111/psyp.13599.

17. Wang, G.; Tian, Y.; Jia, S.; Zhou, W.; Zhang, W. Acupuncture regulates the heart rate variability. *J Acupunct Meridian Stud* **2015**, *8*, 94-98, doi:10.1016/j.jams.2014.10.009.
18. Guangjun, W.; Yuying, T.; Shuyong, J.; Wenting, Z.; Weibo, Z. Bilateral hegu acupoints have the same effect on the heart rate variability of the healthy subjects. *Evidence-based complementary and alternative medicine : eCAM* **2014**, *2014*, 106940, doi:10.1155/2014/106940.
19. Wang, G.; Tian, Y.; Jia, S.; Zhou, W.; Zhang, W. Pilot study of acupuncture point laterality: evidence from heart rate variability. *Evidence-based complementary and alternative medicine : eCAM* **2013**, *2013*, 476064, doi:10.1155/2013/476064.
20. Niskanen, J.P.; Tarvainen, M.P.; Ranta-Aho, P.O.; Karjalainen, P.A. Software for advanced HRV analysis. *Comput Methods Programs Biomed* **2004**, *76*, 73-81, doi:10.1016/j.cmpb.2004.03.004.
21. Hui, K.K.; Liu, J.; Marina, O.; Napadow, V.; Haselgrove, C.; Kwong, K.K.; Kennedy, D.N.; Makris, N. The integrated response of the human cerebro-cerebellar and limbic systems to acupuncture stimulation at ST 36 as evidenced by fMRI. *Neuroimage* **2005**, *27*, 479-496, doi:10.1016/j.neuroimage.2005.04.037.
22. Bernardi, L.; Rossi, M.; Fratino, P.; Finardi, G.; Mevio, E.; Orlandi, C. Relationship between phasic changes in human skin blood flow and autonomic tone. *Microvasc Res* **1989**, *37*, 16-27.
23. Hsiu, H.; Hsu, W.C.; Hsu, C.L.; Huang, S.M. Assessing the effects of acupuncture by comparing needling the hegu acupoint and needling nearby nonacupoints by spectral analysis of microcirculatory laser Doppler signals. *Evid-Based Compl Alt* **2011**, *2011*, 435928, doi:10.1093/ecam/neq073.
24. Thorn, C.E.; Matcher, S.J.; Meglinski, I.V.; Shore, A.C. Is mean blood saturation a useful marker of tissue oxygenation? *Am. J. Physiol.-Heart Circul. Physiol.* **2009**, *296*, H1289-H1295, doi:10.1152/ajpheart.01192.2008.
25. Bernjak, A.; Clarkson, P.B.; McClintock, P.V.; Stefanovska, A. Low-frequency blood flow oscillations in congestive heart failure and after beta1-blockade treatment. *Microvasc Res* **2008**, *76*, 224-232, doi:10.1016/j.mvr.2008.07.006.
26. Kvandal, P.; Landsverk, S.A.; Bernjak, A.; Stefanovska, A.; Kvernmo, H.D.; Kirkeboen, K.A. Low-frequency oscillations of the laser Doppler perfusion signal in human skin. *Microvasc Res* **2006**, *72*, 120-127, doi:10.1016/j.mvr.2006.05.006.
27. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>. **2020**.
28. Klaus, B.; Strimmer, K. fdrtool: Estimation of (Local) False Discovery Rates and Higher Criticism. R package version 1.2.16. <https://CRAN.R-project.org/package=fdrtool>. **2021**.
29. Eicke, B.M.; Milke, K.; Schlereth, T.; Birklein, F. Comparison of continuous wave Doppler ultrasound of the radial artery and laser Doppler flowmetry of the fingertips with sympathetic stimulation. *J Neurol* **2004**, *251*, 958-962, doi:10.1007/s00415-004-0471-7.
30. Gupta, B.N.; Nier, K.; Hensel, H. Cold-sensitive afferents from the abdomen. *Pflugers Arch* **1979**, *380*, 203-204.

31. Wang, G.; Litscher, D.; Tian, Y.; Gaischek, I.; Jia, S.; Wang, L.; Zhang, W.; Litscher, G. Acupuncture Point Laterality: Evidence and Perspective. *Integrative Medicine International* **2014**, *1*, 102-107.
32. Mezentseva, L.V.; Pertsov, S.S. Correlation-and-Spectral Analysis of Time Series for Microcirculatory Parameters in Vessels of Symmetrical Organs. *Bull Exp Biol Med* **2019**, *166*, 301-305, doi:10.1007/s10517-019-04337-y.
33. Scott, E.M.; Greenwood, J.P.; Stoker, J.B.; Gilbey, S.G.; Mary, D.A. Water drinking and sympathetic activation. *Lancet* **2000**, *356*, 2013, doi:10.1016/S0140-6736(05)72987-1.

Figures

Figure 1

EGG power spectra changes. (A) Cutaneous electrode position, red indicates recording electrodes, green indicates reference electrode; black indicates ground electrode; (B) raw data of EGG; (C) power spectra of EGG; (D) EGG peak frequency before and after stimulation. Before stimulation, $F(2,57)=0.38$, $P=0.6884$; after stimulation, $F(2,57)=6.45$, $P=0.003$; for 4 °C stimulation, $t=3.9819$ $\times$ $P_{corr} < 0.001$ (Pre VS Post, paired t test); for 10 °C stimulation, $t=4.9807$ $\times$ $P_{corr} < 0.001$ (Pre VS Post, paired t test); for 30 °C stimulation, $t=1.8492$ $\times$ $P_{corr} = 0.24$ (Pre VS Post, paired t test). P_{corr} : corrected p value.

Figure 2

Change in HRV. (A) ECG signal from channel 1. (B) RR intervals of ECG. (C) Change rate of mean RR, $F(2,57)=8.73$, $P=0.0005$, $**$, $P_{corr}<0.01$ (4 °C VS 30 °C); $##$, $P_{corr}<0.01$ (10 °C VS 30 °C). (D) Change rate of RMSSD, $F(2,57)=3.75$, $P=0.029$. $*$, $P_{corr}<0.05$, (4 °C VS 30 °C). P_{corr} : corrected p value. All values are reported as the mean $\pm$ SE.

Figure 3

Blood perfusion in both sides. (A) Recording position at ST 36. (B) Raw data of blood flux in both sides. (C) The absolute value of the difference between two sides. (D) Laterality index of blood perfusion. Main effect of stimulates: $F(2,57)=5.749$, $P=0.0053$; main effect of time: $F(3,171)=5.268$, $P=0.00169$; interaction effect: $F(6,171)=5.8$, $P<0.0001$. Post hoc analysis revealed a significant increase at 30 °C group from the first session *, $P_{corr}<0.05$, **, $P_{corr}<0.01$ (4 °C VS 30 °C); #, $P_{corr}<0.05$, ##, $P_{corr}<0.01$ (10 °C VS 30 °C). P_{corr} : corrected p value. All values are reported as the mean $\pm$ SE.

Figure 4

Change rate of different frequency intervals transformed by wavelet. (A) Wavelet analysis of blood flux. (B) Frequency interval I (0.0095-0.02 Hz), main effect of stimulates: $F(2,57)=0.723$, $P=0.49$; main effect of time: $F(3,171)=7.923$, $P<0.01$; interaction effect: $F(6,171)=0.954$, $P=0.458$. (C) Frequency interval II (0.02-0.06 Hz), main effect of stimulates: $F(2,57)=0.952$, $P=0.392$; main effect of time: $F(3,171)=9.371$, $P<0.01$; interaction effect: $F(6,171)=0.696$, $P=0.653$. (D) Frequency interval III (0.06-0.15 Hz), main effect of stimulates: $F(2,57)=3.288$, $P=0.0445$; main effect of time: $F(3,171)=2.485$, $P=0.0624$; interaction effect: $F(6,171)=2.7$, $P=0.0157$. (E) Frequency interval IV (0.15-0.4 Hz), main effect of stimulates: $F(2,57)=2.746$, $P=0.0727$; main effect of time: $F(3,171)=2.392$, $P=0.07036$; interaction effect: $F(6,171)=3.172$, $P=0.0056$. (F) Frequency interval V (0.4-1.6 Hz), main effect of stimulates: $F(2,57)=6.085$, $P=0.00403$; main effect of time: $F(3,171)=3.159$, $P=0.0261$; interaction effect: $F(6,171)=5.1$, $P<0.001$. Post hoc analysis revealed a significant increase at 30 °C group from the first session *, $P_{corr}<0.05$, **, $P_{corr}<0.01$ (4 °C VS 30 °C); #, $P_{corr}<0.05$, ##, $P_{corr}<0.01$ (10 °C VS 30 °C). P_{corr} : corrected p value. All values are reported as the mean $\pm$ SE.

Figure 5

Relationship between laterality index of different frequency intervals and change rate of RMSSD. (A). Frequency I at 4 °C. (B). Frequency I at 10 °C. (C). Frequency I at 30 °C. (D) Frequency II at 4 °C. (E) Frequency II at 10 °C. (F) Frequency II at 30 °C. (G) Frequency III at 4 °C. (H) Frequency III at 10 °C. (I) Frequency III at 30 °C. (J) Frequency IV at 4 °C. (K) Frequency IV at 10 °C. (L) Frequency IV at 30 °C. (M) Frequency V at 4 °C. (N) Frequency V at 10 °C. (O) Frequency V at 30 °C. ρ , Spearman's correlation coefficient; P_{corr} , corrected p value.

Figure 6

General illustration of the relationship. There was no significant correlation between the laterality index of lower extremity blood flow and DF or RMSSD. Under the condition of 4 °C stimulation, there was a weak negative correlation between the laterality index of frequency interval I and RMSSD after wavelet transform. There was a weak positive correlation between the laterality index of interval II and RMSSD under the 30 °C stimulation. Red line indicates $P_{corr} > 0.05$; while green line indicates $P_{corr} < 0.05$.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplementary.docx](#)