Lifestyle Factors Associated with Prevalent and Exacerbated Musculoskeletal Pain After the Great East Japan Earthquake: A Cross-Sectional Study from the Fukushima Health Management Survey

Hiroshige Jinnouchi (✉ shigenouchi@gmail.com)

Nippon Medical School <https://orcid.org/0000-0001-7677-7017>

Tetsuya Ohira

Department of Epidemiology, Fukushima Medical University School of Medicine, 1 Hikariga-oka, Fukushima, 960-1295, Japan.

Hironobu Kakihana

Department of Hygiene and Public Health, Osaka Medical College, 2-7 Daigakumachi, Takatsuki, Osaka, 569-8686, Japan.

Ko Matsudaira

Department of Medical Research and Management for Musculoskeletal Pain, 22nd Century Medical and Research Center, Faculty of Medicine, the University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo, 113-8655, Japan.

Masaharu Maeda

Department of Disaster Psychiatry, Fukushima Medical University School of Medicine, 1 Hikariga-oka, Fukushima, 960-1295, Japan.

Hirooki Yabe

Department of Neuropsychiatry, Fukushima Medical University School of Medicine, 1 Hikariga-oka, Fukushima, 960-1295, Japan.

Yuriko Suzuki

Department of Adult Mental Health, National Center of Neurology and Psychiatry, National Institute of Mental Health, 4-1-1 Ogawa-Higashi, Kodaira, 187-8553, Japan.

Mayumi Harigane

Fukushima Medical University

Hiroyasu Iso

Public Health, Department of Social Medicine, Osaka University Graduate School of Medicine, 2-2 Yamadaoka, Suita, Osaka, 565-0871, Japan.

Tomoyuki Kawada

Nippon Medical School

Seiji Yasumura

Department of Public Health, Fukushima Medical University School of Medicine, 1 Hikariga-oka, Fukushima, 960-1295, Japan.

Kenji Kamiya

Research Institute for Radiation Biology and Medicine, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima City, Hiroshima, 734-8553, Japan.

Research article

Keywords: musculoskeletal pain, lifestyle factor, the Great East Japan Earthquake

Posted Date: January 13th, 2020

DOI: <https://doi.org/10.21203/rs.2.20646/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Public Health on May 13th, 2020. See the published version at <https://doi.org/10.1186/s12889-020-08764-9>.

Abstract

Background: While the prevalence of post-disaster musculoskeletal pain has been documented, its associated disaster-related factors have not been investigated. This study determined the prevalence of and lifestyle factors associated with musculoskeletal pain among evacuees of the Great East Japan Earthquake.

Methods: We conducted a cross-sectional study of 34,919 evacuees, aged 40-89 years, without obvious disabilities in activities of daily living at about 1 year after the disaster. The evacuees were asked about their musculoskeletal pain (low back and limb pain) and lifestyle factors: use of evacuation shelters or temporary housing at any point of time, job loss after the disaster, decreased income after the disaster, current smoking status, current drinking status, lack of sleep, regular exercise, and participation in recreational or community activities. Furthermore, psychological factors, such as traumatic reactions, psychological distress, and somatic symptoms, affecting musculoskeletal pain were assessed. We used multinomial logistic regression analysis to calculate odds ratios of each lifestyle factor for prevalent and prevalent plus exacerbated musculoskeletal pain.

Results: Musculoskeletal pain prevalence was 32.8%: 27.6% for prevalent and 5.2% for prevalent plus exacerbated musculoskeletal pain. Multivariable adjusted odds ratios and 95% confidence intervals of lifestyle factors associated with prevalent and prevalent plus exacerbated musculoskeletal pain were as follows: shelter use (prevalent: 1.02, 0.96-1.08; exacerbated: 1.44, 1.29-1.60), job loss (prevalent: 1.03, 0.96-1.10; exacerbated: 1.30, 1.16-1.47), decreased income (prevalent: 1.13, 1.05-1.21; exacerbated: 1.29, 1.14-1.45), current heavy drinking (prevalent: 1.33, 1.21-1.47; exacerbated: 1.38, 1.14-1.68), insomnia (prevalent: 1.22, 1.15-1.29; exacerbated: 1.50, 1.36-1.65), exercising almost daily (prevalent: 0.83, 0.77-0.91; exacerbated: 0.80, 0.68-0.95), and participating in community activities often (prevalent: 0.83, 0.75-0.92; exacerbated: 0.76, 0.61-0.95).

Conclusions: Prevalent and exacerbated musculoskeletal pain were inversely associated with frequencies of exercise and participating in recreational or community activities and positively associated with decreased income, current heavy drinking, and insomnia. The use of evacuation shelters or temporary housing/job loss was positively associated only with exacerbated musculoskeletal pain. These results suggest that post-disaster lifestyle factors are potentially associated with musculoskeletal pain. For better post-disaster pain management, further studies are needed to confirm these results in other disasters and highlight underlying mechanisms.

Background

Disasters cause dramatic impacts on our life; the long-lasting impacts gradually affect people's health. Musculoskeletal pain affects daily activities and quality of life [1, 2]; however, there are insufficient post-disaster epidemiological pain studies. Most previous post-disaster studies have mainly focused on the prevalence of pain. A retrospective study of 958 evacuees conducted about 1 month after the 2009

earthquake in L'Aquila, Italy, showed that 35% of the evacuees had some pain, and, unlike acute pain, chronic pain increased over the 0.5 to 1 month following the disaster [3]. A cross-sectional study of 71 evacuees from the Great East Japan Earthquake conducted about 18 months (1.5 years) after the disaster showed that 62% of evacuees experienced chronic pain in some body part, the most frequent ones being the lower back and limbs [4]. From these findings, it is evident that more than half the evacuees experience musculoskeletal pain after a disaster. Other previous studies, while not conducted in post-disaster situations, have reported that musculoskeletal pain and its prevalence, persistence, and/or exacerbation are associated with lifestyle factors such as physical activity [5–7], drinking status [8], and insomnia [9, 10]. Depressive symptoms have been reported as well [11]; such negative symptoms are involved in persistent and/or exacerbated pain through central neural mechanisms [12].

The Great East Japan Earthquake of 2011 forced approximately 160,000 people to leave their hometowns. The evacuees experienced not only the earthquake itself but also a tsunami and/or the indirect effects of a nuclear power plant accident. These disasters and experiences affected their living environment, relatives and friends, and socioeconomic status [13, 14]. There were also some changes seen between the pre- and post-disaster phases in lifestyle status among evacuees; for example, smoking, drinking, and physical activities [15] as well as psychological distress [16]. However, the relationships between post-disaster lifestyle and musculoskeletal pain, as well as proportions of prevalent, persistent, and/or exacerbated pain in mid-to-long term after the disaster, are not well known. It is therefore important that we identify relationships between lifestyle factors and musculoskeletal pain to achieve better health management to prevent these issues in the mid-to-long term period after the disaster.

The purpose of this study was to determine the prevalence of musculoskeletal pain among the evacuees, and to investigate the association of lifestyle factors with the prevalence and exacerbation of musculoskeletal pain after the Great East Japan Earthquake. We hypothesised that disaster-related lifestyle factors such as shelter use at any point of time may be related to musculoskeletal pain after a disaster. We also hypothesised that disaster-related psychological stress may increase musculoskeletal pain.

Methods

Subjects

This cross-sectional study included 55,727 evacuees, aged 40–89 years (mean age, 61 years), who experienced the Great East Japan Earthquake, and who had taken the Mental Health and Lifestyle Survey as a part of the Fukushima Health Management Survey. Detailed information about the survey is described elsewhere [17, 18]. Briefly, the Mental health and lifestyle Survey is one part of the Fukushima Health Management Survey. The target population were all the residents who lived in the evacuation zones, which is a government-designated area (20 km radius) around the nuclear power plant. The earthquake occurred on 11 March 2011, and the study period was from 18 January 2012 to 31 October

2012. We sent a self-administered questionnaire on various lifestyle and mental health factors to all the residents. The overall response rate of the Mental health and lifestyle Survey was 40.7% [14].

We defined an evacuee as a resident in evacuation areas who reported to have experienced the earthquake, including those who had experienced the tsunami and/or the indirect effects of the nuclear power plant accident. The exclusion criteria were as follows: not having experienced the earthquake (n=2,026); having any major disabilities in activities of daily living such as eating, dressing, going to the toilet, or shopping (n=3,351). We imposed this latter restriction because individuals with disabilities may represent a distinct lifestyle compared to individuals without disabilities, which may lead to misclassification due to their avoiding activity-related pain. We also excluded missing information or abnormal values for basic characteristics and medical history (n=6,256), musculoskeletal pain and lifestyle-related information (n=7,052), and psychological factors (n=2,113). Finally, we used 34,919 people (female, n=18,156; male, n=16,763) without major disabilities at about 1.0 ± 0.1 years (average $\pm$ standard deviation) after the disaster. The Ethics Committee of Fukushima Medical University approved this survey protocol. The participants were determined to have provided their consent to participate by returning the survey questionnaires.

Musculoskeletal pain

We asked all participants to answer the multiple-choice question “In the past few days, have you experienced uncomfortable symptoms due to any diseases or injuries?”. If the participant responded “Yes” to low back pain or limb pain, we regarded that the participant had major musculoskeletal symptoms. The most common areas of musculoskeletal pains are the low back, knee, hip, ankle, foot, shoulder, elbow, wrist, and fingers (the latter eight are in the category of limb pain) [19]. We further asked whether the pain was exacerbated after the disaster. The subjects were categorised into 3 groups: absent musculoskeletal pain (absent), prevalent plus exacerbated musculoskeletal pain (exacerbated), and prevalent musculoskeletal pain (prevalent).

Lifestyle factors

We asked participants about lifestyle factors including the use of evacuation shelters or temporary housing at any point in time (shelter use: no/yes), having lost their job after the disaster (job loss: no/yes), decrease in income after the disaster (decreased income: no/yes), current smoking status (smoking status: never/past/current light smoking/current heavy smoking), current drinking status (drinking status: never/past/current light drinking/current heavy drinking), subjective lack of sleep (insomnia: no/yes), current regular exercise (regular exercise: almost none/1 time per week/2-4 times per week/almost daily), and participation in recreational or community activities (community activities: never or rarely/sometimes/often).

Psychological factors

We assessed psychological factors that may affect musculoskeletal pain such as traumatic reactions (post-traumatic stress disorder check list: 44 points or more) [20, 21], psychological distress (Kessler psychological distress scale: 13 points or more) [22, 23], and somatic symptoms (none/one/two or more subjective symptoms except for musculoskeletal pain). Their subjective symptoms were categorised into 4 groups as follows based on a study by Zijlema et al. (2013): cardiopulmonary/autonomic (palpitations, breathing trouble), gastrointestinal (constipation/diarrhoea, loss of appetite, abdominal pain), general (headaches, dizziness), and others (difficulty urinating) [24].

Other confounding factors

We also gathered information regarding potential confounding factors such as age (40-64 years/65-74 years/75-89 years); sex (male/female); history of hypertension, diabetes, and dyslipidaemia (no/yes); education status (junior high school or less/high school/colleges/university or higher); having experienced the tsunami and/or indirect effects of the nuclear power plant accident (no/yes); house damage (none/partly destroyed/half or more destroyed); and loss of relatives or friends (no/yes).

Statistical analysis

A generalised linear model was used to compare basic characteristics, psychological factors, and lifestyle factors between the groups (absent versus prevalent, absent versus exacerbated, and prevalent versus exacerbated). Because even trivial differences may be statistically significant due to our large sample, we regarded differences of $\pm 3\%$ or more between the groups in addition to p-values less than 0.05 as statistically significant to highlight the significance of our findings. Multinomial logistic regression analysis was used to calculate odds ratios and 95% confidence intervals of each lifestyle factor related to prevalent musculoskeletal pain and exacerbated musculoskeletal pain. Two multivariable adjusted models were constructed. Model 1 was the adjustment for background information (age, sex, history of hypertension/diabetes/dyslipidaemia, and educational status) and disaster experience (tsunami/indirect nuclear power plant accident, house damage, loss of close person) and each lifestyle factor. This model was used to estimate odds ratios of what were considered unmodifiable characteristics. Model 2 was the further adjustment for psychological factors (traumatic reactions and psychological distress) and the number of uncomfortable symptoms (none, one, and two or more) to the model 1. This model aimed to estimate the non-psychometric effects (e.g. mechanical stress) from the psychometric effects.

Results

Prevalence of musculoskeletal pain

The proportions of participants with prevalent and exacerbated musculoskeletal pain were 27.6% and 5.2%, respectively (Table 1). The proportion of participants with both low back pain and limb pain was greater in evacuees with exacerbated musculoskeletal pain than in those with prevalent musculoskeletal pain. According to additional stratified analyses, the prevalence of lower back pain was 25.8% and that of

limb pain was 17.2%. We also found that a greater prevalence was observed in women than in men and that prevalence increased with age, except for exacerbated low back pain (see Supplemental table 1).

	Musculoskeletal pain		
	Absent	Prevalent	Exacerbated
N	23456	9650	1813
Age			
40 to 64 years	15210 (64.8)	5713 (59.2)*	1077 (59.4)* [†]
65 to 74 years	5165 (22.0)	2266 (23.5)	440 (24.3)
75 to 89 years	3081 (13.1)	1671 (17.3)*	296 (16.3)*
Sex			
Male	11690 (49.8)	4375 (45.3)*	698 (38.5)* [†]
Female	11766 (50.2)	5275 (54.7)*	1115 (61.5)* [†]
History of disease			
Hypertension	10358 (44.2)	5146 (53.3)*	988 (54.5)*
Diabetes	4551 (19.4)	2325 (24.1)*	453 (25.0)*
Dyslipidemia	8734 (37.2)	4486 (46.5)*	890 (49.1)*
Educational status			
Junior high school or less	5823 (24.8)	2853 (29.6)*	455 (25.1) [†]
High school	12079 (51.5)	4773 (49.5)	917 (50.6)
Colleges/university or higher	5554 (23.7)	2024 (21.0)	441 (24.3) [†]
Type of disaster experience			
Tsunami	4603 (19.6)	2294 (23.8)*	473 (26.1)*
Indirect nuclear power plant accident	11754 (50.1)	5743 (59.5)*	1198 (66.1)* [†]
House damage			
None	6296 (26.8)	2078 (21.5)*	311 (17.2)* [†]
Partly destroyed	13811 (58.9)	5852 (60.6)	1109 (61.2)
Half or more destroyed	3349 (14.3)	1720 (17.8)*	393 (21.7)*

Loss of close person			
No	19298 (82.3)	7362 (76.3)*	1286 (70.9)*†
Yes	4158 (17.7)	2288 (23.7)*	527 (29.1)*†
Pain region			
Low back pain only	-	4779 (49.5)	685 (37.8)†
Limb pain only	-	2124 (22.0)	319 (17.6)†
Both	-	2747 (28.5)	809 (44.6)†
Percentages are shown in parentheses. Statistically significant and observed $\pm 3\%$ or more difference compared to the Absent group (*), and to the Prevalent group (†); The total percentage does not always add up to 100.			

Table 1

Prevalence of musculoskeletal pain and basic background information of 34 919 evacuees from the earthquake

Frequency analyses of lifestyle factors

Compared to the absence of pain, participants with prevalent or exacerbated musculoskeletal pain showed significant greater proportions of shelter use, job loss, decreased income, never smoking, never drinking, insomnia, almost no regular exercise, and no or rare participation in community activities (Table 2). Proportions of current light drinking, regular exercising almost daily, and occasional or frequent participation in community activities were significantly lower among participants with prevalent or exacerbated musculoskeletal pain (Table 2).

Table 2
Associated lifestyle factors with prevalent and exacerbated musculoskeletal pain

	Musculoskeletal pain		
	Absent	Prevalent	Exacerbated
Shelter use			
No	15931 (67.9)	6012 (62.3)*	923 (50.9) [†]
Yes	7525 (32.1)	3638 (37.7)*	890 (49.1) [†]
Job loss			
No	19057 (81.2)	7438 (77.1)*	1257 (69.3) [†]
Yes	4399 (18.8)	2212 (22.9)*	556 (30.7) [†]
Decreased income			
No	19106 (81.5)	7600 (78.8)	1374 (75.8) [†]
Yes	4350 (18.5)	2050 (21.2)	439 (24.2) [†]
Smoking status			
Never	12862 (54.8)	5380 (55.8)	1086 (59.9) [†]
Past	5848 (24.9)	2438 (25.3)	408 (22.5)
Current light smoking	2374 (10.1)	901 (9.3)	147 (8.1)
Current heavy smoking	2372 (10.1)	931 (9.6)	172 (9.5)
Drinking status			
Never	11075 (47.2)	4664 (48.3)	916 (50.5)*
Past	727 (3.1)	367 (3.8)	71 (3.9)
Current light drinking	8976 (38.3)	3429 (35.5)	622 (34.3)*
Current heavy drinking	2678 (11.4)	1190 (12.3)	204 (11.3)
Insomnia			
No	15728 (67.1)	4677 (48.5)*	645 (35.6) [†]
Yes	7728 (32.9)	4973 (51.5)*	1168 (64.4) [†]
Regular exercise			

	Musculoskeletal pain		
	Absent	Prevalent	Exacerbated
Almost not	11071 (47.2)	4783 (49.6)*	915 (50.5)*
1 time per week	3373 (14.4)	1387 (14.4)	240 (13.2)
2–4 times per week	4994 (21.3)	2117 (21.9)	422 (23.3)
Almost daily	4018 (17.1)	1363 (14.1)*	236 (13.0)*
Community activities			
Never or rarely	13894 (59.2)	6427 (66.6)*	1285 (70.9)*†
Sometimes	7118 (30.3)	2548 (26.4)*	429 (23.7)*
Often	2444 (10.4)	675 (7.0)*	99 (5.5)*
Percentages are shown in parentheses. Statistically significant and observed $\pm$ 3% or more difference compared to the Absent group (*), and to the Prevalent group (†); The total percentage does not always add up to 100.			

Frequency analyses of psychological factors

Participants with prevalent musculoskeletal pain and exacerbated musculoskeletal pain showed significantly greater proportions of traumatic reaction, psychological distress, and two or more of uncomfortable symptoms compared to participants without any pain (Table 3). These point-estimated odds ratios of psychological factors are shown in the Supplemental table 2.

Table 3
Associated psychological factors with prevalent and exacerbated musculoskeletal pain

	Musculoskeletal pain		
	Absent	Prevalent	Exacerbated
Traumatic reaction (PCL)			
Normal	20196 (86.1)	6643 (68.8)*	1000 (55.2)**†
44 points or over	3260 (13.9)	3007 (31.2)*	813 (44.8)**†
Psychological distress (K6)			
Normal	21345 (91.0)	7561 (88.4)*	1223 (67.5)**†
13 points or over	2111 (9.0)	2089 (21.6)*	590 (32.5)**†
Uncomfortable symptoms			
Diarrhea/constipation	2221 (9.5)	2881 (29.9)*	689 (38.0)**†
Headache	2078 (8.9)	3060 (31.7)*	706 (38.9)**†
Dizziness	1329 (5.7)	1950 (20.2)*	448 (24.7)**†
Shortness of breath breathing trouble	1034 (4.4)	1792 (18.6)*	407 (22.4)**†
Palpitations	1043 (4.4)	1656 (17.2)*	416 (22.9)**†
Poor appetite	476 (2.0)	793 (8.2)*	234 (12.9)**†
Pain during urination/ difficulty urinating	281 (1.2)	515 (5.3)*	140 (7.7)*
Number of uncomfortable symptoms			
None	17985 (76.7)	3149 (32.6)*	439 (24.2)**†
One	2889 (12.3)	2599 (26.9)*	471 (26.0)**†
Two or more	2582 (11.0)	3902 (40.4)*	903 (49.8)**†
Percentages are shown in parentheses. Statistically significant and observed $\pm$ 3% or more difference compared to the Absent group (*), and to the Prevalent group (†); The total percentage does not always add up to 100; PCL, post-traumatic stress disorder check list; K6, Kessler psychological distress scale.			

Multivariable adjusted analyses of lifestyle factors

Multinomial logistic regression analysis was used to calculate multivariate adjusted odds ratios and 95% confidence intervals. After adjusting their background information, disaster experience, psychological factors, the number of uncomfortable symptoms, and each lifestyle factor, the lifestyle factors that were significantly associated with prevalent musculoskeletal pain and exacerbated musculoskeletal pain were decreased income (prevalent: 1.13, 1.05–1.21; exacerbated: 1.29, 1.14–1.45), current heavy drinking (prevalent: 1.33, 1.21–1.47; exacerbated: 1.38, 1.14–1.68), insomnia (prevalent: 1.22, 1.15–1.29; exacerbated: 1.50, 1.36–1.65), exercising almost daily (prevalent: 0.83, 0.77–0.91; exacerbated: 0.80, 0.68–0.95), and participating in community activities often (prevalent: 0.83, 0.75–0.92; exacerbated: 0.76, 0.61–0.95) (Table 4). Shelter use (prevalent: 1.02, 0.96–1.08; exacerbated: 1.44, 1.29–1.60) and job loss (prevalent: 1.03, 0.96–1.10; exacerbated: 1.30, 1.16–1.47) were significantly associated only with exacerbated musculoskeletal pain. Adjustment for psychological factors appeared to weaken these associations with prevalent musculoskeletal pain for shelter use, job loss, and past/light smoking; associations were present with prevalent and exacerbated musculoskeletal pains in current heavy smoking, past drinking, and those who regularly exercised 1 or more times per week. Sub-analyses of pain region, sex, and age groups are shown in the Supplemental tables 3–5.

Table 4

Multinomial odds ratios of associated lifestyle factors with prevalent and exacerbated musculoskeletal pain

	Crude		Multivariable adjusted odds ratios ¹		Multivariable adjusted odds ratios ²	
	Prevalent	Exacerbated	Prevalent	Exacerbated	Prevalent	Exacerbated
Shelter use						
Yes	1.28, 1.22–1.35	2.04, 1.85–2.25	1.11, 1.05–1.17	1.57, 1.42–1.74	1.02, 0.96–1.08	1.44, 1.29–1.60
Job loss						
Yes	1.28, 1.22–1.37	1.92, 1.73–2.13	1.19, 1.12–1/27	1.57, 1.40–1.76	1.03, 0.96–1.10	1.30, 1.16–1.47
Decreased income						
Yes	1.19, 1.12–1.26	1.40, 1.25–1.57	1.18, 1.11–1.26	1.36, 1.21–1.54	1.13, 1.05–1.21	1.29, 1.14–1.45
Smoking status						
Past	1.00, 0.94–1.06	0.83, 0.73–0.93	1.15, 1.07–1.24	1.14, 0.98–1.33	1.04, 0.96–1.12	1.02, 0.87–1.19
Current light smoking	0.91, 0.84–0.99	0.73, 0.61–0.88	1.10, 1.00–1.21	0.99, 0.82–1.20	1.00, 0.91–1.11	0.85, 0.70–1.04
Current heavy smoking	0.94, 0.86–1.02	0.86, 0.73–1.02	1.15, 1.04–1.27	1.25, 1.02–1.52	1.06, 0.95–1.18	1.11, 0.90–1.36
Drinking status						
Past	1.19, 1.04–1.35	1.20, 0.93–1.54	1.21, 1.05–1.39	1.37, 1.04–1.37	0.98, 0.84–1.14	1.05, 0.79–1.39
Current light drinking	0.90, 0.85–0.94	0.85, 0.77–0.94	1.05, 0.99–1.11	1.08, 0.96–1.22	1.08, 1.01–1.15	1.11, 0.98–1.25

	Crude		Multivariable adjusted odds ratios ¹		Multivariable adjusted odds ratios ²	
	Prevalent	Exacerbated	Prevalent	Exacerbated	Prevalent	Exacerbated
Current heavy drinking	0.95, 0.88–1.03	0.94, 0.80–1.11	1.26, 1.15–1.38	1.29, 1.07–1.56	1.33, 1.21–1.47	1.38, 1.14–1.68
Insomnia						
Yes	2.16, 2.05–2.26	3.22, 2.94–3.53	2.10, 2.00–2.21	2.96, 2.70–3.25	1.22, 1.15–1.29	1.50, 1.36–1.65
Regular exercise						
1 time per week	0.95, 0.89–1.02	0.86, 0.74–0.99	0.91, 0.85–0.98	0.83, 0.71–0.97	0.94, 0.87–1.02	0.86, 0.73–1.00
2–4 times per week	0.98, 0.92–1.04	1.02, 0.91–1.15	0.89, 0.83–0.95	0.92, 0.81–1.05	0.92, 0.86–0.99	0.96, 0.84–1.10
Almost daily	0.79, 0.73–0.84	0.71, 0.61–0.82	0.74, 0.69–0.80	0.71, 0.60–0.83	0.83, 0.77–0.91	0.80, 0.68–0.95
Community activities						
Sometimes	0.77, 0.73–0.82	0.65, 0.58–0.73	0.81, 0.77–0.86	0.72, 0.64–0.81	0.93, 0.88–0.99	0.86, 0.76–0.98
Often	0.60, 0.55–0.65	0.44, 0.36–0.54	0.66, 0.60–0.72	0.54, 0.44–0.67	0.83, 0.75–0.92	0.76, 0.61–0.95
<p>Odds ratios and 95% confidence intervals were shown. Bolded numbers mean significant association ($p < 0.05$). For each factor, the following categories were used as a reference category: ‘no’ for shelter use, job loss, decreased income, and insomnia; ‘never’ for smoking and drinking status; ‘almost not’ for regular exercise; and ‘never or rarely’ for community activities.</p> <p>¹Adjustment for background information (age, sex, history of hypertension/diabetes/dyslipidemia, and educational status) and disaster experience (tsunami/indirect nuclear power plant accident, house damage, loss of close person) and each lifestyle factors; ²Further adjustment for psychological factors (traumatic reaction and psychological distress) and the number of uncomfortable symptoms (none, one, and two or more).</p>						

Discussion

In this 1-year post-disaster population-based cross-sectional study, the frequency of exercise and participating in recreational or community activities showed inverse associations with both prevalent and exacerbated musculoskeletal pain. In contrast, decreased income, current heavy drinking, and insomnia showed positive associations. Use of evacuation shelters or temporary housing and job loss showed positive association only with exacerbated musculoskeletal pain. These significant associations were observed even after adjustment for psychological factors and each lifestyle factor. Nevertheless, adjustment for psychological factors appeared to weaken most of those associations.

A previous study showed that the prevalence of musculoskeletal pain among evacuees at 1.5 years after the Great East Japan Earthquake disaster was 22% for low back pain and 24% for limb pain [4]. The prevalence of low back pain in our results was similar to that in the previous study, while the prevalence of limb pain seemed to be lower. This discrepancy could be due to differences in population age and the proportion of women to men; our population had a mean age of 61 years and the proportion of women was 52.6%, whereas the population of the previous study had a mean age of 75 years and the proportion of women was 77.5%. Musculoskeletal pain, particularly knee pain, generally increases with age and is more frequently observed in women [25]. This may explain the abovementioned differences.

The frequency of exercise and participation in recreational or community activities showed inverse associations with musculoskeletal pain; several previous studies have supported these associations. Light to moderate intensity of physical activities, possibly mediated by frequency of exercise and community activities, may reduce the risk of chronic musculoskeletal pain [5–7]. We measured physical and community activities, especially those focused on during leisure time. We also hypothesised that the physical and community activities during non-leisure time differed from those in pre-disaster conditions. Shelter use, job loss (or decreased income) was likely rare in pre-disaster conditions; as our results showed, these factors were associated with musculoskeletal pain in post-disaster conditions. According to another 1-year post-disaster cross-sectional study of 4,316 elderly people, the factors of displacement from home, non-working status, and lack of social networking (Lubben's social network scale, 11 or lower) were significantly associated with low physical activity [26]. This partially supports our assumption that shelter use and job loss are associated with musculoskeletal pain via low physical activity.

We also need to discuss the association of insomnia with musculoskeletal pain. According to an annual longitudinal population-based study of 1,746 individuals, musculoskeletal pain may increase the risk for the incidence and persistence of insomnia [9, 10], and insomnia may increase the risk for persistent musculoskeletal pain [10]. Regarding the effects of the direction of musculoskeletal pain increasing the risk of insomnia, path analysis revealed that physical limitation and reduced social participation could explain 68% of the effects [27]. According to a post-disaster cross-sectional study of 4,176 individuals who remained in their homes, significant inverse associations with insomnia were observed in social and community networks such as pleasure in life as a consequence of engaging in social/community networks and interactions with neighbours [28].

Association of high alcohol consumption with prevalent and exacerbated musculoskeletal pain may have been influenced by various factors. For example, one possible reason for the associations observed with alcohol consumption is that people might be injured in a fall due to impaired motor skills, which is likely to occur when blood-alcohol concentration is 0.3 mg/ml or higher [29]. Other possible reasons are chronic alcohol myopathy [30] and gout [31]. Alternatively, we should also consider that some evacuees may use alcohol for pain relief [8]. Alcohol may affect the central nervous system and temporally reduce the intensity of the pain [8].

Throughout the above results, adjustment for psychological factors weakened most of the point-estimated odds ratios (except for current heavy drinking). The changes in psychological stress may influence the prevalent and exacerbated musculoskeletal pain [11, 32]. For example, greater psychological distress was observed among individuals who lived in temporary housing for longer periods [33]. Shelter use tends to create psychological stress due to the change in the available space, neighbours' noise, and lack of a future housing plan [33]. These results suggest that post-disaster lifestyle factors are potentially associated with musculoskeletal pain via both mechanical stress and psychological stress.

The present study had several strengths. First, to the best of our knowledge, this appears to be one of the first studies to use such a large sample size to investigate post-disaster pain epidemiology. Second, systematic recruitment and investigation were conducted after the disaster. We should also discuss the limitations of the present study. First, this cross-sectional study was conducted at one time point (approximately 1 year after the disaster) and different social and living environment changes may have occurred until that period. Thus, these findings could not be generalised to any time point after a disaster. Moreover, it is unclear whether some lifestyle factors, such as regular exercise and community activity, had continued from pre-disaster conditions. In addition, due to the nature of the cross-sectional design, the findings do not allow us to draw conclusions on the time sequence for these lifestyle factors and the musculoskeletal pain. Second, our definition of musculoskeletal pain could not assess cervical and thoracic pain. It also could not discriminate among shoulder, knee, or foot pain. Pain severity (pain intensity or disability), types (nociceptive or neuropathic pain), and related pain experiences in post-disaster conditions were unclear. Third, the low response rate of the overall survey may have influenced the results. The response rate could be different depending on the exposure status of lifestyle factors; according to this, some estimated values in our results could be greater or smaller than the true values in the whole population. For example, if the response rate was higher for those who answered "yes" to participating in recreational or community activities and lower for those with musculoskeletal pain, a stronger inverse association might have been observed between them. However, we could not determine the differences between these response rates. Fourth, we did not gather information about medication use or access to hospitals. There might have been differences between participants in their medical environments. Access to pain medication and/or accessibility to a hospital after the disaster may have differed between participants and, thus, influenced our results.

Conclusion

In this cross-sectional study after a disaster, prevalent and exacerbated musculoskeletal pain were inversely associated with frequencies of exercise and participating in recreational or community activities and were positively associated with decreased income, current heavy drinking, and insomnia. Besides, the use of evacuation shelters or temporary housing and job loss were positively associated only with exacerbated musculoskeletal pain. These results suggest that post-disaster lifestyle factors are potentially associated with musculoskeletal pain. To achieve better post-disaster pain management, further studies are needed to confirm the consistency of these results in other disasters and to highlight the underlying causative mechanisms.

List Of Abbreviations

Not applicable.

Declarations

Ethics approval and consent to participate

Informed consent was obtained from all participants, and the study was approved by the Ethics Committee of the Fukushima Medical University (#1316) and Nippon Medical School (approval code, 29-18).

Consent for publication

Not applicable.

Availability of data and material

No additional data available.

Competing interests

The authors declare that there are no competing interests.

Funding

This survey was supported in part by the national "Health Fund for Children and Adults Affected by the Nuclear Incident".

Authors' contributions

Concept/idea/research design: HJ, TO, and HI

Writing: HJ, TO, HK, HI, KM, TK, and SY

Project management: TO, MM, HY, YS, MH, SY, and KK

Consultation (including review of manuscript before submitting): all authors

Acknowledgements

Medical Center for the Fukushima Health Management Survey; Shunichi Yamashita (Founding Senior Director, Radiation Medical Center for the Fukushima Health Management Survey); Kenji Kamiya (Senior Director, Radiation Medical Center for the Fukushima Health Management Survey); and Koichi Tanigawa (Senior Director, Radiation Medical Center for the Fukushima Health Management Survey).

Authors' information

The findings and conclusions of this article are solely the responsibility of the authors and do not represent the official views of Fukushima Prefecture government.

References

1. Blyth FM, Noguchi N. Chronic Musculoskeletal Pain and its Impact on Older People. *Best Pract Res Clin Rheumatol*. 2017; 31(2):160-8.
2. Shah RC, Buchman AS, Boyle PA, Leurgans SE, Wilson RS, Andersson GB, Bennett DA. Musculoskeletal Pain is Associated with Incident Mobility Disability in Community-Dwelling Elders. *J Gerontol A Biol Sci Med Sci*. 2011; 66(1): 82-8.
3. Angeletti C, Guetti C, Papola R, Petrucci E, Ursini ML, Ciccozzi A, Masi F, Russo MR, Squarcione S, Paladini A, Pergolizzi J, Taylor R Jr, Varrassi G, Marinangeli F. Pain After Earthquake. *Scand J Trauma Resusc Emerg Med*. 2012; 29(20): 43.
4. Yabuki S, Ouchi K, Kikuchi S, Konno S. Pain, Quality of Life and Activity in Aged Evacuees Living in Temporary Housing After the Great East Japan Earthquake of 11 March 2011: A Cross-Sectional Study in Minamisoma City, Fukushima Prefecture. *BMC Musculoskelet Disord*. 2015; 10(16): 246.
5. Shiri R, Falah-Hassani K. Does Leisure Time Physical Activity Protect Against Low Back Pain? Systematic Review and Meta-Analysis of 36 Prospective Cohort Studies. *Br J Sports Med*. 2017; 51(19): 1410-8.
6. Song J, Chang AH, Chang RW, Lee J, Pinto D, Hawker G, Nevitt M, Dunlop DD. Relationship of Knee Pain to Time in Moderate and Light Physical Activities: Data from Osteoarthritis Initiative. *Semin Arthritis Rheum*. 2018; 47(5): 683-8.
7. Landmark T, Romundstad P, Borchgrevink PC, Kaasa S, Dale O. Associations Between Recreational Exercise and Chronic Pain in the General Population: Evidence from the HUNT 3 Study. *Pain*. 2011; 152(10): 2241-7.
8. Riley JL 3rd, King C. Self-Report of Alcohol Use for Pain in a Multi-Ethnic Community Sample. *J Pain*. 2009; 10(9): 944-52.
9. Tang NK, McBeth J, Jordan KP, Blagojevic-Bucknall M, Croft P, Wilkie R. Impact of Musculoskeletal Pain on Insomnia Onset: A Prospective Cohort Study. *Rheumatology (Oxford)*. 2015; 54(2): 248-56.

10. Jansson-Fröjmark M, Boersma K. Bidirectionality Between Pain and Insomnia Symptoms: A Prospective Study. *Br J Health Psychol*. 2012; 17(2): 420-31.
11. Urquhart DM, Bell RJ, Cicuttini FM, Cui J, Forbes A, Davis SR. Negative Beliefs About Low Back Pain Are Associated with High Pain Intensity and High Level Disability in Community-Based Women. *BMC Musculoskelet Disord*. 2008; 9(1): 148.
12. Abdallah CG, Geha P. Chronic Pain and Chronic Stress: Two Sides of the Same Coin. *Chronic Stress* (Thousand Oaks). 2017; 1.
13. Zhang W, Ohira T, Yasumura S, Maeda M, Otsuru A, Harigane M, Horikoshi N, Suzuki Y, Yabe H, Nagai M, Nakano H, Hiroaki M, Uemura M, Takahashi H, Kamiya K, Yamashita S, Abe M. Effects of Socioeconomic Factors on Cardiovascular-Related Symptoms Among Residents in Fukushima After the Great East Japan Earthquake: A Cross-Sectional Study Using Data from the Fukushima Health Management Survey. *BMJ Open*. 2017; 7(6): e014077.
14. Nagai M, Ohira T, Zhang W, Nakano H, Maeda M, Yasumura S, Abe M. Lifestyle-Related Factors that Explain Disaster-Induced Changes in Socioeconomic Status and Poor Subjective Health: A Cross-Sectional Study from the Fukushima Health Management Survey. *BMC Public Health*. 2017; 17(1): 340.
15. Ohira T, Hosoya M, Yasumura S, Satoh H, Suzuki H, Sakai A, Ohtsuru A, Kawasaki Y, Takahashi A, Ozasa K, Kobashi G, Hashimoto S, Kamiya K, Yamashita S, Abe M. Evacuation and Risk of Hypertension After the Great East Japan Earthquake: The Fukushima Health Management Survey. *Hypertension*. 2016; 68(3): 558-64.
16. Yabe H, Suzuki Y, Mashiko H, Nakayama Y, Hisata M, Niwa S, Yasumura S, Yamashita S, Kamiya K, Abe M. Psychological Distress After the Great East Japan Earthquake and Fukushima Daiichi Nuclear Power Plant Accident: Results of a Mental Health and Lifestyle Survey Through the Fukushima Health Management Survey in FY2011 and FY2012. *Fukushima J Med Sci*. 2014; 60(1): 57-67.
17. Yasumura S, Hosoya M, Yamashita S, Kamiya K, Abe M, Akashi M, Kodama K, Ozasa K. Study Protocol for the Fukushima Health Management Survey. *J Epidemiol*. 2012; 22(5): 375-83.
18. Ohira T, Hosoya M, Yasumura S, Satoh H, Suzuki H, Sakai A, Ohtsuru A, Kawasaki Y, Takahashi A, Ozasa K, Kobashi G, Kamiya K, Yamashita S, Abe M. Effect of Evacuation on Body Weight After the Great East Japan Earthquake. *Am J Prev Med*. 2016; 50(5): 553-60.
19. Cunningham LS, Kelsey JL. Epidemiology of Musculoskeletal Impairments and Associated Disability. *Am J Public Health*. 1984; 74(6): 574-9.
20. Weathers FW, Litz BT, Herman DS, Huska JA, Keane TM. The PTSD Checklist (PCL): Reliability, Validity, and Diagnostic Utility. Weathers FW, Litz BT, Herman DS, Hushka JA, Keane TM. The PTSD Checklist (PCL), Reliability, Validity, and Diagnostic Utility. The Annual Meeting of International Society for Traumatic Stress Studies. San Antonio, TX. 1993.
21. Suzuki Y, Yabe H, Horikoshi N, Yasumura S, Kawakami N, Ohtsuru A, Mashiko H, Maeda M. Diagnostic Accuracy of Japanese Posttraumatic Stress Measures After a Complex Disaster: The

- Fukushima Health Management Survey. *Asia Pac Psychiatry*. 2017; 9(1): e12248.
22. Kessler RC, Barker PR, Colpe LJ, Epstein JF, Gfroerer JC, Hiripi E, Howes MJ, Normand SL, Manderscheid RW, Walters EE, Zaslavsky AM. Screening for Serious Mental Illness in the General Population. *Arch Gen Psychiatry*. 2003; 60(2): 184-9.
 23. Furukawa TA, Kawakami N, Saitoh M, Ono Y, Nakane Y, Nakamura Y, Tachimori H, Iwata N, Uda H, Nakane H, Watanabe M, Naganuma Y, Hata Y, Kobayashi M, Miyake Y, Takeshima T, Kikkawa T. The Performance of the Japanese Version of the K6 and K10 in the World Mental Health Survey Japan. *Int J Methods Psychiatr Res*. 2008; 17(3): 152-8.
 24. Zijlema WL, Stolk RPL, Löwe B, Rief W, BioSHaRE., White PD, Rosmalen JG. How to Assess Common Somatic Symptoms in Large-Scale Studies: A Systematic Review of Questionnaires. *J Psychosom Res*. 2013; 74(6): 459-68.
 25. Takahashi A, Kitamura K, Watanabe Y, Kobayashi R, Saito T, Takachi R, Kabasawa K, Oshiki R, Tsugane S, Iki M, Sasaki A, Yamazaki O, Nakamura K. *J Pain Res*. 2018; 11: 3161-9.
 26. Yoshimura E, Ishikawa-Takata K, Murakami H, Tsuboyama-Kasaoka N, Tsubota-Utsugi M, Miyachi M, Yokoyama Y, Sakata K, Kobayashi S, Ogawa A, Nishi N. Relationships Between Social Factors and Physical Activity Among Elderly Survivors of the Great East Japan Earthquake: A Cross-Sectional Study. *BMC Geriatr*. 2016; 16: 30.
 27. Tang NK, McBeth J, Jordan KP, Blagojevic-Bucknall M, Croft P, Wilkie R. Impact of Musculoskeletal Pain on Insomnia Onset: A Prospective Cohort Study. *Rheumatology (Oxford)*. 2015; 54(2): 248-56.
 28. Matsumoto S, Yamaoka K, Inoue M, Muto S. Social Ties May Play a Critical Role in Mitigating Sleep Difficulties in Disaster-Affected Communities: A Cross-Sectional Study in the Ishinomaki Area, Japan. *Sleep*. 2014; 37(1): 137-45.
 29. Johnston JJ, McGovern SJ. Alcohol Related Falls: An Interesting Pattern of Injuries. *Emerg Med J*. 2004; 21(2): 185-8.
 30. Simon L, Jolley SE, Molina PE. Alcoholic Myopathy: Pathophysiologic Mechanisms and Clinical Implications. *Alcohol Res*. 2017; 38(2): 207-17.
 31. Tu HP, Tung YC, Tsai WC, Lin GT, Ko YC, Lee SS. Alcohol-Related Diseases and Alcohol Dependence Syndrome is Associated with Increased Gout Risk: A Nationwide Population-Based Cohort Study. *Joint Bone Spine*. 2017; 84(2): 189-96.
 32. Ng SK, Cicuttini FM, Wang Y, Wluka AE, Fitzgibbon B, Urquhart DM. Negative Beliefs About Low Back Pain Are Associated with Persistent High Intensity Low Back Pain. *Psychol Health Med*. 2017; 22(7): 790-9.
 33. Tanji F, Tomata Y, Sekiguchi T, Tsuji I. Period of Residence in Prefabricated Temporary Housing and Psychological Distress After the Great East Japan Earthquake: A Longitudinal Study. *BMJ Open*. 2018; 8(5): e018211.

Figures

a. Flow diagram

b. Study period

Figure 1

Diagram of the present study. Flow diagram of participant enrollment, and selection.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [MusculoskeletalPainXtheGreatEastJapanEarthquakevrev2suppl.docx](#)