The Association Between Helicobacter Pylori Infection and Erosive Gastroesophageal Reflux Disease; a Cross-Sectional Study

Ramin Niknam

Shiraz University of Medical Sciences

Kamran Bagheri Lankarani

Shiraz University of Medical Sciences

Mohsen Moghadami

Shiraz University of Medical Sciences

Seyed Alireza Taghavi

Shiraz University of Medical Sciences

Leila Zahiri

Shiraz University of Medical Sciences

Mohammad Javad Fallahi (✉ fallahimj@sums.ac.ir)

Shiraz University of Medical Sciences

Research Article

Keywords: Helicobacter pylori, Gastroesophageal reflux disease, GERD, Esophagus, Endoscopic findings

Posted Date: December 17th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1124716/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at BMC Infectious Diseases on March 19th, 2022. See the published version at <https://doi.org/10.1186/s12879-022-07278-6>.

Abstract

Background

The association between *H. pylori* (*Helicobacter pylori*) infection and gastroesophageal reflux disease (GERD) is a complex and confusing subject, so we designed this study.

Method

In a cross-sectional study, all patients referred for endoscopy due to dyspepsia were enrolled. The diagnosis of erosive GERD was made by endoscopy. Patients with normal esophagus were selected as comparison group. Random gastric biopsies were taken from all participants to diagnose *H. pylori* infection.

Result

In total, 1916 patients were included in this study, of whom 45.6% had GERD. The mean age (SD) was 42.95 (16.32). Overall, 1442 (75.3%) patients were positive for *H. pylori* infection. The frequency of *H. pylori* infection in mild GERD patients was higher than the severe GERD, but this difference was not significant ($P=0.214$). Except for sociodemographic status ($P < 0.001$), other variables including gender, age, ethnicity, body mass index (BMI), smoking, and presence of hiatus hernia in patients had no significant association with the frequency of *H. pylori* infection. According to Robust Poisson regression models analysis, the association of *H. pylori* (PR 1.026; 95%CI 0.990-1.064; $P=0.158$) and sociodemographic status were not significantly different between the two groups. But smoking, increased BMI, older age, presence of hiatus hernia, and peptic ulcer diseases were significantly associated with GERD compared with the non-GERD group.

Conclusion

In our results, there was no association between *H. pylori* infection and erosive GERD. Further studies are recommended, especially considering the effects of pathophysiological and other confounding factors on *H. pylori* and GERD.

Background

Gastroesophageal reflux disease (GERD), as a common gastrointestinal (GI) disorder, refers to symptoms or tissue damage caused by retrograde movement of the stomach contents to the esophagus. One of the most common complications of GERD is esophageal inflammation. Factors that may be contributing to the disease include lower esophageal sphincter dysfunction, increased numbers of transient lower esophageal sphincter relaxations, hiatus hernia, delayed gastric emptying, ineffective esophageal clearance, and the presence of an acid pocket (1–3).

Helicobacter pylori (*H. pylori*), as a gram-negative bacterium, plays an important role in the pathogenesis of different GI diseases including gastric ulcer, gastric mucosal lymphoma and gastric cancer (4). However, the relationship between *H. pylori* and GERD is a complex and confusing subject that needs further investigation (5, 6).

Some studies have reported a higher prevalence of *H. pylori* in patients with GERD, while some have observed an inverse relationship. This relationship is difficult to justify because GERD is a disease that is affected by different risk factors include BMI, smoking, lifestyle habits, host factors and more(1–3).

To date, the true relationship of *H. pylori* to GERD is still unclear (4), and recent studies have shown that more research is needed to clarify this association, with a more focus on confounding factors in GERD and *H. pylori* (5–7), so we designed this study.

Methods

Population and study design

In this cross-sectional study, we evaluated the frequency of *H. pylori* infection in cases with endoscopic diagnosis of erosive GERD from March 2013 to November 2020 in Fars province, southern Iran. For this purpose, all consecutive Iranian patients referred to the endoscopy unit for esophagogastroduodenoscopy (EGD) due to dyspepsia were evaluated for erosive GERD. Participants with abnormal esophageal endoscopy other than GERD were excluded. In order to compare the GERD group with a comparison group, we selected other participants from the same referred dyspepsia patients whose esophageal mucosa was completely normal at endoscopy [Figure 1]. Patients with the following conditions were excluded in both groups: history of *H. pylori* eradication, recent treatment with H₂ blocker or proton pump inhibitors or non-steroidal anti-inflammatory drugs(NSAIDs) or medications induced GERD (e.g. anticholinergics, inhaled bronchodilators, and birth control pills), esophageal or gastric surgery, upper GI malignancy, and participants with poor cooperation. A checklist of EGD findings was filled out by gastroenterologist including of esophagus, stomach, and duodenum. An interviewer who was trained before starting the study, collected and recorded different demographic variables including age, sex, height, weight, sociodemographic status, smoking, and also histological reports of *H. pylori* in the checklist. Finally, the GERD group was compared with the non-GERD group in terms of *H. pylori*, considering the effect of confounding factors.

Esophagogastroduodenoscopy

All upper EGD were performed by a gastroenterologist to determine the presence or absence of erosive GERD and to evaluate its severity. In order to fully evaluate the upper GI tract, gastric and duodenal endoscopic findings were also examined. Gastric and duodenal endoscopic findings were divided into three groups: ulcerative, abnormal non-ulcerative (any evidence of mucosal lesion without ulcer), and normal (8). Hiatus hernia was diagnosed when the apparent separation between squamocolumnar junction and diaphragmatic impression was greater than two centimeter during quiet respiration (9).

To detect *H. pylori* infection, random biopsies from the antrum and body of stomach were obtained from all participants in the GERD and non-GERD groups. The samples were transferred to the laboratory in 10% formalin and under appropriate conditions. For histological diagnosis of *H. pylori*, staining was performed with Hematoxylin and eosin and also Giemsa staining.

Diagnosis and definition of GERD severity

The diagnosis of erosive GERD was made by EGD and the Los Angeles classification was used to grade the esophagitis. One or more mucosal breaks confined to the mucosal folds (each no longer than 5 mm), was defined as grade A. At least 1 mucosal break greater than 5 mm long confined to the mucosal folds, was defined as grade B. At least 1 mucosal break continuous between the tops of 2 or more mucosal folds, was defined as grade C. Circumferential mucosal break, was defined as grade D (10, 11).

Ethical approval/statement

This study was performed after obtaining the approval of Shiraz university ethical committee and Institutional Review Board (ID number: 93-01-13-8789) and based on Declaration of Helsinki regarding ethical principles for medical research. Written informed consent was obtained from all patients or their legal guardians to use their medical records in this study.

Statistical Analysis

Continuous data were calculated as means and standard deviations, whereas categorical parameters were expressed as percentages. Chi-squared test, T-test, and One-way ANOVA were used to capture the main differences between subjects, where appropriate. Robust Poisson regression analysis was used for estimating prevalence ratios (PRs) and confidence intervals (CIs) to evaluate the association of various independent variables on the GERD. All analyses were performed with the commercial software "Statistical Package for the Social Sciences" (SPSS version 25.0, IBM, Chicago, USA). A p value of < 0.05 was considered statistically significant.

Results

In total, 1916 patients were included in this study, of whom 874 (45.6%) had GERD and 1042 (54.4%) participants were considered as the non-GERD group (Figure 1). The mean age (SD) was 42.95 (16.32) ranging from 19 to 90. 672 (35.1%) and 1244 (64.9%) patients were male and female, respectively. Mean age in males and females was 46.17 ± 18.15 and 41.22 ± 14.96 , respectively. 1402 (73.2%) patients were rural residents and 183 (9.6%) were smokers. The demographic characteristics of the participants are shown in Table 1.

Overall, 1442 (75.3%) patients were positive and the others were *H. pylori* negative. Except for sociodemographic status ($P < 0.001$), other variables including gender ($P = 0.063$), age ($P = 0.695$), ethnicity ($P = 0.392$), body mass index (BMI) ($P = 0.236$), smoking ($P = 0.682$), and presence of hiatus hernia ($P = 0.601$) had no significant association with the frequency of *H. pylori* infection (Table 2).

Among patients with GERD, the presence of hiatus hernia, age, BMI, and smoking of patients in GERD group were significantly higher than non-GERD group ($P < 0.05$) but there was no significant difference in gender, sociodemographic status, and ethnicity between two groups (Table 2).

H. pylori infection was diagnosed in 673 (77.0%) patients in the GERD group, while 769 (73.8%) in the non-GERD group were positive for *H. pylori* infection. Frequency of *H. pylori* infection in the GERD group was higher than in the non-GERD group but there was no significant difference ($P = 0.106$). On the other hand, *H. pylori* infection was detected in 587 (77.6%) patients in mild GERD patients (LA grade A and B), while 39 (69.6%) in severe GERD (LA grade C and D) were positive for *H. pylori* infection. Although the frequency of *H. pylori* infection in mild GERD patients (LA grade A and B) was higher than the severe GERD (LA grade C and D), this difference was not significant ($P = 0.214$) (Table 3).

Comparison of the frequency of *H. pylori* infection in patients with and without GERD in different types of gastroduodenal endoscopic findings is presented in Table 4. Among patients with GERD, the frequency of *H. pylori* infection was significantly higher in those with gastric abnormal lesions include ulcerative lesions than normal gastric findings ($P = 0.006$) but in non-GERD group, this difference was not significant ($P = 0.068$). On the other hand, the frequency of *H. pylori* infection in both GERD ($P = 0.042$) and non-GERD ($P = 0.006$) groups in duodenal ulcer patients was significantly higher than normal endoscopic findings (Table 4).

The demographic and clinical characteristics and distribution of participants according to the BMI classification are shown in Table 5. Overall, the BMI (SD) was 24.54 (5.48) Kg/m², of which 1126 (58.8%) were in the normal weight group. Although 107 (5.6%) of patients were underweight, 480 (25.1%) and 203 (10.6%) were overweight and obese, respectively. There was a significant relationship between BMI and the presence of GERD ($P = 0.001$) as well as the sociological status ($P = 0.007$) and presence of hiatus hernia ($P < 0.001$) of the participants. There was no significant difference between BMI classification and frequency of other variables such as gender ($P = 0.134$), age ($P = 0.409$), gastric findings ($P = 0.073$), duodenal findings ($P = 0.629$), cigarette smoking ($P = 0.053$), and *H. pylori* infection ($P = 0.345$).

Robust Poisson regression models was used for estimating the PRs and 95% CIs to evaluate the association of various independent variables on the GERD (Table 6). *H. pylori* (PR 1.026; 95% CI 0.990-1.064; $P = 0.158$) and sociodemographic status (PR 1.030; 95% CI 0.998-1.063; $P = 0.064$) were not significantly associated with GERD group than non-GERD group. However, smoking (PR 1.139; 95% CI 1.089-1.192; $P < 0.001$), increased BMI (PR 1.060; 95% CI 1.027-1.093; $P < 0.001$), presence of hiatus hernia (PR 1.140; 95% CI 1.095-1.188; $P < 0.001$), and increased age (PR 1.002; 95% CI 1.001-1.003; $P < 0.001$) were significantly associated with GERD group compared to non-GERD group using regression analysis. In addition, the gastric ulcer (PR 0.875; 95% CI 0.809-0.947; $P = 0.001$) and duodenal ulcer (PR 0.911; 95% CI 0.862-0.963; $P = 0.001$) were also positively associated with presence of GERD.

Discussion

Our study showed that the frequency of *H. pylori* infection in the erosive GERD and non-GERD groups was not significantly different. In addition, there was no significant difference in the frequency of *H. pylori* between mild and severe GERD (Table 3). According to Robust Poisson regression models analysis, some variables including smoking, increased BMI, older age, presence of hiatus hernia, and peptic ulcer diseases (but not *H. pylori* infection) were significantly associated with GERD compared with the non-GERD group (Table 6). These results, in agreement with some studies (12–14), support the hypothesis that there is no association between the frequency of *H. pylori* infection and GERD.

GERD is a common GI disorder with different risk factors including obesity, smoking, alcohol use, pregnancy, scleroderma, and some foods or medications (1–3). Lifestyle modification is recommended as the first step in the treatment of GERD. Proton pump inhibitors are the mainstay of medical treatment for GERD if medication is needed, although a possible link between long-term use of these drugs and an increased risk of some side effects has been shown (1, 2, 15). *H. pylori*, as a common infection, plays an important role in the pathogenesis of various benign and malignant gastroduodenal diseases including gastric ulcer, gastric mucosal lymphoma and gastric cancer (4), however, there is still no agreement on its role in GERD. According to some reports, an inverse relationship between *H. pylori* and GERD has been observed (5, 16–19), but some other studies have not shown this relationship (12–14).

A study in healthy young Japanese volunteers, conducted by Tanaka et al., aimed to determine the prevalence and risk factors of *H. pylori* and GERD and their interrelationship. In this study, similar to our results, *H. pylori* infection had no effect on the prevalence of GERD, but obesity was a risk factor for GERD. They also showed that gender was a risk factor for GERD, but the frequency of smoking or abdominal hernia was not significantly different between groups that was different from our results. (12). Mahdi et al. investigated the association between CagA+ *H. pylori* and GERD and compared them with the healthy group. They concluded that the presence of *H. pylori* in patients with GERD was significantly increased compared to controls group (20).

In a research from Iran, 470 patients with dyspepsia and GERD were studied. The rate of *H. pylori* infection was 78.1%, which was almost similar to our results (Table 3) but the mean age of our patients was lower than their participants. They found no relationship between hiatus hernia and *H. pylori*, which was inconsistent with our results. (21). In another study from Iran, they did not find any association between *H. pylori* in patients with GERD compared to controls (22), which was consistent with our results.

Grand et al. conducted a study to examine 184 patients with reflux symptoms who underwent endoscopy with biopsy, esophageal pH-metry, and manometry. They showed that the role of *H. pylori* infection in the development of GERD as well as in the pathogenesis of esophageal reflux esophagus was not significant but hiatus hernia was significantly associated with the presence of reflux esophagus (23). In a study by Gisbert et al., they used pH-metry and endoscopy to diagnose GERD. In their research, *H. pylori* infection was not associated with GERD based on both procedures (24). Another study based on esophageal manometry, 24-hour pH monitoring, and EGD showed that GERD features, such as abnormal esophageal acid, erosive esophagus, or Barrett's esophagus, were not related to *H. pylori* (25), which is consistent with our results.

A prospective study of 146 patients with GERD, to determine the prevalence of *H. pylori* infection, found that there was no significant evidence for an important role in *H. pylori* infection in causing GERD and erosive esophagitis. In addition, although there was a significant relationship between hiatus hernia and reflux esophagitis, there was no significant correlation between HP and hiatus hernia, which was completely consistent with our results (26). Two other prospective evaluations by O'Connor et al. and Pieramico et al. also did not support the significant association between *H. pylori* infection and GERD (27, 28).

A study of 2508 GERD populations by Mari et al. showed that *H. pylori* infection was observed in 299 (11.9%) patients. Patients with GERD and *H. pylori* in this study were significantly younger, smoked more, and had less severe esophagitis, which was not similar to our study results (19). In Another study by Wang et al., in a non-erosive esophageal esophagus, showed that *H. pylori* infection was inversely associated with GERD, whereas male hiatus hernia were important factors associated with GERD (16). Other than the effect of hiatus hernia, other results of this study were inconsistent with our study.

Two studies from Korea showed that *H. pylori* seropositivity is preventive (17) and absence of *H. pylori* and male gender were associated with reflux esophagitis (18), which is not consistent with our study. But in one of the mentioned studies, reflux esophagitis was significantly associated with hiatal hernia and BMI that was similar to our results (18). A study by Yalaki et al., aimed at comparing and evaluating the relationship between GERD and *H. pylori* in adult patients with gastric localization of *H. pylori* infection and its historical features, the incidence of *H. pylori* has been shown to be significantly lower in patients with GERD than in the control group. This result is not consistent with the results of our study (5).

In a research from Indonesia, 104 patients with dyspepsia was analyzed to determine the prevalence of GERD and its risk factors. 53.8% of their patients had GERD that, similar to our results, smoking was significantly associated with GERD and most participants were classified as LA grade A. They also showed that higher economies increase the risk of GERD (29). In our study, although the frequency of *H. pylori* in the rural was significantly higher than the urban participants, there was no significant difference between the two groups in terms of GERD (Table 2,6).

As shown in this Table 4, among our patients with GERD, the frequency of *H. pylori* infection was significantly higher in those with gastric ulcer than normal gastric findings but in non- GERD group, this difference was not significant. Peptic ulcer disease is commonly associated disease with GERD (2) and EGD plays an important role in the diagnosis and differentiation of benign and malignant GI diseases and its complications include peptic ulcer disease and GERD (8–11). In a retrospective research by Jie et al., 953 peptic ulcer patients, 180 peptic ulcers and GERD patients, and 298 GERD patients were analyzed. They concluded that in patients with GERD, the prevalence of *H. pylori* infection in gastric ulcer patients was higher than without gastric ulcer (30), which was consistent with our results (Table 4). Moreover, gastric and duodenal ulcers, but not *H. pylori* infection, were significantly associated with GERD compared with the non-GERD group, according to Robust Poisson regression models analysis (Table 6).

Different treatment regimens have been suggested for *H. pylori* and some reports have been published on the effect of *H. pylori* eradication on GERD, however, their results have been inconsistent. (31–37). Although some reports have shown an inverse association between *H. pylori* eradication with GERD development (33, 35, 36), others have shown no beneficial effect of *H. pylori* eradication on GERD (31, 32, 34). Finally, there is no consensus on the hypothesis that eradicating *H. pylori* may cause or worsen GERD (4, 38).

The relationship between *H. pylori* and GERD is a complex and confusing issue due to the influence of various pathophysiological factors between them (5, 6). One reason for the heterogeneity of the results of previous researches to find the true relationship between *H. pylori* and GERD may be that the design of many studies was only to find a simple relationship between them, whereas in the final analysis of many of these reports, the effect of confounding factors for this association have not been measured. For instance, The *H. pylori* infection may make people susceptible to GERD by increasing gastric acid secretion, either directly infecting the gastric-type columnar epithelium, or by the action of noxious substances secreted by the infection into refluxed gastric juice (39). *H. pylori* seems to lead to much more complex changes in the gastric mucosa, including the modification of afferent neural signals and the secretion of specific gastric hormones. Ghrelin is a hormone that is mainly produced and released by the stomach with numerous functions. Ghrelin, in addition to enhancing gastric secretion, has a potent prokinetic function in the LES; this phenomenon, together with impaired vagal control, may play a role in the association of *H. pylori* infection with the development of GERD. Therefore, ghrelin and vagal activity may be missing links that partly explain the relationship between GERD and *H. pylori* infection (40).

The strength of our study was to analyze the association between GERD and *H. pylori* infection, taking into account many confounding factors. Other strengths of our study were the size of the considerable sample size, the presence of the comparison group, and the appropriate diagnostic evaluation for all participants. Our research also had limitations. One important limitation was that we included only erosive GERD patients, so the results of this study may not be generalizable to patients with non-erosive reflux disease. Another limitation was that the effects of some pathophysiological factors for both GERD and *H. pylori* were not measured in this study. Endoscopic biopsies to detect *H. pylori* in our study were sent to the laboratory as a mixture of gastric body and antrum in one sample container. It is recommended that in future studies, biopsies of different areas of the stomach be sent to the laboratory in separate sample containers for more accurate evaluation. Finally, this research was conducted in only one center, so a multicenter study is suggested.

Conclusions

Although our results support the hypothesis that there is no association between the frequency of *H. pylori* infection and erosive GERD, the available data do not provide sufficient evidence to define the true relationship between them and this issue remains controversial. We recommend further studies in this area, especially considering the effects of pathophysiological and other confounding factors on *H. pylori* infection and GERD.

Abbreviations

H. pylori: Helicobacter pylori

GERD: gastroesophageal reflux disease

BMI: body mass index

GI: gastrointestinal

EGD: esophagogastroduodenoscopy

NSAIDs: non-steroidal anti-inflammatory drugs

Declarations

Competing interests: all author declare that they have no competing interest.

Ethics approval and consent to participate: Shiraz university ethical committee and Institutional Review Board (ID number: 93-01-13-8789) and based on Declaration of Helsinki regarding ethical principles for medical research. Written informed consent was obtained from all patients or their legal guardians to use their medical records in this study.

Consent for publication: Not applicable

Availability of data and materials: The datasets generated and/or analyzed during the current study are not publicly available due our research center policy but are available from the corresponding author on reasonable request.

Funding: the authors receive no financial support for the research, authorship and/or publication of this article

Authors' contributions: Ramin Niknam, Kamran Bagheri Lankarani, Mohsen Moghadami and Seyed Alireza Taghavi contributed to the study conception and design. All authors performed material preparation, data collection and analysis. The first draft of the manuscript was written by Ramin Niknam and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Acknowledgements: This study was supported by research Council of Shiraz University of Medical Sciences, Shiraz, Iran (ID number: 93-01-13-8789).

References

1. Chen J, Brady P. Gastroesophageal Reflux Disease: Pathophysiology, Diagnosis, and Treatment. Gastroenterol Nurs. 2019;42(1):20-8.

2. Dunlap JJ, Patterson S. Gastroesophageal Reflux Disease. *Gastroenterol Nurs*. 2019;42(2):185-8.
3. Kellerman R, Kintanar T. Gastroesophageal Reflux Disease. *Prim Care*. 2017;44(4):561-73.
4. Jonaitis L, Pellicano R, Kupcinskas L. Helicobacter pylori and nonmalignant upper gastrointestinal diseases. *Helicobacter*. 2018;23 Suppl 1:e12522.
5. Yalaki S, Pulat H, Ilhan A. Localization of Helicobacter pylori gastritis and the relation of existing histopathological features with reflux esophagitis. *Scand J Gastroenterol*. 2020;55(1):27-33.
6. Yucel O. Interactions between Helicobacter pylori and gastroesophageal reflux disease. *Esophagus*. 2019;16(1):52-62.
7. Scida S, Russo M, Miraglia C, Leandro G, Franzoni L, Meschi T, et al. Relationship between Helicobacter pylori infection and GERD. *Acta Biomed*. 2018;89(8-S):40-3.
8. Niknam R, Manafi A, Fattahi MR, Mahmoudi L. The association between gastric endoscopic findings and histologic premalignant lesions in the Iranian rural population. *Medicine (Baltimore)*. 2015;94(17):e715.
9. Kahrilas PJ, Kim HC, Pandolfino JE. Approaches to the diagnosis and grading of hiatal hernia. *Best Pract Res Clin Gastroenterol*. 2008;22(4):601-16.
10. Armstrong D, Bennett JR, Blum AL, Dent J, De Dombal FT, Galmiche JP, et al. The endoscopic assessment of esophagitis: a progress report on observer agreement. *Gastroenterology*. 1996;111(1):85-92.
11. Lundell LR, Dent J, Bennett JR, Blum AL, Armstrong D, Galmiche JP, et al. Endoscopic assessment of oesophagitis: clinical and functional correlates and further validation of the Los Angeles classification. *Gut*. 1999;45(2):172-80.
12. Tanaka Y, Sakata Y, Hara M, Kawakubo H, Tsuruoka N, Yamamoto K, et al. Risk Factors for Helicobacter pylori Infection and Endoscopic Reflux Esophagitis in Healthy Young Japanese Volunteers. *Intern Med*. 2017;56(22):2979-83.
13. Calleja JL, Suarez M, De Tejada AH, Navarro A, Pantogerd G. Helicobacter pylori infection in patients with erosive esophagitis is associated with rapid heartburn relief and lack of relapse after treatment with pantoprazole. *Dig Dis Sci*. 2005;50(3):432-9.
14. Hackelsberger A, Schultze V, Gunther T, von Arnim U, Manes G, Malfertheiner P. The prevalence of Helicobacter pylori gastritis in patients with reflux oesophagitis: a case-control study. *Eur J Gastroenterol Hepatol*. 1998;10(6):465-8.
15. Fattahi MR, Niknam R, Shams M, Anushiravani A, Taghavi SA, Omrani GR, et al. The Association Between Prolonged Proton Pump Inhibitors Use and Bone Mineral Density. *Risk Manag Healthc Policy*. 2019;12:349-55.
16. Wang PC, Hsu CS, Tseng TC, Hsieh TC, Chen CH, Su WC, et al. Male sex, hiatus hernia, and Helicobacter pylori infection associated with asymptomatic erosive esophagitis. *J Gastroenterol Hepatol*. 2012;27(3):586-91.
17. Ko SH, Baeg MK, Jung HS, Kim P, Choi MG. Russian Caucasians have a higher risk of erosive reflux disease compared with East Asians: A direct endoscopic comparison. *Neurogastroenterol Motil*.

2017;29(5).

18. Nam SY, Park BJ, Cho YA, Ryu KH, Choi IJ, Park S, et al. Different effects of dietary factors on reflux esophagitis and non-erosive reflux disease in 11,690 Korean subjects. *J Gastroenterol*. 2017;52(7):818-29.
19. Mari A, Mahroum N, Bragazzi NL, Shalaata M, Khoury T, Watad A, et al. Demographics and Clinical and Endoscopic Characteristics of Patients with Helicobacter pylori Infection and Gastroesophageal Reflux Disease: A Case-Control Study. *Gastroenterol Res Pract*. 2019;2019:3819893.
20. Mahdi BM. The relationship between helicobacter pylori infection and gastro-esophageal reflux disease. *N Am J Med Sci*. 2011;3(3):142-5.
21. Nobakht H, Boghratian A, Sohrabi M, Panahian M, Rakhshani N, Nikkhah M, et al. Association between Pattern of Gastritis and Gastroesophageal Reflux Disease in Patients with Helicobacter Pylori Infection. *Middle East J Dig Dis*. 2016;8(3):206-11.
22. Shavalipour A, Malekpour H, Dabiri H, Kazemian H, Zojaji H, Bahrudi M. Prevalence of cytotoxin-associated genes of Helicobacter pylori among Iranian GERD patients. *Gastroenterol Hepatol Bed Bench*. 2017;10(3):178-83.
23. Grande M, Lisi G, De Sanctis F, Grande S, Esser A, Campanelli M, et al. Does a relationship still exist between gastroesophageal reflux and Helicobacter pylori in patients with reflux symptoms? *World J Surg Oncol*. 2014;12:375.
24. Gisbert JP, de Pedro A, Losa C, Barreiro A, Pajares JM. Helicobacter pylori and gastroesophageal reflux disease: lack of influence of infection on twenty-four-hour esophageal pH monitoring and endoscopic findings. *J Clin Gastroenterol*. 2001;32(3):210-4.
25. Oberg S, Peters JH, Nigro JJ, Theisen J, Hagen JA, DeMeester SR, et al. Helicobacter pylori is not associated with the manifestations of gastroesophageal reflux disease. *Arch Surg*. 1999;134(7):722-6.
26. Grande M, Cadeddu F, Villa M, Attina GM, Muzi MG, Nigro C, et al. Helicobacter pylori and gastroesophageal reflux disease. *World J Surg Oncol*. 2008;6:74.
27. O'Connor HJ, Cunnane K. Helicobacter pylori and gastro-oesophageal reflux disease—a prospective study. *Ir J Med Sci*. 1994;163(8):369-73.
28. Pieramico O, Zanetti MV. Relationship between intestinal metaplasia of the gastro-oesophageal junction, Helicobacter pylori infection and gastro-oesophageal reflux disease: a prospective study. *Dig Liver Dis*. 2000;32(7):567-72.
29. Miftahussurur M, Doohan D, Nusi IA, Adi P, Rezkitha YAA, Waskito LA, et al. Gastroesophageal reflux disease in an area with low Helicobacter pylori infection prevalence. *PLoS One*. 2018;13(11):e0205644.
30. Jie W, Qinghong X, Zhitao C. Association of Helicobacter pylori infection with gastroesophageal reflux disease. *J Int Med Res*. 2019;47(2):748-53.
31. Fallone CA, Barkun AN, Friedman G, Mayrand S, Loo V, Beech R, et al. Is Helicobacter pylori eradication associated with gastroesophageal reflux disease? *Am J Gastroenterol*. 2000;95(4):914-20.
32. McColl KE, Dickson A, El-Nujumi A, El-Omar E, Kelman A. Symptomatic benefit 1-3 years after H. pylori eradication in ulcer patients: impact of gastroesophageal reflux disease. *Am J Gastroenterol*.

2000;95(1):101-5.

33. Rokkas T, Ladas SD, Triantafyllou K, Liatsos C, Petridou E, Papatheodorou G, et al. The association between CagA status and the development of esophagitis after the eradication of *Helicobacter pylori*. *Am J Med*. 2001;110(9):703-7.
34. Schwizer W, Thumshirn M, Dent J, Guldenschuh I, Menne D, Cathomas G, et al. *Helicobacter pylori* and symptomatic relapse of gastro-oesophageal reflux disease: a randomised controlled trial. *Lancet*. 2001;357(9270):1738-42.
35. Take S, Mizuno M, Ishiki K, Nagahara Y, Yoshida T, Yokota K, et al. *Helicobacter pylori* eradication may induce de novo, but transient and mild, reflux esophagitis: Prospective endoscopic evaluation. *J Gastroenterol Hepatol*. 2009;24(1):107-13.
36. Xie T, Cui X, Zheng H, Chen D, He L, Jiang B. Meta-analysis: eradication of *Helicobacter pylori* infection is associated with the development of endoscopic gastroesophageal reflux disease. *Eur J Gastroenterol Hepatol*. 2013;25(10):1195-205.
37. Mahmoudi L, Farshad S, Seddigh M, Mahmoudi P, Ejtehad F, Niknam R. High efficacy of gemifloxacin-containing therapy in *Helicobacter Pylori* eradication: A pilot empirical second-line rescue therapy. *Medicine (Baltimore)*. 2016;95(42):e4410.
38. Malfertheiner P, Megraud F, O'Morain CA, Gisbert JP, Kuipers EJ, Axon AT, et al. Management of *Helicobacter pylori* infection-the Maastricht V/Florence Consensus Report. *Gut*. 2017;66(1):6-30.
39. Spechler SJ. Does *Helicobacter pylori* infection contribute to gastroesophageal reflux disease? *Yale J Biol Med*. 1998;71(2):143-8.
40. Thor PJ, Blaut U. *Helicobacter pylori* infection in pathogenesis of gastroesophageal reflux disease. *J Physiol Pharmacol*. 2006;57 Suppl 3:81-90.

Tables

Table 1. Demographic characteristics of participants with dyspepsia (n=1916)

Gender	
Male	672 (35.1%)
Female	1244 (64.9%)
Age distribution (yrs.)	
<30	486 (25.4%)
30-39	438 (22.9%)
40-49	380 (19.8%)
50-59	293 (15.3%)
≥ 60	319 (16.6%)
Age (yrs.) ; Mean ± SD	42.95 ± 16.32
Ethnicity	
Fars	1460 (76.2%)
Lurs	280 (14.6%)
Turke	108 (5.6%)
Others	68 (3.5%)
Sociodemographic status	1402 (73.2%)
Rural	514 (26.8%)
Urban Age	
Body mass index(Kg/m²);Mean±SD	24.54 ± 5.48
Cigarette smoking	183 (9.6%)
Hiatus hernia	223 (11.6%)

Table 2. Comparison of demographic characteristics of dyspeptic patients considering the presence of H. pylori and GERD (n=1916)

Variable	GERD; N (%)	Non-GERD; N (%)	P value	Positive H pylori; N (%)	Negative H pylori; N (%)	P value
Gender ¹			0.363			0.063
Male	316 (36.2%)	356 (34.2%)		489 (33.9%)	183 (38.6%)	
Female	558 (63.8%)	686 (65.8%)		953 (66.1%)	291 (61.4%)	
Age (yrs.) ; Mean ± SD ²	44.31±16.67	41.81±15.94	0.001	42.87±15.95	43.22±17.41	0.695
Sociodemographic status ¹			0.210			<0.001
	599 (68.5%)	686 (65.8%)		1055 (73.2%)	230 (48.5%)	
Rural	275 (31.5%)	356 (34.2%)			244 (51.5%)	
Urban				387 (26.8%)		
Ethnicity ¹			0.426			0.392
Fars	651 (74.5%)	809 (77.6%)		1111 (77.0%)	349 (73.6%)	
Lurs	139 (15.9%)	56 (13.5%)		207 (14.4%)	73 (15.4%)	
Turke	52(5.9%)	141 (5.4%)		76 (5.3%)	32 (6.8%)	
Others	32 (3.7%)	36 (3.5%)		48 (3.3%)	20 (4.2%)	
Body mass index(Kg/m);Mean ± SD ²	24.86 ± 5.44	24.28 ± 5.51	0.021	24.63 ± 5.48	24.28 ± 5.50	0.236
Cigarette smoking ¹	115 (13.2%)	68 (6.5%)	<0.001	140 (9.7%)	43 (9.1%)	0.682
Hiatus hernia ¹	151 (17.3%)	72 (6.9%)	<0.001	171 (11.9%)	52 (11.0%)	0.601

Notes: ¹ Chi-square Test; ² T-Test

Abbreviations: H. pylori, helicobacter pylori; GERD, gastroesophageal reflux disease

Table 3. Comparison of the frequency of H. pylori infection in dyspeptic patients considering the presence and grades of GERD (n=1916)

Endoscopic findings	Positive H. pylori	Negative H. pylori	P value
Groups; N (%) ¹			0.106
GERD	673 (77.0%)	201 (23.0%)	
Non-GERD	769 (73.8%)	273 (26.2%)	
Grades of GERD; N (%) ^{1,2}			0.214
LA grade A	322 (75.8%)	103 (24.2%)	
LA grade B	265 (80.1%)	66 (19.9%)	
LA grade C	26 (66.7%)	13 (33.3%)	
LA grade D	13 (76.5%)	4 (23.5%)	

Notes: ¹ Chi-square Test; ¹ The Los Angeles Classification of gastroesophageal reflux disease was used for grading Abbreviations: H. pylori, helicobacter pylori; GERD, gastroesophageal reflux disease

Table 4. Comparison of frequency of H. pylori infection in patients with and without GERD in different types of gastroduodenal endoscopic findings in dyspeptic patients

Variables		Gastric endoscopic findings; N (%)			P value ¹
		Normal	Ulcerative	Abnormal non-ulcerative	
GERD	Positive H. pylori	359 (73.0%)	19 (82.6%)	295 (82.2%)	0.006
	Negative H. pylori	133 (27.0%)	4 (17.4%)	64 (17.8%)	
Non- GERD	Positive H. pylori	418 (71.1%)	38 (74.5%)	313 (77.7%)	0.068
	Negative H. pylori	170 (28.9%)	13 (25.5%)	90 (22.3%)	
		Duodenal endoscopic findings; N (%)			
		Normal	Ulcerative	Abnormal non-ulcerative	
GERD	Positive H. pylori	575 (75.7%)	47 (88.7%)	51 (83.6%)	0.042
	Negative H. pylori	185 (24.3%)	6 (11.3%)	10 (16.4%)	
Non-GERD	Positive H. pylori	633 (72.0%)	86 (86.0%)	50 (79.4%)	0.006
	Negative H. pylori	246 (28.0%)	14 (14.0%)	13 (20.6%)	

Notes: ¹ Chi-square Test

Abbreviations: H. pylori, helicobacter pylori; GERD, gastroesophageal reflux disease

Table 5. Distribution of demographic and clinical features of participants (n=1916) according to BMI classification ¹

Variables	Underweight;N(%)	Normal;N(%)	Overweight;N(%)	Obese;N(%)	P value
	107 (5.6%)	1126 (58.8%)	480 (25.1%)	203(10.6%)	
Gender²					0.134
Male	45 (42.1%)	388 (34.5%)	178 (37.1%)	61 (30.0%)	
Female	62 (57.9%)	738 (65.5%)	302 (62.9%)	142 (70.0%)	
Age (yrs.); Mean $\pm$ SD³	40.81 $\pm$ 16.35	43.36 $\pm$ 16.34	42.53 $\pm$ 16.51	42.82 $\pm$ 15.75	0.409
H. pylori infection²					0.345
Positive	77 (72.0%)	841 (74.7%)	375 (78.1%)	149 (73.4%)	
Negative	30 (28.0%)	285 (25.3%)	105 (21.9%)	54 (26.6%)	
Esophageal findings²					0.001
GERD	44 (41.1%)	477(42.4%)	245 (51.0%)	108 (53.2%)	
Non-GERD	63 (58.9%)	649 (57.6%)	235 (49.0%)	95 (46.8%)	
Gastric findings²					0.073
Normal	70 (65.4%)	615 (54.6%)	277 (57.7%)	118 (58.1%)	
Ulcerative	6 (5.6%)	42 (3.7%)	14 (2.9%)	12 (5.9%)	
Abnormal non-ulcerative	31 (29.0%)	469 (41.7%)	189 (39.4%)	73 (36.0%)	
Duodenal findings²					0.629
Normal	95 (88.8%)	954 (84.7%)	414 (86.3%)	176 (86.7%)	
Ulcerative	9 (8.4%)	92 (8.2%)	35 (7.3%)	17 (8.4%)	
Abnormal non-ulcerative	3 (2.8%)	80 (7.1%)	31 (6.5%)	10 (4.9%)	
Cigarette smoking²	18 (16.8%)	107 (9.5%)	39 (8.1%)	19 (9.4%)	0.053
Sociodemographic status²					0.007
Rural	83 (77.6%)	763 (67.8%)	320 (66.7%)	119 (58.6%)	
Urban	24 (22.4%)	363 (32.2%)	160 (33.3%)	84 (41.4%)	

Hiatus hernia ²	9 (8.4%)	108 (9.6%)	69 (14.4%)	37 (18.2%)	<0.001
----------------------------	----------	------------	------------	------------	--------

Notes: ¹ The body mass index (Kg/m2) was classified according to the World Health Organization (WHO) classification into 4 groups of less than 18.5 as under-weight range, 18.5 to < 25 as normal, 25.0 to < 30 as over-weight range, and 30.0 or higher as obese range; ² Test: Chi-squared test ; ³ Test: One-way ANOVA

Abbreviations: BMI, body mass index; H. pylori, helicobacter pylori

Table 6. Robust Poisson regression models estimating prevalence ratio (PR) and 95% confidence interval (CI) to evaluate the association of various independent variables on the GERD

Variable	Crude model		Adjusted model	
	PR (95% CI)	P value	PR (95% CI)	P value
Cigarette smoking		<0.001		<0.001
Yes	1.132 (1.082-1.186)		1.139 (1.089-1.192)	
No	1.0		1.0	
Body mass index ¹		<0.001		<0.001
Overweight range and obese Normal and under-weight	1.066 (1.033-1.100)		1.060 (1.027-1.093)	
	1.0		1.0	
Hiatus hernia		<0.001		<0.001
Yes	1.175 (1.129-1.223)		1.140 (1.095-1.188)	
No	1.0		1.0	
Gastric ulcer		<0.001		0.001
Yes	0.897(0.826-0.973)		0.875 (0.809-0.947)	
No	1.0		1.0	
Duodenal ulcer		0.004		0.001
Yes	0.919(0.867-0.974)		0.911 (0.862-0.963)	
No	1.0		1.0	
Age	1.002 (1.001-1.003)	0.001	1.002 (1.001-1.003)	<0.001
Sociodemographic status		0.211		0.064
Rural	1.021(0.988-1.055)		1.030 (0.998-1.063)	
Urban	1.0		1.0	
Helicobacter pylori		0.106		0.158
Positive	1.030 (0.994-1.068)		1.026 (0.990-1.064)	
Negative	1.0		1.0	

Notes: ¹ The body mass index (Kg/m²) was classified according into 2 groups of less than 25 as normal and under-weight, 25 or higher as overweight range and obese.

Figures

Figure 1

Flow diagram for the patient's selection process

¹ All consecutive patients referred to the endoscopy unit for esophagogastroduodenoscopy because of dyspepsia.

² Patients with the following conditions were excluded: history of H. pylori eradication, recent treatment with H2 blocker or proton pump inhibitors or NSAIDs or medications induced GERD (e.g. anticholinergics, inhaled bronchodilators, and birth control pills), esophageal or gastric surgery, upper gastrointestinal malignancy, and participants with poor cooperation.

³ The diagnosis of erosive gastroesophageal reflux disease was made by esophagogastroduodenoscopy

Abbreviation: GERD, gastroesophageal reflux disease