Vibrational Resonant Inelastic X-Ray Scattering Inliquid Acetic Acid: A Ruler for Molecular Chain Lengths

Viktoriia Savchenko

Royal Institute of Technology

Iulia-Emilia Brumboiu

Royal Institute of Technology

Victor Kimberg

Royal Institute of Technology

Michael Odelius (✉ odelius@fysik.su.se)

Stockholm University

Pavel Krasnov

Siberian Federal University

Ji-Cai Liu

North China Electric Power University

Jan-Erik Rubensson

Uppsala University

Olle Björneholm

Uppsala University

Conny Sâthe

Lund University

Johan Gråsjö

Uppsala University

Minjie Dong

Uppsala University

Annette Pietzsch

Helmholtz-Zentrum Berlin für Materialien und Energie

Alexander Föhlisch

Helmholtz-Zentrum Berlin für Materialien und Energie

Thorsten Schmitt

Paul Scherrer Institute

Daniel McNally

Paul Scherrer Institute

Xingye Lu

Paul Scherrer Institute

Sergey Polyutov

Siberian Federal University

Patrick Norman

Royal Institute of Technology

Marcella Iannuzzi

University of Zurich

Faris Gel'mukhanov

Royal Institute of Technology

Victor Ekholm

Lund University

Research Article

Keywords: hydrogen bond, ACA, XAS

Posted Date: December 3rd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-112475/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on February 18th, 2021. See the published version at <https://doi.org/10.1038/s41598-021-83248-3>.

Vibrational resonant inelastic X-ray scattering in liquid acetic acid: a ruler for molecular chain lengths

Viktoriia Savchenko^{1,2,3,*}, Iulia Emilia Brumboiu^{1,4}, Victor Kimberg^{1,2,3,*}, Michael Odelius^{5,*}, Pavel Krasnov^{1,2}, Ji-Cai Liu^{6,13}, Jan-Erik Rubensson⁷, Olle Björneholm⁷, Conny Sâthe⁸, Johan Gråsjö^{7,14}, Minjie Dong⁷, Annette Pietzsch⁹, Alexander Föhlisch^{9,10}, Thorsten Schmitt¹¹, Daniel McNally¹¹, Xingye Lu¹¹, Sergey Polyutov^{2,3}, Patrick Norman¹, Marcella Iannuzzi¹², Faris Gel'mukhanov^{1,2,3}, and Victor Ekholm^{7,8}

¹Department of Theoretical Chemistry and Biology, KTH Royal Institute of Technology, Stockholm, 10691, Sweden

²Siberian Federal University, Krasnoyarsk, 660041, Russia

³Kirensky Institute of Physics, Federal Research Center KSC SB RAS, Krasnoyarsk, 660036, Russia

⁴Department of Chemistry, Korea Advanced Institute of Science and Technology, Daejeon, 34141, Korea

⁵Department of Physics, AlbaNova University Center, Stockholm University, Stockholm, SE-106 91, Sweden

⁶Department of Mathematics and Physics, North China Electric Power University, Beijing, 102206, China

⁷Department of Physics and Astronomy, Uppsala University, Box 516, Uppsala, SE-751 20, Sweden

⁸MAX IV Laboratory, Lund University, Box 118, Lund, SE-221 00, Sweden

⁹Institute for Methods and Instrumentation in Synchrotron Radiation Research FG-ISRR, Helmholtz-Zentrum Berlin für Materialien und Energie Albert-Einstein-Strasse 15, Berlin, 12489, Germany

¹⁰Institut für Physik und Astronomie, Universität Potsdam, Karl-Liebknecht-Strasse 24-25, Potsdam, 14476, Germany

¹¹Swiss Light Source, Photon Science Division, Paul Scherrer Institut, Villigen PSI, CH-5232, Switzerland

¹²Physical Chemistry Institute, University of Zürich, Zürich, 8057, Switzerland

¹³Collaborative Innovation Center of Light Manipulations and Applications, Shandong Normal University, Jinan, 250358, China

¹⁴Department of Medicinal Chemistry, Uppsala University, Box 574, Uppsala, 75123, Sweden

*Correspondence and requests for materials should be addressed to VS (viksav@kth.se), VK (kimberg@kth.se), and MO (odelius@fysik.su.se).

ABSTRACT

Quenching of vibrational excitations in resonant inelastic X-ray scattering (RIXS) spectra of liquid acetic acid is observed. At the oxygen core resonance associated with localized excitations at the O-H bond, the spectra lack the typical progression of vibrational excitations observed in RIXS spectra of comparable systems. We interpret this phenomenon as due to strong rehybridization of the unoccupied molecular orbitals as a result of hydrogen bonding. This allows us to address the molecular structure of the liquid, and to determine a lower limit for the average molecular chain length.

Introduction

The hydrogen bond (HB) is of central importance in chemistry and biochemistry and it is crucial in chemical reactions, supramolecular structures, molecular assemblies, and even life processes. Consequently, HBs have been immensely studied over the years, exploiting a plethora of spectroscopy and scattering methods, with liquid water as an important showcase¹. The new generation synchrotron radiation sources have allowed for a refinement of the resonant inelastic X-rays scattering (RIXS) technique, which now gives access to detailed information about the nature of the HB, not only in liquid water²⁻⁶, but also liquids like acetone⁷ and methanol⁸.

Vibrational excitations observed in RIXS spectra give information about the local potential surface of the electronic ground state⁹, and about the electronic-vibronic dynamics during the scattering process. The dynamics is especially dramatic when the intermediate core-excited state is dissociative in a bond to a light atom, e.g., a state in which an O-H antibonding orbital is populated. This dynamics is sensitively reflected in extended vibrational progressions in the RIXS spectra, and the exploration of these phenomena for the investigation of HBs in the liquid phase is a new research field which we are currently entering.

Recently, unambiguous experimental evidence for ultrafast proton-transfer dynamics following excitation on the "pre-peak"

resonance of liquid water was provided^{5,6}. Significant nuclear rearrangement could be expected as the pre-peak resonance is associated with the $1s^{-1}4a_1$ excitation in gas-phase water, in which ultrafast dissociation arises from an antibonding O-H σ^* character of the $4a_1$ orbital¹⁰. RIXS spectra of liquid water excited at this resonance show a long vibrational progression (Fig. 1 (a)). Also in corresponding measurements on methanol⁸ and acetone⁷, it was found that the long vibrational progression of the RIXS spectra of the free molecules was largely preserved in the liquid. These observations suggest that the intermolecular interactions, albeit significant, only play a minor role for the RIXS dynamics. Here we will demonstrate that this by no means is a general rule.

Results

In the present work we combine state-of-the-art theory with high-resolution RIXS spectra (Fig. 1 (b,c)) of liquid acetic acid (ACA) to analyze the influence of intermolecular interactions. We show that the HB has a dramatic influence on the molecular dynamics. Whereas ultrafast dissociation, and accompanying extended vibrational progression are expected in RIXS spectra excited at the O-H antibonding intermediate state of O $1s^{-1}\sigma^*$ character for the free molecule, the corresponding dynamics is almost entirely quenched in liquid ACA.

Carboxylic acids have received a lot of experimental and theoretical interest over more than a century. There is now a consensus that the predominant structures of ACA in the gas phase are the monomer and the symmetric cyclic dimer¹¹. In the solid state, the crystalline unit cell consists of non-symmetric dimers which belong to a chain structure¹¹. The local structure of liquid ACA is more complex and is still heavily debated (see Ref. ¹¹⁻¹³ and references therein). The linear cis-cis and cis-trans configurations are theoretically found in less ordered chains and rings of liquid acetic acid¹¹. However, neutron scattering¹⁴ and Raman spectroscopy¹⁵ show a remarkable similarity in the hydrogen bond length when going from the crystal to the liquid structure and it has been concluded that the liquid structure is similar to that of a disordered crystal with predominant chain structures.

The interpretation of spectroscopic and scattering data of disordered liquids is complicated by the difficulty to get direct access to the structure, mainly because of the ambiguity in the analysis and interpretation of structural measurements. A crucial role is played by theoretical modeling, but in spite of theoretical support, the solution of the inverse problem is ambiguous (Refs. ¹²⁻¹⁴). Therefore, complementary investigations are called for, and here we use modeling-supported X-ray absorption spectroscopy (XAS) and RIXS spectroscopy to shed light on the structure and interactions in ACA.

The spectral region of our interest consists of the first two peaks in the oxygen K-edge XAS of acetic acid (Fig. 2). Because of the different chemical shifts of the O $1s$ levels of carbonyl (O_C) and hydroxyl (O_H) oxygen atoms the chemical site can be selected by tuning the excitation energy. Indeed, the XAS spectrum of the acetic acid monomer has two distinct pre-edge peaks and the analysis shows that each peak is related to a distinct oxygen atom (Fig. 2). XAS simulations of the isolated ACA show that the first peak around 532 eV is due to $O_C 1s \rightarrow \pi^*$ core excitation, while the second peak at 535 eV is formed by two close-lying peaks $O_H 1s \rightarrow \pi^*$ and $O_H 1s \rightarrow \sigma^*$ with almost the same intensity¹⁶. The C-O equilibrium distance of the ground state and the core excited state $O_C 1s^{-1}\pi^*$ state differs (Fig. 3) due to the strong interaction between the $2p$ orbitals of carbon and oxygen. Consequently, a long C=O vibrational progression is predicted for the gas-phase RIXS spectrum (Fig. 1 (b)).

The O-H equilibrium distance of the core excited $O_H 1s^{-1}\pi^*$ state of the ACA monomer is almost the same as the ground state, largely because the oxygen $2p$ orbital does not interact with the hydrogen $1s$ orbital (Fig. 3), and consequently, RIXS spectra excited at this resonance are predicted to show almost no OH vibrations (Fig. S3¹⁷). In contrast, the $O_H 1s^{-1}\sigma^*$ is dissociative (Fig. 3), resulting in an extensive vibrational progression (Fig. 1 (c)) in the corresponding monomer RIXS spectrum.

Experimentally, we find that the vibrational progressions predicted for free molecules to a large extent are preserved in the RIXS spectra of the liquids. The sensitivity to the chemical surrounding is typically accentuated for highly excited vibrational states, where potential surfaces are modified by intermolecular interactions. Therefore, vibrational progressions are often smeared out and attenuated at large energy losses. For lower energy losses, on the other hand, the behavior simulates gas-phase predictions. This is illustrated for the O $1s \rightarrow 4a_1$ resonance in water (Fig. 1 (a)), and for the $O_C 1s \rightarrow \pi^*$ resonance in ACA (Fig. 1 (b)). The predicted O-H vibrations for gas-phase water and O-C vibrations for ACA, are both retained in the RIXS spectra of the liquids, at least up to $v = 5$, before intermolecular interactions smear out the intensity.

For the second resonance of liquid ACA the situation is dramatically different (Fig. 1 (c)). Whereas a long progression of O-H stretch vibrations with a $\omega_{\text{vib}}^{\text{exp}} \approx 0.44$ eV is predicted for the free molecule, vibrational excitations are virtually absent in the experimental RIXS spectrum. This observation suggests that intermolecular interactions dramatically quench the ultrafast dissociation, and that the HB has a major impact on the dynamics. Notable, however, is a weak vibrational peak structure, where the vibrational peaks are separated 0.16-0.17 eV. These could be attributed to the OH bending mode and/or carboxyl C-O stretch^{18,19}. To shed light on the unusual effect of complete quenching of the O-H stretch progression, we analyze the core-excitations of ACA molecules in gas phase and in liquid. For comparison with liquid phase, an experimental gas-phase spectrum would be ideal. However, the spectral quality needed to make a comparison is state-of-the-art, and presently there is no instrument available where such a gas-phase measurement can be done. Instead, we use a high-quality theoretical calculation

of the RIXS spectrum of the ACA monomer (Fig. 1 (c)) which is expected to accurately predict an experimental gas-phase spectrum (compare to our previous study of RIXS of free water molecule²⁰). We use this as a reference to compare with the measured RIXS spectra of the liquid.

We represent the gas-phase ACA by the monomer, and assume that a chain of hydrogen bonded molecules is the representative structure of liquid ACA (Fig. 4 (d)). Here, we identify two distinct OH groups: one at the end of the chain, not involved in HB, which we denote "end", and one inside the chain, involved in HB, which we denote "in". These two OH groups are expected to have significantly different XAS spectra, which we have computed using the shortest possible chains, i.e. the dimers. Two types of dimers are included, namely the regular dimer (centrosymmetric cyclic dimer, which also exist in the gas phase) and the inverted dimer (asymmetric noncyclic dimer which is the shortest representative of the chain present in the liquid). The XAS spectra, computed in the same way as for the monomer, are shown in Fig. 2.

Discussion

Note that while π^* and σ^* resonances are predicted to contribute almost equally to the 535-eV peak in the XAS spectrum of the monomer (Fig. 2 (a)), the σ^* resonance *does not contribute at all* in the regular dimer, as previously discussed in the literature¹⁶ (Fig. 2 (b)). This dramatic change is an obvious effect of the HB, which affects both "in" OH groups in the regular dimer. For the inverted dimer the XAS spectrum comprises transitions in both "in" and "end" OH groups (Fig. 2 (c)), and whereas the "in" OH group has similar features as the OH groups in the regular dimer the XAS signal of the "end" OH resembles the signal of the OH group in the ACA monomer. This is a manifestation of the building block principle, according to which the occurrence of specific groups leaves a fingerprint in the spectrum. The effect of the HBs is, however, not very large in the total XAS, and condensation results only in slight shifts of the spectral features (Fig. 2 (d-e)). In contrast, a strong effect is expected in RIXS: Whereas the "end" OH groups are almost unaffected by the HB, and the 535-eV peak comprises transitions to the dissociative σ^* core-excited state a long vibrational progression is expected in the RIXS spectrum, the peak in the spectra of the "in" OH groups has no contribution from the σ^* state, and consequently no significant vibrational excitations are expected. In this way a RIXS measurement allows us to clearly distinguish between "in" and "end" OH groups, and the observation that vibrational excitations are virtually absent shows that HBs are significant, and puts an upper limit to the occurrence of "end" OH groups in the liquid, which we estimate in the following.

The total number of ACA units in a liquid can be written approximately as

$$\mathcal{N} = \ell_{\text{ch}} N_{\text{ch}} + \ell_{\text{cycl}} N_{\text{cycl}} \quad (1)$$

Here ℓ_{ch} and ℓ_{cycl} are the average number of ACA monomers (or length in ACA units) in chain and cyclic structures, respectively, while N_{ch} and N_{cycl} are the average number of chains and cyclic structures, respectively. Here we assume that the chain is the representative chain structure from crystalline ACA with one "end" OH group (see Fig. 4 (d)). The cyclic structures have only "in" OH groups (see Fig. 4 (c)). The relative abundance of the "end" OH groups is $N_{\text{ch}}/\mathcal{N}$. This ratio represents the relative intensity of the RIXS profile with the OH vibrational progression caused by the "end" OH group.

A detailed analysis of peak-to-noise ratio in the RIXS spectra of liquid ACA and the calculated $\nu = 1$ peak intensities in the ACA monomer is performed in the Supplementary information¹⁷. From this analysis, we conclude that for a vibrational progression to exceed the experimental noise level (more precisely the level of detectability), the fraction of chains must obey the following relation $\mathcal{N}/N_{\text{ch}} = \ell_{\text{eff}} = \ell_{\text{ch}} + \ell_{\text{cycl}} N_{\text{cycl}}/N_{\text{ch}} \gtrsim 3.5$, where ℓ_{eff} denotes the effective length of chain or cycle unit.

Since our measurements do not display a OH vibrational progression, this means that the effective length ℓ_{eff} should be longer than 3.5 monomer units.

$$\frac{\mathcal{N}}{N_{\text{ch}}} = \ell_{\text{ch}} + \ell_{\text{cycl}} \frac{N_{\text{cycl}}}{N_{\text{ch}}} > 3.5. \quad (2)$$

Let us take the recommended structure from a recent analysis¹⁴ of neutron scattering data, where it is claimed that the dominant structure is the trimer chain. If trimer chains were the only motif then $N_{\text{cycl}} = 0$ and $\ell_{\text{ch}} = 3$, so that the condition for observing vibrational excitation would be fulfilled. Our results, indicate that the effective chain length is longer. Our estimate corresponds to a lower bound for ℓ_{eff} within the current *signal/noise* ratio. One can expect to refine the estimate, possibly giving an increased ℓ_{eff} , in future experiments with improved statistics.

So far, we have discussed RIXS only from the perspective of the $O_H 1s^{-1}\sigma^*$ core-excited state of the ACA monomer and dimers. However, one may also expect vibrational structure in RIXS due to the $O_H 1s^{-1}\pi^*$ core-excited states which are also populated when the photon energy is tuned near the second XAS peak (Fig. 2). One cannot exclude that the HB changes the potential energy curve of the π^* core-excited states and opens inelastic scattering channels, but, as shown in the Supplementary information¹⁷, this does not occur.

Additionally, ab initio molecular dynamics (AIMD) simulations of the XAS spectrum were performed to investigate the sensitivity to solvation and HBs, using the transition-potential DFT approach (see the Supplementary information¹⁷). The results of the simulations (Fig. 4 (b)) confirm that the σ^* core-excited state only contributes little to the second XAS peak in liquid ACA. The individual contributions of the C=O and O-H oxygen atoms to the *O* 1s XAS spectra (displayed in blue and red lines, respectively) show that liquid interaction have a profoundly different influence on the two sets of core-excited states. In Fig. 4 (a,b), we also show the decomposition of the XAS spectrum contributions into in-plane and out-of-plane components which allows us to understand the influence of the HB on the O-H σ^* core-excited state. In Fig. S1 (b,c) of the Supplementary information¹⁷ we further investigate the hydrogen bonding by ordering the out-of-plane and in-plane contributions according to hydrogen bond distance in the donating direction of the OH group, and we identify the trend that the σ^* character only barely reaches down to the O-H pre-edge peak even for very distorted hydrogen bonding configurations.

In conclusion, our RIXS measurements of liquid ACA show the absence of a vibrational progression of the OH stretch mode under pre-edge core-excitation of the hydroxyl oxygen contrary to liquid water and methanol. We attribute this behavior to an unusually strong influence of the HB, which changes the character of the intermediate state, and quenches the vibrational progression. The effect allows us to put an upper limit to the average concentration of "end" OH groups in the liquid that are not strongly involved in HBs, implying that the average length of chains must be larger than 3 and/or there must be a significant abundance of cyclic structures. Finally we note that the method exploited here, based on the quenching of the vibrational progression in RIXS, can be used for structure investigations with some generality: We have found that the intensity of the OH vibrational progression can be used as the "ruler" of the chain length. X-ray diffraction²¹ shows that the ratio between dimers and chains is shifted towards dimers in the case of liquid propionic acids compared to ACA, and the proposed structure may be validated with this new method. We envision applications to a variety of liquids, such as formic acid, methyl acetate, and methyl propionate in the near future.

Methods

Experiment. The experiment was performed at room temperature with the SAXES spectrometer²² at the RIXS end station of the ADRESS beam line²³ at the Swiss Light Source at the Paul Scherrer Institut. The liquid acetic acid acquired from Sigma-Aldrich had a purity level $\geq 99\%$ and the water sample was Milli-Q purified and de-ionized. We utilized a flow-cell separating the sample from the vacuum by a 100 nm thick Si_3N_4 window. Due to the risk of window rupture under irradiation, the cell was moved every 5 min. The resonantly scattered photons were detected at a 90° angle from the incoming horizontally polarized X-ray beam by three different detectors with an experimental resolution of ≈ 45 meV for liquid ACA and water. The spectra were measured at several times (3-6 times) at each excitation energy where the duration of each measurement was 5 minutes. The presented spectra were calculated as the sum of these spectra and normalized by dividing by the number of measurements times the number of detectors. To avoid errors from this procedure, the spectra of these individual scans were shifted to same energy scale by using a fit to the elastic line before joining them for further data processing. The energy calibration was based on the O_2 RIXS spectrum²⁴. We compare the intensities of RIXS spectra for the scattering through the carbonyl (O_C) and hydroxyl (O_H) oxygen atoms. The normalization of intensities of these scattering channels was refined by adjusting the spectral intensity to achieve the noise (by the root mean square error) in the signal-free anti-Stokes range of -6 to -2 eV to be equal for these spectra (See Supplementary information¹⁷ and Fig. S5).

Theory. The main part of simulations were performed using post Hartree-Fock ab initio methods accompanied by the techniques based on density functional theory (DFT). We computed the X-ray absorption profile of the ACA monomer using the core-valence separation (CVS) approximation at the algebraic diagrammatic construction (ADC) level of theory. Specifically, the CVS-ADC(2)-x variant was used in combination with the 6-31++G(p,d) basis set, as outlined in Refs.^{8,25}. It is noted that the XAS profile of the ACA monomer as computed using the multi-reference RASPT2 approach displays a good agreement with the ADC(2)-x simulation shown in Fig.2. The RIXS profile was computed using the wave packet software as in Refs.^{6,8,20,26} using the potential energy curves (PECs) along the O-H and C-O stretching modes. The ground and core-excited PECs were determined at the RASSCF/ANO-RCC-VTZP level (with 10 electrons in 13 active molecular orbitals) followed by RASPT2 calculations to account for dynamic electron correlation effects. The full computational details can be found in the Supplementary information¹⁷.

References

1. Soper, A. K. Is water one liquid or two? *The J. Chem. Phys.* **150**, 234503, DOI: <https://aip.scitation.org/doi/full/10.1063/1.5096460> (2019).
2. Fuchs, O. *et al.* Isotope and temperature effects in liquid water probed by x-ray absorption and resonant x-ray emission spectroscopy. *Phys. review letters* **100**, 027801, DOI: <https://doi.org/10.1103/PhysRevLett.100.027801> (2008).

3. Pietzsch, A. *et al.* Snapshots of the Fluctuating Hydrogen Bond Network in Liquid Water on the Sub-Femtosecond Timescale with Vibrational Resonant Inelastic x-ray Scattering. *Phys. Rev. Lett.* **114**, 088302, DOI: [10.1103/PhysRevLett.114.088302](https://doi.org/10.1103/PhysRevLett.114.088302) (2015).
4. Nilsson, A. & Pettersson, L. G. The structural origin of anomalous properties of liquid water. *Nat. communications* **6**, 1–11, DOI: [10.1038/ncomms9998](https://doi.org/10.1038/ncomms9998) (2015).
5. Niskanen, J. *et al.* Compatibility of quantitative X-ray spectroscopy with continuous distribution models of water at ambient conditions. *Proc. Natl. Acad. Sci. United States Am.* **116**, 4058–4063, DOI: [10.1073/pnas.1815701116](https://doi.org/10.1073/pnas.1815701116) (2019).
6. Vaz da Cruz, V. *et al.* Probing hydrogen bond strength in liquid water by resonant inelastic X-ray scattering. *Nat. Commun.* **10**, 1013, DOI: [10.1038/s41467-019-08979-4](https://doi.org/10.1038/s41467-019-08979-4) (2019).
7. Sun, Y.-P. *et al.* Intramolecular soft modes and intermolecular interactions in liquid acetone. *Phys. Rev. B* **84**, 132202, DOI: <https://doi.org/10.1103/PhysRevB.84.132202> (2011).
8. Vaz da Cruz, V. *et al.* Nuclear dynamics in resonant inelastic X-ray scattering and X-ray absorption of methanol. *The J. Chem. Phys.* **150**, 234301, DOI: [10.1063/1.5092174](https://doi.org/10.1063/1.5092174) (2019).
9. Schreck, S. *et al.* Ground state potential energy surfaces around selected atoms from resonant inelastic x-ray scattering. *Sci. reports* **6**, 1–8, DOI: [10.1038/srep20054](https://doi.org/10.1038/srep20054) (2016).
10. Hjelte, I. *et al.* Evidence for ultra-fast dissociation of molecular water from resonant auger spectroscopy. *Chem. physics letters* **334**, 151–158, DOI: [https://doi.org/10.1016/S0009-2614\(00\)01434-2](https://doi.org/10.1016/S0009-2614(00)01434-2) (2001).
11. Riniker, S. *et al.* Temperature dependence of the dielectric permittivity of acetic acid, propionic acid and their methyl esters: a molecular dynamics simulation study. *ChemPhysChem* **13**, 1182–1190, DOI: <https://doi.org/10.1002/cphc.201100949> (2012).
12. Imberti, S. & Bowron, D. T. Formic and acetic acid aggregation in the liquid state. *J. Physics: Condens. Matter* **22**, 404212, DOI: <http://dx.doi.org/10.1088/0953-8984/22/40/404212> (2010).
13. Zhang, M., Chen, L., Yang, H. & Ma, J. Theoretical study of acetic acid association based on hydrogen bonding mechanism. *The J. Phys. Chem. A* **121**, 4560–4568, DOI: <https://doi.org/10.1021/acs.jpca.7b03324> (2017).
14. Chebaane, A., Trabelsi, S., Nasr, S. & Bellissent-Funel, M.-C. Local order in fully deuterated liquid acetic acid as studied by neutron scattering. complementarity to x-ray results. *J. Mol. Liq.* **198**, 204–210, DOI: <https://doi.org/10.1016/j.molliq.2014.06.029> (2014).
15. Nakabayashi, T., Kosugi, K. & Nishi, N. Liquid structure of acetic acid studied by raman spectroscopy and ab initio molecular orbital calculations. *The J. Phys. Chem. A* **103**, 8595–8603, DOI: <https://doi.org/10.1021/jp991501d> (1999).
16. Tabayashi, K., Takahashi, O., Namatame, H. & Taniguchi, M. Substituent r-effects on the core–electron excitation spectra of hydrogen-bonded carboxylic-acid (r-cooh) clusters: Comparison between acetic-acid and formic-acid clusters. *Chem. Phys. Lett.* **557**, 1–9, DOI: <https://doi.org/10.1016/j.cplett.2012.10.043> (2013).
17. See supplementary information at <http://www.nature.com/srep> for AIMD simulation and spectral calculation details.
18. Olbert-Majkut, A., Ahokas, J., Lundell, J. & Pettersson, M. Raman spectroscopy of acetic acid monomer and dimers isolated in solid argon. *J. Raman Spectrosc.* **42**, 1670–1681, DOI: <https://doi.org/10.1002/jrs.2909> (2011).
19. See assignment in Olbert-Majkut *et al.*¹⁸ e.g. Table 1 (HAc monomer) ν_7 : $\nu_{\text{C-O}} + \delta_{\text{COH}} (\text{A}') 1325.0 \text{ cm}^{-1} = 0.164 \text{ eV}$ and Table 2 (HAc dimer) $\nu_{\text{C-O}} + \delta_{\text{COH}} (\text{B}_u) 1309.3 \text{ cm}^{-1} = 0.162 \text{ eV}$.
20. Couto, R. C. *et al.* Selective gating to vibrational modes through resonant X-ray scattering. *Nat. communications* **8**, 12725–12734, DOI: [10.1038/ncomms14165](https://doi.org/10.1038/ncomms14165) (2017).
21. Gorbunova, T., Shilov, V. & Batalin, G. X-ray diffraction study of formic, acetic, and propionic acids in the liquid state. *J. Struct. Chem.* **14**, 388–392, DOI: <https://doi.org/10.1007/BF00746986> (1973).
22. Ghiringhelli, G. *et al.* SAXES, a high resolution spectrometer for resonant x-ray emission in the 400–1600eV energy range. *Rev. Sci. Instruments* **77**, 113108, DOI: [10.1063/1.2372731](https://doi.org/10.1063/1.2372731) (2006).
23. Strocov, V. N. *et al.* Synchrotron Radiation High-resolution soft X-ray beamline ADRESS at the Swiss Light Source for resonant inelastic X-ray scattering and angle-resolved photoelectron spectroscopies. *J. Synchrotron Rad* **17**, 631–643, DOI: [10.1107/S0909049510019862](https://doi.org/10.1107/S0909049510019862) (2010).
24. Hennies, F. *et al.* Resonant inelastic scattering spectra of free molecules with vibrational resolution. *Phys. review letters* **104**, 193002, DOI: <https://doi.org/10.1103/PhysRevLett.104.193002> (2010).

25. Wenzel, J., Holzer, A., Wormit, M. & Dreuw, A. Analysis and comparison of cvs-adc approaches up to third order for the calculation of core-excited states. *The J. chemical physics* **142**, 214104, DOI: <https://doi.org/10.1063/1.4921841> (2015).
26. Gel'mukhanov, F. & Ågren, H. Resonant X-ray Raman scattering. *Phys. reports* **312**, 87–330, DOI: [https://doi.org/10.1016/S0370-1573\(99\)00003-4](https://doi.org/10.1016/S0370-1573(99)00003-4) (1999).
27. Robin, M., Ishii, I., McLaren, R. & Hitchcock, A. Fluorination effects on the inner-shell spectra of unsaturated molecules. *J. Electron Spectrosc. Relat. Phenom.* **47**, 53–92, DOI: [10.1016/0368-2048\(88\)85005-9](https://doi.org/10.1016/0368-2048(88)85005-9) (1988).
28. Tokushima, T. *et al.* Selective observation of the two oxygen atoms at different sites in the carboxyl group (-COOH) of liquid acetic acid. *Phys. Chem. Chem. Phys.* **11**, 1679–1682, DOI: [10.1039/b818812b](https://doi.org/10.1039/b818812b) (2009).

Acknowledgements

The work is supported by the Swedish Research Council (VR). The research work of V.S. is funded by a Swedish Institute scholarship. M.O. acknowledges financial support from the Carl Tryggers Foundation. J.-C.L. thanks for the support by the National Science Foundation of China under grant Nos. 11574082, 11974108 and the Fundamental Research Funds for the Central Universities (No. 2018MS050). F.G., V.K., P.K. and S.P. acknowledge support within the Russian Science Foundation (Project No. 16-12-10109). F.G. acknowledges also the support from the Helmholtz Virtual Institute VI419 "Dynamic Pathways in Multidimensional Landscapes". The synchrotron experiments have been performed at the ADRESS beamline of the Swiss Light Source at the Paul Scherrer Institut (PSI). The work at PSI is supported by the Swiss National Science Foundation through the NCCR MARVEL and the Sinergia project "Mott Physics Beyond the Heisenberg (MPBH) model" (SNSF Research grant numbers CRSII2:141962 and CRSII2:1607651). The research leading to these results has received funding from the European Community's Seventh Framework Programme (FP7/2007–2013) under Grant Agreement No. 290605 (COFUND: PSIFELLOW). The computations were performed on resources provided by the Swedish National Infrastructure for Computing (SNIC) at the Swedish National Supercomputer Center (NSC), the High Performance Computer Center North (HPC2N), and Chalmers Centre for Computational Science and Engineering (C3SE).

Author contributions statement

V.S. performed theoretical simulations of nuclear dynamics, prepared the text of manuscript and figures; V.E. suggested and planned the experiment, collected the data, carried out the data analysis; J.E.R., O.B., C.S., J.G., M.D., A.P., A.F., T.S., D.M., and X.L. participated in the experiment and results discussion; I.E.B., M.O., P.K. performed quantum chemical calculations; J.C.L., S.P., P.N., M.I. participated in theoretical analysis and results discussion; F.G., M.O., and V.K. proposed theoretical model and wrote the paper. All authors reviewed the manuscript.

Additional information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing interests: The authors declare no competing financial interests.

Figure 1. Liquid phase and gas phase RIXS spectra of water⁵ (a) and acetic acid (b,c). The experimental spectra of liquid phase (blue) are compared to computed gas phase spectra (red), obtained using *ab initio* methods as outlined in the text and normalized to $\nu = 1$. The inserts shows the experimental XAS spectra of gas phase (dashed blue) water⁵ (a), acetic acid²⁷ (b,c) and liquid (blue) water⁵ (a), acetic acid²⁸ (b, c), where the vertical arrows point out the corresponding core-excitation energy in XAS. (In gas phase, on top of the resonances. In (a) for liquid water at 535.0 eV. In (b,c) for liquid ACA at 532.0 and 534.0 eV for the $O_C 1s \rightarrow \pi^*$ and $O_H 1s \rightarrow \pi^*$ resonances, respectively). The RIXS intensities of ACA in panels b and c are matched to each other, as it is shown in Fig. S5¹⁷. The energy loss is the difference between the energies of incoming and emitted X-ray photons.

Figure 2. CVS-ADC(2)-x calculated XAS spectra of (a) the acetic acid monomer, (b) regular dimer and (c) inverted dimer, showed in comparison to measured XAS spectra of (d) acetic acid liquid from Ref.²⁸ and (e) gas phase from Ref.²⁷. The experimental and theoretical XAS spectra are shown in blue and red, respectively. The calculated oscillator strengths are shown as bar graphs, where each bar is coloured according to the atom of origin depicted in the molecular structure next to each spectrum. The XAS spectra have been obtained by broadening the bar graphs using Gaussian functions of 0.5 eV full width at half maximum. The high photon energy region (537 eV and above) of the dimer spectra is missing. This is due to the high computational cost of the CVS-ADC(2)-x method, where for the dimers we could only compute a limited number of excitation vectors, allowing the full description of only the first two experimental XAS peaks. In the case of the monomer, we could instead include enough excitation vectors to cover the full photon energy window used in the experiment.

Figure 3. The potential energy curves from RASPT2 calculations of acetic acid in the ground, $1s_{O_C}^{-1}$ (a) and $1s_{O_H}^{-1}$ (b) core-excited states along the C=O and O-H stretching normal modes, respectively. Only the lowest bound $1s_{O_C}^{-1}\pi^*$ contributes to the first XAS peak, and both close lying bound $1s_{O_H}^{-1}\pi^*$ and dissociative $1s_{O_H}^{-1}\sigma^*$ states form the second XAS peak of acetic acid monomer (Fig. 2). The oscillatory strengths are almost the same for the $1s_{O_H}^{-1}$ core-excited states and they are about four times smaller as compared to the $1s_{O_C} \rightarrow \pi^*$ core-excitation (Fig. 2).

Figure 4. XAS from DFT calculations of the O_C (blue) and O_H (red) K-edges in gas phase (a) and liquid (b) ACA are shown and further decomposed into out-of-plane (π^* -type, dotted) and in-plane (σ^* -type, dashed-dotted) contributions. The spectra are computed using the XFH transition potential method and AIMD approach (see for more details the Supplementary information¹⁷). (c) A cyclic structure without "end" OH group. (d) A chain structure of liquid acetic acid contains an "end" OH group.

Figures

Figure 1

Liquid phase and gas phase RIXS spectra of water⁵ (a) and acetic acid (b,c). The experimental spectra of liquid phase (blue) are compared to computed gas phase spectra (red), obtained using ab initio methods as outlined in the text and normalized to $v = 1$. The inserts show the experimental XAS spectra of gas phase (dashed blue) water⁵ (a), acetic acid²⁷ (b,c) and liquid (blue) water⁵ (a), acetic acid²⁸ (b, c), where the vertical arrows point out the corresponding core-excitation energy in XAS. (In gas phase, on top of the resonances. In (a) for liquid water at 535.0 eV. In (b,c) for liquid ACA at 532.0 and 534.0 eV for the OC $1s \rightarrow \pi^*$ and OH $1s \rightarrow \pi^*$ resonances, respectively). The RIXS intensities of ACA in panels b and c are matched to each other, as it is shown in Fig. S517. The energy loss is the difference between the energies of incoming and emitted X-ray photons.

Figure 2

CVS-ADC(2)-x calculated XAS spectra of (a) the acetic acid monomer, (b) regular dimer and (c) inverted dimer, showed in comparison to measured XAS spectra of (d) acetic acid liquid from Ref.28 and (e) gas phase from Ref.27. The experimental and theoretical XAS spectra are shown in blue and red, respectively. The calculated oscillator strengths are shown as bar graphs, where each bar is coloured according to the atom of origin depicted in the molecular structure next to each spectrum. The XAS spectra have been

obtained by broadening the bar graphs using Gaussian functions of 0.5 eV full width at half maximum. The high photon energy region (537 eV and above) of the dimer spectra is missing. This is due to the high computational cost of the CVS-ADC(2)-x method, where for the dimers we could only compute a limited number of excitation vectors, allowing the full description of only the first two experimental XAS peaks. In the case of the monomer, we could instead include enough excitation vectors to cover the full photon energy window used in the experiment.

Figure 3

The potential energy curves from RASPT2 calculations of acetic acid in the ground, $1s-1O_C$ (a) and $1s-1O_H$ (b) core-excited states along the C=O and O-H stretching normal modes, respectively. Only the lowest bound $1s-1O_C \pi^*$ contributes to the first XAS peak, and both close lying bound $1s-1O_H \pi^*$ and dissociative $1s-1O_H \sigma^*$ states form the second XAS peak of acetic acid monomer (Fig. 2). The oscillatory strengths are almost the same for the $1s-1O_H$ core-excited states and they are about four times smaller as compared to the $1s-1O_C \pi^*$ core-excitation (Fig. 2).

Figure 4

XAS from DFT calculations of the OC (blue) and OH (red) K-edges in gas phase (a) and liquid (b) ACA are shown and further decomposed into out-of-plane (π^* -type, dotted) and in-plane (σ^* -type, dashed-dotted) contributions. The spectra are computed using the XFH transition potential method and AIMD approach (see for more details the Supplementary information17). (c) A cyclic structure without "end" OH group. (d) A chain structure of liquid acetic acid contains an "end" OH group.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SciRepACAMS.pdf](#)