

Separation of Phenyl Acetic Acid and 6-Aminopenicillanic Acid Applying Aqueous Two-Phase Systems Based on Copolymers and Salts

Farzaneh Ghazizadeh Ahsaie

Amirkabir University of Technology

Gholamreza Pazuki (✉ ghpazuki@aut.ac.ir)

Amirkabir University of Technology

Research Article

Keywords: Aqueous two-phase systems, Copolymers, Partitioning, 6-APA, 6-Aminopenicillanic acid, Phenyl acetic acid

Posted Date: December 3rd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-113249/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on February 10th, 2021. See the published version at <https://doi.org/10.1038/s41598-021-82476-x>.

Separation of Phenyl Acetic Acid and 6-Aminopenicillanic Acid Applying Aqueous Two-Phase Systems based on Copolymers and Salts

Farzaneh Ghazizadeh Ahsaie¹, Gholamreza Pazuki¹

¹Department of Chemical Engineering, Amirkabir University of Technology (Tehran Polytechnic),
Tehran, Iran

* Corresponding author: Tel: +98-021-64543159. Fax: +98-021-66405847. E-mail address:
ghpazuki@aut.ac.ir (G.R. Pazuki)

Abstract

6-Aminopenicillanic acid (6-APA) is used for synthesis of semisynthetic antibiotics. Polymer-salt aqueous two-phase systems (ATPSs) were applied for separation of 6-aminopenicillanic acid (6-APA) and phenyl acetic acid (PAA), as the products of hydrolyzation reaction of Penicillin G / Penicillin V. The binodal curves of ATPS composed of a copolymer (reverse Pluronic 10R5, Pluronic L35 and PEG-ran-PPG) and a salt (Tri-sodium citrate, tri-potassium citrate, di-potassium phosphate, sodium sulphate and magnesium sulphate) were obtained. The results show that, at a fixed PPG/PEG ratio, block copolymers have larger two-phase region compared with random copolymer. After screening on the partition coefficient of PAA and 6-APA separately, Na_2SO_4 was selected for studying the effect of the copolymer structure and the composition of salt and copolymer on partitioning, considering higher selectivity of PAA and 6-APA. 10R5- Na_2SO_4 ATPS was selected as the most appropriate system for separation of 6-APA and PAA. This system was used for separation of mixture of 6-APA and PAA. The results show that selectivity was ≈ 53 and smaller in a system, containing a mixture of 6-APA and PAA. This observation can be justified by the interaction between 6-APA and PAA. Molecular interaction between these two molecules were investigated by the Flory-Huggins interaction parameter.

Keywords: Aqueous two-phase systems, Copolymers, Partitioning, 6-APA, 6-Aminopenicillanic acid, Phenyl acetic acid

1. Introduction

Penicillin was firstly discovered by Fleming in 1932. They are widely used in antimicrobial therapy due to their high bacterial activity, low toxicity and good medical efficacy [1]. In order to decelerate the existence of drug-resistance bacteria and ensure the drug influence, now 6-aminopenicillanic acid (6-APA) is used as basic raw material for synthesis of semisynthetic antibiotics [1–4]. 6-APA was discovered in late 1950s [5]. It can be produced by chemical and enzymatic methods. Toxic materials such as pyridine, NOCl and PCl_5 are used in chemical method which can cause environmental problems. Compare to chemical method, the enzymatic method has several advantages: mild environment, high capacity, low cost and safety [1]. In this method Penicillin G / Penicillin V are hydrolyzed by penicillin acylase to produce 6-APA [4]. The by-product of this synthesis is phenyl acetic acid. Despite the advantage of this method, solubility of Penicillin G, penicillin acylase, 6-APA and PAA cause problems for purification [4].

In 1896, Beijerinck first observed a phase separation in aqueous solution of starch and gelatin. For the first time, Albertsson applied aqueous two phase system (ATPS), for partitioning of plant cell particles [6–8]. ATPSs can be formed by mixing of two compounds (polymer, salt, ionic liquid, surfactant, to name but a few) in water above a critical concentration [6]. These systems are environment-friendly, non-flammable, scalable, efficient for many purifications and separations and low energy consumption [6,7,9,10]. 6-APA and phenylacetic acid are usually separated from the hydrolysate by precipitation at its pI (isoelectric pH) and n-butyl acetate solvent extraction, respectively. Instead of conventional solvent extraction, aqueous two-phase systems can be an alternative.

The polymer-salt ATPSs are more attractive due to their low viscosity, fast phase separation, low cost and higher selectivity [11]. Based on the components of ATPS and their composition, several mechanisms for phase separation was suggested, like: salting-out, hydrophilic and hydrophobic

interactions. In polymer-salt ATPS, salting out and hydrophobic interactions are considered as dominant mechanisms [12].

The difference in physicochemical properties of the top and the bottom phases, cause an uneven distribution of biomolecules in ATPS. One of the factors affecting the partitioning of the solute is the phase-forming properties [13]. Copolymers are consisted of two or more monomers. Their properties could be adjusted by manipulating in number of monomers [14]. One of the most widely used copolymers is ethylene glycol (PEG) - propylene glycol (PPG). These copolymers can be in a random (UCON) or a block (normal Pluronic and reverse Pluronic (R-pluronic)) form. Using copolymers as phase forming compounds in ATPSs causes an extreme partitioning, hence selectivity and efficiency increase [15]. Self-assembly of amphiphilic copolymers (block copolymers) in aqueous solutions, enhance the interaction between copolymer and hydrophobic compounds [16,17]. Svensson and et al. indicated that self-assembly of copolymers is not a dominant factor in the partitioning of hydrophilic proteins [18].

In this paper, the application of three different PEG-PPG copolymers (random, normal and R-Pluronic) in 6-APA and PAA purification were investigated. Phase behaviors of these copolymers with five different salts were studied. To study the capacity of these ATPSs for separation of both biomolecules, we started by studying the partition behavior of each drug. The effect of salt (anion and cation) and copolymer structure (random, normal and reverse) on binodal curves and partitioning of 6-APA and PAA were investigated. The influence of concentration of salt and copolymers were studied in Na₂SO₄-copolymer ATPS. The partition behavior and separation of 6-APA and PAA were investigated for a mixture of them. In order to study the interaction between 6-APA and PAA, ¹H-NMR and Flory-Huggins interaction parameter (χ_{12}) were applied. The Flory-Huggins interaction parameter was calculated based on the Hansen solubility parameter.

2. Materials and methods

2.1. Materials

Pluronic L35 (Mw=1900), R-Pluronic 10R5 (Mw=2000) (PO-EO-block polymers with approximately 50% EO), PEG-ran-PPG-ran-PEG (UCON) from *Leuconostoc spp*, Tri-sodium citrate, tri-potassium citrate and di-potassium phosphate were purchased from Sigma. Sodium sulphate and magnesium sulphate were acquired from Merck. Phenyl acetic acid and 6-aminopenicillanic acid were acquired from TITRACHEM with mass fraction purity of 98%, prepared in double distilled water.

2.2. Methods

2.2.1 Phase diagrams and tie-lines

All phase diagrams were determined at two temperatures (298 and 283 K) and atmospheric pressure through cloud-point titration method described in ref [19]. The experimental data were correlated with Merchuk equation (1). The adjustable parameters (A, B, C) were determined by regression of experimental data.

$$[\text{copolymer}] = A \times \exp[(B \times [\text{Salt}]^{0.5}) - (C \times [\text{salt}]^3)] \quad (1)$$

The composition of each phase was calculated by Merchuck equation and applying gravimetric method [20] (equations (2-5)).

$$[\text{copolymer}]_{Top} = A \times \exp[(B \times [\text{salt}]_{Top}^{0.5}) - (C \times [\text{salt}]_{Top}^3)] \quad (2)$$

$$[\text{copolymer}]_{Bot} = A \times \exp[(B \times [\text{salt}]_{Bot}^{0.5}) - (C \times [\text{salt}]_{Bot}^3)] \quad (3)$$

$$[\text{copolymer}]_{Top} = \frac{[\text{copolymer}]_F}{\alpha} - \frac{1-\alpha}{\alpha} [\text{copolymer}]_{Bot} \quad (4)$$

$$[\text{salt}]_{Top} = \frac{[\text{salt}]_F}{\alpha} - \frac{1-\alpha}{\alpha} [\text{salt}]_{Bot} \quad (5)$$

The subscripts F, Top and Bot stand for feed, top and bottom phases, respectively. α is the ratio of top and bottom phase mass. Tie-line length and slope (TLL and STL, respectively) are obtained through equations (6-7).

$$TLL = \sqrt{([copolymer]_{TOP} - [copolymer]_{BOT})^2 + ([salt]_{TOP} - [salt]_{BOT})^2} \quad (6)$$

$$STL = \frac{[copolymer]_{TOP} - [copolymer]_{BOT}}{[salt]_{TOP} - [salt]_{BOT}} \quad (7)$$

2.2.2 Partitioning of PAA and 6-APA

The salt-copolymer ATPSs were prepared gravimetrically ($u(m) = 10^{-4}$ g). The mixtures were stirred vigorously and left to reach thermodynamic equilibrium and complete phase separation for at least 24 h at 298 K. Both phases (copolymer-rich and salt -rich phase) were carefully separated and weighed to determine biomolecule concentrations. In this work, the partition coefficient (K_{drug}), the percentage extraction efficiency ($EE\%$) and selectivity (Sel) were defined following Eqs. (8-10):

$$K_{drug} = \frac{[drug]_{TOP}}{[drug]_{BOT}} \quad (8)$$

$$EE\% = \frac{K_{drug}V_r}{K_{drug}V_r + 1} \times 100 \quad (9)$$

$$Sel = \frac{K_{drug_{PAA}}}{K_{drug_{6-APA}}} \quad (10)$$

where $[drug]$ and V_r represent the concentration of drug and the volume ratio between top and bottom phases.

Phenyl acetic acid (PAA) and 6-aminopenicillanic acid (6-APA) were determined by UV Spectroscopy at 254 and 207 nm, respectively, using previously determined calibration curves. In order to eliminate the interference of phase forming components, ternary mixtures at specific

concentrations of each component were prepared, using pure water instead of the aqueous solutions of drugs, to be used as blank solution. Three replicates were prepared for each assay, the results were reported as average of triplicated calculated partition coefficient. In order to study the partition coefficient in system contains both 6-APA and PAA, HPLC has been applied. The HPLC procedure is explained in ref. [21].

2.2.3 Intermolecular Interaction

In order to investigate the molecular interaction between PAA and 6-APA, the Hansen solubility parameter has been applied.

The Hansen Solubility parameter has been used to calculate the Flory-Huggins interaction parameter between PAA and 6-APA as an important parameter for interaction estimation [22]. The Hansen Solubility parameter includes three contributions: dispersive (δ_d), polar (δ_p), and hydrogen bonding (δ_h); as defined in the following:

$$\delta_d = \frac{\sum_i F_{d_i}}{\sum_i V_i} \quad (11)$$

$$\delta_p = \frac{\left(\sum_i F_{p_i}^2 \right)^{0.5}}{\sum_i V_i} \quad (12)$$

$$\delta_h = \frac{\left(\sum_i E_{h_i} \right)^{0.5}}{\sum_i V_i} \quad (13)$$

Where $F_{d,i}$, $F_{p,i}$ and $F_{h,i}$ are the group contribution of dispersion forces, polar forces and hydrogen bond energy, respectively and V_i is the group contribution of structural group i to molar volume.

The solubility parameter can be obtained applying equations (11) to (13). The Flory-Huggins

interaction parameter can be obtained following equation (15):

$$\Delta\delta = \sqrt{(\delta_{d,1} - \delta_{d,2})^2 + 4(\delta_{p,1} - \delta_{p,2})^2 + (\delta_{h,1} - \delta_{h,2})^2} \quad (14)$$

$$\chi_{12} = (\Delta\delta)^2 V_1 / RT \quad (15)$$

3. Results

3.1. Binodal curves determination

An aqueous mixture of copolymer and salt split into two phases above certain concentrations of components. The nature of each phase forming can affect the two-phase region. In this work, the effect of the structure of copolymer, temperature and salt anion/cation on the binodal curves were studied. The experimental data were fitted by Merchuk equation (Eq. 1). The Merchuk adjustable parameters are reported in tables 1 and 2.

Table 1. Merchuk parameters and correlation coefficient (R^2) obtained by fitting Eq. 1. (298 K)

Salt	Copolymer	$A \pm \sigma$	$B \pm \sigma$	$10^3(C \pm \sigma)$	R^2
$MgSO_4$	Pluronic L35	52.1 ± 6.7	-0.382 ± 0.085	0.34 ± 0.2	0.997
	UCON	45.1 ± 6.8	-0.279 ± 0.090	0.20 ± 0.11	0.998
	Pluronic 10R5	64.4 ± 5.4	-0.472 ± 0.060	0.39 ± 0.14	0.993
$Na_3C_6H_5O_7$	Pluronic L35	61.1 ± 8.2	-0.393 ± 0.122	1.22 ± 0.83	0.991
	UCON	66.6 ± 5.3	-0.393 ± 0.066	0.50 ± 0.18	0.998
	Pluronic 10R5	71.4 ± 6.4	-0.520 ± 0.079	0.89 ± 0.09	0.996
Na_2SO_4	Pluronic L35	77.1 ± 3.5	-0.541 ± 0.049	0.32 ± 0.01	0.995
	UCON	86.0 ± 13.8	-0.655 ± 0.193	1.35 ± 0.11	0.999
	Pluronic 10R5	83.6 ± 9.0	-0.740 ± 0.126	2.5 ± 0.17	0.998
K_2HPO_4	Pluronic L35	89.9 ± 14.3	-0.775 ± 0.145	2.64 ± 0.17	0.988
	UCON	68.9 ± 8.0	-0.499 ± 0.106	1.10 ± 0.59	0.998

	Pluronic 10R5	88.2 ± 13.2	-0.740 ± 0.123	1.78 ± 0.92	0.994
K₃C₆H₅O₇	Pluronic L35	71.9 ± 3.8	-0.481 ± 0.038	0.60 ± 0.12	0.999
	UCON	75.6 ± 9.1	-0.458 ± 0.118	0.80 ± 0.64	0.984
	Pluronic 10R5	99.0 ± 8.1	-0.627 ± 0.054	0.66 ± 0.19	0.998


Table 2. Merchuk parameters and correlation coefficient (R^2) obtained by fitting Eq. 1. (283 K)

Salt	Copolymer	$A \pm \sigma$	$B \pm \sigma$	$10^3(C \pm \sigma)$	R^2
MgSO₄	Pluronic L35	73.7 ± 2.2	-0.380 ± 0.018	0.70 ± 0.02	0.994
	UCON	69.1 ± 1.2	-0.438 ± 0.009	0.13 ± 0.01	0.999
	Pluronic 10R5	72.4 ± 1.7	-0.487 ± 0.0130	0.18 ± 0.01	0.998
Na₃C₆H₅O₇	Pluronic L35	66.5 ± 1.2	-0.368 ± 0.011	0.72 ± 0.02	0.998
	UCON	68.1 ± 1.3	-0.408 ± 0.012	0.37 ± 0.01	0.997
	Pluronic 10R5	74.4 ± 1.6	-0.484 ± 0.014	0.6 ± 0.03	0.997
Na₂SO₄	Pluronic L35	64.1 ± 10.1	-468 ± 0.103	2.19 ± 0.44	0.997
	UCON	81.7 ± 2.7	-0.634 ± 0.021	1.2 ± 0.00	1.000
	Pluronic 10R5	74.4 ± 2.9	-0.642 ± 0.025	2.61 ± 0.10	1.000
K₂HPO₄	Pluronic L35	65.6 ± 2.0	-0.405 ± 0.018	1.50 ± 0.05	0.999
	UCON	85.6 ± 2.3	-0.571 ± 0.016	0.86 ± 0.04	0.998
	Pluronic 10R5	82.2 ± 2.5	-0.622 ± 0.019	1.10 ± 0.05	0.999
K₃C₆H₅O₇	Pluronic L35	62.9 ± 1.0	-0.328 ± 0.008	0.38 ± 0.00	0.999
	UCON	84.8 ± 7.6	-0.402 ± 0.036	0.20 ± 0.02	0.998
	Pluronic 10R5	69.4 ± 1.7	-0.385 ± 0.014	0.48 ± 0.02	0.998

Figures 1 and 2 represent the experimental binodal curves. To eliminate the interference of molecular weight of the different phase forming components on the analysis, the binodal curves are represented in molality scale (figure S1 and S2). The results show that two-phase region expands by increasing the temperature. The solubility and hydrophilicity of the copolymers decrease by increasing the temperature. At higher temperature lower amount of copolymer is needed to form two-phase system. In order to study the effect of salt cation on the binodal curve of copolymer/salt ATPS, $\text{MgSO}_4 / \text{Na}_2\text{SO}_4$ and $\text{Na}_3\text{C}_6\text{H}_5\text{O}_7 / \text{K}_3\text{C}_6\text{H}_5\text{O}_7$ were selected. ATPSs based on MgSO_4 formed at higher concentration of phase forming components compared to Na_2SO_4 . This shows that Na^+ ion is more able to form two phases compared to Mg^{2+} . The salting-out ability of ions can be described based on Hofmeister series and free energy hydration (ΔG_{hyd}). The ΔG_{hyd} values of Mg^{2+} and Na^+ are $-1830 \text{ kJ mol}^{-1}$ and -365 kJ mol^{-1} , respectively [23]. More negative ΔG_{hyd} value for ions leading to more salting-out ability. Although the Hofmeister series and ΔG_{hyd} show that Mg^{2+} has more salting-out effect compared to Na^+ but the obtained results depicted a reverse trend. It should be mentioned that cations with higher valance have more interactions with ether oxygen of PEG monomers in the copolymer structure leading to more salting-in [19]. Moreover, several authors had reported that Na_2SO_4 is more able than MgSO_4 to form ATPSs [19,23–25]. According to experimental binodal curves of $\text{Na}_3\text{C}_6\text{H}_5\text{O}_7$ and $\text{K}_3\text{C}_6\text{H}_5\text{O}_7$ salts, Na^+ shows more salting-out effect than K^+ . This observation is in agreement with Hofmeister series and ΔG_{hyd} ($\Delta G_{hyd,K^+} = -295 \text{ kJmol}^{-1}$) [26–29]. In order to investigate the effect of anion on the binodal curves, $\text{Na}_3\text{C}_6\text{H}_5\text{O}_7 / \text{Na}_2\text{SO}_4$ and $\text{K}_3\text{C}_6\text{H}_5\text{O}_7 / \text{K}_2\text{HPO}_4$ were chosen. The Hofmeister series for anions follows as [29]: $\text{HPO}_4^{2-} < \text{SO}_4^{2-} < \text{C}_6\text{H}_5\text{O}_7^{3-}$. The reported values of ΔG_{hyd} of HPO_4^{2-} , $\text{C}_6\text{H}_5\text{O}_7^{3-}$ and SO_4^{2-} are -1125 , -2793 and -1080 kJmol^{-1} , respectively. The obtained results depicted that both SO_4^{2-} and HPO_4^{2-} has larger salting out effect than $\text{C}_6\text{H}_5\text{O}_7^{3-}$. It was expected that $\text{C}_6\text{H}_5\text{O}_7^{3-}$ salt, to be more effective in salting-out due to its position in

Hofmeister series and more negative ΔG_{hyd} , but the experimental results show that $C_6H_5O_7^{3-}$ provided smaller two phase region. This behavior was observed in several literatures [26,27,30] .

Figure 1. Experimental binodal curves determined for the ATPSs composed of different copolymer + salt at 298 K






Figure 2. Experimental binodal curves determined for the ATPSs composed of different copolymer + salt at 283 K

Various structures of PEG-PPG copolymers (Pluronic L35, R-Pluronic 10R5 and UCON) contain 50% wt of PEG were investigated. The tendency of phase separation follows as: R-Pluronic 10R5 \approx Pluronic L35 > UCON. This trend shows that UCON is more hydrophilic, while the hydrophilicity of R-Pluronic 10R5 and Pluronic L35 is similar. Wu et al studied the effect of position of PEG and PPG in block copolymer structure on hydrophilic - lipophilic balance (HLB) and relative solubility number (RSN) [31]. RSN and HLB depend on molecular weight, $\frac{\text{PEG}}{\text{PPG}}$ ratio and copolymer structure. They showed that Pluronic copolymers has higher HLB and RSN than

corresponding R-Pluronic copolymers. The higher values of HLB and RSN the larger hydrophilicity of copolymers. They also reported that the difference between RSN value is not significant for small copolymers (molecular weight < 3000). Reported RSN and HLB values for Pluronic L35, R-Pluronic 10R5 are reported in table 3. As they have reported the studied block copolymers are similar in hydrophilicity and molecular weight leading to little difference in their binodal curve.

Table 3. RSN and HLB values of Pluronic L35 and R-Pluronic 10R5

Copolymer	RSN	HLB
Pluronic L35	22.3*	18-23**
R-Pluronic 10R5	21.0*	12-18**

* ref. [31]; ** ref.[32]

Several tie-lines were determined for each system at 298 K. The equilibrium concentrations, TLL and STL are reported in table S1-S3. The changes in STL are due to the difference hydrophobicity of the copolymers. In the ATPSs of Pluronic L35, R-Pluronic 10R5 and UCON the absolute value of STL follows: Pluronic L35 \approx R-Pluronic 10R5 > UCON.

3.2. Distribution of biomolecule

In order to evaluate the capacity of these ATPS in separation of 6APA and PAA, the partition behavior of 6APA and PAA were studied separately. According to the binodal curves at two different temperature, higher temperature was selected due to bigger two-phase region. The mixture point of 9 and 20 wt% of salt and copolymer, respectively were selected. Since this mixture point in potassium citrate- UCON ATPS is located in one phase region, (13, 20) mixture point was studied for this salt. The partition coefficients were reported in table 4. PAA and 6-APA are considered as hydrophobic and hydrophilic molecules, respectively, due to their octanol-water

partition coefficients $\log K_{OW,PAA} = 1.43$ [33] and $\log K_{OW,6-APA} = -0.8$ [34]. As it is obvious, PAA has more affinity to the top phase (hydrophobic phase). In all studied ATPSs, except in K_2HPO_4 and $K_3C_6H_5O_7$ based one, the hydrophilic molecule (6-APA) is more concentrated in the bottom phase as Vobecka and co-workers reported that partition coefficient of 6-APA was greater than one in K_2HPO_4 -PEG ATPS [35]. Since K_2HPO_4 and $K_3C_6H_5O_7$ are two basic salts, pH value of the ATPSs based on them is greater than one. 6-APA has no net charge at its Isoelectric pH ($pI=3.9$) [36], but it is negatively charged in K_2HPO_4 and $K_3C_6H_5O_7$ ($pH > pI$). 6-APA shows more affinity to the top phase in K_2HPO_4 and $K_3C_6H_5O_7$ based ATPSs, despite the more hydrophobicity of top phase. Haghtalab et al. observed similar behavior for the partitioning of α -amylase in PEG - K_2HPO_4 ATPS. They reported that increasing pH value, causes an enhancement in the partition coefficient of α -amylase ($pH > pI=5.4$) [37]. Several authors have reported that negatively charged molecules show more affinity to the (PEG/ copolymer PEG-PPG)-rich phase since PEG polymer behaves as positively charged molecule [38–40]. In all studied systems, Na_2SO_4 -ATPSs have highest partition coefficient and lowest water ratio between top and bottom phases. Water ratio values are reported in table S4. All the reported values are lower than one (less water in the top phase) hence the top phase is more hydrophobic.

Table 4. Partition coefficient (K), extraction efficiency (EE%) and selectivity (Sel.) of 6-APA and PAA in systems based in copolymers + salts + water.

Copolymer	Salt	$K_{6-APA} \pm \sigma$	$K_{PAA} \pm \sigma$	$EE_{6-APA} \pm \sigma$	$EE_{PAA} \pm \sigma$	Sel.
Pluronic 10R5	K ₃ C ₆ H ₅ O ₇	1.04 ± 0.07	4.4 ± 0.20	66.62 ± 1.68	75.95 ± 0.95	4.60
	K ₂ HPO ₄	1.49 ± 0.01	5.99 ± 0.40	49.85 ± 0.17	80.20 ± 1.27	4.05
	Na ₃ C ₆ H ₅ O ₇	0.2 ± 0.10	2.26 ± 0.38	89.43 ± 5.67	54.60 ± 8.29	10.42
	MgSO ₄	0.52 ± 0.06	4.36 ± 0.12	89.09 ± 1.23	60.88 ± 0.66	8.47
	Na ₂ SO ₄	0.78 ± 0.08	19.95 ± 0.44	60.89 ± 2.51	94.14 ± 0.14	24.92
Pluronic L35	K ₃ C ₆ H ₅ O ₇	1.01 ± 0.08	6.24 ± 0.40	66.62 ± 1.76	76.89 ± 1.21	6.64
	K ₂ HPO ₄	1.7 ± 0.01	4.8 ± 0.08	49.26 ± 0.07	74.27 ± 0.33	2.80
	Na ₃ C ₆ H ₅ O ₇	0.47 ± 0.20	1.79 ± 0.38	82.21 ± 8.05	46.24 ± 5.20	3.85
	MgSO ₄	0.44 ± 0.00	5.90 ± 0.90	90.61 ± 0.01	55.82 ± 3.01	12.30
	Na ₂ SO ₄	0.52 ± 0.08	49.00 ± 8.60	72.98 ± 0.70	97.06 ± 0.49	91.52
UCON	K ₃ C ₆ H ₅ O ₇	3.30 ± 0.01	4.97 ± 0.13	35.39 ± 3.79	73.45 ± 1.57	1.49
	K ₂ HPO ₄	2.12 ± 0.19	6.14 ± 0.28	45.66 ± 2.40	77.05 ± 1.28	2.81
	Na ₃ C ₆ H ₅ O ₇	0.40 ± 0.10	1.20 ± 0.02	82.47 ± 0.00	47.05 ± 11.47	4.70
	MgSO ₄	0.58 ± 0.01	6.80 ± 0.20	89.91 ± 0.16	56.80 ± 2.09	11.76
	Na ₂ SO ₄	0.70 ± 0.01	16.6 ± 0.50	62.73 ± 0.50	93.81 ± 2.55	30.07

After the investigation of the effect of salt, Na_2SO_4 was selected for further study, considering higher partition coefficient for PAA and higher selectivity. 5 mixture points (table 5) were chosen to study the effect of salt and copolymer weight fraction. The results are shown in table 5. An enhancement in copolymer and salt weight fraction causes an increase in partition coefficient of PAA.

Although Pluronic L35 has higher value of HLB and RSN which lead to more hydrophilicity but the hydrophobic molecule (PAA) showed more tendency to the top phase in ATPS based on Pluronic L35 compared with Pluronic 10R5. This observation can be explained by the higher ability of Pluronic L35 in micellization than Pluronic 10R5. Self-aggregation behavior in reverse and normal Pluronic is different [41]. Normal Pluronic forms star-like micelles, while R-Pluronic can form flower-like or random network micelles [42,43]. Micellization is a more favorable process in normal Pluronic solution compared with reverse one [42]. Aggregation number and CMC in R-Pluronic solution is smaller and higher than normal Pluronic [44], respectively. The obtained results show that R-Pluronic has lower ability in increasing the solubility of hydrophobic molecules than normal Pluronic. Similar results were reported for solubility and partitioning of toluene and 1,2-dichloroethane (hydrophobic molecules) in normal and R-Pluronic solutions [45,46].

Table 5. Weight percentage composition of each mixture point tested ($W_t\%$), partition coefficient (K), extraction efficiency (EE%) and selectivity (Sel.) for systems based in copolymers and Na₂SO₄.

[salt copolymer] $W_t\%$	Pluronic 10R5			Pluronic L35			UCON		
	$K_{PAA} \pm \sigma$	$K_{6-APA} \pm \sigma$	Sel.	$K_{PAA} \pm \sigma$	$K_{6-APA} \pm \sigma$	Sel.	$K_{PAA} \pm \sigma$	$K_{6-APA} \pm \sigma$	Sel.
[9 20]	19.54 ± 0.65	0.78 ± 0.08	25	38.83 ± 0.85	0.43 ± 0.02	91	21.00 ± 1.63	0.71 ± 0.02	30
[9 25]	26.15 ± 0.12	0.25 ± 0.02	104	40.87 ± 0.62	0.47 ± 0.03	87	27.42 ± 1.11	0.63 ± 0.03	44
[9 15]	18.92 ± 1.27	0.15 ± 0.02	123	11.45 ± 0.22	0.25 ± 0.09	46	13.05 ± 0.41	0.54 ± 0.02	24
[7 20]	16.73 ± 5.57	0.58 ± 0.11	29	20.20 ± 1.18	0.54 ± 0.00	37	11.35 ± 0.58	0.43 ± 0.05	26
[11 20]	41.44 ± 1.42	0.33 ± 0.01	124	39.67 ± 0.94	0.64 ± 0.01	62	27.07 ± 2.32	0.39 ± 0.10	69
	$EE_{PAA} \pm \sigma$	$EE_{6-APA} \pm \sigma$		$EE_{PAA} \pm \sigma$	$EE_{6-APA} \pm \sigma$		$EE_{PAA} \pm \sigma$	$EE_{6-APA} \pm \sigma$	
[9 20]	94.14 ± 0.14	60.89 ± 2.51		97.06 ± 0.49	72.98 ± 0.70		93.81 ± 2.55	62.73 ± 0.50	
[9 25]	96.61 ± 0.08	78.61 ± 1.47		97.79 ± 0.18	66.42 ± 1.19		96.48 ± 0.13	61.37 ± 1.23	
[9 15]	91.20 ± 0.55	92.23 ± 0.96		86.71 ± 0.55	87.86 ± 4.03		89.73 ± 0.29	73.56 ± 0.69	
[7 20]	94.19 ± 0.15	64.14 ± 4.20		95.68 ± 0.15	62.59 ± 0.20		91.89 ± 0.37	70.01 ± 2.40	
[11 20]	96.52 ± 0.12	81.75 ± 0.42		96.09 ± 0.12	71.49 ± 0.39		94.74 ± 0.43	79.35 ± 4.04	

According to the reported partition coefficient for PAA and 6-APA in table 5, the maximum selectivity is greater than 100. To study the capacity of these ATPS for simultaneous partitioning and separation of PAA and 6-APA in mixture, Pluronic 10R5 – Na₂SO₄ ATPS was selected considering higher selectivity (≈ 120). The obtained results are reported in table 6. The results show that selectivity of PAA and 6-APAA in this ATPS is approximately 53. Reduction in selectivity shows that there is an interaction between 6-APA and PAA.

Table 6. Weight percentage composition ($W_t\%$), partition coefficient (K), extraction efficiency (EE%) and selectivity (Sel.) for a mixture of PAA and 6-APA in 10R5- Na₂SO₄

ATPS (wt%)	K_{PAA}	K_{6-APA}	Sel.
10R5 – Na₂SO₄ [11 20]	39.62	0.74	53.54

3.3 Intermolecular Interaction

The Flory-Huggins interaction parameter (χ_{ij}) can be used to study the compatibility between two components. Table 7 represents the values of χ_{12} . The lower value of χ_{12} , the higher compatibility and more favorable interaction for PAA-X pair (X: copolymer, water, 6-APA) [47,48]. The compatibility with PAA follows the order: 6-APA > copolymer > water. It can be concluded that $K_{PAA} > 1$ and higher concentration of PAA in copolymer-rich phase is due to the higher affinity of PAA to copolymer. The smaller selectivity for separation of a concentrated mixture of PAA and 6-APA can be justified by the affinity of 6-APA with PAA.

Table 7. Hansen solubility parameters and χ_{12} values for all compounds at 298 K

	δ_d	δ_p	δ_h	δ_t	χ_{12}
PAA	21.23	4.45	10.12	23.94	-
6-APA	27.09	10.74	16.14	33.31	0.21
Copolymer	17.22	0.92	9.39	19.63	0.74
water	15.5	16.0	42.3	47.8	7.7

4. Conclusion

In this study, the phase behavior of three different structures of copolymers and five salts were studied. The respective binodal curves and tie lines were determined. The results show that block copolymers (Pluronic L35 and R-Pluronic 10R5) have more ability for phase separation. These systems were applied for partitioning of PAA and 6-APA. The hydrophobic interaction and salting-out ability of salts have significant effects on partition coefficient of both PAA and 6-APA. It was concluded that the most hydrophobic solute has partitioned preferentially towards the most hydrophobic phase and the most hydrophilic solute for the most hydrophilic phase. According to the values of selectivity, these ATPSs shows high potential for separation of 6-APA and PAA.

5. References

- [1] K. Li, M.A.A. Mohammed, Y. Zhou, H. Tu, J. Zhang, C. Liu, Z. Chen, R. Burns, D. Hu, J.M. Ruso, Z. Tang, Z. Liu, Recent progress in the development of immobilized penicillin G acylase for chemical and industrial applications: A mini-review, *Polym. Adv. Technol.* 31 (2020) 368–388. <https://doi.org/10.1002/pat.4791>.
- [2] D. -H Chen, S. -S Wang, T. -C Huang, Separation of phenylacetic acid, 6-aminopenicillanic acid and penicillin G with electrodialysis under constant current, *J. Chem. Technol. Biotechnol.* 64 (1995) 284–292. <https://doi.org/10.1002/jctb.280640311>.
- [3] A.H. Demirçelik, I. Perçin, A. Denizli, Supermacroporous hydrophobic affinity sorbents for penicillin acylase purification, *J. Macromol. Sci. Part A Pure Appl. Chem.* 54 (2017) 71–79. <https://doi.org/10.1080/10601325.2017.1261618>.
- [4] S.R. Ayakar, G.D. Yadav, Development of novel support for penicillin acylase and its application in 6-aminopenicillanic acid production, *Mol. Catal.* 476 (2019) 110484. <https://doi.org/10.1016/j.mcat.2019.110484>.
- [5] R.P. Elander, Industrial production of β -lactam antibiotics, *Appl Microbiol Biotechnol.* 61 (2003) 385–392. <https://doi.org/10.1007/s00253-003-1274-y>.

- [6] M. Iqbal, Y. Tao, S. Xie, Y. Zhu, D. Chen, X. Wang, L. Huang, D. Peng, A. Sattar, M.A.B. Shabbir, H.I. Hussain, S. Ahmed, Z. Yuan, Aqueous two-phase system (ATPS): an overview and advances in its applications, *Biol. Proced. Online*. 18 (2016) 1–18. <https://doi.org/10.1186/s12575-016-0048-8>.
- [7] N.L. Abbott, D. Blankschtein, T.A. Hatton, On protein partitioning in two-phase aqueous polymer systems., *Bioseparation*. 1 (1990) 191–225.
- [8] V.P. Priyanka, R.L. Gardas, Mono- and di- cationic ionic liquids based aqueous biphasic systems for the extraction of diclofenac sodium, *Sep. Purif. Technol.* 234 (2020) 116048. <https://doi.org/10.1016/j.seppur.2019.116048>.
- [9] R. Sadeghi, N. Ebrahimi, M.D. Tehrani, Investigation of carbohydrates as non-charged, non-toxic and renewable soluting-out agent for polymer based aqueous biphasic systems implementation, *Polymer (Guildf)*. 98 (2016) 365–377. <https://doi.org/10.1016/j.polymer.2016.06.050>.
- [10] Y.P. Jiménez, C.R. Freijeiro, A. Soto, O. Rodríguez, Phase equilibrium for polymer/ionic liquid aqueous two-phase systems, *Fluid Phase Equilib.* 506 (2020). <https://doi.org/10.1016/j.fluid.2019.112387>.
- [11] M.T. Zafarani-Moattar, S. Emamian, S. Hamzehzadeh, Effect of temperature on the phase equilibrium of the aqueous two-phase poly(propylene glycol) + tripotassium citrate system, *J. Chem. Eng. Data*. 53 (2008) 456–461.

<https://doi.org/10.1021/je700549u>.

- [12] A. Chakraborty, K. Sen, RSC Advances Phase separation in aqueous systems for realizing virtually significant extractions †, RSC Adv. 4 (2014) 64328–64335. <https://doi.org/10.1039/C4RA06798C>.
- [13] M. Svensson, F. Joabsson, P. Linse, F. Tjerneld, Partitioning of hydrophobic amino acids and oligopeptides in aqueous two-phase system containing self-aggregating block copolymer. Effects of temperature, salts and surfactants, J. Chromatogr. A. 761 (1997) 91–101. [https://doi.org/10.1016/S0021-9673\(96\)00830-8](https://doi.org/10.1016/S0021-9673(96)00830-8).
- [14] M.S. Bakshi, S. Sachar, T. Yoshimura, K. Esumi, Association behavior of poly(ethylene oxide)-poly(propylene oxide)-poly(ethylene oxide) block copolymers with cationic surfactants in aqueous solution, J. Colloid Interface Sci. 278 (2004) 224–233. <https://doi.org/10.1016/j.jcis.2004.05.025>.
- [15] M.C. de Oliveira, M.A.N. de Abreu Filho, P. de A. Pessoa Filho, Phase equilibrium and protein partitioning in aqueous two-phase systems containing ammonium carbamate and block copolymers PEO-PPO-PEO, Biochem. Eng. J. 37 (2007) 311–318. <https://doi.org/10.1016/j.bej.2007.05.010>.
- [16] L.H. Haraguchi, R.S. Mohamed, W. Loh, P.A. Pess[^], P.A. Pessoa Filho, Phase equilibrium and insulin partitioning in aqueous two-phase systems containing

- block copolymers and potassium phosphate, *Fluid Phase Equilib.* 215 (2004) 1–15. [https://doi.org/10.1016/S0378-3812\(03\)00368-6](https://doi.org/10.1016/S0378-3812(03)00368-6).
- [17] F.G. Ahsaie, G. Pazuki, T.E. Sintra, P. Carvalho, S.P.M. Ventura, Study of the Partition of Sodium Diclofenac and Norfloxacin in Aqueous Two-Phase Systems based on Copolymers and Dextran, *Fluid Phase Equilib.* (2020) 112868. <https://doi.org/https://doi.org/10.1016/j.fluid.2020.112868>.
- [18] M. Svensson, K. Berggren, A. Veide, F. Tjerneld, Aqueous two-phase systems containing self-associating block copolymers. Partitioning of hydrophilic and hydrophobic biomolecules, *J. Chromatogr. A.* 839 (1999) 71–83. [https://doi.org/10.1016/S0021-9673\(99\)00038-2](https://doi.org/10.1016/S0021-9673(99)00038-2).
- [19] H. Shaker Shiran, M. Baghbanbashi, F. Ghazizadeh Ahsaie, G. Pazuki, Study of curcumin partitioning in polymer-salt aqueous two phase systems, Elsevier B.V, 2020. <https://doi.org/10.1016/j.molliq.2020.112629>.
- [20] J.A.A. Jose C. Merchuk , Barbara A. Andrews, Aqueous two-phase systems for protein separation: Phase separation and applications, *J. Chromatogr. A.* 711 (1998) 285–293. <https://doi.org/10.1016/j.chroma.2012.03.049>.
- [21] S. Miao, J. Chen, X. Cao, Preparation of a novel thermo-sensitive copolymer forming recyclable aqueous two-phase systems and its application in bioconversion of Penicillin G, *Sep. Purif. Technol.* 75 (2010) 156–164.

<https://doi.org/10.1016/j.seppur.2010.08.003>.

- [22] A.F.M. Barton, Handbook of Solubility Parameters and Other Cohesion Parameters, 1991.
- [23] J.G. Pimentel, S.F. Bicalho, O.R.R. Gandolfi, L.A.A. Verissimo, S. de S. Castro, E.A. Souza, C.M. Veloso, R. da C.I. Fontan, V.S. Sampaio, R.C.F. Bonomo, Evaluation of salting-out effect in the liquid–liquid equilibrium of aqueous two-phase systems composed of 2-propanol and Na₂SO₄/MgSO₄ at different temperatures, Fluid Phase Equilib. 450 (2017) 184–193. <https://doi.org/10.1016/j.fluid.2017.08.001>.
- [24] M. Do Carmo Hespanhol Da Silva, L.H. Mendes Da Silva, J.A. Júnior, R.O. Guimarães, J.P. Martins, Liquid-liquid equilibrium of aqueous mixture of triblock copolymers L35 and F68 with Na₂SO₄, Li₂SO₄, or MgSO₄, J. Chem. Eng. Data. 51 (2006) 2260–2264. <https://doi.org/10.1021/je060340l>.
- [25] J.P. Martins, C. De Paula Carvalho, L.H.M. Da Silva, J.S. Dos Reis Coimbra, M. Do Carmo Hespanhol Da Silva, G. Dias Rodrigues, L.A. Minim, Liquid-liquid equilibria of an aqueous two-phase system containing poly(ethylene) glycol 1500 and sulfate salts at different temperatures, J. Chem. Eng. Data. 53 (2008) 238–241. <https://doi.org/10.1021/je700538z>.
- [26] and E.A.M. Sara C. Silvério, Oscar Rodríguez José A. Teixeira, The Effect

- of Salts on the Liquid – Liquid Phase Equilibria of PEG600 + Salt Aqueous Two-Phase Systems, *J. Chem. Eng. Data.* 58 (2013) 3528–3535.
- [27] and E.A.M. Sara C. Silve´rio, Oscar Rodri´guez, Jose´ A. Teixeira, LSRE-Laboratory, Liquid - Liquid Equilibria of UCON + (Sodium or Potassium) Phosphate Salt Aqueous Two-Phase Systems at 23 ° C, *J. Chem. Eng. Data.* 55 (2010) 1285–1288.
- [28] Y. Wang, Y. Mao, J. Han, Y. Liu, Y. Yan, Liquid - Liquid equilibrium of potassium phosphate/potassium citrate/sodium citrate + ethanol aqueous two-phase systems at (298.15 and 313.15) K and correlation, *J. Chem. Eng. Data.* 55 (2010) 5621–5626. <https://doi.org/10.1021/je100501f>.
- [29] L.N. Yun Wang , Yuanyuan Li, Juan Han b, Jinchen Xia, Xu Tang, Tong Chen, Cloudy behavior and equilibrium phase behavior of triblock copolymer L64 þ salt þ water two-phase systems, *Fluid Phase Equilib.* 409 (2016) 439–446. <https://doi.org/10.1016/j.fluid.2015.10.046>.
- [30] E.A.M. Estela Lladosa , Sara C. Silv´erio b, Oscar Rodr´ıguez a, Jos´e A. Teixeira, (Liquid + liquid) equilibria of polymer-salt aqueous two-phase systems for laccase partitioning : UCON 50-HB-5100 with potassium citrate and (sodium or potassium) formate at 23 ° C, *J. Chem. Thermodyn.* 55 (2012) 166–171. <https://doi.org/10.1016/j.jct.2012.06.002>.

- [31] H.H. Jiangying Wu, Yuming Xu, Tadeusz Dabros, CANMETEnergy, Effect of EO and PO positions in nonionic surfactants on surfactant properties and demulsification performance, *Colloids Surfaces A Physicochem. Eng. Asp.* 252 (2005) 79–85. <https://doi.org/10.1016/j.colsurfa.2004.09.034>.
- [32] Y. Zhao, M.B. Brown, S.A. Jones, Engineering novel topical foams using hydrofluoroalkane emulsions stabilised with pluronic surfactants, *Eur. J. Pharm. Sci.* 37 (2009) 370–377. <https://doi.org/10.1016/j.ejps.2009.03.007>.
- [33] A.M. Api, D. Belsito, D. Botelho, M. Bruze, G.A. Burton, J. Buschmann, M.L. Dagli, M. Date, W. Dekant, C. Deodhar, M. Francis, A.D. Fryer, L. Jones, K. Joshi, S. La Cava, A. Lapczynski, D.C. Liebler, D. O'Brien, A. Patel, T.M. Penning, G. Ritacco, J. Romine, N. Sadekar, D. Salvito, T.W. Schultz, I.G. Sipes, G. Sullivan, Y. Thakkar, Y. Tokura, S. Tsang, RIFM fragrance ingredient safety assessment, linalyl phenylacetate, CAS Registry Number 7143-69-3, *Food Chem. Toxicol.* 122 (2018) S703–S713. <https://doi.org/10.1016/j.fct.2018.11.028>.
- [34] C.Y. Kao, T.C. Lo, W.C. Lee, Influence of polyvinylpyrrolidone on the hydrophobic properties of partially porous poly(styrene-divinylbenzene) particles for biological applications, *J. Appl. Polym. Sci.* 87 (2003) 1818–1824. <https://doi.org/10.1002/app.11653>.

- [35] M.P. Lucie Vobecká, Alexandr Romanov, Zdeněk Slouka, Pavel Hasal, Optimization of aqueous two-phase systems for the production of 6-aminopenicillanic acid in integrated microfluidic reactors-separators, *N. Biotechnol.* 47 (2018) 73–79. <https://doi.org/10.1016/j.nbt.2018.03.005>.
- [36] M. Su, J. Wang, J. Gong, H. Sun, Isoelectrical points and solubility of 6-aminopenicillanic acid in water + 1-butanol + butyl acetate, *J. Chem. Eng. Data.* 54 (2009) 373–375. <https://doi.org/10.1021/je800403t>.
- [37] A. Haghtalab, B. Mokhtarani, G. Maurer, Experimental results and thermodynamic modeling of the partitioning of lysozyme, bovine serum albumin, and α -amylase in aqueous two-phase systems of PEG and (K_2HPO_4 or Na_2SO_4), *J. Chem. Eng. Data.* 48 (2003) 1170–1177. <https://doi.org/10.1021/je0340102>.
- [38] S.M. Waziri, B.F. Abu-Sharkh, S.A. Ali, The effect of pH and salt concentration on the coexistence curves of aqueous two-phase systems containing a pH responsive copolymer and polyethylene glycol, *Fluid Phase Equilib.* 205 (2003) 275–290. [https://doi.org/10.1016/S0378-3812\(02\)00297-2](https://doi.org/10.1016/S0378-3812(02)00297-2).
- [39] S. Saravanan, J.R. Rao, B.U. Nair, T. Ramasami, Aqueous two-phase poly(ethylene glycol)-poly(acrylic acid) system for protein partitioning:

- Influence of molecular weight, pH and temperature, *Process Biochem.* 43 (2008) 905–911. <https://doi.org/10.1016/j.procbio.2008.04.011>.
- [40] S.F. Darani, F.G. Ahsaie, G. Pazuki, S. Abdolrahimi, Aqueous two-phase systems based on thermo-separating copolymer for partitioning of doxorubicin, *J. Mol. Liq.* (2020) 114542. <https://doi.org/https://doi.org/10.1016/j.molliq.2020.114542>.
- [41] B.V.N. Phani Kumar, R.R. Reddy, A. Pan, V.K. Aswal, K. Tsuchiya, G.K.S. Prameela, M. Abe, A.B. Mandal, S.P. Moulik, Physicochemical Understanding of Self-Aggregation and Microstructure of a Surface-Active Ionic Liquid [C4mim] [C8OSO3] Mixed with a Reverse Pluronic 10R5 (PO8EO22PO8), *ACS Omega.* 3 (2018) 5155–5164. <https://doi.org/10.1021/acsomega.8b00267>.
- [42] E. Larrañeta, J.R. Isasi, Phase behavior of reverse poloxamers and poloxamines in water, *Langmuir.* 29 (2013) 1045–1053. <https://doi.org/10.1021/la304245p>.
- [43] T. Liu, V.M. Nace, B. Chu, Cloud-Point Temperatures of B n E m B n and P n E m P n Type Triblock Copolymers in Aqueous Solution, *J. Phys. Chem. B.* 102 (1998) 1065–1065. <https://doi.org/10.1021/jp9801119>.
- [44] T. Patel, P. Bahadur, J. Mata, The clouding behaviour of PEO-PPO based

- triblock copolymers in aqueous ionic surfactant solutions: A new approach for cloud point measurements, *J. Colloid Interface Sci.* 345 (2010) 346–350. <https://doi.org/10.1016/j.jcis.2010.01.079>.
- [45] G. Lazzara, S. Milioto, M. Gradzielski, The solubilisation behaviour of some dichloroalkanes in aqueous solutions of PEO-PPO-PEO triblock copolymers: A dynamic light scattering, fluorescence spectroscopy, and SANS study, *Phys. Chem. Chem. Phys.* 8 (2006) 2299–2312. <https://doi.org/10.1039/b516242b>.
- [46] F. Gabelle, W.J. Koros, R.S. Schechter, Solubilization of Aromatic Solutes in Block Copolymers, *Macromolecules.* 28 (1995) 4883–4892. <https://doi.org/10.1021/ma00118a014>.
- [47] Y. Tian, C. Shi, Y. Sun, C. Zhu, C.C. Sun, S. Mao, Designing micellar Nanocarriers with improved drug loading and stability based on solubility parameter, *Mol. Pharm.* 12 (2015) 816–825. <https://doi.org/10.1021/mp5006504>.
- [48] M.M. Lübtow, M.S. Haider, M. Kirsch, S. Klisch, R. Luxenhofer, Like Dissolves Like? A Comprehensive Evaluation of Partial Solubility Parameters to Predict Polymer-Drug Compatibility in Ultrahigh Drug-Loaded Polymer Micelles, *Biomacromolecules.* 20 (2019) 3041–3056. <https://doi.org/10.1021/acs.biomac.9b00618>.

Author Contribution statement

Farzaneh Ghazizadeh Ahsaie: Conceptualization, Methodology, Software, Writing.

Gholamreza Pazuki:, Writing, Methodology, Supervision.

Competing financial interests:

The authors declare no competing financial and/or non-financial interests.

Figures




Figure 1

Experimental binodal curves determined for the ATPSs composed of different copolymer + salt at 298 K




Figure 2

Experimental binodal curves determined for the ATPSs composed of different copolymer + salt at 283 K

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [sup6apa.pdf](#)