Medication adherence and Survival prospects among hospitalized heart failure patients in a tertiary hospital in Tanzania: A Prospective Cohort Study.

Pedro Pallangyo (✉ pedro.pallangyo@gmail.com)

The Jakaya Kikwete Cardiac Institute

Jalack Millinga

Jakaya Kikwete Cardiac Institute

Smita Bhalia

JAKAYA KIKWETE CARDIAC INSTITUTE

Zabella Mkojera

JAKAYA KIKWETE CARDIAC INSTITUTE

Nsajigwa Misidai

JAKAYA KIKWETE CARDIAC INSTITUTE

Happiness J Swai

JAKAYA KIKWETE CARDIAC INSTITUTE

Naairah R Hemed

JAKAYA KIKWETE CARDIAC INSTITUTE

Alice Kaijage

JAKAYA KIKWETE CARDIAC INSTITUTE

Mohamed Janabi

JAKAYA KIKWETE CARDIAC INSTITUTE

Research note

Keywords: heart failure, nonadherence, poor adherence, low adherence, drug adherence, medication adherence, medication compliance, noncompliance, Tanzania

Posted Date: January 14th, 2020

DOI: <https://doi.org/10.21203/rs.2.20768/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Research Notes on February 21st, 2020. See the published version at <https://doi.org/10.1186/s13104-020-04959-w>.

Abstract

Objective : Management of heart failure is complex and multifaceted but adherence to medications remains the cornerstone of preventing avoidable readmissions, premature deaths, and unnecessary healthcare expenses. Despite of evidence-based efficacy on anti-failure drugs, poor adherence is pervasive and remains a significant barrier to improving clinical outcomes in heart failure population.

Results : We enrolled 459 patients with established diagnosis of heart failure. The mean age was 46.4 years and participants aged ≤ 50 years constituted 55.4% of the cohort. There was a female predominance (56.5%), 67.5% resided in urban areas and 74.2% had primary education. Of the 419 participants eligible for assessment of medication adherence, 313 (74.7%) had poor adherence and 106 (25.3%) had good adherence. Possession of a health insurance was found to be the strongest associated factor for adherence (OR 8.7, 95% CI 4.7-16.0, $P < 0.001$). Participants with poor adherence displayed a 70% increased risk for rehospitalization compared to their counterparts with good adherence (RR 1.7, 95% CI 1.2-2.9, $p = 0.04$). Poor adherence was found to be the strongest predictor of early mortality (HR 2.5, 95% CI 1.3-4.6, $p < 0.01$). Poor medication adherence in patients with heart failure is associated with increased readmissions and mortality.

Introduction

Cardiovascular disorders (CVD) are responsible for about one-third of all global mortalities with over three-quarters of deaths transpiring in the developing world.¹ In spite of the remarkable advances in novel screening techniques and therapeutic directions, the prognosis of heart failure (HF) remains strikingly poor around the globe particularly in the developing nations.²⁻⁷ Owing to its chronic nature, clinical management of HF necessitate long-term use of several drugs to reduce morbidity (digoxin, diuretics, and nitrates)⁸⁻¹⁰ and mortality (angiotensin converting enzyme [ACE] inhibitors, aldosterone receptor blockers [ARBs], and beta-blockers).¹¹⁻¹³ However, universally low prescription rates (19.8% in low-income, 30.7% in lower middle-income, 54.9% in upper middle-income and 88.8% in high-income countries) of such drugs among patients who require them is observed.¹⁴

Despite of all developments in heart-failure management, adherence plays a pivotal role in attaining maximal therapeutic benefits (i.e. better clinical outcomes). Nevertheless, regardless of the assessment tool used or population studied, adherence rates are consistently suboptimal across studies making it a significant public health issue.¹⁵⁻²⁵ Poor adherence to prescribed regimens is pervasive and results in preventable hospitalizations, premature deaths and unnecessary health care expenditure regardless of the underlying cardiovascular etiology.¹⁵⁻²⁶ In this prospective cohort study, we sought to explore the adherence pattern, associated factors and outcomes among hospitalized heart failure patients in a tertiary hospital in Tanzania.

Methods

Recruitment process and Definition of terms

All patients who were hospitalized at our institution between March and October 2018 with established diagnosis of heart failure (for at least 3 months' prior enrollment) were consecutively enrolled for this study. Sociodemographic, clinical, laboratory, echocardiographic, and adherence data were gathered using a structured questionnaire during the hospital admission of enrollment. Participants underwent a 2-dimensional echocardiography for diagnosis reconfirmation. Renal functions were estimated using the Modification of Diet in Renal Disease (MDRD) equation and estimated glomerular filtration rate (eGFR) value of $< 60 \text{ mL/min/1.73 m}^2$ was used to define renal dysfunction. Diagnosis of anemia utilized the WHO criteria i.e. Hemoglobin (Hb) concentration of $< 13.0 \text{ g/dL}$ and $< 12.0 \text{ g/dL}$ for males and females respectively. Diabetes was defined by fasting blood glucose (FBG) levels $\geq 7.0 \text{ mmol/L}$ or use of glucose lowering agents. Hypertension was defined as systolic blood pressure (SBP) $> 140 \text{ mmHg}$ and/or diastolic blood pressure (DBP) $> 90 \text{ mmHg}$ or use of antihypertensive medications. Total cholesterol level greater than 6.2 mmol/L was used to define dyslipidemia. Hyponatremia, hypokalemia, hypocalcemia, and hypomagnesemia were defined by concentrations $< 135 \text{ mmol/L}$, $< 3.5 \text{ mmol/L}$, $< 2.1 \text{ mmol/L}$ and $< 0.7 \text{ mmol/L}$ respectively. Potassium levels $> 5.0 \text{ mmol/L}$ was used to denote hyperkalemia. We assessed adherence based on the last time a participant last took her heart failure medications. Participants were regarded to have good adherence if they took all of their prescribed heart failure medications within 72 hours before their index admission and poor adherence was defined as not taking all medications for 72 hours or more.

Follow-up and Study Outcomes

Follow-up was conducted through scheduled weekly phone calls and continued through April 2019 with a predetermined stopping point providing a maximum of 180 days of follow-up for each patient after enrollment. Data was censored after completion of follow-up or death, whichever occurred first. A participant was deemed lost to follow-up when despite all attempts couldn't be reached through phone numbers provided. Our primary outcome measures were rehospitalization and all-cause mortality during the study duration. We defined rehospitalization as any cardiovascular-related hospital admission following a successful discharge from the hospitalization of enrollment. Early mortality was defined as death during the hospitalization of enrollment.

Statistical analysis

All statistical analyses utilized STATA v11.0 software. Pearson Chi square and Student's T-test were used to compare categorical and continuous variables respectively. Linear logistic regression analyses were used to assess for factors associated with adherence and predictors of rehospitalization. Based on their adherence status, participants were compared with respect to survival using Cox proportional-hazards regression model. Differences in survival between the low- and high-adherence groups were compared using the log-rank test. We report Odds ratio (OR), Relative risk (RR) and Hazard ratio (HR) with 95% confidence intervals (CI) and p-values where appropriate. All tests were 2-sided and $p < 0.05$ was used to denote statistical significance.

Results

Study Population.

A total of 459 heart failure patients met the inclusion criteria and were enrolled into this study. During follow-up, 40 (8.7%) participants exited; 5 due to incomplete key data and 35 were lost to follow-up. Table 1 displays the baseline characteristics of participants. The mean age of our heart failure cohort was 46.4 ± 18.9 years and those aged ≤ 50 years constituted 55.4% of all participants. There was a female predominance (56.6%) and over two-thirds of all participants resided in urban areas. The mean BMI was 25.1 ± 5.2 and 39.4% of patients were overweight or obese. About 7.2% of participants were in NYHA functional class II while classes III and IV constituted 36.5% and 56.3% respectively. Fifty two point seven percent (52.7%) had a history of hypertension, 13.6% had diabetes, 6.7% were infected with HIV, 51.3% had renal insufficiency and 72.1% were anemic. Echocardiography revealed HHD was the predominant cause of heart failure (40.1%) followed by DCM (27.0%) and RHD (23.2%).

Table 1
Baseline characteristics of participants (N = 419)

Characteristic	All (N = 419)	Poor adherence (n = 313)	Good adherence (n = 106)	p-value
Age	46.4 (18.9)	45.5 (19.0)	49.1 (18.6)	0.09
Age groups	103 (24.6%)	82 (26.2%)	21 (19.8%)	0.19
<30	129 (30.8%)	96 (30.7%)	33 (31.1%)	0.94
30–50	187 (44.6%)	135 (43.1%)	52 (49.1%)	0.28
>50				
Sex	182 (43.4%)	139 (44.4%)	43 (40.6%)	0.50
Male	237 (56.6%)	174 (55.6%)	63 (59.4%)	
Female				
Residence	283 (67.5%)	197 (62.9%)	86 (81.1%)	0.001
Urban	136 (32.5%)	116 (37.1%)	20 (18.9%)	
Rural				
Marital status	100 (23.9%)	82 (26.2%)	18 (17.0%)	0.05
Single	296 (70.6%)	213 (68.1%)	83 (78.3%)	0.05
Married	23 (05.5%)	18 (05.7%)	5 (04.7%)	0.67
Divorced/Widowed				
Education	16 (03.8%)	12 (03.9%)	4 (03.8%)	0.96
None	295 (70.4%)	248 (79.2%)	47 (44.3%)	<0.001
Primary	68 (16.2%)	36 (11.5%)	32 (30.2%)	<0.001
Secondary	40 (09.6%)	17 (05.4%)	23 (21.7%)	<0.001
University				
Occupation	76 (18.1%)	47 (15.0%)	29 (27.3%)	<0.01
None	311 (74.3%)	250 (79.9%)	61 (57.6%)	<0.001
Employed/self-employed	32 (07.6%)	16 (05.1%)	16 (15.1%)	0.001
Retired				
Body Mass Index	25.1 (05.2)	24.8 (04.2)	26.0 (07.4)	0.04
BMI Categories	11 (02.6%) 243 (58.0%)	7 (02.2%) 188 (60.1%)	4 (03.8%) 55 (51.9%)	0.37
Underweight				0.14
Normal	105 (25.1%)	79 (25.2%)	26 (24.5%)	0.89
Overweight	60 (14.3%)	39 (12.5%)	21 (19.8%)	0.06
Obese				
Health insured	93 (22.2%)	32 (10.2%)	61 (57.6%)	<0.001
Yes	326 (77.8%)	281 (89.8%)	45 (42.4%)	
No				
HF etiology	113 (27.0%)	78 (24.9%)	34 (32.1%)	0.15
DCM	168 (40.1%)	134 (42.8%)	35 (33.0%)	0.08
HHD	97 (23.2%)	72 (23.0%)	25 (23.6%)	0.90
RHD	41 (09.8%)	29 (09.3%)	12 (11.3%)	0.55
Others				

Characteristic	All (N = 419)	Poor adherence (n = 313)	Good adherence (n = 106)	p-value
Comorbidities	221 (52.7%)	171 (54.6%)	50 (47.2%)	0.19
Hypertension	57 (13.6%)	39 (12.5%)	18 (17.0%)	0.24
Diabetes	28 (6.7%)	15 (4.8%)	13 (12.3%)	0.01
HIV/AIDS	215 (51.3%)	163 (52.1%)	52 (49.1%)	0.59
Renal insufficiency	100 (23.9%)	80 (25.6%)	20 (18.9%)	0.16
eGFR < 15	302 (72.1%)	234 (74.8%)	68 (64.2%)	0.04
Anemia	99 (23.6%)	75 (24.0%)	24 (22.6%)	0.77
Hb < 8 g/dL				
NYHA class	30 (7.2%)	19 (6.0%)	11 (10.4%)	0.13
II	153 (36.5%)	112 (35.8%)	41 (38.7%)	0.59
III	236 (56.3%)	182 (58.2%)	54 (50.9%)	0.19
IV				
Admission days	14.0 (13.3)	13.8 (13.4)	14.3 (12.8)	0.74
HF-related	211 (50.4%)	167 (53.3%)	44 (41.5%)	0.04
hospitalization	208 (49.6%)	146 (46.7%)	62 (58.5%)	
1st				
>1				

Medication adherence

Overall, 337 (73.4%) were on ACEI, 122 (26.6%) on ARB, 386 (84.1%) on beta-blockers, 432 (94.1%) on diuretics, 395 (86.1%) on aldosterone antagonists, 166 (36.2%) on inotropes and 36 (7.8%) were on digoxin. Of the 419 participants eligible for assessment of medication adherence, 313 (74.7%) had poor adherence and 106 (25.3%) had good adherence. The mean number of days' participants last took medications before the index hospitalization was 17.7 ($\pm$ 6.9) days. Among participants with poor adherence, 254 (81.2%) had not taken any of their anti-failure medications within the past one-week prior admission. Inability to afford medications was the most (87.3%) reported reason for nonadherence. Other reported factors affecting adherence in this cohort included; medication side effects (8.1%), forgetfulness (53.9%), negligence (26.0%), local unavailability of drugs (18.9%) and pill burden (34.4%). Differences in age, sex, marital status, and BMI displayed similar medication adherence patterns, Table 1. However, during bivariate analyses four characteristics including education level, residence, employment status, and health insurance possession showed significant associations with adherence, Table 2. Significant variables then underwent multivariate logistic regression analysis where possession of a health insurance was found to be the strongest associated factor for adherence (OR 8.7, 95% CI 4.7–16.0, $P < 0.001$), Table 2.

Table 2
Factors Associated with Adherence

Control group	Comparative group	OR	95% CI	P-value	Adj. OR	Adj. 95% CI	Adj. P-value
age < 50	age ≥ 50	0.8	0.5–1.2	0.3	-	-	-
Female	Male	1.2	0.7–1.8	0.5	-	-	-
≥ secondary education	≤ primary education	5.3	3.3–8.6	< 0.001	1.9	0.9-4.0	0.07
married	Single	1.7	1.0-2.8	0.05	-	-	-
Employed	no employment	0.3	0.2–0.5	< 0.001	1.2	0.6–2.4	0.6
Urban	Rural	2.5	1.5–4.3	0.001	2.0	1.1–3.7	0.03
no comorbidity	≥ 1 comorbidity	0.9	0.5–1.4	0.56	-	-	-
health insurance	not insured	11.9	7.0-20.2	< 0.001	8.7	4.7–16.0	< 0.001

Rehospitalization and Mortality

Overall, 208 (49.6%) patients had a history of a prior cardiovascular-related hospitalization. During follow-up, rehospitalization rates were 32.8%, 48.1% and 53.0% at 30, 90 and 180 days respectively. Despite of similar rehospitalization rates among poor and good-adherence participants at 30-days (35.4% vs 27.2%, $p = 0.12$) and 90-days (51.8% vs 40.2%, $p = 0.07$), patients with poor adherence had significantly higher rates of rehospitalization at 180 days (57.5% vs 43.5%, $p = 0.03$). Participants with poor adherence displayed a 70% increased risk for rehospitalization compared to their counterparts with good adherence (RR 1.7, 95% CI 1.2–2.9, $p = 0.04$).

Overall, 177 (42.2%) patients survived the 180-days of follow-up. The overall mean survival days was 103.3 ± 74.8 days and participants with good adherence (140.5 ± 63.1 days) displayed a longer survival compared to their poor adherence (90.8 ± 74.3 days) counterparts, $p < 0.001$. Mortality rate was 30.8%, 48.7% and 57.8% at 30, 90 and 180 days respectively. Regardless of the assessment time, participants with poor adherence displayed superior mortality compared to those with good adherence i.e. 37.1% vs 12.3% at 30 days, 56.6% vs 25.5% at 90 days, and 65.5% vs 34.9% at 180 days; all $p < 0.001$). Additionally, we performed subgroup analyses to assess for all-cause mortality by adherence status. In all 19 characteristics involved in subgroup analyses, participants with poor adherence had inferior survival rates compared to their counterparts with good adherence, Fig. 1. More interestingly, even within the subgroup

of those who possessed a health insurance, it was observed that poor adherence participants fared worse compared to good adherence controls, (HR 1.6, 95% CI 1.0-2.4, $p = 0.05$).

Discussion

Management of heart failure is complex and multifaceted but adherence to medications remains a fundamental measure to prevent acute exacerbations.^{27,28} Despite of unwavering evidence on the efficacy of anti-failure drugs, poor adherence is common and remains a significant barrier to improving clinical outcomes in heart failure population. Estimates of nonadherence in heart failure patients have varied widely (22%-90%)^{18,25,29-36} in the literature. In this present study, less than one-fifth of participants were categorized as having high adherence. Our rate of nonadherence is skewed to the extreme undesired end of the reported range in the literature.

With regards to reasons for poor adherence, numerous factors have predominated in various studies. For instance, in studies by Toh et al (71%) and Mujtaba et al (72.7%), poor medication instructions was the most reported factor.^{25,36} On the other hand, studies by Aggarwal et al and Dickson et al found forgetfulness and comorbidities respectively as the leading factors for nonadherence.^{29,32} In this present study, nearly 90% of nonadherent participants reported medication cost as the major barrier to their adherence. These findings are in unison with Dunlay et al study as far as cost being the most reported factor is concerned, however it was a barrier in a significantly lesser proportion (22%) compared to what we observed.³⁷ While majority of known risk factors for nonadherence are potentially modifiable, inability to comply due to poverty is not. Owing to this, improving medication adherence in impoverished societies continues to be a very difficult undertaking. It should not be forgotten that such poor societies and their already overwhelmed health sectors continue to struggle with prevention and management of the ever present infectious diseases.

Several studies have demonstrated the repercussions of poor adherence on prognosis of heart failure.^{16,32,33,38,39} Moreover, numerous studies have established the prognostic benefits of interventions to improve adherence.^{24, 40-50} In this present study, nearly 60% of participants with poor adherence were rehospitalized within 6-months of enrollment. Our findings are in consonance with several other prospective studies which have produced rehospitalization rates ranging between 20% and 69%.^{16,32,33,38,39} Additionally, intervention studies have uniformly shown that improved adherence is associated with reduction (3%-96%) in readmission risk.^{40,41,43,44,46,47,48,49} Furthermore, systematic reviews and meta-analyses by Ruppert et al and Unverzagt et al revealed a 21% and 10% decreased odds of rehospitalization respectively in the adherence intervention arm.^{24,50}

Survival prospects among heart failure patients remain poor all over the globe. Overall, less than half of patients in this study survived the 6-months of follow-up. Nonadherent participants displayed about three times mortality hazard compared to their adherent counterparts. Similar to our findings, intervention studies have shown mortality reduction (2%-84%) in favor of adherent participants.^{40,41,42,44,45,46,48}

Moreover, two meta-analyses that included over 50 studies each showed a 2% and 11% mortality reduction in favor of the adherence intervention arm.^{24,50} Poor adherence was found to be the strongest predictor of early mortality in this study. To solidify on the significance of adherence in heart failure prognostication, participants with low adherence displayed significantly higher rates of primary outcomes compared to their high adherence counterparts in all subgroup analyses we conducted.

Conclusions

In conclusion, findings of this present study provide important insight pertaining to medication adherence and its potential in dictating the prognosis of heart failure patients residing in resource-limited settings. Poor adherence in patients with heart failure contributes to a considerable burden on the healthcare system above all increased rehospitalizations and mortality. These findings call for deliberate efforts to ensure that measures to assess and improve adherence are incorporated and become an integral component in routine clinical practice. Furthermore, strategies to improve health insurance acquisition including endeavours to make it a right rather than a privilege is fundamental in improving adherence especially among persons living in impoverished societies.

Limitations

Medication adherence was ascertained by self-report and thus reporting bias and recall bias could have in some way affected our findings. Prospective comparison of patients receiving adherence intervention versus control would allow a more rigorous evaluation of adherence potential in prognosticating heart failure and should be considered in the future studies in this setting.

List Of Abbreviations

ACE, Angiotensin Converting Enzyme; ARB, Angiotensin Receptor Blocker; BMI, Body Mass Index; CI, Confidence Interval; CVD, Cardiovascular Disorders; DBP, Diastolic Blood Pressure; DCM, Dilated Cardiomyopathy; eGFR, estimated Glomerular Filtration Rate; FBG, Fasting Blood Glucose; Hb, Hemoglobin; HF, Heart Failure; HHD, Hypertensive Heart Disease; HR, Hazard Ratio; MDRD, Modification of Diet in Renal Disease; MMAS-8, 8-item Morisky Medication Adherence Scale; NYHA, New York Heart Association; OR, Odd Ratio; RHD, Rheumatic Heart Disease; RR, Relative Risk; SBP, Systolic Blood Pressure; WHO, World Health Organization.

Declarations

Ethics approval and consent to participate

The study was approved by the Directorate of Research and Publications of the Muhimbili University of Health and Allied Sciences and the permission to conduct the study was granted by the Jakaya Kikwete Cardiac Institute. All the participants or caretakers in capacity to consent on behalf provided written informed consent.

Consent to Publish

Not applicable

Availability of Data and Materials

The final version of data set supporting the findings of this paper is submitted together with this manuscript to the editorial committee. All the raw data is included in this manuscript. There are no ethics restrictions preventing the sharing of the raw data.

Competing interest

The authors have no conflict of interest to declare

Funding

This study was possible through a research grant from the Commission for Science and Technology (COSTECH) of Tanzania. The investigators and the study sponsor never had confidentiality agreement.

Author contributions

PP conceived the study. JM, ZM, NM, HJS and NRH conducted all the interviews and physical examinations. Echocardiography was performed by MJ, AK, and SB. PP performed data entry and analysis. The corresponding author wrote the first draft of the manuscript, and other authors contributed to and approved it. All authors made the decision to submit the manuscript for publication. All authors assume responsibility for the accuracy and integrity of the analysis.

Acknowledgement

We thank the nursing and medical staff of the Jakaya Kikwete Cardiac Institute for their cooperation and active participation during this study. We extend our gratitude to all the study participants and their relatives for their willingness, tolerance and cooperation offered during the entire study duration.

Correspondence

Pedro Pallangyo, Jakaya Kikwete Cardiac Institute, P.O Box 65141, Dar es Salaam, email: pedro.pallangyo@gmail.com, cell: +255 783 351 165.

References

1. World Health Organization. Cardiovascular diseases Fact sheet reviewed September 2016. <http://www.who.int/mediacentre/factsheets/fs317/en/>.
2. Kraus S, Ogunbanjo G, Sliwa K, Ntusi NAB. Heart failure in sub-Saharan Africa: A clinical approach. S Afr Med J 2016;106(1):23-31.

3. Ntusi NB, Mayosi BM. Epidemiology of heart failure in sub-Saharan Africa. *Expert Rev Cardiovasc Ther.* 2009 Feb;7(2):169-80.
4. Makubi A, Hage C, Lwakatare J, et al. Contemporary aetiology, clinical characteristics and prognosis of adults with heart failure observed in a tertiary hospital in Tanzania: the prospective Tanzania Heart Failure (TaHeF) study. *Heart.* 2014 Aug; 100(16): 1235-41.
5. Pallangyo P, Fredrick F, Bhalia S, et al. Cardiorenal Anemia Syndrome and Survival among Heart Failure Patients in Tanzania: A Prospective Cohort Study. *BMC Cardiovasc Disord.* 2017; **17**:59.
6. Parmar KR, Xiu PY, Chowdhury MR, et al. In-hospital treatment and outcomes of heart failure in specialist and non-specialist services: a retrospective cohort study in the elderly Open Heart 2015; 2:
7. Karaye KM, Sani MU. Factors associated with poor prognosis among patients admitted with heart failure in a Nigerian tertiary medical centre: a cross-sectional study. *BMC Cardiovasc Disord.* 2008; 8: 16.
8. Gheorghiade M, Patel K, Filippatos G, et al. Effect of oral digoxin in high-risk heart failure patients: a pre-specified subgroup analysis of the DIG trial. *Eur J Heart Fail.* 2013 May; 15(5): 551–59.
9. Zannad F, McMurray JJ, Krum H, van Veldhuisen DJ, Swedberg K, Shi H, et al. Eplerenone in patients with systolic heart failure and mild symptoms. *N Engl J Med.* 2011; 364(1):11-21.
10. Metra M, Teerlink JR, Voors AA, et al. Vasodilators in the treatment of acute heart failure: what we know, what we don't. *Heart Fail Rev.* 2009; 14:299–307.
11. Konstam MA, Rousseau MF, Kronenberg MW, Udelson JE, Melin J, Stewart D, et al. Effects of the angiotensin converting enzyme inhibitor enalapril on the long-term progression of left ventricular dysfunction in patients with heart failure. SOLVD Investigators. *Circulation.* 1992; 86(2):431-8.
12. Pfeffer MA, Swedberg K, Granger CB, Held P, McMurray JJ, Michelson EL, et al. Effects of candesartan on mortality and morbidity in patients with chronic heart failure: the CHARM-Overall programme. *Lancet.* 2003; 362(9386):759-66.
13. Haas SJ, Vos T, Gilbert RE, Krum H. Are beta-blockers as efficacious in patients with diabetes mellitus as in patients without diabetes mellitus who have chronic heart failure? A meta-analysis of large-scale clinical trials. *Am Heart J.* 2003; 146:848–53.
14. Yusuf S, Islam S, Chow CK et al., Use of secondary prevention drugs for cardiovascular disease in the community in high-income, middle-income, and low-income countries (the PURE Study): a prospective epidemiological survey. *The Lancet* 2011; 378 (9798): 1231–43.
15. Simpson SH, Eurich DT, Majumdar SR, et al. A metaanalysis of the association between adherence to drug therapy and mortality. 1006; 333:18 –26.
16. Osterberg L, Blaschke T. Adherence to medication. *N Engl J Med.* 2005; 353:487– 497.
17. Ho PM, Bryson CL, Rumsfeld JS. Medication Adherence: Its Importance in Cardiovascular Outcomes. *Circulation.* 2009; 119:3028-3035.
18. Oosterom-Calo R, van Ballegooijen AJ, Terwee CB, et al. Determinants of adherence to heart failure medication: a systematic literature review. *Heart Fail Rev* 2013; 18:409–427.

19. Juarez DT, Williams AE, Chen C, et al. Factors Affecting Medication Adherence Trajectories for Patients With Heart Failure. *Am J Manag Care*. 2015; 21(3):197-205.
20. Molloy GJ, O'Carroll RE, Witham MD, et al. Interventions to Enhance Adherence to Medications in Patients With Heart Failure: A Systematic Review. *Circ Heart Fail*. 2012; 5:126-133.
21. Fitzgerald AA, Powers JD, Ho PM, et al. Impact of medication nonadherence on hospitalizations and mortality in heart failure. *J Card Fail*. 2011;17(8):664–669.
22. Shah D, Simms K, Barksdale DJ, Wu Improving medication adherence of patients with chronic heart failure: challenges and solutions. *Research Reports in Clinical Cardiology* 2015;6 87–95.
23. [Ruppar TM](#), [Delgado JM](#), [Temple J](#). Medication adherence interventions for heart failure patients: A meta-analysis. *Eur J Cardiovasc Nurs*. 2015 Oct; 14(5):395-404.
24. Ruppar TM, Cooper PS, Mehr DR, et al. Medication Adherence Interventions Improve Heart Failure Mortality and Readmission Rates: Systematic Review and Meta-Analysis of Controlled Trials. *J Am Heart Assoc*. 2016 June; 5(6).
25. Toh CT, Jackson B, Gascard DJ, et al. Barriers to Medication Adherence in Chronic Heart Failure Patients during Home Visits. *J Pharm Pract Res* 2010; 40: 27-30.
26. Sabaté E. Adherence to long-term therapies: evidence for action. World Health Organization, Geneva, Switzerland, 2003.
27. Kemp C, Conte J. The pathophysiology of heart failure. *Cardiovasc Pathol*. 2012; 21(5):365–371.
28. O'Connor CM, Stough WG, Gallup DS, Hasselblad V, Gheorghiade M. Demographics, clinical characteristics, and outcomes of patients hospitalized for decompensated heart failure: observations from the IMPACT-HF registry. *J Card Fail*. 2005;11(3):200–205.
29. [Aggarwal B](#), [Pender A](#), [Mosca L](#), [Mochari-Greenberger H](#). Factors associated with medication adherence among heart failure patients and their caregivers. *J Nurs Educ Pract*. 2015; 5(3):22-27.
30. Zhang Y, Wu SH, Fendrick AM, Baicker K. Variation in Medication Adherence in Heart Failure. *JAMA Intern Med*. 2013;173(6):468-470.
31. [Knafl GJ](#), [Riegel B](#). What puts heart failure patients at risk for poor medication adherence? [Patient Prefer Adherence](#). 2014 Jul 17;8:1007-18.
32. Dickson VV, Knafl GJ, Riegel B. Predictors of Medication Nonadherence Differ Among Black and White Patients with Heart Failure. *Res Nurs Health*. 2015 Aug; 38(4): 289–300.
33. Wu JR, Frazier SK, Rayens MK, et al. Medication Adherence, Social Support, and Event-Free Survival in Patients with Heart Failure. *Health Psychol*. 2013 Jun; 32(6): 637–646.
34. Alakhali KM, Daniel PS, Noohu AM, Sirajudeen SA. Patient Medication Adherence and Physician Prescribing among Congestive Heart Failure Patients of Yemen. *Indian J Pharm Sci*. 2013 Sep-Oct; 75(5): 557–562.

35. Lee D, Mansi I, Bhushan S, Parish R. Non-Adherence in At-Risk Heart Failure Patients: Characteristics and Outcomes. *Journal of Nature and Science*, 2015; 1(5):e95
36. Mujtaba SF, Masood T, Saad M. Reasons of Medical Noncompliance in Heart Failure Patients. *Pakistan Heart Journal*. 2010 July-Dec; 43 (3-4)
37. Dunlay SM, Eveleth JM, Shah ND, et al. Medication Adherence Among Community-Dwelling Patients With Heart Failure. *Mayo Clin Proc*. 2011 Apr; 86(4): 273–281.
38. Albert NM. Improving medication adherence in chronic cardiovascular disease. *Crit Care Nurse*. 2008 Oct;28(5):54-64; quiz 65.
39. Leventhal MJ, Riegel B, Carlson B, De GS. Negotiating compliance in heart failure: remaining issues and questions. *Eur J Cardiovasc Nurs*. 2005; 4:298-307.
40. Antonicelli R, Testarmata P, Spazzafumo L, et al. Impact of telemonitoring at home on the management of elderly patients with congestive heart failure. *J Telemed Telecare*. 2008; 14:300–305.
41. Antonicelli R, Mazzanti I, Abbatecola AM, Parati G. Impact of home patient telemonitoring on use of b-blockers in congestive heart failure. *Drugs Aging*. 2010;27:801–805.
42. Assyag P, Renaud T, Cohen-Solal A, et al. RESICARD: East Paris network for the management of heart failure: absence of effect on mortality and rehospitalization in patients with severe heart failure admitted following severe decompensation. *Arch Cardiovasc Dis*. 2009;102:29–41.
43. Bisharat B, Hafi L, Baron-Epel O, et al. Pharmacist counselling to cardiac patients in Israel prior to discharge from hospital contribute to increasing patient's medication adherence closing gaps and improving outcomes. *J Transl Med*. 2012; 10:34.
44. Bocchi EA, Cruz F, Guimaraes G, et al. Long-term prospective, randomized, controlled study using repetitive education at six-month intervals and monitoring for adherence in heart failure outpatients: the REMADHE trial. *Circ Heart Fail*. 2008;1:115–124.
45. DeWalt DA, Schillinger D, Ruo B, et al. Multisite randomized trial of a single-session versus multisession literacy sensitive self-care intervention for patients with heart failure. *Clin Trial Regist*. 2012;125:2854–2862.
46. Falces C, Lopez-Cabezas C, Andrea R, et al. Intervencion educativa para mejorar el cumplimiento del tratamiento y prevenir reingresos en pacientes de edad avanzada con insuficiencia cardiaca [An educative intervention to improve treatment compliance and to prevent readmissions of elderly patients with heart failure]. *Med Clin (Barc)*. 2008;131:452–456.
47. Ferrante D, Varini S, Macchia A, et al; on behalf of the GESICA Investigators. Long-term results after a telephone intervention in chronic heart failure: DIAL (randomized trial of phone intervention in chronic heart failure) follow-up. *J Am Coll Cardiol*. 2010;56:372–378.
48. Powell LH, Calvin JE Jr, Richardson D, et al. Self-management counselling in patients with heart failure: the heart failure adherence and retention randomized behavioral trial. *JAMA*.

2010;304:1331–1338.

49. Shively MJ, Gardetto NJ, Kodiath MF, et al. Effect of patient activation on self-management in patients with heart failure. *J Cardiovasc Nurs*. 2013;28:20–34.
50. Unverzagt S, Meyer G, Mittmann S, et al. Improving Treatment Adherence in Heart Failure: A Systematic Review and Meta-analysis of Pharmacological and Lifestyle Interventions. *Dtsch Arztebl Int*. 2016 Jun; 113(25): 423–430.

Figures

Figure 1

Forest plot displaying subgroup analyses for all-cause mortality by adherence status.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [DataSheetAdhFinal.xlsx](#)
- [DatasetCodes.docx](#)