

# An increase in psychosomatic symptoms among youth: Is it connected to adult mental health?

Fabrizia Giannotta (✉ [fabrizia.giannotta@su.se](mailto:fabrizia.giannotta@su.se))

Stockholms Universitet <https://orcid.org/0000-0002-3005-1840>

Kent W. Nilsson

Uppsala University: Uppsala Universitet

Cecilia Åslund

Uppsala Universitet

Peter Larm

Stockholm University: Stockholms Universitet

---

## Research Article

**Keywords:** Psychosomatic symptoms, somatic symptoms, musculoskeletal symptoms, psychological symptoms, adolescents, young adults.

**Posted Date:** March 18th, 2022

**DOI:** <https://doi.org/10.21203/rs.3.rs-1137822/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

**Background.** Despite an increase in mental health problems, with psychosomatic symptoms having been observed in new generations of youth, the extent to which these problems correspond to an increase in adult mental problems is unknown. The present study investigates whether adolescents with high levels of psychosomatic symptoms are at risk of developing depression and anxiety problems in adulthood and whether sex moderates any association. Moreover, we aim to understand whether different clusters of youth psychosomatic symptoms – somatic, psychological and musculoskeletal – have different impacts on adult mental health.

**Methods.** 1545 Swedish adolescents – aged 13 (49%) and 15 (51%) – completed surveys at baseline (T1) and 3 years later (T2); of them, 1174 (61% females) also participated after 6 years (T3). Multivariate logistic models were run.

**Results.** Youth with high levels of psychosomatic symptoms had higher odds of high levels of depressive symptoms at T2 and T3. Moreover, psychosomatic symptoms at T1 predicted a high level of anxiety symptoms and diagnoses of anxiety disorders at T3. When analyzed separately, musculoskeletal symptoms predicted higher odds of having high levels of depressive symptoms at T2 and T3 and anxiety symptoms at T3, while psychological symptoms predicted high levels of anxiety symptoms at T2. Moreover, somatic symptoms at T1 predicted diagnoses of depression and anxiety disorders at T3. Sex did not moderate any of the relationships.

**Conclusions.** The study supports the idea that the decline of mental health in youth can seriously impact the psychological health of new generations of young adults.

## Background

In the last few decades, there has been an observed increase in adolescents' mental health problems. In a systematic review, Bor et al. (1) pointed out that while externalising problems were rather stable during the transition between the 20th and 21st centuries, internalising symptoms increased, especially among adolescent girls. Consistently, cross-cohort comparison studies, conducted in the UK, Greece, Iceland, the Netherlands and Norway, showed a long-term increase in adolescent emotional problems from 1970 to 2000 (2–5). These trends were only partly confirmed in the period between 2000 and 2018, when the increase in mental health problems concerned mainly the Nordic countries (6–8). All in all, these studies support the idea that the mental health of new generations of youth, especially those living in the Nordic countries, may become a more serious public health concern than it has been in the past.

Despite the general consensus that adolescents' mental health is in decline, especially in the Nordic countries, the extent to which youth mental health problems evolve into subsequently diagnosed disorders, such as depression or anxiety, in adulthood is still unknown. In this regard, Baxter et al. (9) have demonstrated that depression and anxiety disorders in adults did not increase during the period between 2000 and 2010. The authors have concluded that the observed decline of mental health in youth might be

explained by an increase in awareness of mental health rather than a real increase in mental health problems in adolescence. However, the discrepancy between the rates of youth mental health problems and adults' depression and anxiety disorders might also be due to the types of measures used to assess youth mental health problems. Most studies, especially in the last two decades, have not used standardized measures of depression and anxiety, but have focused on psychological distress and psychosomatic symptoms/complaints (5, 6, 8, 10–12).

To what extent are psychosomatic symptoms in adolescence related to the development of depressive and anxiety disorders in adulthood? A clear relation has been extensively demonstrated in cross-sectional studies (for a review, see 13). However, given the interrelation between mental health problems – i.e., psychosomatic symptoms, depression and anxiety – cross-sectional studies are limited in that they cannot establish what predicts what. In other words, the extent to which adolescent psychosomatic symptoms predict the development of depressive and anxiety disorders in adulthood cannot be understood with a cross-sectional design. Consequently, to disentangle the relation between psychosomatic symptoms and mental health problems, longitudinal studies are needed.

To our knowledge, longitudinal studies with population samples that examine the extent to which youth psychosomatic symptoms are related to adults' psychiatric disorders are scarce. Moreover, some of these studies, which do not control for baseline levels of psychiatric disorders (e.g., 14), cannot give conclusive answers in any case. Among the few exceptions, Bohman et al. (15) and Shanahan et al. (16) have reported that somatic symptoms, such as headaches, stomachaches, or muscular/joint aches in adolescence or childhood, predict psychiatric/mental disorders in adulthood, while Shelby et al. (17) have found that functional abdominal pain in adolescence is related to anxiety in early adulthood. However, these studies focus on specific somatic complaints, while most of the studies that show an increase in youth mental health problems include a more comprehensive set of symptoms, both psychological and physical, in line with the definition provided by WHO (18). Therefore, in order to investigate whether the observed increase in mental health problems among adolescents in the last two decades might lead to increases in adults' mental disorders, studies that focus on similar psychosomatic symptoms and follow adolescents into adulthood are needed. This study aims at filling this gap.

In this study, we focus on a cohort within the recent generation of Swedish adolescents that have shown an increase in psychosomatic symptoms and follow them into young adulthood. Given the concern raised about increased psychosomatic symptoms, our primary goal is to investigate whether elevated levels of psychosomatic symptoms in adolescence are related to subsequent depressive and/or anxiety disorders in young adulthood. Moreover, as previous research has highlighted the finding that mental health problems increase more in girls (e.g., 6, 8), we investigate whether psychosomatic symptoms elevate adult mental health problems in different ways in girls and boys. Finally, given that some of the above-mentioned studies have used only self-report measures to assess psychiatric disorders (16) while others have used diagnostic instruments (19), and that comparison between them is difficult, we used both self-report validated instruments and official diagnoses of disorders to increase the construct

validity of the study. In short, using two cohorts of adolescents, 13 and 15 years-old at baseline, the aims of the study were:

1. To investigate the extent to which high numbers of psychosomatic symptoms, i.e., psychological musculoskeletal, and somatic complaints, predict high of depressive and anxiety symptoms and diagnoses of depression and/or anxiety disorder 3 years and 6 years after the baseline.
2. To investigate the moderating roles of sex and SES in the associations between psychosomatic symptoms at baseline and depression and anxiety 3 and 6 years after baseline.

## Methods

### Study design

### Procedure

The longitudinal dataset comes from the “SALVE- Cohort” study, which aims at following youngsters from two cohorts, born in 1997 and in 1999. The youngsters were contacted when they were 12-13 and 14-15 years-old (T1), 15-16 and 17-18 years-old (T2), and 18-19 and 20-21 years-old (T3). At T1, participants were contacted by regular mail and invited to participate in the longitudinal study. They were informed that their participation was voluntary and that they could interrupt it at any time. They returned a self-reported questionnaire at T1, T2 and T3. The study was approved by the Ethical Review Board in Uppsala (Dnr. 2012/187).

### Participants

The original eligible adolescents were N=4712, of whom N=1868 (38.46%), responded at wave 1 (T1). At wave 2 (T2), the adolescents were contacted again, and 1575 of the original sample filled in the questionnaire (see 20). Finally, 1174 of the adolescents who participated at T1 also returned the questionnaire at T3.

In sum, the final sample that participated at T1 and T2 consisted of 1575 young people, of whom 58% (N=664) were female; 20% (N=319) had non-Scandinavian parents; 49% (N=804) were born in 1999 and 51% (N=774) in 1997. The final sample that participated at T1 and T3 consisted of 1174 young people, of whom 61% (N=722) were female; 20% (N=229) had non-Scandinavian parents; 51% (N=601) were born in 1999 and 49% (N=573) in 1997.

### Measures

Psychosomatic symptoms were measured using eight items from the WHO scale assessing the frequency of symptoms in the last three months (18). The answers range from never (0) to always (4). A

Confirmatory Factor Analysis (CFA) showed that the model best fitting these items was from a second-order confirmatory factor analysis with a latent dimension of psychosomatic symptoms that comprised three subdimensions of symptoms, namely psychological (i.e., feeling nervous, feeling irritable, feeling sleepy, 3 items), somatic (headache, stomachache, 2 items) and musculoskeletal (pain in the shoulders/neck, pain in the back/hips, pain in the hands/knees/legs/feet, 3 items) ( $\chi^2=77.03$ ,  $p>.01$ ,  $df=17$ ,  $RMSEA=.04$ ,  $CFI=.98$ ,  $SRMR=.034$ ). Accordingly, in the analysis we used both the total index of *Psychosomatic symptoms* ( $\alpha=.71$ ) and its three dimensions, namely *Psychological*, *Somatic*, and *Musculoskeletal symptoms*. Since our interest was in high levels of psychosomatic symptoms, the total index and the three subdimensions were dichotomized into high (top 25th percentile) and low.

Depressive symptoms. The Depression Self-Rating Scale Adolescent version, DSRSA (21) was used. The scale comprises 15 items based on the DSM-IV criteria for a major depressive disorder. The adolescents were asked about their depressive feelings in the last two weeks, with a yes/no response alternative. In accordance with the DSRSA scale, the index used in the analysis was calculated by adding reported symptoms, where each set of symptoms was counted only once (0-9 points). Cronbach's alphas were 0.81, 0.77 and 0.87, for T1, T2 and T3 respectively. Moreover, a dichotomous variable was created, in which adolescent boys and girls fulfilling the DSM-IV A-criterion were classified as having high symptoms of depression.

Anxiety. Anxiety symptoms were measured using the Spence Children's Anxiety Scale (SCAS, 22) at waves 1 and 2, while a short version of the same scale for adults was used at wave 3. The SCAS consists of 44 items, of which 38 cover all the six categories of anxiety disorder highlighted in the Diagnostic and Statistical Manual of Mental Disorders (23), while 6 are used as filler items to reduce negative bias. Alternative responses go from 0 (never) to 3 (always). The score of 33 has been identified as the cut-off for a diagnosis of generalized anxiety disorder (24) for children and adolescents. At wave 3, an adapted version with 15 items of the scale was used to assess the anxiety symptoms of the young adults. In the Swedish population, the cut-off for total anxiety disorder was  $>18$ . Cronbach alphas were 0.87, 0.89 and 0.91 and respectively. We used the above-mentioned cut-offs to classify the adolescents with high vs low symptoms.

Diagnoses of depressive and anxiety disorders. At T3, the young adults were asked whether they had received a diagnosis of depression and/or or some anxiety disorder, with yes/no as response alternatives.

Covariates. Sex (male, female), age (born in 1997 or 1999) and parents' country of birth (Scandinavian parents vs at least one of the parents born outside Scandinavia) were used as covariates in the analyses.

## Statistical Analyses

To investigate the effects of psychosomatic symptoms at T1 on levels of depression and anxiety at T2 and T3 and on diagnoses of depressive and/or anxiety disorder at T3, multivariate logistic models were constructed, controlling for initial levels of depression and anxiety respectively, and also for sex, age and

SES. First, we ran models assessing the effect of the dichotomized total index of psychosomatic symptoms (low vs high, based on the top 25th percentile at T1) on depressive or anxiety symptoms (low vs high, based on DSRS and RCMA cutoffs) at T2 and T3, and on diagnoses of depression or anxiety disorder at T3. Then, we ran models using the subscales of psychosomatic symptoms, i.e., high somatic, high musculoskeletal, and high psychological symptoms (high=top 25th percentile). Finally, to test whether sex was a moderator of the effects of psychosomatic symptoms, all the models were re-run adding an interaction term, i.e., psychosomatic symptoms \* sex.

## Results

### Attrition analyses

Attrition analyses were conducted to investigate whether adolescents with mental health problems were more likely to drop out at wave 2 or at wave 3. We compared the means at T1 of adolescents who participated at T1 and T2 or at T1 and T3 with the means at T1 of adolescents who participated at T1 and not at T2, or at T1 and not at T3, respectively. We found no differences in psychosomatic (T1-T2:  $F(1, 1803) = .592$ , n.s., T1-T3:  $F(1, 1803) = .013$ , n.s.), somatic (T1-T3:  $F(1, 1826) = .863$ , n.s., T1-T3:  $F(1, 1826) = .177$ , n.s.), musculoskeletal (T1-T2:  $F(1, 1800) = .275$ , n.s., T1-T3:  $F(1, 1800) = .005$ , n.s.), or psychological symptoms (T1-T2:  $F(1, 1815) = .123$ , n.s., T1-T3:  $F(1, 1815) = .027$ , n.s.). Moreover, adolescents who dropped out at T2 or at T3 were not more depressed (based on the DSRS cutoff) (T1-T2:  $\chi^2 = 1.11$ , n.s.; T1-T3:  $\chi^2 = 3.32$ , n.s) or more anxious (based on the RCMA cutoffs) (T1-T2:  $\chi^2 = .09$ , n.s.; T1-T3:  $\chi^2 = .968$ , n.s.) at T1 than those that did not drop out.

### Descriptive Analyses

The descriptive data are presented in Tables 1 and 2. At all the waves, girls reported higher psychosomatic symptoms in general, and also more specific somatic, psychological and musculoskeletal symptoms, than boys. They also reported higher percentages of depression and anxiety (see Tables 1 and 2).

Table 1

Means (standard deviations) and proportions (N) of the main variables of the study for the participants at T1 and T2.

<b>Samples T1 and T2</b>	<b>Male</b> % (N)//M(Sd)	<b>Female</b> % (N)//M(Sd)	<b>F or <math>\chi^2</math>, df, and p</b>	<b>Total sample</b> % (N)//M(Sd)
<b>Independent variables</b>				
Psychosomatic symptoms T1	8.05 (4.50)	10.65 (4.95)	108.547 (1, 1552), p<.001	9.53 (4.92)
Somatic symptoms T1	2.10 (1.44)	2.97 (1.53)	132.47 (1, 1570), p<.001	2.60 (1.51)
Psychological symptoms T1	2.33 (1.38)	3.00 (1.51)	79.27 (1, 1563), p<.001	2.71 (1.49)
Musculoskeletal symptoms T1	1.66 (1.55)	2.10 (1.66)	28.24 (1, 1550), p<.001	1.91 (1.63)
High anxiety symptoms (RCMAS) T1	5% (35)	19% (169)	60.31, df=1, p<.001	13% (204)
High depressive symptoms (DSRS) T1	6% (42)	16%(148)	318.84; df=1, p<.001	12% (190)
<b>Dependent variables</b>				
High anxiety symptoms (RCMAS) T2	6.5% (43)	33% (293)	152.22, df=1, p<.001	21% (336)
High depressive symptoms (DSRS) T2	14% (91)	33% (296)	74.64 (1), p<.001	25% (387)

Table 2

Means (standard deviations) and proportions (N) of the main variables of the study for the participants at waves 1 and 3.

<b>Samples T1 and T3</b>	<b>Male</b> % (N)//M(Sd)	<b>Female</b> %/(N)//M(Sd)	<b>F or <math>\chi^2</math>, df, and p</b>	<b>Total sample</b> %/(N)//M(Sd)
<b>Independent variables</b>				
Psychosomatic symptoms T1	8.07 (4.62)	10.52 (4.97)	69.84 (1, 1151), p<.001	9.57 (4.97)
Somatic symptoms T1	2.12 (1,44)	2.95 (1.53)	81.91 (1, 1169), p<.001	2.63 (1.57)
Psychological symptoms T1	2.98 (1.49)	3.00 (1.51)	59.15 (1, 1157), p<.001	2.71 (1.49)
Musculoskeletal symptoms T1	1.66 (1.58)	2.09 (1.66)	18.51 (1, 1148), p<.001	1.93 (1.64)
High anxiety symptoms (RCMAS) T1	5% (24)	17% (122)	33.99, df=1, p<.001	13% (147)
High depressive symptoms (DSRS) T1	6% (27)	15% (106)	20.95; df=1, p<.001	11% (133)
<b>Dependent variables</b>				
High anxiety symptoms (RCMAS) T3	16% (71)	37% (264)	59.52, df=1, p<.001	29% (336)
High depressive symptoms (DSRS) T3	25% (112)	44% (317)	44.09 (1), p<.001	37% (430)
Diagnosis of depression T3	3% (16)	9% (68)	14.57 (1), p<.001	7% (83)
Diagnosis of anxiety T3	3% (15)	9% (68)	15.86, df=1, p<.001	7% (83)

### Psychosomatic symptoms at T1 and depression and anxiety at T2 and T3

The participants who had a high number of psychosomatic symptoms at T1 had 77% (OR=1.79, 95% CI: 1.31-2.45) increased odds at T2, and 56% (OR=1.56, 95% CI: 1.12-2.17) increased odds at T3, of showing high depressive symptoms, after controlling for the initial values of depressive symptoms, sex, parents' country of birth and age. When the symptoms were analyzed separately, it emerged that musculoskeletal symptoms drove this relation. That is, youth with a high number of musculoskeletal symptoms had a 52% higher odds ratio at T2 (OR=1.52, 95% CI: 1.14-2.04), and a 49% higher one at T3 (OR=1.49, 95% CI: 1.11-2.03), of showing high depressive symptoms, after controlling for initial values of depressive symptoms, sex, parents' country of birth and age. Somatic and psychological symptoms were not significant (see Table 3).

Table 3

Odds ratios (OR) for the associations between mental health problems at baseline and depression at T2 and T3, controlling for sex, age and SES.

	High depressive symptoms T2 OR (95% CI)	High depressive symptoms T3 OR (95% CI)	Diagnosis of depressive disorder at T3 OR (95% CI)
<b>Model 1</b>			
Psychosomatic symptoms (1=top 25th percentile)	1.79 (1.31-2.45)	1.56 (1.12-2.17)	2.67 (1.55-4.62)
Number of depressive symptoms at T1	1.32 (1.24-1.41)	1.16 (1.08-1.24)	1.21 (1.09-1.35)
Sex	2.37 (1.79-3.13)	2.00 (1.53-2.62)	2.12 (1.18-3.76)
Age	.95 (.83-1.08)	1.09 (.97-1.25)	1.16 (.91-1.48)
Parents' country of birth	1.14 (.83-1.56)	.93 (.67-1.29)	.50 (.24-1.05)
<b>Model 2</b>			
Somatic symptoms (1=top 25th percentile)	.99 (.68-1.47)	1.12 (.75-1.67)	2.36 (1.36-4.11)
Musculoskeletal symptoms (1=top 25th percentile)	1.52 (1.14-2.04)	1.49 (1.11-2.03)	1.24 (.74-2.09)
Psychological symptoms (1=top 25th percentile)	1.45 (1.02-2.15)	1.31 (.87-1.96)	1.25 (.68-2.30)
Number of depressive symptoms (1=top 25th percentile)	1.32 (1.24-1.41)	1.16 (1.08-1.24)	1.23 (1.10-1.38)
Sex	2.46 (1.86-3.25)	1.99 (1.52-2.61)	2.11 (1.18-3.76)
Age	.94 (.83-1.07)	1.10 (.97-1.25)	1.17 (.92-1.49)
Parents' country of birth	.92 (.66-1.28)	.95 (.85-1.06)	.48 (.23-1.02)

Regarding the diagnosis of depressive disorders, adolescents who reported high levels of somatic symptoms at T1 showed an increased probability (OR=2.36, 95% CI: 1.36-4.11) of a diagnosis of depression at T3, even after controlling for depressive symptoms at baseline.

Psychosomatic symptoms at T1 did not increase the odds of showing high anxiety symptoms at T2, but they did at T3 (OR=1.72, CI=1.23-2.41) (see Table 4). When divided into subscales of symptoms, youth with high levels of psychological symptoms at T1 had 63% (OR=1.63, 95% CI: 1.07-2.64) higher odds of developing high anxiety symptoms at T2, but the relationship was no longer significant at T3 (see Table 4). However, youth with high levels of musculoskeletal symptoms at T1 had 51% higher odds (OR=1.51, 95% CI: 1.09-2.09) of suffering from a high level of anxiety symptoms at T3 (see Table 4).

Table 4

Odds Ratios (OR) for the associations between mental health problems at baseline and anxiety at T2 and T3, controlling for sex, age and SES.

	High anxiety symptoms T2 OR (95% CI)	High anxiety symptoms T3 OR (95% CI)	Diagnosis of anxiety T3
<b>Model 1</b>			
Psychosomatic symptoms (1=top 25th percentile)	1.21 (.86-1.71)	1.72 (1.23-2.41)	3.26 (1.91-5.56)
Number of anxiety symptoms at T1	1.08 (1.06-1.09)	1.05 (1.04-1.06)	1.02 (1.00-1.03)
Sex	4.35 (3.00-6.32)	2.09 (1.52-2.86)	2.21 (1.19-4.08)
Age	1.13 (.98-1.31)	1.11 (.97-1.28)	1.08 (.85-1.38)
Parents' country of birth	1.15 (.81-1.66)	.76 (.52-1.11)	.53 (.25-1.10)
<b>Model 2</b>			
Somatic symptoms (1=top 25th percentile)	1.01 (.66-1.54)	1.46 (.96-2.21)	2.02 (1.12-3.61)
Musculoskeletal symptoms (1=top 25th percentile)	1.15 (.82-1.60)	1.51 (1.09-2.09)	1.55 (.92-2.61)
Psychological symptoms (1=top 25th percentile)	1.63 (1.09-2.43)	1.11 (.72-1.70)	1.49 (.80-2.79)
Number of anxiety symptoms	1.08 (1.06-1.09)	1.05 (1.03-1.06)	1.02 (1.00-1.04)
Sex	4.47 (3.07-6.53)	2.08 (1.51-2.85)	2.44 (1.30-4.60)
Age	1.12 (.97-1.30)	1.12 (.98-1.29)	1.07 (.84-1.37)
Parents' country of birth	1.12 (.78-1.62)	.76 (.52-1.10)	.44 (.20-.96)

Regarding the diagnosis of anxiety disorder, adolescents with high psychosomatic symptoms at T1 had a higher probability (OR=3.26, CI=1.91-5.56) of an anxiety disorder at T3. Moreover, reporting somatic symptoms at T1 was associated with an increased probability (OR=2.02, 95% CI: 1.12-3.61) of a diagnosed anxiety disorder at T3.

## The Moderating Effect Of Sex

Sex did not moderate the relation between psychosomatic symptoms at T1 and depression at T2 or at T3. When it comes to the different subscales of symptoms, none of them had a different effect on depression in boys and girls (see Table 5).

Table 5

Odd ratios (OR) of the interaction terms used to assess the moderating role of sex in predicting the impact of mental health problems at baseline on depression and anxiety at T2 and T3.

<b>Models with depression as outcome</b>	<b>High depressive symptoms T2</b>	<b>High depressive symptoms T3</b>	<b>Diagnosis of depressive disorder T3</b>
	<b>OR (95% CI)</b>	<b>OR (95% CI)</b>	<b>OR (95% CI)</b>
Psychosomatic symptoms x sex	1.13 (.58-2.20)	1.26 (.64-2.50)	1.25 (.36-4.34)
Somatic symptoms x sex	3.17 (.99-10.16)	1.35 (.54-3.32)	.68 (.17-2.67)
Musculoskeletal symptoms x sex	.86 (.46-1.56)	.76 (.41-1.39)	2.40 (.59-9.74)
Psychological symptoms x sex	.71 (.33-1.52)	2.20 (.93-5.20)	.55 (.14-2.14)
<b>Models with anxiety as outcome</b>	<b>High anxiety symptoms T2</b>	<b>High anxiety symptoms T3</b>	<b>Diagnosis of anxiety disorder at T3</b>
	<b>OR (95% CI)</b>	<b>OR (95% CI)</b>	<b>OR (95% CI)</b>
Psychosomatic symptoms x sex	.68 (.29-1.60)	.90 (.43-1.89)	1.07 (.30-3.76)
Somatic symptoms x sex	2.66 (.62-11.50)	1.28 (.47-3.50)	1.45 (.27-7.68)
Musculoskeletal symptoms x sex	1.00 (.43-2.30)	.95 (.48-1.91)	3.47 (.69-17.51)
Psychological symptoms x sex	.46 (.19-1.17)	1.31 (.51-3.37)	.84 (.19-3.63)

Sex did not moderate the relation between psychosomatic symptoms at T1 and anxiety at T2 or T3. Moreover, none of the different subscales of symptoms had a different relation to anxiety in boys and girls (see Table 5).

## Discussion

The aim of the current study was to investigate whether an adolescent's mental health problems, in term of psychosomatic symptoms, can predict an increased risk of developing depressive and anxiety symptoms and/or receiving the diagnosis of a depressive or anxiety disorder in adulthood. We found that psychosomatic symptoms in adolescence increased the risk of both depression and anxiety three and six years later. Moreover, when looking into the types of symptoms that were most predictive, we found that adolescents who suffer from musculoskeletal symptoms are more at risk of developing high depressive and anxiety symptoms, while adolescents who complain of somatic symptoms, e.g., headaches and

stomachaches, have an increased risk of being diagnosed with an anxiety and/or depressive disorder six years later. Finally, these effects did not differ between girls and boys.

Our study shows that somatic symptoms, such as headaches and stomachaches, can predict depression and/or an anxiety disorder up to 6 years later. This result is in line with many cross-sectional studies that have pointed to relations between such symptoms and depression and anxiety (13, 25), even to the extent that it has been argued that these symptoms can be used as predictors of suicide attempts (26). There are also some longitudinal studies that have come to the same conclusion, showing that somatic symptoms independently predict both depression and anxiety, after controlling for depression and anxiety at baseline and using both diagnostic interviews and registry data on hospital-based mental health care (15, 16, 19). The present study corroborates this assertion, while adding that the contributions of both psychosomatic symptoms in general and of somatic symptoms in particular to mental health are the same for girls and boys. To the best of our knowledge, this is the first study to determine that, despite the different levels of mental health problems in girls and boys, somatic symptoms play a similar role in the development of depressive and anxiety disorders in adulthood.

Another important contribution of the study lies in its focus on the role of psychosomatic symptoms other than the somatic, i.e., on musculoskeletal symptoms. Although some studies have shown that musculoskeletal pain in children is associated with internalizing symptoms (27), to our knowledge, the long-term effects of symptoms of this kind have not been investigated before. We found that they may also contribute to a decline in mental health in adulthood, in both the short and the long term. When assessed via self-report measures, musculoskeletal symptoms were the only symptoms in the current study that were able to predict depression and anxiety three and six years later. This result highlights the importance of not underestimating the role of these symptoms. While somatic symptoms have been investigated extensively, and many hypotheses have been advanced regarding their relations with psychiatric disorders (for a review, see 28), the reasons why musculoskeletal symptoms might be associated with mental health problems are unknown. The symptoms may be a direct consequence of increased computer time in new generations of youth (29) entailing decreased physical activity, which in turn is associated with reduced mental health, whether or not in the form of diagnosed problems (30). This might provide an explanation for why this association has not involved real-life diagnoses of depressive and anxiety disorders of the kind our study has highlighted. However, the hypothesis needs to be confirmed in future studies.

## Limitations And Strengths

The study has some limitations. First, the psychosomatic symptoms were assessed only at baseline and not at the follow-ups. This limits the opportunity to understand how they might develop in late adolescence and young adulthood, and their contribution to possible mental ill-health. Moreover, as in all the other study population studies in this arena, initial acceptance of participation in the study was somewhat low (around 40%). Although this rate is quite common in this type of study, we cannot exclude the possibility that low participation impacted the external validity of the results, especially when

considering that non-participants are often more at risk of the negative outcomes considered (e.g., depression). Attrition, however, is unlikely to have influenced the results as the adolescents who stayed in the project did not differ from those who dropped out at T1.

This study also has some strengths. First, to the best of our knowledge, it is the first study to examine the longitudinal effects of different subcategories of psychosomatic symptoms on mental health. Second, the use of different measures to assess mental health problems, i.e., self-reported symptoms and clinical diagnoses, provide good support for the validity of the results.

## **Conclusions**

Many recent studies have pointed to an increase in mental health problems, often in terms of psychosomatic symptoms, in the new generations of youth. This study demonstrates that high levels of psychosomatic symptoms in adolescence increase the risk of developing high levels of both depressive and anxiety symptoms and depressive and anxiety disorders. Therefore, it provides a rationale closely to monitor adolescents, boys and girls to the same extent, and especially those with somatic and musculoskeletal symptoms, in order to prevent the development of serious disorders in adulthood.

## **Declarations**

### **Ethics approval and consent to participate**

The study was approved by the Ethical Review Board in Uppsala (Dnr. 2012/187). All the adolescents and their parents gave written informed consent to participate in the study.

### **Consent for publication**

Not applicable.

### **Availability of data and materials**

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

### **Competing interests**

All the authors declare they have no competing interests.

### **Funding**

The study was supported by a grant from the Swedish Research Council for Health, Working Life and Welfare (FORTE) (2019-00492) to F.G. Also, funding of the data collection for the study was supported by grants from the Söderström König Foundation (SLS-559921, SLS-655791, SLS-745221), Åke Wiberg's

Foundation (M15-0239), and the Swedish Research Council for Health, Working Life and Welfare (FORTE) (2015-00897) to CÅ.

None of these organizations played any role in the study design, collection, analysis or interpretation of the data, writing the manuscript, or the decision to submit the paper for publication.

### **Authors' contributions:**

F.G. developed the concept and design of the study, and also drafted the manuscript. P.L. collaborated with F.G. in drafting the manuscript. K.N. and C.Å. developed the study concept and design for the whole SALVe cohort project, and helped with interpretation of the results of this study. P.L., K.N. and C.Å. critically revised the text. All authors read and approved the final manuscript.

### **Acknowledgements**

Not applicable

### **Authors' information**

FG is an Associate Professor of Psychology at the Department of Public Health, Stockholm University. Her research interests cover the study of the protective and risk factors for youth development, with a recent focus on youth mental health. She is also interested in prevention programs that focus on changing negative trajectories of children with disruptive behaviors and on improving well-being of normative children.

KN is a Professor of Psychiatry at the Centre for Clinical Research, Uppsala University. His major research interests include determinants and correlates to psychiatric health problems in adolescence and gene-environmental interactions.

CÅ is a Professor of Social Medical Research at the Department of Public Health and Caring Sciences, Uppsala University. Her main research interests are depression and antisocial behaviour in adolescents, focusing on the influence of social factors and gene-environment interaction.

PL is an Associate Professor and Senior Lecturer at the Department of Public Health Sciences at Stockholm University. His major research interest includes determinants and correlates of alcohol use in adolescence.

## **References**

1. Bor W, Dean AJ, Najman J, Hayatbakhsh R. Are child and adolescent mental health problems increasing in the 21st century? A systematic review. Australian New Zealand journal of psychiatry. 2014;48(7):606–16.

2. Collishaw S. Annual research review: secular trends in child and adolescent mental health. *J Child Psychol Psychiatry*. 2015;56(3):370–93.
3. Sigfusdottir ID, Asgeirsdottir BB, Sigurdsson JF, Gudjonsson GH. Trends in depressive symptoms, anxiety symptoms and visits to healthcare specialists: a national study among Icelandic adolescents. *Scandinavian journal of public health*. 2008;36(4):361–8.
4. Sweeting H, Young R, West P. GHQ increases among Scottish 15 year olds 1987–2006. *Soc Psychiatry Psychiatr Epidemiol*. 2009;44(7):579.
5. Hagquist C. Discrepant trends in mental health complaints among younger and older adolescents in Sweden: an analysis of WHO data 1985–2005. *J Adolesc Health*. 2010;46(3):258–64.
6. Potrebny T, Wiium N, Lundegård MM-I. Temporal trends in adolescents' self-reported psychosomatic health complaints from 1980-2016: A systematic review and meta-analysis. *PLOS one*. 2017;12(11):e0188374.
7. De Looze M, Cosma A, Vollebergh W, Duinhof E, de Roos S, van Dorsselaer S, et al. Trends over time in adolescent emotional wellbeing in the Netherlands, 2005-2017: links with perceived schoolwork pressure, parent-adolescent communication and bullying victimization. *J Youth Adolesc*. 2020;49(10):2124–35.
8. Cosma A, Stevens G, Martin G, Duinhof EL, Walsh SD, Garcia-Moya I, et al. Cross-national time trends in adolescent mental well-being from 2002 to 2018 and the explanatory role of schoolwork pressure. *J Adolesc Health*. 2020;66(6):50-S8.
9. Baxter AJ, Scott KM, Ferrari AJ, Norman RE, Vos T, Whiteford HA. Challenging the myth of an “epidemic” of common mental disorders: trends in the global prevalence of anxiety and depression between 1990 and 2010. *Depress Anxiety*. 2014;31(6):506–16.
10. Bremberg S. Mental health problems are rising more in Swedish adolescents than in other Nordic countries and the Netherlands. *Acta Paediatr*. 2015;104(10):997–1004.
11. Potrebny T, Wiium N, Haugstvedt A, Sollesnes R, Torsheim T, Wold B, et al. Health complaints among adolescents in Norway: A twenty-year perspective on trends. *PloS one*. 2019;14(1):e0210509.
12. Ottova-Jordan V, Smith OR, Gobina I, Mazur J, Augustine L, Cavallo F, et al. Trends in multiple recurrent health complaints in 15-year-olds in 35 countries in Europe, North America and Israel from 1994 to 2010. *The European Journal of Public Health*. 2015;25(suppl\_2):24–7.
13. Campo JV. Annual Research Review: Functional somatic symptoms and associated anxiety and depression—developmental psychopathology in pediatric practice. *J Child Psychol Psychiatry*. 2012;53(5):575–92.
14. Hotopf M, Mayou R, Wadsworth M, Wessely S. Temporal relationships between physical symptoms and psychiatric disorder. *The British journal of psychiatry*. 1998;173(3):255–61.
15. Bohman H, Låftman SB, Cleland N, Lundberg M, Päären A, Jonsson U. Somatic symptoms in adolescence as a predictor of severe mental illness in adulthood: a long-term community-based follow-up study. *Child Adolesc Psychiatry Mental Health*. 2018;12(1):1–12.

16. Shanahan L, Zucker N, Copeland WE, Bondy C, Egger HL, Costello EJ. Childhood somatic complaints predict generalized anxiety and depressive disorders during adulthood in a community sample. *Psychological medicine*. 2015;45(8):1721.
17. Shelby GD, Shirkey KC, Sherman AL, Beck JE, Haman K, Shears AR, et al. Functional abdominal pain in childhood and long-term vulnerability to anxiety disorders. *Pediatrics*. 2013;132(3):475–82.
18. Åslund C, Starrin B, Nilsson KW. Social capital in relation to depression, musculoskeletal pain, and psychosomatic symptoms: a cross-sectional study of a large population-based cohort of Swedish adolescents. *BMC Public Health*. 2010;10(1):1–10.
19. Bohman H, Jonsson U, Päären A, von Knorring L, Olsson G, von Knorring A-L. Prognostic significance of functional somatic symptoms in adolescence: a 15-year community-based follow-up study of adolescents with depression compared with healthy peers. *BMC Psychiatry*. 2012;12(1):1–10.
20. Vadlin S, Åslund C, Nilsson KW. A longitudinal study of the individual-and group-level problematic gaming and associations with problem gambling among Swedish adolescents. *Brain behavior*. 2018;8(4):e00949.
21. Svanborg P, Ekselius L. Self-assessment of DSM-IV criteria for major depression in psychiatric out- and inpatients. *Nord J Psychiatry*. 2003;57(4):291–6.
22. Spence SH. Structure of anxiety symptoms among children: a confirmatory factor-analytic study. *J Abnorm Psychol*. 1997;106(2):280–97.
23. Association AP. *Diagnostic and statistical manual of mental disorders*. 4th ed. Washington, DC: Author.; 1994.
24. Olofsdotter S, Sonnby K, Vadlin S, Furmark T, Nilsson KW. Assessing adolescent anxiety in general psychiatric care: Diagnostic accuracy of the Swedish self-report and parent versions of the Spence Children's Anxiety Scale. *Assessment*. 2016;23(6):744–57.
25. Campo JV, Fritsch SL. Somatization in children and adolescents. *Journal of the American Academy of Child Adolescent Psychiatry*. 1994;33(9):1223–35.
26. Heinz A, Catunda C, van Duin C, Willems H. Suicide prevention: Using the number of health complaints as an indirect alternative for screening suicidal adolescents. *J Affect Disord*. 2020;260:61–6.
27. Mikkelsen M, Sourander A, Piha J, Salminen JJ. Psychiatric symptoms in preadolescents with musculoskeletal pain and fibromyalgia. *Pediatrics*. 1997;100(2):220–7.
28. Beck JE. A developmental perspective on functional somatic symptoms. *J Pediatr Psychol*. 2008;33(5):547–62.
29. Sigmund E, Sigmundová D, Badura P, Kalman M, Hamrik Z, Pavelka J. Temporal trends in overweight and obesity, physical activity and screen time among Czech adolescents from 2002 to 2014: A national health behaviour in school-aged children study. *Int J Environ Res Public Health*. 2015;12(9):11848–68.
30. Cao H, Qian Q, Weng T, Yuan C, Sun Y, Wang H, et al. Screen time, physical activity and mental health among urban adolescents in China. *Preventive medicine*. 2011;53(4-5):316–20.