

# Commitment to protective measures during COVID-19 pandemic in Syria: a nationwide cross-sectional study.

**Mosa Shibani** (✉ [Moosa.Shibani@gmail.com](mailto:Moosa.Shibani@gmail.com))

Faculty of Medicine, Syrian Private University, Damascus, Syria <https://orcid.org/0000-0002-4424-2728>

**Mhd Amin Alzabibi**

Faculty of Medicine, Syrian Private University, Damascus, Syria

**Abdul Fattah Mohandes**

Faculty of Medicine, University of Aleppo, Aleppo, Syria. <https://orcid.org/0000-0002-6506-0851>

**Humam Armashi**

Faculty of Medicine, Syrian Private University, Damascus, Syria

**Tamim Alsuliman**

Hematology and Cell Therapy Department, Saint-Antoine Hospital, AP-HP Sorbonne University, Paris, France.

**Angie Mouki**

Faculty of Pharmacy, Maykop State Technological University, Maykop, Russia.

**Hlma Ismail**

Faculty of Medicine, Syrian Private University, Damascus, Syria

**Shahd Alhayk**

Faculty of Medicine, Syrian Private University, Damascus, Syria

**Ahmad abdulateef Rmman**

Faculty of Medicine, Syrian Private University, Damascus, Syria

**Hala Adel Almohi Alsaid Mushaweh**

Faculty of Medicine, University of Aleppo, Aleppo, Syria.

**Marah Mansour**

Faculty of Medicine, Tartous University, Tartous, Syria.

**Elias Battikh**

Department of Internal Medicine, Hamad Medical Corporation, Doha, Qatar.

**Naram Khalayli**

Faculty of Medicine, Damascus University, Damascus, Syria.

**Bisher Sawaf**

Department of Internal Medicine, Hamad Medical Corporation, Doha, Qatar.

**Mayssoun Kudsi**

Department of Rheumatology, Almowasa University Hospital, College of Medicine, Damascus University, Damascus, Syria

---

## Research Article

**Keywords:** COVID-19, Prevention, Syria, Commitment

**Posted Date:** December 6th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-1140688/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

## Background

Severe acute respiratory syndrome coronavirus 2 continues to impose itself on all populations of the world. Given the slow pace of vaccination in the developing world and the absence of effective treatments, adherence to precautionary infection control measures remains the best way to prevent the COVID-19 pandemic from spiraling out of control. In this study, we aim to evaluate the extent to which the Syrian population adheres to these measures and analyze the relationship between demographic variables and adherence.

## Methods

This cross-sectional study took place in Syria between January 17 and March 17, 2021. A structured self-administered questionnaire was used to collect the data. The questionnaire was distributed in both electronic and printed versions. Collected data were analyzed using SPSS v.25. The chi-square test was used to address the correlation between adherence and demographic variables.

## Results

Of the 10083 reached out, only (74.6%) responded. Of them, 4026 (53.5%) were women, 3984 (52.9%) were single, and 1908 (25.3%) had earned university degrees. 5286 (70.25) were classified as the good adherers to protective measures. Statistically significant differences across age, sex, marital status, financial status, employment, and educational attainment when correlated against commitment to preventive measures. Young age, female sex, good financial status, and having a full-time job and post-graduate education were positively correlated to the stronger commitment to preventive measures. Furthermore, those who believed that COVID-19 poses a major risk to them or society were more committed to preventive measures than those who did not.

## Conclusion

The participants in this study generally showed a high level of adherence to the preventive measures compared to participants in other studies from around the world, with some concerns regarding risk perception and the sources of information they depend on. Public health and community education efforts should focus on maintaining, if not expanding, this level of commitment, which would mitigate the pandemic's impact on Syrian society.

## Background:

The continued spread of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) - the causative agent of coronavirus disease 2019 (COVID-19) - has impacted all aspects of life worldwide, with over 250 million cases and over 5 million deaths (1). On 11 March 2020, the World Health Organization (WHO) officially declared COVID-19 to be a global pandemic (2), and recommended comprehensive strategies to prevent the spread of the virus (3). Since person-to-person transmission mainly occurs via respiratory droplets, close contact with infected individuals during talking, sneezing, coughing, and indirect contact with contaminated objects or surfaces (4, 5), the most important recommendations include self-isolation, physical distancing, wearing face masks, and practicing hand hygiene (3, 6). Many governments have implemented these virus-mitigation measures to contain the spread of the virus and protect vulnerable populations from infection. These cooperative efforts are important in lowering mortality rates and preventing health care systems from being overburdened. However, populations must be highly committed to these measures to ensure their success. In the absence of effective treatments and in light of recent evidence showing decreased vaccine-induced immunity after 5-7 months (7), preventive infection-control measures remain the best hope for containing the disease. The Syrian government began implementing precautionary measures to pre-empt the spread of the disease before the first case was even reported on 22 March 2020 (8, 9). All schools, colleges, commercial and leisure centers, gyms, and places of worship were closed, and a 6pm-to-6am curfew was put into effect. However, these measures lasted only for two months (March to May 2020) (8). A recent nationwide study in Syria reported good levels of awareness among the Syrian population regarding COVID-19 in general and preventive measures in particular (10). Almost all participants (99%) were aware that proper hand hygiene, avoidance of crowded places, isolation at home, and wearing masks in public places are the main preventive measures (10). Information about the Syrian population's general knowledge regarding infection control will not only help to inform policy-makers as they make important decisions on how to best confront this pandemic, but it is also important to measure the extent to which the population adheres to these measures so that gaps between knowledge and practice can be addressed. As of this writing, 45,468 laboratory-confirmed cases and 2,637 casualties of COVID-19 have been reported by the Syrian Ministry of Health (11). However, considering that testing has been limited in scale and that the cost of test kits is relatively high, it is likely that official numbers are deceptively low and do not reflect the severity of the pandemic in Syria. The war in Syria has raged for over 10 years and continues to impose massive burdens on the population, including economic, social, and educational challenges (12). A consequence of this widespread conflict is the largest refugee crisis since World War II (13). The Syrian healthcare system, already devastated by the war and suffering staffing, supply, and funding shortages, has all but collapsed in the face of the pandemic. The Syrian economy has been particularly hard hit, and due to recent inflation, the number of people in need (PIN) is expected to increase from 11 million in 2020 to 13.3 million in 2021 (12). The rising price of personal protective equipment and medical supplies, as well as other COVID-19 related factors that have increased the cost of living, has resulted in the majority of Syrian families being unable to afford leaving their jobs and sacrificing income to self-isolation and physical distancing measures (14). In this study, we aim to measure the commitment of the Syrian population to infection prevention and control (IPC) measures (such as hand washing, wearing masks

and gloves, and avoidance of handshaking and face-touching), measure perceived risk regarding COVID-19, and study the correlation between commitment and some demographic variables.

## **Methods:**

### **Study design, setting, and participants:**

A nationwide cross-sectional study was performed between January 17 to March 17, 2021. Data was collected using a structured self-administered questionnaire which was distributed to a nationally representative sample of Syrian people. The questionnaire, based on a previous study, was modified to be suitable for Syrian society (15–17). It was then piloted on a sample of 15 people to ensure clarity, and adjustments were made based on their feedback. Chain-referral and convenience sampling methods were employed by distributing the questionnaire in two formats: electronically as a Google Form survey via social media and messaging platforms (Facebook, Whatsapp, Twitter), and physically as hard copies to patients, their companions, and workers in public hospitals in each of Damascus, Aleppo, Homs, Tartous, Hama, and Sweida governorates. The sample size was calculated using OpenEpi online software available at "<https://www.openepi.com/SampleSize/SSPropor.htm>". According to data from the United Nations, the estimated population of Syria in 2019 was about 18 million (18); based on this figure, the sample required to represent the total population was calculated to be 7336, with a confidence level of 95% and a confidence interval of 1.14.

Inclusion criteria were that the person is: (1) 18 years old or above, (2) literate, (3) a Syrian citizen living in Syria, and (4) willing to complete the questionnaire. Of the 8083 participants who responded to the questionnaire, 551 were excluded for not meeting the inclusion criteria (17 withdrew their consent to participate, 543 were not Syrians or lived outside of Syria), yielding a final sample size of 7,531 participants.

## **Measures:**

The questionnaire consisted of 32 questions divided into 3 sections:

### **1. Socio-demographic characteristics :**

13 questions about age, gender, marital status, nationality, governorate of origin, place of residence (urban or rural), financial status, employment status, educational level, father's and mother's educational level, health insurance coverage, and work or study in a healthcare-related field. Age was divided into 4 groups: 18-24, 25-44, 45-65, and >65 years. Governorates were divided into 5 categories based on geographical location: 1- Central governorates (Damascus, Rif Dimashq, Hama, Homs), 2- Eastern governorates (Deir ez-Zor, Al-Hasakah, Ar-Raqqah), 3- Western governorates (Latakia, Tartous), 4- Northern governorates (Aleppo, Idlib), 5-Southern governorates (Daraa, Quneitra, As-Suwayda).

### **2. COVID-19 general information:**

5 questions about prior infection with SARS-CoV-2, the risk this virus poses to the individual and to Syrian society as a whole, and participants' source of information.

### **3. Commitment to infection prevention and control (IPC) measures.**

12 yes-no statements about various IPC measures including: wearing a face mask, social event cancellation or postponement, self-isolation, cleaning or disinfecting touched items, carrying sanitizing hand-gel, reduced face-touching, healthy diet, avoiding people who have cold or flu-like symptoms, using tissues when sneezing or coughing, and washing hands with soap and water.

#### **Statistical analysis:**

Data from the hard copy questionnaires were entered manually by the investigators to the original Google Forms online questionnaire that was used to collect online data, after which it was exported to a Microsoft Excel spreadsheet. The raw data was then encoded in Excel to make it compatible with the statistics software. A 13 point scale was used to measure the level of commitment to IPC measures. Each individual measure was given one point, then each participant was categorized into one of three categories based on how many protective measures he/she applied: 1- Low commitment (0 - 3 protective measures), 2- Moderate commitment (4 - 8 protective measures) and 3- High commitment (9 - 12 protective measures). We used Statistical Package for Social Sciences version 25.0 ( SPSS Inc., Chicago, IL, United States) to analyze the data. Categorical variables were reported as frequencies and percentages. Pearson's chi-square test was used to study the associations between categorical groups. A  $p$ -value  $< 0.05$  was considered statistically significant.

#### **Ethical considerations:**

The study protocol was approved by the respective Research Ethics Committee at each of Damascus, Aleppo, Tartous, and Syrian Private Universities, and the ethics committees of each hospital from which data was collected. Informed consent was obtained from every participant prior to participation.

#### **Results:**

##### ***Participant characteristics:***

Of 10083 participants, 7531 agreed to participate and fully completed the questionnaire (response rate = 74.7%), of which 3505 (46.5%) were males and 4026 (53.5%) were females. The dominant age group was 18-24 years old: 3124 (41.5%), followed by 25 - 44 years old: 2338 (31%), 45-65 years old: 1686 (22.4%), and  $> 65$  years: 383 (5.1%). Over the half of the study population originated from the central governorates: 4376 (58.1%), and 1165 (15.5%) from the western governorates. Regarding marital status, 3984 (52.9%) were single, 2825 (37.5%) were married, and 500 (6.6%) were in a relationship. The majority of participants were financially middle class: 3241 (43%), and only 371 (4.9%) had excellent financial status. University students (2906 ,38.6%) and university graduates (1908, 25.3%) represented the majority

of responders, while only 324 (4.3%) did not receive any formal education. When asked if employed or studying in a healthcare-related field, 4936 (65.5%) of the participants answered “no”. (Table 1)

### ***COVID-19 related information:***

When asked about a previous infection with COVID-19, the majority of participants (4549 ,60.4%) answered “no”, 2742 (36.4%) said they experienced symptoms, and only 240 (3.2%) had a PCR-confirmed infection. On the other hand, when the participants were asked if they know someone who has had a PCR-confirmed infection, 4571 (60.7%) said “yes”. Regarding the imposed risk of COVID-19 on Syrian society, 4639 (61.6%) believe it poses a major risk, 1621 (21.5%) believe it poses a minor risk, and 314 (4.2%) said there is no risk at all. However, when asked about the extent to which the virus poses a personal risk, 3224 (42.8%) replied “minor risk”, 2218 (29.5%) replied “major risk”, 834 (11.1%) replied “no risk at all”, and 1255 (16.7%) did not know.(Table 2). Healthcare workers and social media (Facebook,Whatsapp, Youtube, Telegram, Instagram, etc.) were the main source of COVID-19 related information for (50.9%) and (46.3%) participants respectively. (Figure 1)

### ***Commitment to preventive measures:***

The vast majority of participants (5286, 70.2%) showed high commitment, 1888 (25.1%) showed moderate commitment, and 357 (4.7%) showed poor commitment. (Table 3).

The most-practiced preventive measure among the study population was “covering the mouth/nose when coughing or sneezing” (6887, 91.4%) followed by “hand washing with soap and water more often than usual” (6637, 88.1%), “avoid people with cold or flu-like symptoms” (6645, 88.2%), “wearing a face mask in public or crowded places” (6577, 87.3%) and “avoiding crowded places” (6317, 83.9%). Half of the study population started to follow a healthy diet 3747 (49.8%) and 4950 (65.7%) reduced their attendance at school, college, university or work.(Table 4)

### ***Correlations between commitment to preventive measures and participants characteristics:***

Chi-square univariate analysis showed a statistically significant difference between males and females regarding commitment to preventive measures. 76.4% of females and 63.1% of males were categorized as highly committed, while only 3.4% of females and 6.2% of males were categorized as poorly committed ( $\chi^2=160.683$ ,  $p$ -value<0.0001). A significant association was found between age groups and adherence to preventive measures: 25-44 year old participants were most committed to IPC measures with 1780(76.1%) categorized as ‘high’, followed by 18–24, 45–65 and >65 age groups: 2269(72.6%), 1022(60.6%) and 215(56.1%) respectively ( $\chi^2 = 204.974$ ,  $p$ -value <0.0001). Participants in relationships (76.6%,  $\chi^2 = 92.002$ ,  $p$ -value <0.0001) and those from the western governorates of Syria (78.8%,  $\chi^2 = 184.079$ ,  $p$ -value <0.0001) were more committed to IPC measures than their single counterparts and those in other parts of the country. Commitment to preventive measures was significantly associated with residency and financial status, with urban residents (71.6% vs. 35.4%,  $\chi^2 = 59.106$ ,  $p$ -value <0.0001) and those in good financial status (76.6%,  $\chi^2 = 279.195$ ,  $p$ -value <0.0001) were the most committed groups.

Participants with post-graduate education (80.6%,  $\chi^2 = 640.976$ ,  $p$ -value  $<0.0001$ ) and students with full time jobs (77.1%,  $\chi^2 = 129.431$ ,  $p$ -value  $<0.0001$ ) were the most committed to preventive measures. (Table 5) Our results revealed that people who believe that COVID-19 poses a major risk to Syrian society were more committed to IPC measures, with 77.4% being highly committed compared to 59.5% from the 'minor risk' group and 43.0% from the 'no risk at all' group. Similarly, those who believe that COVID-19 poses a major risk to them personally were more committed to preventive measures, as (79.5%) of them were in the high commitment category. (Table 6)

## Discussion:

The COVID-19 pandemic has significantly impacted humanity and forced governments across the world to adopt extensive infection prevention and control measures with varying degrees of severity (19). Syria is a low-income country that has been ravaged by civil war for over a decade, diminishing the ability to adequately respond to the pandemic and impose meaningful quarantines. As such, results of measures adopted by other countries and regions cannot be relied upon to predict the course of the pandemic in Syria, and the extraordinary difficulties facing the country and the realities on the ground must be taken into account when uniquely assessing the situation in Syria. The efficiency and impact of infection prevention and control (IPC) measures can be optimized by obtaining insights into the population's current commitment to such measures. To the best of our knowledge, this is the first nationally representative study to offer insights into people's adherence to preventive measures during the COVID-19 pandemic in Syria. Our findings revealed that 70.2% of the population claim to adhere to most of the preventive measures asked about in the questionnaire. This level of adherence is similar to that found in a Belgian study, and better than that from an Ethiopian study (20, 21). The majority of our sample were young, and the age distribution of our population was generally consistent with the demographic data reported by the Central Bureau of Statistics (CBS), Damascus, Syria (22). According to the latest CBS report, 40% of the Syrian population were below 24 years old, and 25.5% were 25–44 years old (compared to 41.5% and 31%, respectively, of our study population) (22). The importance of face masks in reducing the spread of the virus is supported by numerous studies (23, 24). One study suggests that complete eradication of the disease can be achieved if 80% of the population uses face masks effectively (25). The vast majority of our participants (87.3%) were committed to wearing face masks in public spaces. This proportion is predictably lower than those from studies in China (98.0%) and Hong Kong (98.8%), where mask-wearing has been ingrained in the culture for decades, but considerably higher than in studies from Northwest Ethiopia (32.42%), Ethiopia (13.9%), Saudi Arabia (56.4%), and the United Kingdom (3.1%) (15, 21, 26–28).

Low income and unemployment were correlated with lower adherence to IPC measures, while higher income and gainful employment was correlated with higher adherence; only 55% of low income responders were highly committed to IPC measures, compared to 76.6% of those with good financial status. This may owe to the high cost of commitment to protective measures, which is prohibitively

expensive for a significant proportion of the Syrian population. The Syrian pound has lost 35% of its value against the US dollar in the last year alone (14), and the percentage of the population living in poverty is 90% and rising (29). While screening tests and social distancing might be considered cost-effective elsewhere in the world, this is not the case for the Syrian population (PCR screens cost \$50 each and are not subsidized). Infection control measures such as sweeping lockdowns have only recently affected most of the world's economies, whereas the Syrian economy, already suffering from a decade of war and crippling sanctions, has been devastated by lockdowns and other pandemic-related economic pressures. As such, Syria's population and healthcare system are in desperate need of international support in the form of financial grants and donations of personal protective equipment, drugs, medical supplies, and vaccines. Income and employment-related results are in line with studies from China and Ethiopia, but do not align with those of a study from Saudi Arabia, likely due to vastly different socio-economic dynamics (15, 26, 28).

UNICEF reported that after a decade of war in Syria, more than half of children continue to be deprived of education (30). The enormous scale of the education crisis is extremely worrying, as it threatens not only the future of an entire generation of children and the country as a whole, but also the important role of schools as conduits for health literacy and education about diseases and the importance of infection control. Our study revealed an important correlation between education and adherence to IPC measures, with commitment increasing significantly as the level of education increases. On one end of the spectrum, only 35.2% of uneducated participants adhere to protective measures, compared to 80.6% of participants with postgraduate education on the other end. These findings are consistent with studies from China, Ethiopia, and Germany (16, 26, 28).

Numerous studies have shown that risk perception can be considered a determinant of individual behavior during a disease outbreak (31–33). The more risk perceived, the more likely people are to adhere to preventive measures. Some studies go even further and suggest that it is important to differentiate between the 'experiential' and 'affective' components of risk perception (34, 35). Earlier research demonstrated that experiential risk perception, "the gut feeling of being vulnerable to risk", was positively associated with applying personal protective actions, such as vaccination and sun protection (36, 37). Our study seems to support this assumption, and other studies in Italy, Northwest Ethiopia reported the same observation (21, 38).

Several previous studies showed that the level of knowledge correlates directly with adherence to preventive measures (28, 39). Ideally, the public should be well-informed by reliable sources of information. Unfortunately, our participants' reliance on untrusted sources on social media is part of a global trend in which misinformation is rampant and pervasive. Social media tends to be the most expedient means of obtaining information for many people, and studies have shown that social media is a fertile and target-rich environment for spreading misinformation and conspiracy theories that negatively affect the quality of the public's knowledge (40–42). Since it is impossible to fully control what is published on social media, local and global health authorities must enhance their presence on these platforms and use engaging content and effective methods to spread awareness and accurate

information. Studying the public's perception and behavior toward COVID-19 provides valuable insight which can help policymakers and healthcare providers to address the knowledge gaps that negatively affect people's perception and behavior, thereby improving the national response to this pandemic. We encourage all concerned institutions to invest the time, resources, and expertise necessary to successfully and significantly leverage social media platforms to drive public health education and COVID-19 awareness campaigns. The rebuilding and rehabilitation of schools must be prioritized, and infection control measures incorporated into the curriculum. Special accommodations should be made for low-income people and families, in the form of distributing infection control kits (composed of a reusable face mask and hand sanitizer) and securing their income when proven to be sick to encourage them to self-quarantine.

## **Conclusion:**

Despite the high level of commitment to infection prevention and control (IPC) measures demonstrated by the participants in our study, it is necessary to stress the importance of continuing this commitment throughout the pandemic. It is recommended that local and international health authorities carry out continuous awareness campaigns with the aim of reminding the population of the importance of consistently applying IPC measures. Moreover, the population should be educated about how to identify and avoid misinformation on social media and to rely on reliable sources of information. Because of the economic and humanitarian situation in war-torn Syria, it is necessary for all concerned bodies and organizations to take serious action and provide appropriate assistance to the healthcare system to help contain this pandemic.

## **Limitations:**

This study is subject to some limitations. First, as a cross-sectional study it may not be able to determine causation, therefore more longitudinal studies are recommended. Second, distributing the questionnaire online only will lead to selection bias, as most people with internet access tend to be younger and wealthier. To minimize this bias we distributed the questionnaire both online and as hard copies. Third, many questions were subject to recall bias. Finally, the economic status question was subjective since the value of the Syrian pound is not stable and the exchange rate continues to fluctuate. This continues to affect the purchasing power of the local currency, with many families whose income was once adequate falling below the poverty line.

## **Declarations:**

### **Ethics approval and consent to participate:**

The study protocol was approved by the Research Ethics Committee in each of Damascus, Aleppo, Tartous, and Syrian Private Universities, and the ethical committees in the concerned hospitals. Informed consent was obtained from every participant prior to participation.

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

### **Availability of data and materials:**

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

### **Competing interest:**

The authors declare that they have no competing interests

### **Funding:**

This project did not receive any funding from any agencies in the public, commercial, or non-profit sectors.

### **Authors' contributions:**

### **Acknowledgments:**

We would like to express our thanks to all the people and organizations that helped in distributing this survey, especially: Syrian Researchers Organization, Impact Makers Team, Syrian Drugs Up To Date Platform, MedDose Organization (NGO).

## **References:**

1. World Health Organization (WHO). WHO Coronavirus (COVID-19) Dashboard 2021 [cited 2021 December]. Available from: <https://covid19.who.int/table>.
2. WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020 [press release]. 11 March 2020.
3. World Health Organization (WHO). Advice for the public: Coronavirus disease (COVID-19) 2021 [updated 1 October 2021; cited 2021 4 December]. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/advice-for-public>.
4. Liu J, Liao X, Qian S, Yuan J, Wang F, Liu Y, et al. Community Transmission of Severe Acute Respiratory Syndrome Coronavirus 2, Shenzhen, China, 2020. *Emerging infectious diseases*. 2020;26(6):1320–3.
5. Zhao S, Lin Q, Ran J, Musa SS, Yang G, Wang W, et al. Preliminary estimation of the basic reproduction number of novel coronavirus (2019-nCoV) in China, from 2019 to 2020: A data-driven

- analysis in the early phase of the outbreak. *International journal of infectious diseases: IJID : official publication of the International Society for Infectious Diseases*. 2020;92:214–7.
6. Centers for Disease Control and Prevention (CDC). COVID-19: How to Protect Yourself & Others 2021 [updated Nov. 29, 2021; cited 2021 4 December]. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/prevention.html>.
  7. Chemaitelly H, Tang P, Hasan MR, AlMukdad S, Yassine HM, Benslimane FM, et al. Waning of BNT162b2 Vaccine Protection against SARS-CoV-2 Infection in Qatar. *The New England journal of medicine*. 2021.
  8. United Nations High Commissioner for Refugees (UNHCR). Syrian Arab Republic: COVID-19 Response 2020 Fact Sheet. 2020.
  9. Gharibah M, Mehchy Z. COVID-19 pandemic: Syria's response and healthcare capacity. *Conflict Research Programme*; 2020.
  10. Mohsen F, Bakkar B, Armashi H, Aldaher N. Crisis within a crisis, COVID-19 knowledge and awareness among the Syrian population: a cross-sectional study. *BMJ open*. 2021;11(4):e043305.
  11. World Health Organization (WHO). WHO COVID-19 Dashboard, Syria 2021 [cited 2021 4 December]. Available from: <https://covid19.who.int/region/emro/country/sy>.
  12. United Nations Office for the Coordination of Humanitarian Affairs (OCHA). Syrian Arab Republic: 2021 Needs and Response Summary (February 2021) 2021 [cited 2021 4 December]. Available from: <https://reliefweb.int/report/syrian-arab-republic/syrian-arab-republic-2021-needs-and-response-summary-february-2021>.
  13. McNatt Z, Boothby NG, Al-Shannaq H, Chandler H, Freels P, Mahmoud AS, et al. Impact of separation on refugee families: Syrian refugees in Jordan. *Columbia University Libraries*; 2018.
  14. UN Office for the Coordination of Humanitarian Affairs. Syrian Arab Republic: COVID-19 Humanitarian Update No. 24 As of 3 March 2021. 2021.
  15. Alshammari AS, Alshammari H, Alshammari S. Factors Associated With Adherence to COVID-19 Preventive Measures Among Saudi Arabians. *Cureus*. 2021;13(4):e14623.
  16. Tomczyk S, Rahn M, Schmidt S. Social Distancing and Stigma: Association Between Compliance With Behavioral Recommendations, Risk Perception, and Stigmatizing Attitudes During the COVID-19 Outbreak. *Frontiers in psychology*. 2020;11:1821.
  17. Vally Z. Public perceptions, anxiety and the perceived efficacy of health-protective behaviours to mitigate the spread of the SARS-Cov-2/ COVID-19 pandemic. *Public health*. 2020;187:67–73.
  18. United Nations Department of Economic and Social Affairs. World Population Prospects 2019 2019 [cited 2021 4 December 2021]. Available from: <https://population.un.org/wpp/>.
  19. Pak A, Adegboye OA, Adekunle AI, Rahman KM, McBryde ES, Eisen DP. Economic Consequences of the COVID-19 Outbreak: the Need for Epidemic Preparedness. *Frontiers in public health*. 2020;8:241.
  20. van Loenhout JAF, Vanderplanken K, Scheen B, Van den Broucke S, Aujoulat I. Determinants of adherence to COVID-19 measures among the Belgian population: an application of the protection

- motivation theory. Archives of public health = Archives belges de sante publique. 2021;79(1):74.
21. Azene ZN, Merid MW, Muluneh AG, Geberu DM, Kassa GM, Yenit MK, et al. Adherence towards COVID-19 mitigation measures and its associated factors among Gondar City residents: A community-based cross-sectional study in Northwest Ethiopia. PloS one. 2020;15(12):e0244265.
  22. Central Bureau Of Statistics (CBS). Statistical Population 2019 [cited 2021 4 December]. Available from: <http://cbssyr.sy/index-EN.htm>.
  23. Li T, Liu Y, Li M, Qian X, Dai SY. Mask or no mask for COVID-19: A public health and market study. PloS one. 2020;15(8):e0237691.
  24. Kisielinski K, Giboni P, Prescher A, Klosterhalfen B, Graessel D, Funken S, et al. Is a Mask That Covers the Mouth and Nose Free from Undesirable Side Effects in Everyday Use and Free of Potential Hazards? International journal of environmental research and public health. 2021;18(8).
  25. Ngonghala CN, Iboi E, Eikenberry S, Scotch M, MacIntyre CR, Bonds MH, et al. Mathematical assessment of the impact of non-pharmaceutical interventions on curtailing the 2019 novel Coronavirus. Mathematical biosciences. 2020;325:108364.
  26. Zhong BL, Luo W, Li HM, Zhang QQ, Liu XG, Li WT, et al. Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: a quick online cross-sectional survey. International journal of biological sciences. 2020;16(10):1745–52.
  27. Bowman L, Kwok KO, Redd R, Yi Y, Ward H, Wei WI, et al. Comparing Public Perceptions and Preventive Behaviors During the Early Phase of the COVID-19 Pandemic in Hong Kong and the United Kingdom: Cross-sectional Survey Study. Journal of medical Internet research. 2021;23(3):e23231.
  28. Dessu S, Tsehay T, Girum T, Timerga A, Solomon M, Tsegaye B, et al. The applicability of basic preventive measures of the pandemic COVID-19 and associated factors among residents in Guraghe Zone. PloS one. 2021;16(8):e0256598.
  29. As Plight of Syrians Worsens, Hunger Reaches Record High, International Community Must Fully Commit to Ending Decade-Old War, Secretary-General Tells General Assembly [press release]. 2021.
  30. United Nations Children's Fund (UNICEF). After almost ten years of war in Syria, more than half of children continue to be deprived of education. Joint statement by Regional Humanitarian Coordinator for the Syria Crisis Muhannad Hadi and UNICEF Regional Director for the Middle East and North Africa Ted Chaiban 2021 [updated 24 January 2021; cited 2021 4 December]. Available from: <https://www.unicef.org/press-releases/after-almost-ten-years-war-syria-more-half-children-continue-be-deprived-education>.
  31. Brewer NT, Chapman GB, Gibbons FX, Gerrard M, McCaul KD, Weinstein ND. Meta-analysis of the relationship between risk perception and health behavior: the example of vaccination. Health psychology: official journal of the Division of Health Psychology, American Psychological Association. 2007;26(2):136–45.
  32. Harrison JA, Mullen PD, Green LW. A meta-analysis of studies of the Health Belief Model with adults. Health education research. 1992;7(1):107–16.

33. FLOYD DL, PRENTICE-DUNN S, ROGERS RW. A Meta-Analysis of Research on Protection Motivation Theory. *Journal of Applied Social Psychology*,. 2000;30(2):407–29.
34. Ferrer RA, Klein WM, Persoskie A, Avishai-Yitshak A, Sheeran P. The Tripartite Model of Risk Perception (TRIRISK): Distinguishing Deliberative, Affective, and Experiential Components of Perceived Risk. *Annals of behavioral medicine: a publication of the Society of Behavioral Medicine*. 2016;50(5):653–63.
35. Kaufman AR, Twesten JE, Suls J, McCaul KD, Ostroff JS, Ferrer RA, et al. Measuring Cigarette Smoking Risk Perceptions. *Nicotine & tobacco research: official journal of the Society for Research on Nicotine and Tobacco*. 2020;22(11):1937–45.
36. Weinstein ND, Kwitel A, McCaul KD, Magnan RE, Gerrard M, Gibbons FX. Risk perceptions: assessment and relationship to influenza vaccination. *Health psychology: official journal of the Division of Health Psychology, American Psychological Association*. 2007;26(2):146–51.
37. Janssen E, van Osch L, de Vries H, Lechner L. Measuring risk perceptions of skin cancer: reliability and validity of different operationalizations. *British journal of health psychology*. 2011;16(Pt 1):92–112.
38. Savadori L, Lauriola M. Risk Perception and Protective Behaviors During the Rise of the COVID-19 Outbreak in Italy. *Frontiers in psychology*. 2020;11:577331.
39. Abate H, Mekonnen CK. Knowledge, Attitude, and Precautionary Measures Towards COVID-19 Among Medical Visitors at the University of Gondar Comprehensive Specialized Hospital Northwest Ethiopia. *Infection and drug resistance*. 2020;13:4355–66.
40. Abdul-Baki MN, Dhemes Y, Alasadi L, Alsuliman T. The substantial awareness role of web-based and social media platforms in developing countries during a pandemic scenario: the example of COVID-19. *Journal of public health (Oxford, England)*. 2020;42(3):642–3.
41. Gabarron E, Oyeyemi SO, Wynn R. COVID-19-related misinformation on social media: a systematic review. *Bulletin of the World Health Organization*. 2021;99(6):455-63a.
42. Wang X, Zhang M, Fan W, Zhao K. Understanding the spread of COVID-19 misinformation on social media: The effects of topics and a political leader's nudge. *Journal of the Association for Information Science and Technology*,.n/a(n/a).

## Tables:

**Table 1.** Participant's Characteristics :

Variables		Total (%) n = 7531
Age range (years)	18 -24	3124 (41.5%)
	25 - 44	2338 (31%)
	45 -65	1686 (22.4%)
	> 65	383 (5.1%)
Gender	Male	3505 (46.5%)
	Female	4026 (53.5%)
Marital status	Single	3984 (52.9%)
	Married	2825 (37.5%)
	In relationship	500 (6.6%)
	Widow	222 (2.9%)
Do you have health insurance?	Yes	1489 (19.8%)
	No	6042 (80.2%)
Educational level	No formal education	324 (4.3%)
	Elementary school	422 (5.6%)
	Secondary school	550 (7.3%)
	Highschool	782 (10.4%)
	University student	2906 (38.6%)
	University graduate	1908 (25.3%)
	Postgraduate degree	639 (8.5%)
Do you work or study in the healthcare system	Yes	2595 (34.5%)
	No	4936 (65.5%)
Mother's educational level	No formal education	1225 (16.3%)
	Primary school	1834 (24.4%)
	Secondary school	1656 (22%)
	University degree	2625 (34.9%)
	Postgraduate degree	191 (2.5%)

Father's educational level	No formal education	760 (10.1%)
	Primary school	1971 (26.2%)
	Secondary school	1512 (20.1%)
	University degree	2776 (36.9%)
	Postgraduate degree	512 (6.8%)
Residency	City	5711 (75.8%)
	Countryside	1820 (24.2%)
Geographical origin	Eastern Syria	365 (4.8%)
	Northern Syria	1272 (16.9%)
	Middle Syria	4376 (58.1%)
	Southern Syria	353 (4.7%)
	Western Syria	1165 (15.5%)
Financial status	Bad	1268 (16.8%)
	Middle	3241 (43%)
	Good	2651 (35.2%)
	Excellent	371 (4.9%)
Employment	Unemployed	1635 (21.7%)
	Part-time job	1114 (14.8%)
	Full-time job	1688 (22.4%)
	Student	2653 (35.2%)
	Student + Part-time job	290 (3.9%)
	Student + Full-time job	131 (1.7%)
	Part-time + Full-time jobs	20 (0.3%)

<b>Table 2.</b> COVID-19 related information:		N(%)
Do you believe you have had coronavirus?	No	4549 (60.4)
	yes with PCR	240 (3.2)
	Yes with symptoms	2742 (36.4)
Do you personally know anyone (excluding yourself) who has had a PCR-confirmed COVID-19 ?	No	2960 (39.3)
	Yes	4571 (60.7)
To what extent do you think COVID-19 poses a risk to people in Syria?	No risk at all	314 (4.2)
	Minor risk	1621 (21.5)
	Major risk	464 (61.6)
	Do not know	956 (12.7)
To what extent do you think COVID-19 poses a risk to you personally?	No risk at all	834 (11.1)
	Minor risk	3224 (42.8)
	Major risk	2218 (29.5)
	Do not know	1255 (16.7)

<b>Table 3.</b> Level of commitment to protective measures.			
The number of preventive measures used by the participants			
Number of measures	N(%)	Level of commitment	N(%)
0	91(1.2%)	Bad commitment	357(4.7%)
1	67(0.9%)		
2	93(1.2%)		
3	106(1.4%)		
4	159(2.1%)	Moderate commitment	1888(25.1%)
5	221(2.9%)		
6	328(4.4%)		
7	461(6.1%)		
8	719(9.5%)		
9	932(12.4%)	Good commitment	5286(70.2%)
10	1278(17.0%)		
11	1511(20.1%)		
12	1565(20.8%)		

<b>Table 4.</b> Specific preventive measures.	Yes (N%)	No (N%)
wearing a face mask when in crowded places or when public places	6577 (87.3)	954 (12.7)
Reduce the amount you go to school, college, university or work	4950 (65.7)	2581 (34.3)
cancelled or postponed a social event such as meeting friends, eating out or going to a sporting event	5508 (73.1)	2023 (26.9)
reduce the amount you go to shops	5764 (76.5)	1767 (23.5)
kept away from crowded places?	6317 (83.9)	1214 (16.1)
cleaned or disinfected things you might touch ( doorknob or hard surfaces)	5517 (73.3)	2014 (26.7)
carried sanitizing hand gel with you when you were out?	5557 (73.8)	1974 (26.2)
reduce the amount you touch your eyes, nose, and\or mouth?	5722 (76)	1809 (24)
followed a healthy diet or took vitamins supplements	3747 (49.8)	3784 (50.2)
tried to avoid people who have cold or flu-like symptoms?	6645 (88.2)	886 (11.8)
usually used tissues when sneezing or coughing	6887 (91.4)	644 (8.6)
washed your hands with soap and water more often than usual?	6637 (88.1)	894 (11.9)

**Table 5.** Correlation between demographic characteristics and Level of commitment to protective measures

Demographic characteristics		Level of commitment			Chi-Square value	P-value
		Bad	Moderate	Good		
Age (years)	18 -24 (n=3124)	82 (2.6%)	773(24.7%)	2269(72.6%)	204.974	0.000*
	25 -44 (n=2338)	101(4.3%)	457(19.5%)	1780(76.1%)		
	45 – 65 (n=1686)	137(8.1%)	527(31.3%)	1022(60.6%)		
	> 65 (n=383)	37(9.7%)	131(34.2%)	215(56.1%)		
Gender	Male (n=3505)	219(6.2%)	1075(30.7%)	2211(63.1%)	160.683	0.000*
	Female (n=4026)	138(3.4%)	813(20.2%)	3075(76.4%)		
marital status	Single (n=3984)	134(3.4%)	978(24.5%)	2872(72.1%)	92.002	0.000*
	Married (n=2825)	202(7.2%)	719(25.5%)	1904(67.4%)		
	in relationship (n=500)	6(1.2%)	111(22.2%)	383(76.6%)		
	Widow (n=222)	15(6.8%)	80(36.0%)	127(57.2%)		
origin	Eastern (n=365)	25(6.8%)	121(33.2%)	219(60.0%)	184.079	0.000*
	Northern (n=1272)	128(10.1%)	384(30.2%)	760(59.7%)		
	Middle (n=4376)	164(3.7%)	1072(24.5%)	3140(71.8%)		
	Southern (n=353)	12(3.4%)	92(26.1%)	249(70.5%)		

	Western (n=1165)	28(2.4%)	219(18.8%)	918(78.8%)		
residency	City (n=5711)	213(3.7%)	1409(24.7%)	4089(71.6%)	59.106	0.000*
	Countryside (n=1820)	144(7.9%)	479(26.3%)	1197(65.8%)		
financial status	Bad (n=1268)	150(11.8%)	420(33.1%)	698(55.0%)	279.195	0.000*
	Middle (n=3241)	134(4.1%)	814(25.1%)	2293(70.7%)		
	Good (n=2654)	59(2.2%)	561(21.2%)	2031(76.6%)		
	Excellent (n=371)	14(3.8%)	93(25.1%)	264(71.2%)		
employment	Unemployed (n=1635)	148(9.1%)	427(26.1%)	1060(64.8%)	129.431	0.000*
	part-time (n=1114)	75(6.7%)	274(24.6%)	765(68.7%)		
	full-time (n=1688)	60(3.6%)	404(23.9%)	1224(72.5%)		
	Student (n=2653)	64(2.4%)	675(25.4%)	1914(72.1%)		
	part-time + student (n=290)	5(1.7%)	77(26.6%)	208(71.7%)		
	full-time + student (n=131)	4(3.1%)	26(19.8%)	101(77.1%)		
	part-time + full-time (n=20)	1(5.0%)	5(25.0%)	14(70.0%)		
academic level	no education	81(25.0%)	129(39.8%)	114(35.2%)	640.976	0.0000*

	(n=324)					
	Elementary	61(14.5%)	151(35.8%)	210(49.8%)		
	(n=422)					
	Secondary	35(6.4%)	182(33.1%)	333(60.5%)		
	(n=550)					
	Highschool	40(5.1%)	212(27.1%)	530(67.8%)		
	(n=782)					
	university student	78(2.7%)	721(24.8%)	2107(72.5%)		
	(n=2906)					
	university graduate (n=1908)	39(2.0%)	392(20.5%)	1477(77.4%)		
	post-university study (n=639)	23(3.6%)	101(15.8%)	515(80.6%)		
father's educational level	no education (n=760)	142(18.7%)	257(33.8%)	361(47.5%)	482.949	0.000*
	primary education (n=1971)	95(4.8%)	543(27.5%)	1333(67.6%)		
	secondary education (n=1521)	40(2.6%)	368(24.3%)	1104(73.0%)		
	university degree (n=2776)	69(2.5%)	628(22.6%)	2079(74.9%)		
	post-uni (n=512)	11(2.1%)	92(18.0%)	409(79.9%)		
mother's educational level	no education (n=1225)	170(13.9%)	408(33.3%)	647(52.8%)	391.734	0.000*
	primary education (n=1834)	72(3.9%)	494(26.9%)	1268(69.1%)		
	secondary education	52(3.1%)	391(23.6%)	1213(73.2%)		

(n=1656)				
university degree	58(2.2%)	560(21.3%)	2007(76.5%)	
(n=2625)				
post-uni (191)	5(2.6%)	35(18.3%)	151(79.1%)	

\* P-value  $\leq 0.05$  considered statistically significant

**Table 6.** Correlation between risk perception and commitment to preventive measures

		Level of commitment			Chi-Square	p.value
		Bad	Moderate	Good		
To what extent do you think coronavirus poses a risk to people in Syria?	No risk at all	54(17.2%)	125(39.8%)	135(43.0%)	433.437	0.000*
	Minor risk	126(7.8%)	531(32.8%)	964(59.5%)		
	Major risk	102(2.2%)	947(20.4%)	3591(77.4%)		
	Do not know	75(7.8%)	285(29.8%)	596(62.3%)		
To what extent do you think coronavirus poses a risk to you personally?	No risk at all	119(14.3%)	311(37.3%)	404(48.4%)	402.271	0.000*
	Minor risk	98(3.0%)	858(26.6%)	2268(70.3%)		
	Major risk	50(2.3%)	405(18.3%)	1763(79.5%)		
	Do not know	90(7.2%)	314(25.0%)	851(67.8%)		

\* P-value  $\leq 0.05$  considered statistically significant

## Figures

Figure .1 Participants sources of information




Figure 1

Participants' sources of information