

# The relative importance of work-related and non-work-related stressors and perceived social support on perceived stress in a cross-sectional population-based sample

Jes Sørensen (✉ [jesbso@rm.dk](mailto:jesbso@rm.dk))

DEFACTUM, Central Denmark Region

Mathias Lasgaard

DEFACTUM, Central Denmark Region

Morten Willert

Department of Occupational Medicine, Danish Ramazzini Centre, Aarhus University Hospital

Finn Larsen

DEFACTUM, Central Denmark Region

---

## Research Article

**Keywords:** Stress, perceived stress scale, survey, population-based, stressors, work, non-work, social support, dominance analysis

**Posted Date:** December 1st, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-114221/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on March 19th, 2021. See the published version at <https://doi.org/10.1186/s12889-021-10594-2>.

# Abstract

**Background:** High levels of perceived stress have a negative bearing on health and well-being, and stress is a major public health issue. According to the Stress Process Model, stressors are socially patterned and combine to produce strain. Despite this, most studies on stress have focused on work-related stressors leaving non-work determinants underinvestigated. The aim of the present study was to determine the relative importance of work-related and non-work-related stressors and perceived social support for the perceived stress level.

**Methods:** Self-reported data were drawn from the 2017 population-based health survey "How are you?" conducted in the Central Denmark Region (N=32,417). Data were linked with data drawn from national administrative registers. Work- and non-work-related stressors assessed included major life events, chronic stressors and daily hassles. Perceived social support was assessed using a single question. Perceived stress was assessed by the 10-item Perceived Stress Scale. We conducted dominance analyses based on a multiple linear regression model to determine the most important contributors to perceived stress. Analyses were weighted and adjusted.

**Results:** Work- and non-work-related stressors along with perceived social support explained 42.5% of the total variance ( $R^2$ ) in perceived stress. The most important predictors were disease, perceived social support and work situation. The stratified analyses produced slightly varying predictor profiles of perceived stress between subgroups. Work situation was the most important predictor in the employed group. However, adding non-work-related predictors to the analysis tripled the explained variance.

**Conclusions:** Level of perceived stress is affected by a combination of work- and non-work-related stressors and perceived social support both at population level and in subgroups. The most important predictors of perceived stress are disease, perceived social support and work situation. Results indicate that public health strategies aiming to reduce stress should take a holistic approach and address a variety of stressor domains rather than focus on a single domain.

**Trial registration:** The study was approved by the Danish Data Protection Agency (r. no. 2012-58-0006) and Central Denmark Region (r. no. 1-16-02-593-16).

## Background

Stress has been defined by Lazarus and Folkman as a state of emotional strain and pressure arising in an individual when demands in the environment are perceived as "taxing or exceeding his or her resources and endangering his or her wellbeing" (1). High levels of perceived stress have a negative impact on health and well-being (2). Stress increases the risk of unemployment (3), and long-term exposure to stressors is a risk factor for developing mental and physical diseases including depression (4-7) and anxiety disorders (8), substance abuse (9, 10), coronary heart disease (11, 12), type 2 diabetes (13, 14) and metabolic syndrome (15). Furthermore, high levels of perceived stress are associated with an increased risk of premature death (16).

Stress is a major public health issue and has long been a topic attracting much interest in a wide range of research fields (2, 17, 18). A particularly active area of research has been work-related stress (17, 19, 20). This area has predominantly been inspired by the Demand-Control-Support (21) and Effort-Reward Imbalance (22) models, focusing on the interplay between the psychosocial work environment and workers' mental health. With the exception of work-life balance, most research in this area has analysed the workplace as a closed system, where high levels of stress among employees are linked primarily to occupational stressors leaving non-work determinants underinvestigated (18, 23).

However, the sociological Stress Process Model introduced in 1981 by Pearlin and colleagues (24) recognized that stressors are socially patterned and seldomly act in isolation but combine in an additive or multiplicative manner to produce strain. The Stress Process Model has served as a paradigm for research into an expanding range of stressors (25). Pearlin argued strongly that stress research should address the full spectrum of stressors found in a person's life rather than single stressors in isolation (25, 26). Despite this, the understanding of the complexity of the stress process remains deficient in the empirical literature (27). According to a review of trends in stress research, literature on specific stressors has become more abundant and research has become increasingly segmented (28). This trend may result in a return to pre-Stress Process Model approaches addressing one stressor at a time and thus ignoring the additive or multiplicative effect of stressors.

One of the gains of the extensive sociological, psychological and epidemiological research into stressors is a conceptual division of the stress continuum into life events, chronic stressors and daily hassles (29). "Life events" are sudden, discrete occurrences such as the death of a near relative or being laid off from work unexpectedly. "Chronic stressors" are enduring strains such as discrimination, permanently compromised health or conflicting demands of multiple social roles. "Daily hassles" include episodic – expected or unexpected – minor events which, as they accumulate, may constitute stressors such as a parent and a child struggling over bedtime or delays in morning traffic. In addition, these very diverse stressors vary in respect to duration, severity and the context in which they occur. It is acknowledged that these different types of stressors all contribute to the overall burden of stress in the population (29). Thus, the measures used for exposure to stressors should tap into a broad spectrum from discrete life events to chronic stressors and daily hassles. The consequences of chronic or recurring stressors may be particularly severe when they surface within major social domains, such as breadwinning, job and family (30). A better understanding of domain-specific stressors and their relative contribution to the overall perceived burden of stress is important for effective public health strategies (31, 32). Prevention and reduction of stress at population level and in target groups facing special needs should be based on empirical knowledge of the main sources of stress.

The other major component of the Stress Process Model is resources available to cope with the challenges experienced (33). Central to this component is access to social support from one's surroundings. In particular, empirical research has established that it is not the actual availability of support that promotes coping but its perceived availability (27, 34). It has been theorised that social support buffers the effects of stressors (24, 26, 35). However, a review indicates that the stress-buffering

effects of social support are “less dramatic and consistent” than the direct effects of social support on mental health (36). This underscores the relevance of investigating the direct effect of perceived social support on the overall stress burden on par with stressors within various social domains.

In this study, we return to the holistic approach to stress suggested by Pearlin (30) by investigating the relative importance of stressors occurring within major social domains and perceived social support on perceived stress in a general adult population.

## **Aim**

The aim of the study was to determine the relative importance and contribution of work-related and non-work-related stressors and perceived social support on the perceived stress level. The study examines stressors and perceived social support at population level and in socio-demographic subgroups in order to generate knowledge for public health strategies.

## **Methods**

### **Study design and data collection**

Self-reported data were drawn from the 2017 population-based health survey “How are you?” conducted in the Central Denmark Region. In 2017, approximately 23% of the Danish population were living in the Central Denmark Region, and the demographic composition (gender, age, and educational attainment) of the population was similar to that of the total Danish population (37). The survey included a representative population sample of 52,000 citizens aged 16 years and above drawn from the Danish Civil Registration System (38). Participants were invited to complete a web-based or postal questionnaire (39). Those who failed to respond to the survey received up to four reminders. The response rate was 62% (32,417), and 80% of the respondents completed the web questionnaire. The study was approved by the Danish Data Protection Agency (r. no. 2012-58-0006) and the Central Denmark Region (r. no. 1-16-02-593-16). Each participant received written information about the purpose of the survey, and informed consent was obtained from all subjects. All methods were carried out in accordance with relevant guidelines and regulations along with the approval. Data from the survey were linked with national administrative registers using the unique personal identification number assigned to all Danish citizens (38). Register data included age, gender, ethnic background and labour market status.

## **Variables**

### **Perceived stress scale**

The level of perceived stress was assessed by the 10-item Perceived Stress Scale (PSS) (40). Based on Lazarus' stress model (41, 42), the PSS is a generic measure of stress in which stress relies upon the person's perception of the stressor as stressful or not. The ten items ask how often in the past month life

was appraised as unpredictable, uncontrollable and overloaded (e.g., "In the last month, how often have you been upset because of something that happened unexpectedly?") (40). The items were scored from 0 to 4 ("never", "almost never", "sometimes", "fairly often", or "very often"). An increasing sum score (range 0 to 40) indicates an increasing perceived stress level (40). PSS has satisfying psychometric properties (43, 44) and has been used in a number of population-based studies (e.g. 3, 45-48). In the present study, Cronbach's  $\alpha$  indicated that the scale had a high internal consistency ( $\alpha=0.87$ ).

## **Stressors**

Work-related and non-work-related stressors were assessed using nine questions covering seven categories from the Life Event Questionnaire (49). The selection of questions was inspired by The Danish National Birth Cohort (50-52). The questions cover major life events, chronic stressors and daily hassles. Stressors were assessed asking "Within the past 12 months have you felt burdened by some of the following things?" The questionnaire covers work situation, financial circumstances, housing conditions, relationship with partner, relationship with family and friends, own disease, disease among close relatives, deaths among close relatives and other types of burdens. The response categories were "no", "yes, a little", "yes, partly" or "yes, a lot". In the present study, all four response categories of each variable were used.

The survey also includes a more comprehensive questionnaire regarding work that covers quantitative demands, job influence, satisfaction and recognition, work-life conflicts, physical and emotional wear and tear, and physical load (53). However, introducing the more comprehensive questionnaire did not alter the analyses, which is in line with a previous study (54). Thus, in order to keep the model simple, the single question about work situation was preferred.

## **Perceived social support**

Perceived social support was assessed using a single item: "Do you have anyone to talk to if you have problems or need support?" The response format was "yes, always", "yes, mostly", "yes, sometimes" or "no, never or almost never". In the present study, all four response categories of each variable were used. The question was inspired by The MOS Social Support Instrument (55).

## **Socio-demographic variables**

Age and gender were assessed using a combination of self-reported and register data. Ethnic background was defined using the Danish Civil Registration System (38). Educational attainment was self-reported and categorised as low (primary school, no further education), medium (upper secondary education, vocational education and/or short higher education) or high (bachelor's degree or higher level of education) according to the Danish version of the International Standard Classification of Education (56). Students were categorised according to their expected graduation level. Labour market status was assessed using a combination of self-reported data and data from the Danish Register for Evaluation of Marginalization (57).

## Data analysis

We conducted dominance analysis, a relatively new computer-intensive method, to determine the most important contributors to perceived stress in the population and in stratified analyses (i.e., gender, age, educational attainment and labour market status) (58, 59). The analyses were based on a multiple linear regression model with PSS as the dependent variable and the nine different stressors and perceived social support as predictor variables. Dominance analysis is a method of ranking predictors according to how much each variable contributes to the total variance of the dependent variable in a model. The method is particularly suitable when predictors are intercorrelated, as may be expected in this study. Dominance analysis is an ensemble method based on estimation of all possible regression models (all subset regressions) (60). The average increase in total variance explained by the model ( $R^2$ ) when adding a variable to all possible sub-models quantifies the importance of the variable. Dominance analysis is used for decomposition of the total  $R^2$ , but it has rarely been used in stress research (61). The dominance analyses in the present study consisted of 1,023 ( $10^2-1$ ) regression models containing all possible combinations of predictors. In addition to the ten predictors, gender, age, ethnic background and educational attainment were included in all models.

Weights constructed by Statistics Denmark using a model-based calibration approach and including socio-demographic characteristics, income, social benefits and healthcare utilisation were applied in order to reduce sampling and non-response bias (39, 62).

Prior to the analyses, data were screened for missing values. The percentage of missing values was acceptable (0-13%). Missing PSS items ranged from 4.3% (item 1) to 5.1% (item 8). If one, two or three items of the PSS scale were missing, the mean of the available items was used to calculate the scale score (63). If responses to more than three items were missing, the PSS score was regarded as missing. Hence, 1,392 observations (4.3%) were excluded.

For stressors, missing values were treated as "no" if respondents had answered at least one of the nine stressor questions. A similar approach was adopted in the "How are you?" health survey with regards to measures of chronic diseases (64).

PSS was obtained in 95.7% of the respondent (Table 1), and stressors were obtained in 93.3% of the respondents (Table 2). Respondents with missing PSS or stressors were more likely to be 16-24 years old, have an ethnic background other than Danish and an unknown educational attainment or work situation (results not shown).

Stata/SE v16.1 (StataCorp, College Station, TX) was used to prepare the data and perform the descriptive and inferential analyses including dominance analysis using the community-contributed extension DOMIN (65).

## Results

## Descriptive analyses

The mean perceived stress level in the population was 12.2 with a range from 0 to 40 (Table 1). The highest stress levels were found among women, 16-24-year-olds, respondents with an ethnic background other than Danish, respondents with a low educational attainment and respondents who were either students or outside the labour market (unemployed, receiving cash or sickness benefits, or receiving a disability pension). The three groups outside the labour market had the highest perceived stress levels of all subgroups.

Prevalence of stressors varied from 42.2% having been burdened by disease during the past 12 months to 13.0% having experienced other types of burdens (Table 2). Absence of perceived social support was found in 4.9%.

## Main findings

Work- and non-work-related stressors along with perceived social support explained 42.5% of the variance in perceived stress level in the adjusted analysis (Table 3 and Figure 1). The three predictors explaining most of the variance in PSS were disease (9.5%), perceived social support (5.9%) and work situation (4.9%) (Table 3 and Figure 1). Adding the non-work-related stressors and perceived social support to the dominance analysis increased the explained variance from 12.1% (work situation 4.9% and sociodemographic characteristics 7.2%) to 42.5%.

## Stratified analyses

The predictors explained 41.9% and 42.1% of the variance in PSS in men and women, respectively (Table 3). In both men and women, the most important predictors were disease, perceived social support and work situation. In the analyses stratified by age group, the predictors explained 42.8%, 44.8% and 34.0% of the variance in PSS, respectively (Table 3). In the young age group (16-24 years), the most important predictors were disease, relationship with family and friends and perceived social support. In the middle-aged group (25-64 years), the most important predictors were disease, work situation and perceived social support. In the elderly group (65 years and above), the most important predictors were disease and perceived social support.

In the analyses stratified by educational attainment, the predictors explained 38.7% of the variance in PSS in the low, 42.6% in the medium and 41.9% in the high educational groups (Table 3). In the low educational attainment group, the most important predictors were disease and perceived social support. In the medium educational attainment group, the most important predictors were disease, perceived social support and work situation. Finally, in the high educational attainment group, the most important predictors were disease, work situation and perceived social support.

In the analyses stratified by employment status, the predictors explained 41.6% for students, 38.4% for employed, 46.1% for unemployed, 51.8% for respondents receiving cash or sickness benefits, 42.6% for respondents receiving disability pension, 43.9% for respondents receiving early retirement pension and 33.3% for respondents receiving retirement pension (Table 3). The top three predictors included disease in all seven subgroups and perceived social support in five subgroups. Among students, the most important predictors were disease, perceived social support, relationship with family and friends and financial circumstances. Among employed respondents, the most important predictors were work situation, perceived social support and disease. Among unemployed respondents, it was disease, work situation, financial circumstances and relationship with family and friends. Among those receiving cash or sickness benefits, it was disease, relationship with family and friends and financial circumstances. Among those receiving disability pensions, it was disease, perceived social support and relationship with family and friends. Furthermore, among those receiving early retirement pensions it was disease, perceived social support and relationship with family and friends. Finally, among those receiving retirement pensions, it was disease and perceived social support.

## Discussion

### Main findings and stratified analyses

We applied a holistic approach to stress, investigating the relative importance of work- and non-work-related stressors and perceived social support on perceived stress in a representative adult population and in sociodemographic subgroups. As recommended in a number of previous studies (3, 18, 23, 31, 32), we investigated both work- and non-work-related predictors of stress. Most noteworthy, the perceived stress level was affected by a combination of work- and non-work-related stressors and perceived social support. The most important predictors of perceived stress were disease, perceived social support and work situation. In fact, including non-work-related stressors and perceived social support along with work-related stressors more than tripled the explained variance in perceived stress compared with work-related stressors alone.

The stratified analyses (including different subgroups) resulted in slightly varying predictor profiles of perceived stress. However, disease was the most important predictor of perceived stress in 14 of the 15 subgroups. Perceived social support was among the top three predictors in 13 subgroups, and work situation was among the top three predictors in seven subgroups (only relevant in eight subgroups). Other important predictors in the stratified analyses were relationship with family and friends, and financial circumstances. Work situation was the single most important predictor in the employed group, and the second most important predictor in the unemployed group. However, even in the employed group, adding non-work-related stressors and perceived social support tripled the explained variance.

Importantly, although a specific stressor explains only a small part of the overall variation in stress level, it may well play an important role in combination with other stressors (32). As mentioned, stressors rarely occur in isolation. The combined effect of several stressors can therefore have a negative effect on the

stress level. In addition, a stressor can be part of a chain of stressors triggered by a primary stressor (*stress proliferation*), or there may be a spill over of stressors across roles or life domains (conflicts between work and family, etc.) (24). This demonstrates the importance of examining multiple stressors simultaneously.

Our results are in accordance with the Stress Process Model, which finds that stressors combine to produce strain (24), and with previous studies reporting that combinations of stressors from both work and non-work domains contribute to mental health (18, 23) and perceived stress in particular (3, 31, 32, 66, 67). However, our results are contrary to the vast majority of studies on stress that have focused on different aspects of the work situation and have concluded that the work situation causes stress and that related negative health effects are explained solely by the work situation (5, 12, 17, 20, 61, 68, 69). Our results are supported by a review reporting lack of knowledge about the relationship between work and non-work domains and their contribution to mental health (18). The authors concluded that since the majority of studies do not include non-work domains, they may overlook the contribution from non-work predictors. Marchand and colleagues concluded that including both work and non-work domains is necessary to avoid erroneous conclusions about the relationship between work and mental health (23). For example, recent years have seen an increased focus on the implications of managing work and family caregiving roles (child caregiving, elder caregiving or "sandwiched" caregiving) with a special focus on gender differences (70). In line herewith, our results – both at population level and in subgroups – indicate that in order to understand the complex interaction of stressors, both non-work-related stressors and perceived social support should be addressed along with the work situation. Thus, our results are in favour of a holistic approach to understanding the relationship between work- and non-work-related stressors and perceived social support, and therefore also of developing public health strategies and interventions aiming at reducing perceived stress at the population level and in high-risk groups.

## Predictors

The three most important predictors of perceived stress in this study were disease, perceived social support and work situation. An American study of older adults identified loneliness, neighbourhood and financial strain as particularly important predictors of perceived stress (31). Different groups of respondents may have different predictors of perceived stress, which is indicated in our study.

Disease as a stressor has received far less attention than stress as a risk factor for developing diseases. Even so, an understanding of the pathway from disease to stress is beginning to emerge (71). This pathway includes loss of control and uncertainty leading to a feeling of threat, which may lead to hopelessness and stress if coping mechanisms are inadequate. A recent review concludes that it may be useful to distinguish between primary mental health disorders and stress caused by disease (72). Four pathways from chronic disease to stress have been suggested (71): 1) Disease often causes pain, reduces physical and mental functioning and can be life-threatening, making it difficult to meet the demands of everyday life. 2) The process of diagnosis and treatment in itself gives rise to concern and

uncertainty and can be both financially burdensome and time consuming. 3) Patients may have concerns whether possible harms outweigh potential benefits of treatment and may experience adverse drug interactions if treated for multiple conditions. 4) Treatment can disrupt social roles and relationships and cause lower self-efficacy and self-worth, thereby affecting mental well-being negatively.

Lack of social support is a well-known predictor of poor mental health (27, 33, 34, 73). Non-work social support can come from a number of sources including a partner, close relatives, friends, neighbours etc. A study of workplace-perceived social support finds that job performance was strengthened by perceived supervisor and co-worker support (74). Our results are in line with a Danish study reporting that a low level of private-life social support increases the risk of symptoms of depression (73), and a review concluding that social support is associated with mental health (34). Even though we do not distinguish between work- and non-work-related social support, our results are in accordance with a recent meta-analysis showing that a low level of support at work increases the risk of stress-related mental disorders (17). However, it remains to be elucidated how social support produces positive mental health outcomes (36).

The work situation is a well-known stressor (17, 19, 20), which is also confirmed in our study. Different aspects of the work situation have been identified as predictors of stress, e.g. effort reward imbalance, high job demands, organisational justice, social support, emotional demands and decision authority (17). A study reports that stress from work-related strain may be transferred from parents to children (75). This may also hold true for transfer to one's partner.

Although The Stress Process Model has served as a paradigm for stressor research since the 1980s, a lack of understanding remains of how work- and non-work-related stressors, coping, social support and personal resources interact with each other (25). Gaining an enhanced understanding of this interaction is mandatory to develop effective strategies against stress encompassing people's entire life situation (31, 32). An American study found that similar combinations of chronic stressors and life events may result in different levels of perceived stress (*multifinality*) and that different combinations of stressors may produce similar levels of perceived stress (*equifinality*) (32).

## Strengths And Limitations

The present study has several strengths. We use a combination of self-reported and register data from a representative population sample with a large sample size and a reasonably high response rate. Register data are of high validity and accuracy. Perceived stress is assessed using the PSS, which has satisfying psychometric properties. Predictors of perceived stress covered a wide range of sources. Furthermore, the dominance analysis allows us to rank the importance and contribution of predictors and eases the interpretation of several regression coefficients simultaneously (60, 76). Finally, data are weighted, and analyses are adjusted for a number of factors, which is expected to reduce residual confounding.

The study also has some limitations. Non-respondents are more likely to be 16-24-years and of an ethnic background other than Danish. Therefore, non-response bias cannot be excluded. Perceived stress,

stressors and social support are assessed within different time frames, which may weaken the variance explained by the predictors. This may partly explain why the dominance analyses explain only around 40% of the total variance. In the 16-24-year-old group, educational level is based on assumptions about future educational level resulting in a larger proportion having a low and medium-high educational level compared with the 25-64-year-old group.

This is a cross-sectional study in which it is not possible to investigate causal relationships between predictors and stress levels.

## **Conclusions**

The results of this population-based study demonstrate that perceived stress level is affected by a combination of work- and non-work-related stressors and perceived social support. This is the case both at population level and in stratified subgroups. The three most important predictors of perceived stress are disease, perceived social support and work situation. Hence, the study indicates that we should address a variety of stressor domains rather than focus on a single domain like work-related stressors. Thus, our results point towards a return to a holistic approach to stress as suggested by Pearlin and colleagues (24).

Work situation is the most important predictor of perceived stress in the employed group. However, even in this group, adding the non-work-related predictors increased the explained variance noticeably, suggesting that work-related stressors only capture part of the total amount of stressors affecting perceived stress. Thus, even among those employed, stress preventions should not focus solely on work-related stressors.

This study provides novel insight into the predictors of perceived stress level applicable to public health strategies aiming to reduce stress both at population level and in high-risk subgroups. Stratified subgroup analyses revealed varying predictor profiles suggesting a need for target group-specific stress reduction strategies.

## **Declarations**

### **Ethics approval and consent to participate**

The study was approved by the Danish Data Protection Agency (r. no. 2012-58-0006) and Central Denmark Region (r. no. 1-16-02-593-16).

Each participant received written information about the purpose of the survey, and informed consent was obtained from all subjects. All methods were carried out in accordance with relevant guidelines and regulations along with the approval.

# Consent for publication

Not applicable.

# Availability of data and materials

The datasets used and analysed during the current study are available from the corresponding author on reasonable request.

# Competing interests

The authors declare that they have no competing interests

# Funding

The "How are you?" health survey was funded by Central Denmark Region. The present study (based on secondary data analysis) was funded by Velliv Foreningen (grant number 19-0176).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

# Authors' contributions

FBL and JBS conceptualized the study, acquired funding, performed analyses and authored the manuscript. ML and MVW contributed to the manuscript, and read and approved the final manuscript.

# Acknowledgements

Not applicable.

# References

1. Lazarus RS, Folkman S. Stress, Appraisal, and Coping: Springer Publishing Company; 1984.
2. O'Connor DB, Thayer JF, Vedhara K. Stress and Health: A Review of Psychobiological Processes. Annual Review of Psychology. 2020.
3. Maehlisen MH, Pasgaard AA, Mortensen RN, Vardinghus-Nielsen H, Torp-Pedersen C, Boggild H. Perceived stress as a risk factor of unemployment: a register-based cohort study. BMC Pub Health. 2018;18(1):728.

4. Liu RT, Alloy LB. Stress generation in depression: A systematic review of the empirical literature and recommendations for future study. *Clin Psychol Rev.* 2010;30(5):582-93.
5. Bonde JPE. Psychosocial factors at work and risk of depression: a systematic review of the epidemiological evidence. *Occup Environ Med.* 2008;65(7):438.
6. Kendler KS, Karkowski LM, Prescott CA. Stressful life events and major depression: Risk period, long-term contextual threat and diagnostic specificity. *J Nerv Mental Dis.* 1998;186(11):661-9.
7. Assari S, Lankarani MM. Stressful Life Events and Risk of Depression 25 Years Later: Race and Gender Differences. *Front Pub Health.* 2016;4:49.
8. Niedhammer I, Malard L, Chastang J-F. Occupational factors and subsequent major depressive and generalized anxiety disorders in the prospective French national SIP study. *BMC Pub Health.* 2015;15(1):200.
9. Spencer RL, Hutchison KE. Alcohol, aging, and the stress response. *Alcohol Res Health.* 1999;23(4):272-83.
10. Keyes KM, Hatzenbuehler ML, Grant BF, Hasin DS. Stress and alcohol: epidemiologic evidence. *Alcohol Res.* 2012;34(4):391-400.
11. Steptoe A, Kivimäki M. Stress and Cardiovascular Disease: An Update on Current Knowledge. *Ann Rev Pub Health.* 2013;34(1):337-54.
12. Sara JD, Prasad M, Eleid MF, Zhang M, Widmer RJ, Lerman A. Association Between Work-Related Stress and Coronary Heart Disease: A Review of Prospective Studies Through the Job Strain, Effort-Reward Balance, and Organizational Justice Models. *J Am Heart Assoc.* 2018;7(9):e008073.
13. Kelly SJ, Ismail M. Stress and Type 2 Diabetes: A Review of How Stress Contributes to the Development of Type 2 Diabetes. *Ann Rev Pub Health.* 2015;36(1):441-62.
14. Harris M, Oldmeadow C, Hure A, Luu J, Loxton D, Attia J. Stress increases the risk of type 2 diabetes onset in women: A 12-year longitudinal study using causal modelling. *PLOS ONE.* 2017;12:e0172126.
15. Kuo W-c, Bratzke LC, Oakley LD, Kuo F, Wang H, Brown RL. The association between psychological stress and metabolic syndrome: A systematic review and meta-analysis. *Obes Rev.* 2019;20(11):1651-64.
16. Prior A, Fenger-Grøn M, Larsen KK, Larsen FB, Robinson KM, Nielsen MG, et al. The Association Between Perceived Stress and Mortality Among People With Multimorbidity: A Prospective Population-Based Cohort Study. *Am J Epidemiol.* 2016;184(3):199-210.
17. van der Molen HF, Nieuwenhuijsen K, Frings-Dresen MHW, de Groene G. Work-related psychosocial risk factors for stress-related mental disorders: an updated systematic review and meta-analysis. *BMJ Open.* 2020;10(7):e034849.
18. Beauregard N, Marchand A, Blanc M-E. What do we know about the non-work determinants of workers' mental health? A systematic review of longitudinal studies. *BMC Pub Health.* 2011;11(1):439.

19. Limonic L, Lennon MC. Work and Unemployment as Stressors. In: Wright ER, Scheid TL, editors. *A Handbook for the Study of Mental Health: Social Contexts, Theories, and Systems*. 3 ed. Cambridge: Cambridge University Press; 2017. p. 224-38.
20. Dewe P, Cooper, C. *Work and Stress: A Research Overview*. London: Routledge; 2020.
21. Karasek R, Theorell T. *Healthy work : stress, productivity and the reconstruction of working life*: New York (N.Y.) : Basic books; 1990.
22. Siegrist J. Adverse health effects of high-effort/low-reward conditions. *J Occup Health Psyc*. 1996;1(1):27-41.
23. Marchand A, Durand P, Haines V, 3rd, Harvey S. The multilevel determinants of workers' mental health: results from the SALVEO study. *Soc Psyc Psychiatr Epidemiol*. 2015;50(3):445-59.
24. Pearlin LI, Menaghan EG, Lieberman MA, Mullan JT. The stress process. *J Health Soc Behav*. 1981;22(4):337-56.
25. Aneshensel CS. Sociological Inquiry into Mental Health: The Legacy of Leonard I. Pearlin. *J Health Soc Behav*. 2015;56(2):166-78.
26. Pearlin LI. The Sociological Study of Stress. *J Health Soc Behav*. 1989;30(3):241-56.
27. Brown LL, Mitchell UA, Ailshire JA. Disentangling the Stress Process: Race/Ethnic Differences in the Exposure and Appraisal of Chronic Stressors Among Older Adults. *J Gerontol: Series B*. 2018;75(3):650-60.
28. Wheaton B, Young M, Montazer S, Stuart-Lahman K. Social stress in the twenty-first century. *Handbook of the sociology of mental health*, 2nd ed. *Handbooks of sociology and social research*. New York, NY, US: Springer Science + Business Media; 2013. p. 299-323.
29. Wheaton B, Montazer S. Stressors, Stress, and Distress. In: Scheid TL, Brown TN, editors. *A Handbook for the Study of Mental Health: Social Contexts, Theories, and Systems*. 2 ed. Cambridge: Cambridge University Press; 2009. p. 171-99.
30. Pearlin LI, Schieman S, Fazio EM, Meersman SC. Stress, Health, and the Life Course: Some Conceptual Perspectives. *J Health Soc Behav*. 2005;46(2):205-19.
31. Scott SB, Jackson BR, Bergeman CS. What contributes to perceived stress in later life? A recursive partitioning approach. *Psychol Aging*. 2011;26(4):830-43.
32. Scott SB, Whitehead BR, Bergeman CS, Pitzer L. Combinations of stressors in midlife: examining role and domain stressors using regression trees and random forests. *J Gerontol B Psychol Sci Soc Sci*. 2013;68(3):464-75.
33. Glanz K, Schwartz MD. Stress, coping, and health behavior. *Health behavior and health education: Theory, research, and practice*, 4th ed. San Francisco, CA, US: Jossey-Bass; 2008. p. 211-36.
34. Brown RL, Ciciurkaite G. Understanding the Connection between Social Support and Mental Health. In: Wright ER, Scheid TL, editors. *A Handbook for the Study of Mental Health: Social Contexts, Theories, and Systems*. 3 ed. Cambridge: Cambridge University Press; 2017. p. 207-23.

35. Cohen S, Wills TA. Stress, social support, and the buffering hypothesis. *Psych Bull.* 1985;98(2):310-57.
36. Thoits PA. Mechanisms Linking Social Ties and Support to Physical and Mental Health. *J Health Soc Behav.* 2011;52(2):145-61.
37. Statistics Denmark. FOLK1 - Population 2017 [Available from: <https://www.statistikbanken.dk/statbank5a/default.asp?w=1920>].
38. Pedersen CB. The Danish Civil Registration System. *Scand J Public Health.* 2011;39(7 Suppl):22-5.
39. Christensen AI, Lau CJ, Kristensen PL, Johnsen SB, Wingstrand A, Friis K, et al. The Danish National Health Survey: Study design, response rate and respondent characteristics in 2010, 2013 and 2017. *Scand J Pub Health.* 2020:1403494820966534.
40. Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *J Health Soc Behav.* 1983;24(4):385-96.
41. Lazarus RS. *Psychological stress and the coping process.* New York, NY, US: McGraw-Hill; 1966.
42. Lazarus RS. Psychological stress and coping in adaption and illness. In: Lipowski ZJ, Lipsitt DR, Whybrow PC, editors. *Psychosomatic medicine : current trends and clinical applications.* New York: Oxford Univ. Press; 1977. p. 14-26.
43. Lee EH. Review of the psychometric evidence of the perceived stress scale. *Asian Nurs Res (Korean Soc Nurs Sci).* 2012;6(4):121-7.
44. Nielsen MG, Ornbol E, Vestergaard M, Bech P, Larsen FB, Lasgaard M, et al. The construct validity of the Perceived Stress Scale. *J Psychosom Res.* 2016;84:22-30.
45. Stubbs B, Vancampfort D, Veronese N, Schofield P, Lin PY, Tseng PT, et al. Multimorbidity and perceived stress: a population-based cross-sectional study among older adults across six low- and middle-income countries. *Maturitas.* 2018;107:84-91.
46. Pedersen AB, Baggesen LM, Ehrenstein V, Pedersen L, Lasgaard M, Mikkelsen EM. Perceived stress and risk of any osteoporotic fracture. *Osteoporos Int.* 2016;27(6):2035-45.
47. Graff S, Prior A, Fenger-Gron M, Christensen B, Glumer C, Larsen FB, et al. Does perceived stress increase the risk of atrial fibrillation? A population-based cohort study in Denmark. *Am Heart J.* 2017;188:26-34.
48. Gawlik KS, Melnyk BM, Tan A. Associations Between Stress and Cardiovascular Disease Risk Factors Among Million Hearts Priority Populations. *Am J Health Promot.* 2019;33(7):1063-6.
49. Norbeck JS. Modification of life event questionnaires for use with female respondents. *Res Nurs Health.* 1984;7(1):61-71.
50. Sondergaard C, Olsen J, Friis-Hasche E, Dirdal M, Thrane N, Sorensen HT. Psychosocial distress during pregnancy and the risk of infantile colic: a follow-up study. *Acta Paediatr.* 2003;92(7):811-6.
51. Tegethoff M, Greene N, Olsen J, Meyer AH, Meinlschmidt G. Maternal psychosocial stress during pregnancy and placenta weight: evidence from a national cohort study. *PloS one.* 2010;5(12):e14478-e.

52. Tegethoff M, Greene N, Olsen J, Schaffner E, Meinlschmidt G. Stress during pregnancy and offspring pediatric disease: A National Cohort Study. *Environ Health Perspect.* 2011;119(11):1647-52.
53. Kristensen TS, Hannerz H, Høgh A, Borg V. The Copenhagen Psychosocial Questionnaire—a tool for the assessment and improvement of the psychosocial work environment. *Scand J Work Environ Health.* 2005;31(6):438-49.
54. Munch-Hansen T, Wieclaw J, Agerbo E, Westergaard-Nielsen N, Bonde JP. Global measure of satisfaction with psychosocial work conditions versus measures of specific aspects of psychosocial work conditions in explaining sickness absence. *BMC Pub Health.* 2008;8(1):270.
55. Sherbourne CD, Stewart AL. The MOS social support survey. *Soc Sci Med.* 1991;32(6):705-14.
56. Eurostat. International Standard Classification of Education (ISCED) 2018 [updated 28-08-2018]. Available from: [https://ec.europa.eu/eurostat/statistics-explained/index.php/International\\_Standard\\_Classification\\_of\\_Education\\_\(ISCED\)](https://ec.europa.eu/eurostat/statistics-explained/index.php/International_Standard_Classification_of_Education_(ISCED)).
57. Hjollund NH, Larsen FB, Andersen JH. Register-based follow-up of social benefits and other transfer payments: accuracy and degree of completeness in a Danish interdepartmental administrative database compared with a population-based survey. *Scand J Pub Health.* 2007;35(5):497-502.
58. Azen R, Budescu D. The Dominance Analysis Approach for Comparing Predictors in Multiple Regression. *Psych Meth.* 2003;8:129-48.
59. Budescu D. Dominance Analysis: A New Approach to the Problem of Relative Importance of Predictors in Multiple Regression. *Psychological Bulletin.* 1993;114:542-51.
60. Luchman J. Determining Subgroup Difference Importance with Complex Survey Designs: An Application of Weighted Dominance Analysis. *Survey Practice.* 2015;8:1-10.
61. Sun J-W, Bai H-Y, Li J-H, Lin P-Z, Zhang H-H, Cao F-L. Predictors of occupational burnout among nurses: a dominance analysis of job stressors. *J Clin Nurs.* 2017;26(23-24):4286-92.
62. Särndal C-E, Lundström S. Estimation in Survey with Nonresponse. 2005.
63. Hawthorne G, Hawthorne G, Elliott P. Imputing Cross-Sectional Missing Data: Comparison of Common Techniques. *Aus New Zea J Psych.* 2005;39(7):583-90.
64. Larsen FB, Pedersen MH, Friis K, Glumer C, Lasgaard M. A Latent Class Analysis of Multimorbidity and the Relationship to Socio-Demographic Factors and Health-Related Quality of Life. A National Population-Based Study of 162,283 Danish Adults. *PLoS One.* 2017;12(1):e0169426.
65. Stata modul by Joseph N. Luchman, Senior Scientist, Fors Marsh Group LLC, Arlington, VA
66. Evans D, Mallet L, Flahault A, Cothureau C, Velázquez S, Capron L, et al. The importance of both workplace and private life factors in psychological distress: a large cross-sectional survey of French railway company employees. *Soc Psych Psych Epidemiol.* 2012;48:1211-24.
67. Hasselberg K, Jonsdottir IH, Ellbin S, Skagert K. Self-reported stressors among patients with Exhaustion Disorder: an exploratory study of patient records. *BMC Psych.* 2014;14(1):66.
68. Nilsen C, Andel R, Fritzell J, Kåreholt I. Work-related stress in midlife and all-cause mortality: can sense of coherence modify this association? *Eur J Pub Health.* 2016;26(6):1055-61.

69. Pandey A, Quick JC, Rossi AM, Nelson DL, Martin W. 11 Stress and the Workplace: 10 Years of Science, 1997–2007. *The handbook of stress science: Biology, psychology, and health*. 2010:137.
70. DePasquale N, Polenick CA, Davis KD, Moen P, Hammer LB, Almeida DM. The Psychosocial Implications of Managing Work and Family Caregiving Roles: Gender Differences Among Information Technology Professionals. *J Fam Issues*. 2017;38(11):1495-519.
71. Garrido MM, Hash-Converse JM, Leventhal H, Leventhal EA. Stress and chronic disease management. *The handbook of stress science: Biology, psychology, and health*. 2011:487-500.
72. Hudson JL, Moss-Morris R. Treating Illness Distress in Chronic Illness. *Eur Psychol*. 2019;24(1):26-37.
73. Madsen IE, Jorgensen AF, Borritz M, Nielsen ML, Rugulies R. Is the association between high strain work and depressive symptoms modified by private life social support: a cohort study of 1,074 Danish employees? *BMC Pub Health*. 2014;14:698.
74. Kim HJ, Hur W-M, Moon T-W, Jun J-K. Is all support equal? The moderating effects of supervisor, coworker, and organizational support on the link between emotional labor and job performance. *BRQ Business Res Quart*. 2017;20(2):124-36.
75. Almeida DM, Davis KD. Workplace Flexibility and Daily Stress Processes in Hotel Employees and their Children. *Ann Am Acad Pol Soc Sci*. 2011;638(1):123-40.
76. Yu TC, Zhou H, Suh K, Arcona S. Assessing the importance of predictors in unplanned hospital readmissions for chronic obstructive pulmonary disease. *Clinicoecon Outcomes Res*. 2015;7:37-51.

## Tables

Table 1. Characteristics of respondents with perceived stress scale (PSS) score (N=31,025).

	PSS mean (SD)	N	% <sup>a</sup>
<b>All</b>	12.2 (7.2)	31,025	—
<b>Gender</b>			
Men	11.4 (6.7)	14,447	49.7
Women	13.1 (7.6)	16,578	50.3
<b>Age</b>			
16-24	14.2 (6.5)	3,505	14.8
25-64	12.2 (7.2)	19,122	62.7
≥65	11.2 (7.5)	8,398	22.5
<b>Ethnic background</b>			
Danish	12.0 (7.4)	29,271	89.9
Other Western <sup>b</sup>	13.4 (5.3)	784	4.2
Non-Western <sup>c</sup>	15.6 (5.2)	970	5.9
<b>Educational attainment</b>			
Low (0-10 years)	14.4 (7.4)	4,437	15.2
Medium (11-15 years)	12.0 (7.3)	16,923	51.8
High (15- years)	11.2 (6.6)	8,497	28.3
Unknown	14.5 (6.5)	1,168	4.7
<b>Employment status</b>			
Students	13.7 (6.2)	2,546	11.2
Employed	11.1 (6.7)	16,288	51.1
Unemployed	15.9 (7.2)	539	2.0
Cash or sickness benefits, etc.	17.9 (7.8)	1,414	5.6
Disability pension	17.8 (7.0)	1,086	4.1

Early retirement pension	9.7 (7.0)	730	2.0
Retirement pension	11.4 (7.4)	7,576	20.7
Unknown	13.7 (6.6)	846	3.3

<sup>a</sup> Weighted percentage. <sup>b</sup> Other Western countries: all 27 EU countries, Andorra, Iceland, Liechtenstein, Monaco, Norway, San Marino, Switzerland, Vatican State, Canada, USA, UK, Australia and New Zealand. <sup>c</sup> Non-Western: all other countries.

Table 2. Prevalence of stressors and perceived social support (N=30,261).

	N	% <sup>a</sup>
<b>Financial circumstances</b>		
No	19,120	58.9
Yes, a little	7,121	25.2
Yes, partly	2,305	9.0
Yes, a lot	1,715	6.9
<b>Housing conditions<sup>b</sup></b>		
No	24,527	77.2
Yes, a little	3,616	13.8
Yes, partly	1,279	5.3
Yes, a lot	838	3.6
<b>Work situation</b>		
No	18,923	59.8
Yes, a little	6,587	22.9
Yes, partly	2,708	9.7
Yes, a lot	2,040	7.7
<b>Relationship with partner</b>		
No	22,927	75.1
Yes, a little	5,196	17.1
Yes, partly	1,366	4.9
Yes, a lot	768	3.0
<b>Relationship with family and friends</b>		
No	22,639	72.9
Yes, a little	6,030	21.1
Yes, partly	1,174	4.4

Yes, a lot	413	1.7
<b>Disease</b>		
No	17,252	57.8
Yes, a little	7,781	24.7
Yes, partly	3,151	10.4
Yes, a lot	2,072	7.1
<b>Disease among close relatives</b>		
No	17,853	60.4
Yes, a little	8,127	25.7
Yes, partly	2,842	9.1
Yes, a lot	1,435	4.7
<b>Deaths among close relatives</b>		
No	24,581	81.2
Yes, a little	3,235	10.6
Yes, partly	1,293	4.3
Yes, a lot	1,152	3.9
<b>Other types of distress</b>		
No	26,531	87.0
Yes, a little	1,913	6.4
Yes, partly	1,004	3.6
Yes, a lot	812	3.0
<b>Perceived social support</b>		
Yes, always	18,374	59.2
Yes, mostly	7,707	25.8

Yes, sometimes	2,825	10.1
No, never or almost never	1,334	4.9

<sup>a</sup> Weighted percentage.

Table 3. Unadjusted and adjusted dominance analyses for all respondents and subgroups stratified by gender, age, educational attainment and employment status (N=29,860). Predictors contributing with more than 10% of the adjusted R<sup>2</sup> are stated in bold.

	Unadjusted R <sup>2</sup>	Adjusted R <sup>2</sup>	Adjusted rank and R <sup>2</sup> of stressors and perceived social support	
<b>All<sup>a</sup></b>	38.7% (N=29,860)	42.5% (N=29,541)	<b>1. Disease (9.5%)</b> <b>2. Perceived social support (5.9%)</b> <b>3. Work situation (4.9%)</b> 4. Relationship with family and friends (4.1%) 5. Financial circumstances (3.2%)	6. Other types of distress (2.3%) 7. Relationship with partner (1.9%) 8. Housing conditions (1.8%) 9. Disease among close relatives (1.3%) 10. Deaths among close relatives (0.5%)
<b>Gender<sup>b</sup></b>				
Men	39.1% (N=13,903)	41.9% (N=13,770)	<b>1. Disease (10.0%)</b> <b>2. Perceived social support (5.9%)</b> <b>3. Work situation (5.4%)</b> 4. Relationship with family and friends (4.0%) 5. Financial circumstances (3.2%)	6. Other types of distress (2.3%) 7. Housing conditions (2.2%) 8. Relationship with partner (1.9%) 9. Disease among close relatives (1.3%) 10. Deaths among close relatives (0.5%)
Women	38.2% (N=15,957)	42.1% (N=15,771)	<b>1. Disease (9.2%)</b> <b>2. Perceived social support (6.0%)</b> <b>3. Work situation (4.6%)</b> <b>4. Relationship with family and friends (4.2%)</b> 5. Financial circumstances (3.2%)	6. Other types of distress (2.5%) 7. Relationship with partner (1.9%) 8. Housing conditions (1.6%) 9. Disease among close relatives (1.4%) 10. Deaths among close relatives (0.5%)

	Unadjusted R <sup>2</sup>	Adjusted R <sup>2</sup>	Adjusted rank and R <sup>2</sup> of stressors and perceived social support	
<b>Age<sup>c</sup></b>				
16-24	38.4% (N=3,302)	42.8% (N=3,272)	<b>1. Disease (6.7%)</b> <b>2. Relationship with family and friends (6.3%)</b> <b>3. Perceived social support (5.3%)</b> 4. Work situation (3.6%) 5. Financial circumstances (3.6%)	6. Other types of distress (3.0%) 7. Housing conditions (1.9%) 8. Relationship with partner (1.5%) 9. Disease among close relatives (1.2%) 10. Deaths among close relatives (0.1%)
25-64	43.1% (N=18,447)	44.8% (N=18,271)	<b>1. Disease (10.0%)</b> <b>2. Work situation (6.9%)</b> <b>3. Perceived social support (6.4%)</b> 4. Relationship with family and friends (4.0%) 5. Financial circumstances (3.8%)	6. Other types of distress (2.6%) 7. Housing conditions (2.1%) 8. Relationship with partner (2.1%) 9. Disease among close relatives (1.3%) 10. Deaths among close relatives (0.5%)

	Unadjusted R <sup>2</sup>	Adjusted R <sup>2</sup>	Adjusted rank and R <sup>2</sup> of stressors and perceived social support	
≥65	29.3% (N=8,111)	34.0% (N=7,998)	<b>1. Disease (10.9%)</b> <b>2. Perceived social support (5.4%)</b> 3. Relationship with family and friends (3.0%) 4. Relationship with partner (2.1%) 5. Disease among close relatives (1.6%)	6. Other types of distress (1.3%) 7. Financial circumstances (1.2%) 8. Housing conditions (1.1%) 9. Deaths among close relatives (0.8%) 10. Work situation (0.3%)
<b>Educational attainment<sup>d</sup></b>				
Low	36.6% (N=4,337)	38.7% (N=4,337)	<b>1. Disease (10.2%)</b> <b>2. Perceived social support (5.6%)</b> 3. Relationship with family and friends (3.7%) 4. Financial circumstances (2.6%) 5. Other types of distress (2.4%)	6. Work situation (2.0%) 7. Relationship with partner (1.8%) 8. Disease among close relatives (1.6%) 9. Housing conditions (1.6%) 10. Deaths among close relatives (0.7%)
Medium	40.3% (N=16,766)	42.6% (N=16,766)	<b>1. Disease (10.3%)</b> <b>2. Perceived social support (6.5%)</b> <b>3. Work situation (5.0%)</b> 4. Relationship with family and friends (4.2%) 5. Financial circumstances (3.6%)	6. Other types of distress (2.1%) 7. Relationship with partner (1.9%) 8. Housing conditions (1.9%) 9. Disease among close relatives (1.3%) 10. Deaths among close relatives (0.5%)


	Unadjusted R <sup>2</sup>	Adjusted R <sup>2</sup>	Adjusted rank and R <sup>2</sup> of stressors and perceived social support	
High	39.8% (N=8,438)	41.9% (N=8,438)	<b>1. Disease (8.2%)</b> <b>2. Work situation (7.4%)</b> <b>3. Perceived social support (5.1%)</b> 4. Relationship with family and friends (4.5%) 5. Financial circumstances (3.2%)	6. Other types of distress (3.0%) 7. Relationship with partner (2.0%) 8. Housing conditions (2.0%) 9. Disease among close relatives (1.4%) 10. Deaths among close relatives (0.3%)
<b>Employment status<sup>a</sup></b>				
Students	37.0% (N=2,535)	41.6% (N=2,526)	<b>1. Disease (6.8%)</b> <b>2. Perceived social support (6.0%)</b> <b>3. Relationship with family and friends (6.0%)</b> <b>4. Financial circumstances (4.3%)</b> 5. Work situation (3.4%)	6. Other types of distress (3.3%) 7. Housing conditions (1.7%) 8. Relationship with partner (1.7%) 9. Disease among close relatives (0.8%) 10. Deaths among close relatives (0.4%)

	Unadjusted R <sup>2</sup>	Adjusted R <sup>2</sup>	Adjusted rank and R <sup>2</sup> of stressors and perceived social support	
Employed (N=16,167)	36.1%	38.4% (N=16,068)	<b>1. Work situation (9.0%)</b> <b>2. Perceived social support (6.1%)</b> <b>3. Disease (6.0%)</b> 4. Relationship with family and friends (3.2%) 5. Relationship with partner (2.5%)	6. Financial circumstances (2.3%) 7. Other types of distress (1.9%) 8. Housing conditions (1.5%) 9. Disease among close relatives (1.4%) 10. Deaths among close relatives (0.2%)
Unemployed (N=539)	43.6%	46.1% (N=536)	<b>1. Disease (10.2%)</b> <b>2. Work situation (7.1%)</b> <b>3. Financial circumstances (5.4%)</b> <b>4. Relationship with family and friends (4.7%)</b> 5. Perceived social support (4.2%)	6. Relationship with partner (2.7%) 7. Housing conditions (2.5%) 8. Other types of distress (1.4%) 9. Disease among close relatives (0.7%) 10. Deaths among close relatives (0.4%)
Cash or sickness benefits (N=1,401)	47.2%	51.8% (N=1,362)	<b>1. Disease (13.8%)</b> <b>2. Relationship with family and friends (6.2%)</b> <b>3. Financial circumstances (5.3%)</b> 4. Perceived social support (4.4%) 5. Other types of distress (3.8%)	6. Work situation (3.3%) 7. Housing conditions (2.4%) 8. Disease among close relatives (2.1%) 9. Relationship with partner (1.4%) 10. Deaths among close relatives (0.4%)

	Unadjusted R <sup>2</sup>	Adjusted R <sup>2</sup>	Adjusted rank and R <sup>2</sup> of stressors and perceived social support	
Disability pension (N=1,029)	38.7%	42.6% (N=986)	<b>1. Disease (8.4%)</b> <b>2. Perceived social support (6.8%)</b> <b>3. Relationship with family and friends (5.8%)</b> 4. Other types of distress (3.2%) 5. Financial circumstances (3.0%)	6. Housing conditions (2.6%) 7. Deaths among close relatives (2.1%) 8. Disease among close relatives (1.9%) 9. Relationship with partner (1.5%) 10. Work situation (1.4%)
Early retirement pension (N=714)	41.4%	43.9% (N=712)	<b>1. Disease (11.7%)</b> <b>2. Perceived social support (9.6%)</b> <b>3. Relationship with family and friends (4.9%)</b> 4. Housing conditions (2.9%) 5. Financial circumstances (2.9%)	6. Other types of distress (2.2%) 7. Work situation (2.0%) 8. Relationship with partner (1.4%) 9. Disease among close relatives (1.3%) 10. Deaths among close relatives (1.2%)
Retirement pension (N=7,299)	29.6%	33.3% (N=7,187)	<b>1. Disease (11.0%)</b> <b>2. Perceived social support (5.6%)</b> 3. Relationship with family and friends (3.2%) 4. Relationship with partner (2.4%) 5. Disease among close relatives (1.6%)	6. Other types of distress (1.4%) 7. Financial circumstances (1.2%) 8. Housing conditions (1.2%) 9. Deaths among close relatives (0.8%) 10. Work situation (0.2%)


<sup>a</sup> Adjusted for gender, age, ethnic background and educational attainment. <sup>b</sup> Adjusted for age, ethnic background and educational attainment. <sup>c</sup> Adjusted for gender, ethnic background and educational attainment. <sup>d</sup> Adjusted for gender, age and ethnic background.

## Figures



**Figure 1**

Adjusted dominance analyses for all respondents (N=29,541). Rank of stressors and social support based on average increase in total variance explained by the model (R<sup>2</sup>) when adding a variable to all possible sub-models. Contribution to total variance is indicated next to the bar in absolute and relative (in parentheses) terms. Sociodemographic characteristics were included in all models.



**Figure 1**

Adjusted dominance analyses for all respondents (N=29,541). Rank of stressors and social support based on average increase in total variance explained by the model (R<sup>2</sup>) when adding a variable to all possible sub-models. Contribution to total variance is indicated next to the bar in absolute and relative (in parentheses) terms. Sociodemographic characteristics were included in all models.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SorensenetalDeterminantsofstressTablesBMCPublicHealth.docx](#)
- [SorensenetalDeterminantsofstressTablesBMCPublicHealth.docx](#)