Tissue Adhesive for Wound Closure in enhanced-recovery Total Hip Arthroplasty: A Prospective, Randomized and Controlled Study

Xiangpeng Kong

Chinese PLA General Hospital

Minzhi Yang

Nankai University

Zheng Cao

Nankai University

Jiying Chen

Chinese PLA General Hospital

Wei Chai (✉ chaiweiguanjie@sina.com)

Chinese PLA General Hospital

Yan Wang

Chinese PLA General Hospital

Research article

Keywords: total hip arthroplasty, enhanced recovery after surgery, tissue adhesive, wound closure, dressing change, prospective, randomized and controlled study

Posted Date: February 12th, 2020

DOI: <https://doi.org/10.21203/rs.2.20875/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: We aimed to present our experience of adopting tissue adhesive as adjunct to standard wound closure in total hip arthroplasty (THA) and evaluate its performance.

Methods: From September 2019 to November 2019, we prospectively enrolled consecutive patients who underwent simultaneous bilateral THA in this randomized and controlled study. Standard wound closure was applied on one side of hip and additional tissue adhesive on the other side at random. We collected and analyzed patients' information, including age, gender, body mass index (BMI), diagnosis, postoperative length of stay (LOS), dressing changes, wound evaluation scores, wound-related cost and complications.

Results: Thirty patients with simultaneous bilateral THA were enrolled in this study. During the hospital stay, the dressing change in hips with tissue adhesive was significantly less than that in the other hips ($p=0.000$). However, the wound-related cost in hips with tissue adhesive was significantly higher ($p=0.000$). At one-month follow up, patient-reported wound measurement of hips with tissue adhesive was significantly better than the other hips ($p=0.004$). Seventeen patients preferred tissue adhesive and only five patients preferred standard wound closure.

Conclusions: Tissue adhesive could significantly reduce wound drainage and increase patient satisfaction, which can be an ideal adjunct to standard wound closure in enhanced-recovery THA.

Background

At present, enhanced recovery after surgery (ERAS) has developed rapidly in the field of joint replacement [1-3]. In some institutes, the length of stay (LOS) after total hip arthroplasty (THA) has been shortened to less than 48 hours and some even became daytime surgery [4,5]. The advent of ERAS also raised the stringent requirements of surgical techniques and perioperative management [6,7]. Wound closure is one of the most important aspects in perioperative management. However, compared with surgical techniques, there were relatively fewer studies on wound closure and care.

Prolonged wound drainage, which is a common complication after joint replacement, could result in delayed wound healing, limited postoperative activity and even periprosthetic joint infection (PJI) [8,9]. One ideal wound closure, which permits rapid rehabilitation without wound drainage, should be design simple and easy to use.

In recent years, tissue adhesive has been introduced and adopted in orthopedic surgery [10-15]. According to the previous studies, tissue adhesive may be an ideal supplement to standard wound closure following total knee arthroplasty (TKA) [8]. It exists as liquid and polymerizes rapidly when contacting with skin tissue. The protective film produced by the tissue adhesive could quarantine with the external environment and prevent the foreign substances from invading the wound in the early postoperative period [16-19]. However, as we know, there was no study using tissue adhesive as the supplement to subcuticular suture in THA.

Thus in this prospective study, we aimed to present our experience of adopting tissue adhesive as adjunct to standard wound closure in THA and evaluate its role and cost performance in reducing wound drainage.

Methods

Study population and design

From September 2019 to November 2019, we prospectively enrolled consecutive patients who underwent simultaneous bilateral THA in this randomized self-control study. The study was approved by the local ethics committee.

Inclusion criteria: 1. age between 18 and 60 years old; 2. the bilateral THA with the same prosthesis through the posterolateral approach; 3. the written informed consent obtained prior to participating in this study. Exclusion criteria: 1. previous open surgery or major trauma or infection in either hip; 2. eloid, psoriasis, eczema or other skin diseases; 3. allergy to the ingredients of the tissue adhesive; 4. underlying malignant tumors; 5. regular anticoagulation therapy; 6. peripheral vascular disease; 7. active inflammatory arthropathy; 8. bilateral surgeries performed in stages.

Sample size calculation: according to the previous study and the preliminary results of our pre-experiment, we set $\alpha=0.05$, $\beta=0.10$, the mean difference of dressing change was 1.0. An estimated 24 patients would be needed to provide 90% power. Finally, we decide to enroll 30 patients, which allowed for 20% loss to follow up.

Surgical procedures of wound closure

All patients underwent the right THA firstly and left secondly. The fixed surgical team performed the surgeries and two fixed residents performed wound closure. Tranexamic acid (TXA) was given intravenously twice before incision and wound closure.

Standard wound closure for different layers: 1. the joint capsule and external rotator muscles were reconstructed with 2-0 Ethibond non-absorbable suture W4843 (Ethicon, Somerville, NJ, USA). 2. the deep fascia and superficial fascia were closed with 2-0 absorbable knotless barbed running suture (Quill, Surgical Specialties Corporation, IL, USA) and 4-0 coated Vicryl Plus antibacterial interrupted suture (Ethicon, Somerville, NJ, USA). 3. the subcuticular tissue was closed with 4-0 absorbable knotless barbed running suture (Ethicon, Somerville, NJ, USA).

The right hip was randomized by computer-generated method in the opaque envelopes after standard wound closure. Standard wound closure was applied on one side of hip and additional tissue adhesive on the other side at random. (Figure 1). The residents remained blinded before the allocations.

The HISTOACRYL® tissue adhesive (B.Braun, Melsungen, German) was applied evenly on both sides of the wound and waited for air-drying of 30 seconds. When the tissue adhesive got dry, a wound dressing was covered on surface. During the hospital stay, patients and caregivers were told to notify the surgeon or nurse for dressing change if blood or exudate soaked the dressing, which was the standard protocol of postoperative dressing change. If dressing could keep dry and clean before discharge, patient wound received one dressing change. The dressing change caused by discharge was not recorded. All patients had antibiotics within 24 hours, shower after 14 days and aspirin in 35 days postoperatively.

Follow-up and wound evaluation

Each patient's preoperative, intraoperative and postoperative information, including age, gender, body mass index (BMI), diagnosis, postoperative LOS, times of dressing changes and wound-related cost, were collected prospectively.

Wound-related cost was a sum cost of the suture, tissue adhesive, wound dressing, and other additional materials during the hospital stay.

At one-month follow up after surgery, wound-related complications and wound-related evaluation scores were recorded. Wound-related complications were defined as redness, dehiscence, subcutaneous hematoma, prolong wound drainage (>5 days), surgical site infection (SSI) and re-suture caused by any reasons. Wound-related evaluation scores included patient scar assessment score (PSAS), Hollander wound evaluation score (HWES) and Vancouver scar score (VSS). Both two residents received professional training and stay satisfying consistency of wound evaluation in the pre-experiment. The mean scores evaluated by two resident were used as final scores. In addition, all patients would be asked to choose their prefer wound closure.

PSAS [20]: the scoring system mainly refers to the patient's own feeling and evaluation of wound. Six represents normal skin and sixty represents worst imaginable scar in the patient's scale (Table 1).

Table 1. Patient scar assessment score (PSAS).

*	1	2	3	4	5	6	7	8	9	10
Is the scar painful?										
Is the scar itching?										
#	1	2	3	4	5	6	7	8	9	10
Is the color of the scar different?										
Is the scar more stiff?										
Is the thickness of the scar different?										
Is the scar irregular?										
Total score Patient Scar Score										
* 0 means “no, no complains”,10 means “yes, more imaginable”										
# 0 means “no, as normal skin”,10 means “yes, very different”										

HWES [21]: the scoring system includes 6 items, which are step-off of borders, contour irregularities, margin separation, edge inversion, excessive distortion and overall appearance. It was evaluated by two independent orthopedic residents unknown to the result of allocation. One point is for each item. The lower the score, the better the wound healing (Table 2).

Table 2. Hollander wound evaluation score (HWES).

Incision attribute	Score if absent	Score if present
Step-off borders	0	1
Contour Irregularities	0	1
Margin Separation	0	1
Edge inversion	0	1
Excessive Distortion	0	1
Overall appearance	0 (satisfactory)	1 (unsatisfactory)
Total Hollander score	0 (best)	6 (worse)

VSS [22]: the scoring system includes 4 items. They are vascularity, pliability, height and pigmentation. It was evaluated by two independent orthopedic residents unknown to the result of allocation. Lower scores represent a more normal appearance (Table 3).

Table 3. Vancouver scar score (VSS).

Score	Vascularity	Pliability	Height	Pigmentation
0	Normal	Normal	Flat	Normal
1	Pink	Supple	<2mm	Hypopigmentation
2	Red	Yielding	2-4mm	Mixed
3	Purple	Firm	>4mm	Hyperpigmentation
4	-	Banding	-	-

Statistical analysis

All statistical analyses were performed by SPSS version 22 (Inc., Chicago, IL, USA). Data was showed as median, mode and interquartile range (IQR) (skewed distribution) or mean $\pm$ standard deviation (SD) (normal distribution). Measurement data was analyzed by student's tests or rank-sum test. Count data was analyzed by rank-sum test or Fisher exact test. A value of $\alpha=0.05$ suits all tests. The intraclass correlation coefficient (ICC) was used to assess the observers' agreement: 0.81 to 1.00, nearly perfect reliability; 0.61 to 0.80, strong reliability; 0.41 to 0.60, moderate reliability; 0.21 to 0.40, fair reliability; and 0 to 0.20, poor reliability.

Results

Thirty-three patients were enrolled in this study. Two patients failed to complete the scheduled follow-up and one patient had re-operation because of recurrent dislocation. Finally, 30 patients were analyzed, which included 23 patients with osteonecrosis of femoral head (ONFH), 5 patients with developmental dysplasia of hip (DDH) and 2 patients with ankylosing spondylitis (AS). The two AS patients have or have had the non-steroidal anti-inflammatory drugs (NSAIDs) regularly for pain relief, and none of them had the immunosuppressants or glucocorticoids between preoperative 1 month and postoperative 1 month. The basic information was shown in Table 4.

Table 4. The demographics of thirty patients.

Basic information	Data
Age (Median, mode, IQR) (years)	30.5, 32, 11
Male:Female	17:13
BMI (Mean $\pm$ SD) (kg/m ²)	22.94 $\pm$ 3.62
Postoperative LOS (Mean, mode, IQR) (day)	4, 4, 1

Three patients (four hips) had prolonged wound drainage and their basic information were shown in Table 5. The BMI of three patients was 28.50 ± 1.94 kg/m² and that of remaining patients was 22.32 ± 3.21 kg/m². The difference was significant ($p=0.003$). There was no PJI during the follow-up period in this cohort.

Table 5. The information of three patients who had prolonged wound drainage.

Sex	Age (years)	Diagnosis	BMI (kg/m ²)	Hip	Wound closure	Treatment
Female	34	ONFH	27.13	Right	Standard	Re-suture and oral antibiotics
Male	31	DDH	27.65	Left	Standard	Re-suture and oral antibiotics
Male	37	ONFH	30.72	Right	Standard	Re-suture and oral antibiotics
-	-	-	-	Left	Tissue adhesive	Partial pressure bandage and oral antibiotics

During the hospital stay, the dressing change in hips with tissue adhesive was significantly less than that in the other hips ($p=0.000$) (Table 6, Figure 2). Wound-related cost in hips with tissue adhesive was significantly higher than that in the other hips ($p=0.000$).

The postoperative one-month appearances of bilateral wound in one patient were shown in Figure 3. From the view of patients, the PSAS in hips with tissue adhesive was significantly better than the other hips ($p=0.004$). From the view of observers, there were no significant differences in the HWES or VSS between two methods (Table 6). These scores assessed by two observers had nearly perfect reliability with each other ($ICC>0.81$) (Table 7).

Based on the subjective feeling from patients. seventeen patients (57%) preferred the side with tissue adhesive, eight patients (27%) had no preference and only five patients (16%) preferred the other side (Figure 4).

Table 6. Dressing change, wound-related cost, complications and evaluation scores between two methods of wound closure in thirty patients.

Data	Tissue adhesive	Standard wound closure	P
Dressing change (Median, mode, IQR)	0, 0, 1	2, 2, 1	0.000
Wound-related cost (Mean±SD) (US dollar)	272.39±10.12	221.83±13.55	0.000
Wound-related complications	1/29	3/27	0.306
PSAS (Mean±SD)	22.83±9.48	30.57±9.54	0.004
HWES (Median, mode, IQR)	0, 0, 0	0, 0, 0	0.414
VSS (Mean±SD)	5.13±1.16	5.77±1.16	0.057

Table 7. The inter-observers' agreements between two observers.

Evaluation system	PSAS, 95% CI	HWES, 95% CI	VSS, 95% CI
Inter-observer agreement	0.962 (0.931 to 0.965)	0.927 (0.892 to 0.964)	0.854 (0.755 to 0.910)

Discussion

This prospective, randomized and self-controlled study found that the supplementary tissue adhesive to standard wound closure in hip replacement could reduce wound drainage and increase patient satisfaction, despite of increasing medical expense. So tissue adhesive combining with standard wound closure can be one reliable and effective choice for enhanced- recovery THA.

Since the development of ERAS in orthopedic surgeries aims to improve wound healing, increase patient satisfaction and reduce hospitalization days and complications, wound closure has become increasingly important.

In hip replacement, the most common wound closure methods are staples or sutures. Yet, it seems to be no consensus in the literature as to which closure method is superior [23]. From the perspective of usage time, skin staples requires less time than suture [24]. From the perspective of wound complications, one meta-analysis including several high-quality studies found that there was a significantly higher risk of developing a wound infection, when the wound was closed with skin staples rather than sutures. Furthermore, this risk was especially greater in patients who underwent hip surgery [25]. From the perspective of wound healing, running subcuticular suture proved to ensure more physiologic robust blood flow than vertical suture and skin staples, which was one important factor in determining wound healing [26].

In our institute, the patients who underwent THA were relatively young, and had the higher requirement of wound appearance and enhanced recovery.

We usually adopted subcuticular suture, rather than skin interrupted suture and staple, for cosmetic appearance. Many studies had also conformed the role of subcuticular suture in improving incision appearance after joint surgery [27-29]. However, this standard wound closure failed to solve the fundamental problem of wound drainage.

Prolonged wound drainage could be detrimental to clinical outcomes [11,30]. Wound management without drainage is an important partway to ERAS, which urgently need new techniques and materials.

How to avoid postoperative wound drainage remains unknown. Maybe tissue adhesive is a good choice. The tissue adhesive in this study is composed by N-butyl-2-cyanoacrylate and early application in humans can be traced back to 1980s [31]. It would form a protective film on the surface of wound and provide enough adhesive strength in 30 seconds. Furthermore, it has certain bacteriostatic action. Therefore, these characteristics and advantages contribute to its extensive clinical application in surgery.

In this study, dressing change in thirty hips with tissue adhesive decreased by 50% when comparing to the other thirty hips. The patient-reported evaluation scores of wounds with tissue adhesive were better than those without tissue adhesive. In the group of standard wound closure, the patients who had bilateral posterolateral incisions needed to lie on their side for 3-5 minutes to finish disinfection and dressing change, which would aggravate the pain of the other incision. But in the group of tissue adhesive, less dressing change could reduce surgeons' workload and inconvenience to patients. This might explain why patients preferred tissue adhesive were three times as many as standard wound closure.

Wood et al reported that the time taken for wounds to stop oozing following hip replacement was significantly related to the BMI [32]. Zhang et al found that patients who were obese (BMI > 30 kg/m²) had higher odds of wound complications in both nonurgent and urgent orthopedic hip surgery [33]. In this

study, there were significantly different BMI between the patients with and without prolonged wound drainage, which also showed that obesity might be a risk factor of wound complications.

Some surgeons have also reported their result of using tissue adhesive in arthroplasty. Gromov and El-Gazzar proved its role in reducing wound drainage when tissue adhesive worked as the supplement to staples [8,11]. However, another two randomized controlled trials showed no significant differences in cosmetic appearance of scars, incidence of complications, and patient satisfaction between tissue adhesive and standard wound closure [10-12]. These authors just analyzed the clinical outcomes, without taking dressing change and medical expenses into consideration. According to the above studies and our experience, tissue adhesive should be the adjunct, not substitution, to standard wound closure in THA.

There is no denying that some patients were allergic to tissue adhesive, though no allergy was found in this study [34,35]. Dunnett performed 912 knee replacement and discovered allergy in 1.7% patients [36]. It usually manifested as erythematous pruritic papular rash surrounding incision site within postoperative 1 to 3 weeks and cleared with topical steroids.

This study has several strengths. Firstly, the randomized controlled trial could reduce selection bias, information bias and confounding bias, which was unavoidable in case-control study. Secondly, the self-controlled study could balance the bias between comorbidities and physical conditions among individuals. Thirdly, patients and residents evaluated wound respectively. Fourthly, the study took economic factors into account.

This study is not without limitations. Firstly, since this study aimed to investigate the effect of wound closure on enhanced-recovery THA, we only followed up patients in postoperative one month. Small sample size and short follow-up period were not sufficient for detailed and comprehensive analysis. Secondly, patients who can bear bilateral hip replacements simultaneously usually have good physical conditions. It is unknown the result when tissue adhesive used in a less healthy population. Thirdly, we didn't conduct mechanical tests to investigate the adhesive strength of tissue adhesive, which influence the persuasiveness of this study. Fourthly, using tissue adhesive generally took only one minute or less and wound closure time weren't recorded, which made it impossible to compare their influence on operating time. Fifthly, because medical consumptive material pricing systems in other institutes were different and wound-related costs after discharge were not analyzed, the cost-performance of tissue adhesive may be different from this study.

Conclusions

Tissue adhesive could significantly reduce wound drainage and increase patient satisfaction, which can be an ideal adjunct to standard wound closure in enhanced-recovery THA.

Abbreviations

THA: total hip arthroplasty; BMI: body mass index; LOS: length of stay; ERAS: enhanced recovery after surgery; PJI: periprosthetic joint infection; TKA: total knee arthroplasty; TXA: tranexamic acid; SSI: surgical site infection; PSAS: patient scar assessment score; HWES: Hollander wound evaluation score; VSS: Vancouver scar score; IQR: interquartile range; SD: standard deviation; ICC: intraclass correlation coefficient; ONFH: osteonecrosis of femoral head; DDH: developmental dysplasia of hip; AS: ankylosing spondylitis.

Declarations

- **Ethics approval and consent to participate:** The study was approved by the medical ethics committee of Chinese PLA General Hospital (S2019-166-01) and registered at Chinese Clinical Trial Registry (the clinical trial registration number: ChiCTR1900025730). All procedures were conducted in compliance with the guidelines of the Declaration of Helsinki. The patients were informed consent prior to their participation in the study.
- **Consent for publication:** We have obtained written consent for publication.
 - **Availability of data and material:** All data generated or analyzed during this study are included in this published article.
 - **Competing interests:** The authors declare that they have no competing interests.
 - **Funding:** None.
- **Author Contributions:** All authors have made substantial contributions to: (1) the conception and design of the study, acquisition of data, or analysis and interpretation of data; (2) drafting the article or revising it critically for important intellectual content; (3) final approval of the version to be submitted. WC: primarily responsible for oversight of the research project, including all data acquisition and analysis, and manuscript preparation and approval. XK and MY: primarily responsible for all computational analyses in the article and the drafting of the manuscript. XK and MY were the co-first authors. ZC, JC and YW: revised the manuscript and helped perform the analysis with constructive discussions. All authors have read and approved the final submitted manuscript.
- **Acknowledgement:** Not applicable.
- **Authors information**

First author: Xiangpeng Kong, MD

- mail:18810999609@163.com.

Address: No.28 Fuxing Road, Haidian, Beijing, China. Mail:100853.

Co-first author: Minzhi Yang, postgraduate

E-mail:Yangminzhi1995@163.com.

Address: No.94 Weijin Road, Nankai, Tianjin, China. Mail:300071.

Author: Zheng Cao, postgraduate

E-mail: 361782992@qq.com.

Address: No.94 Weijin Road, Nankai, Tianjin, China. Mail:300071.

Author: Jiying Chen, MD

E-mail: chenjiyingjoint@sina.com.

Address: No.28 Fuxing Road, Haidian, Beijing, China. Mail:100853.

Corresponding author: Wei Chai, MD

E-mail: chaiweiguanjie@sina.com.

Address: No.28 Fuxing Road, Haidian, Beijing, China. Mail:100853.

Author: Yan Wang, MD

E-mail: wangyanguanjie@sina.com.

Address: No.28 Fuxing Road, Haidian, Beijing, China. Mail:100853.

References

1. Kehlet Henrik, Fast-track hip and knee arthroplasty.[J] .Lancet, 2013, 381: 1600-2.
2. Stambough Jeffrey B, Nunley Ryan M, Curry Madelyn C et al. Rapid recovery protocols for primary total hip arthroplasty can safely reduce length of stay without increasing readmissions.[J] .J Arthroplasty, 2015, 30: 521-6.
3. Husted Henrik, Holm Gitte, Jacobsen Steffen, Predictors of length of stay and patient satisfaction after hip and knee replacement surgery: fast-track experience in 712 patients.[J] .Acta Orthop, 2008, 79: 168-73.
4. Raphael Michael, Jaeger Melanie, van Vlymen Janet, Easily adoptable total joint arthroplasty program allows discharge home in two days.[J] .Can J Anaesth, 2011, 58: 902-10.
5. Petersen Pelle B, Kehlet Henrik, Jørgensen Christoffer C et al. Incidence and Risk Factors for Stroke in Fast-Track Hip and Knee Arthroplasty-A Clinical Registry Study of 24,862 Procedures.[J] .J Arthroplasty, 2019, 34: 743-749.e2.
6. Khan Sameer K, Malviya Ajay, Muller Scott D et al. Reduced short-term complications and mortality following Enhanced Recovery primary hip and knee arthroplasty: results from 6,000 consecutive procedures.[J] .Acta Orthop, 2014, 85: 26-31.
7. Malek I A, Royce G, Bhatti S U et al. A comparison between the direct anterior and posterior approaches for total hip arthroplasty: the role of an 'Enhanced Recovery' pathway.[J] .Bone Joint J, 2016, null: 754-

60.

8. Gromov Kirill, Troelsen Anders, Raaschou Sofie et al. Tissue Adhesive for Wound Closure Reduces Immediate Postoperative Wound Dressing Changes After Primary TKA: A Randomized Controlled Study in Simultaneous Bilateral TKA.[J] .Clin. Orthop. Relat. Res., 2019, undefined: undefined.
9. Scuderi Giles R, Avoiding Postoperative Wound Complications in Total Joint Arthroplasty.[J] .J Arthroplasty, 2018, 33: 3109-3112.
10. Khan R J K, Fick D, Yao F et al. A comparison of three methods of wound closure following arthroplasty: a prospective, randomised, controlled trial.[J] .J Bone Joint Surg Br, 2006, 88: 238-42.
11. El-Gazzar Yaser, Smith Daniel C, Kim Sun Jin et al. The use of dermabond® as an adjunct to wound closure after total knee arthroplasty: examining immediate post-operative wound drainage.[J] .J Arthroplasty, 2013, 28: 553-6.
12. Livesey C, Wylde V, Descamps S et al. Skin closure after total hip replacement: a randomised controlled trial of skin adhesive versus surgical staples.[J] .J Bone Joint Surg Br, 2009, 91: 725-9.
13. Miller Adam G, Swank Michael L, Dermabond efficacy in total joint arthroplasty wounds.[J] .Am J. Orthop., 2010, 39: 476-8.
14. Garbuz Donald, 2-octylcyanoacrylate, staples, and sutures had similar wound closure outcomes after total hip or knee replacement.[J] .J Bone Joint Surg Am, 2006, 88: 1895.
15. Lieberman Jay, Skin adhesive or staples were not different for skin closure after total hip replacement. [J] .J Bone Joint Surg Am, 2010, 92: 476.
16. Pelissier P, Casoli V, Le Bail B et al. Internal use of n-butyl 2-cyanoacrylate (Indermil) for wound closure: an experimental study.[J] .Plast. Reconstr. Surg., 2001, 108: 1661-6.
17. Anderson J M, Gibbons D F, The new generation of biomedical polymers.[J] .Biomater Med Devices Artif Organs, 1974, 2: 235-48.
18. Braginsky Lena, Javellana Melissa, Cleveland Emily et al. Tissue Adhesive Compared With Sterile Strips After Cesarean Delivery: A Randomized Controlled Trial.[J] .Obstet Gynecol, 2019, 134: 295-301.
19. Dumville Jo C, Coulthard Paul, Worthington Helen V et al. Tissue adhesives for closure of surgical incisions.[J] .Cochrane Database Syst Rev, 2014, undefined: CD004287.
20. Raklyar Eduard, Zloty David M, Use of a patient and observer scar assessment scale to evaluate the V-Y advancement flap for reconstruction of medial cheek defects.[J] .Dermatol Surg, 2012, 38: 1968-74.
21. Hollander J E, Singer A J, Valentine S et al. Wound registry: development and validation.[J] .Ann Emerg Med, 1995, 25: 675-85.
22. Baryza M J, Baryza G A, The Vancouver Scar Scale: an administration tool and its interrater reliability. [J] .J Burn Care Rehabil, 1995, 16: 535-8.
23. Krishnan Rohin, MacNeil S Danielle, Malvankar-Mehta Monali S, Comparing sutures versus staples for skin closure after orthopaedic surgery: systematic review and meta-analysis.[J] .BMJ Open, 2016, 6: e009257.
24. Shantz Jesse A, Vernon James, Leiter Jeff et al. Sutures versus staples for wound closure in orthopaedic surgery: a randomized controlled trial.[J] .BMC Musculoskelet Disord, 2012, 13: 89.

25. Smith Toby O,Sexton Debbie,Mann Charles et al. Sutures versus staples for skin closure in orthopaedic surgery: meta-analysis.[J] .BMJ, 2010, 340: c1199.
26. Wyles Cody C,Jacobson Steven R,Houdek Matthew T et al. The Chitranjan Ranawat Award: Running Subcuticular Closure Enables the Most Robust Perfusion After TKA: A Randomized Clinical Trial.[J] .Clin. Orthop. Relat. Res., 2016, 474: 47-56.
27. Parvizi J, Ghanem E, Sharkey P et al. Does Excessive Anticoagulation Predispose the Patient to Periprosthetic Infection?[J]. Journal of Arthroplasty, 2007, 22(2):0-305.
28. Agarwala Sanjay,Vijayvargiya Mayank,Concealed cosmetic closure in total knee replacement surgery - A prospective audit assessing appearance and patient satisfaction.[J] .J Clin Orthop Trauma, 2019, 10: 111-116.
29. Rui Min,Zheng Xin,Sun Shao-Song et al. A prospective randomised comparison of 2 skin closure techniques in primary total hip arthroplasty surgery.[J] .Hip Int, 2018, 28: 101-105.
30. Shetty A A,Kumar V S,Morgan-Hough C et al. Comparing wound complication rates following closure of hip wounds with metallic skin staples or subcuticular vicryl suture: a prospective randomised trial. [J] .J Orthop Surg (Hong Kong), 2004, 12: 191-3.
31. Messi G,Canciani G,Marchi A G,Costs and benefits of the use of tissue adhesives in wounds in children.[J] .Pediatr Med Chir, 1990, 12: 185-8.
32. Wood James J,Bevis Paul M,Bannister Gordon C,Wound oozing after total hip arthroplasty.[J] .Ann R Coll Surg Engl, 2007, 89: 140-2.
33. Zhang Joyce C,Matelski John,Gandhi Rajiv et al. Can Patient Selection Explain the Obesity Paradox in Orthopaedic Hip Surgery? An Analysis of the ACS-NSQIP Registry.[J] .Clin. Orthop. Relat. Res., 2018, 476: 964-973.
34. El-Dars L D,Chaudhury W,Hughes T M et al. Allergic contact dermatitis to Dermabond after orthopaedic joint replacement.[J] .Contact Derm., 2010, 62: 315-7.
35. Gonzalo-Garijo,MAngeles,Pérez-Calderón,Remedios,Pérez-Rangel. Inmaculada et al. Contact dermatitis after orthopaedic surgery.[J] .Contact Derm., 2009, 61: 299-300.
36. Durando Dunnett,Porubsky Caitlin,Winter Suzanne et al. Allergic contact dermatitis to dermabond (2-octyl cyanoacrylate) after total knee arthroplasty.[J] .Dermatitis, 2014, 25: 99-100.

Figures

Figure 1

The appearances of bilateral wounds in the operating room (left: tissue adhesive; right: standard wound closure).

Figure 2

The times of dressing change between two methods in thirty patients.

Figure 3

The preference distribution for wound closure in thirty patients.

Figure 4

The appearances of bilateral wounds at postoperative one month (left: tissue adhesive; right: standard wound closure).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [CONSORT2010Checklist.pdf](#)