

Choroidal vascularity index changes in chronic migraine patients

Taha Sezer (✉ drtahasezer@gmail.com)

Duzce University School of Medicine <https://orcid.org/0000-0002-4888-4293>

Research Article

Keywords: Choroidal vascularity index, Choroidal thickness, Migraine, Optical coherence tomography

Posted Date: March 7th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1147803/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Purpose: Migraine is a neurovascular disorder characterized by recurrent headaches. The relationship between migraine disease and the choroid, has been examined in an attempt to elucidate the underlying pathophysiological mechanisms. This study evaluated choroidal vascularity index (CVI) in chronic migraine patients.

Methods: In this prospective study, we compared CT and CVI values of 36 chronic migraine patients (30 women and 6 men) during an attack-free period with those of 36 healthy individuals (30 women and 6 men) with no systemic or ocular disease, including headache. All patients underwent a detailed eye examination before enhanced depth imaging optical coherence tomography (EDI-OCT) imaging. Migraine patients were grouped as those with and without aura and were asked to rate their headache severity on visual analog scale (VAS; range 1-10) and estimate their monthly migraine frequency.

Results: The mean subfoveal CT (SFCT) was $300.52 \pm 88.30 \mu\text{m}$ in the migraine group and $262.85 \pm 70.68 \mu\text{m}$ in the control group. The mean CVI was $71.8\% \pm 6.2\%$ in the migraine group and $70.7\% \pm 5.3\%$ in the control group. SFCT and CVI did not differ significantly between the migraine and control groups ($p > 0.05$). VAS pain score was 8.17 ± 0.33 in the migraine group and was not correlated with SFCT ($r = 0$, $p = 0.998$) or CVI ($r = -0.06$, $p = 0.731$). The monthly migraine frequency was 5.60 ± 3.60 and was not correlated with SFCT ($r = -0.17$, $p = 0.328$) or CVI ($r = -0.06$, $p = 0.731$).

Conclusion: CT and CVI showed no significant differences from controls in chronic migraine patients during an attack-free period.

Introduction

Migraine is a neurological disorder characterized by severe, recurrent unilateral headaches.[1] The diagnosis is based on the characteristics of the headache and associated neurological symptoms such as gastrointestinal and autonomic nervous system symptoms. These symptoms include photophobia, phonophobia, and vomiting, and the pain is usually aggravated by physical activity. One-third of migraine patients also experience transient visual, sensory, language, or motor disturbances before (or rarely, during) the headache, which are referred to as the aura.[2]

The pathophysiology of migraine is not well understood and there is no consensus on existing theories. Vasogenic, neurogenic, and cortical spreading depression theories have been proposed to explain migraine pathophysiology.[3] The vasogenic theory attributes the headache to prolonged vasospasm followed by vasodilation,[4, 5] whereas the neurogenic theory suggests that vascular changes in migraine occur as a result of neuronal dysfunction. In particular, the release of numerous vasogenic neuropeptides in the trigeminovascular region and the triggering of nociceptive impulses support that the pathophysiology of migraine may be of neurovascular origin.[6] Therefore, migraine is currently considered a neurovascular disease.

The choroid receives most of the ocular blood flow, and understanding changes in its structure may provide insight into choroidal and ocular blood flow.[7, 8] Choroidal imaging was generally performed with indocyanine green angiography (ICGA) and contact B-scan ultrasound (US) before the introduction of spectral domain optical coherence tomography (SD-OCT) into clinical use.[9] Although choroidal imaging with SD-OCT was inadequate at first, the enhanced depth imaging technique (EDI-OCT) uses longer wavelengths and has enabled detailed visualization of the luminal and stromal structures of the choroidal layer.[10] Most recently, the increasing clinical use of optical coherence tomography angiography (OCTA) has allowed non-invasive examination of the deep and superficial retinal, choriocapillaris, and choroidal circulation.[11]

Numerous studies using EDI-OCT to identify vascular changes in the pathophysiology of migraine have investigated choroidal thickness (CT).[12–17] While measurement of CT may be helpful in clinical research, it is not a reliable parameter because it can be influenced by multiple factors, including diurnal variation, age, gender, and axial length. Therefore, research focus has shifted to choroidal vascularity index (CVI), which is not affected by physiological factors and is determined as the ratio of luminal area to total choroidal area using special software.[18–20] In this study, we aimed to examine differences in CT and CVI in patients with chronic migraine during an attack-free period compared to healthy controls.

Materials And Methods

Thirty-six migraine patients (30 women and 6 men) who were being followed due to chronic migraine and were referred from the neurology clinic of Düzce University and 36 control subjects (30 women and 6 men) with no ocular or systemic disease and no headache complaints were compared. The study was approved by the Düzce University Institutional Review Board and Ethics Committee and adhered to the Declaration of Helsinki. All participants provided informed consent to use their clinical data for this study.

The migraine patients were grouped as those with and without aura according to the Headache International Society criteria.[21] They were asked to rate their headache pain severity using a visual analogue scale (between 0-10 points) and estimate their monthly headache frequency. All patients used nonsteroidal anti-inflammatory drugs for their migraine attacks.

Patients with any disease that may affect choroidal flow (e.g., hypertension, diabetes mellitus, vasculitis, renal failure), smoking history, and use of drugs likely to affect CT (e.g., sildenafil, triptan, ergot alkaloids, antihistamines, decongestants) were not included in the study.

Ophthalmologic Examination

All participants in the control and migraine groups underwent a detailed ophthalmological examination by the same physician (M.B.). Best-corrected visual acuity (BCVA), fundoscopy, slit-lamp examination, and intraocular pressure measurement were performed in both groups.

Exclusion criteria included spherical and cylindrical refractive errors greater than +/-3 diopters (D), amblyopia, retinal or choroidal pathology, intraocular surgery, and media opacity that would prevent OCT imaging. The right eyes of all participants were evaluated in the study. SD-OCT scans were performed at the same time of day (9:00-10:00 am) to minimize the effect of diurnal variation on the choroid.[22]

Choroidal Thickness Measurement

CT measurements were performed by the same experienced ophthalmologist (S.T.) using SD-OCT (Heidelberg Engineering, Heidelberg, Germany). A Spectralis OCT (Heidelberg Engineering) was used with a standardized imaging protocol. A 9-mm horizontal image centered on the fovea was obtained with an average of 100 B-scans in each section to improve the signal-to-noise ratio. Eye-tracking mode was also used. All subjects were examined with pupil dilation. CT was measured from the outer edge of the hyperreflective line corresponding to the retinal pigment epithelium (RPE) to the hyporeflexive line corresponding to the choroidal-scleral interface. Measurements were made in three different regions, at the foveal center and 1000 μm nasal and temporal of the fovea.

Choroid Vasculature Index Assessment

Sonoda et al. used the image binarization technique to calculate CVI.[23] In this study we used the slightly modified technique described by Agrawal et al.[18] Open-source ImageJ software was used for image processing (version 1.47; provided in the public domain by the National Institutes of Health, Bethesda, MD, USA; <http://imagej.nih.gov/ij/>). Briefly, 1×1 pixel EDI-OCT images were opened in ImageJ and the scale was set to 200 μm . A total choroidal area (TCA) 1.5 mm in width and centered on the fovea was selected and marked using with the manual plotting polygonal tool (Fig. 1). The upper border of the choroid was marked at the RPE and the lower border was marked at the choroid-sclera junction. The entire length of the OCT B-scan was used for analysis. Then the EDI-OCT B scan was converted to 8-bit images using the default setting. Niblack's automated local threshold tool was applied to delineate the luminal area (LA) and stromal area (SA). The image was then converted back to an RGB (red, green, blue) image to enable computation of LA with the color threshold tool (Fig. 2). Lastly, CVI was calculated as the ratio of LA to TCA. CVI assessment was performed by the same physician who measured CT (S.T.).

Statistical Analysis

Statistical analysis was performed using SPSS software version 16.0 (SPSS, Inc., Chicago, IL, USA). The normality of data distributions was tested using Shapiro–Wilk test. Continuous variables were shown as median (min–max). Nominal data were analyzed by Pearson's chi-square or Fisher's exact test as appropriate. Differences between values in the three groups were analyzed using Kruskal-Wallis test. Mann–Whitney U test was used for pairwise comparisons of the groups. The correlation between SFCT and migraine variables was evaluated using Spearman's correlation coefficient. Statistical significance was defined as $p < 0.05$.

Results

Thirty-six right eyes of 36 participants in the migraine group (30 women, 6 men) and 36 right eyes of 36 participants in the control group (30 women, 6 men) were included in the study. The mean ages in the migraine and control groups were 34.7 ± 1.5 years (range 20-52) and 35.1 ± 1.4 years (range 23-51), respectively ($p=0.868$). The groups were also similar in gender distribution. The migraine group included 5 patients who experienced migraine with aura and 31 patients with migraine without aura. BCVA was 0.00 logMAR (20/20 Snellen equivalent) in all eyes. The mean refractive error was -0.07 ± 0.72 D (range -2.25 to $+1.50$ D) in the migraine group and -0.15 ± 0.74 D (range -2.50 to $+1.50$ D) in the control group (Table 1).

Table1. Demographic data of patients and controls

Characteristics	Migraine group (n= 36)	Control group (n= 36)	<i>P</i> value
Age, years	34.7 ± 1.5	35.1 ± 1.4	0.868*
Gender, n (male/female)	6/30	6/30	
Refractive error, D	-0.07 ± 0.72	-0.15 ± 0.74	0.677**
VAS score (range)	8.17 ± 0.33 (1-10)		
Attacks per month (range)	5.60 ± 3.60 (1-14)		

* Student's t-test

** Mann-Whitney U test

VAS: Visual analogue scale

SFCT was 300.52 ± 88.30 μm in the migraine group and 262.85 ± 70.68 μm in the control group. There were no significant differences between the groups in SFCT or CT 1000 μm temporal and nasal of the fovea ($p>0.05$). The mean CVI was $71.8\% \pm 6.2\%$ in the migraine group and $70.7\% \pm 5.3\%$ in the control group ($p>0.05$). Mean CVI, TCA, LA, and SA did not differ significantly between the migraine and control groups (Table 2). There was also no significant difference in CVI between migraine patients with and without aura ($p>0.05$) (Table 3).

Table 2
Comparison of choroidal parameters between patients with migraine and the control group

Parameter	Migraine group mean \pm SD	Control group mean \pm SD	<i>P</i> value
Subfoveal CT (μm)	300.52 \pm 88.30	262.85 \pm 70.68	0.107**
N1000 CT (μm)	295.44 \pm 91.25	263.84 \pm 78.54	0.205**
T1000 CT (μm)	289.33 \pm 93.48	263.92 \pm 66.75	0.333**
Total Choroidal Area (mm^2)	0.715 \pm 0.219	0.689 \pm 0.183	0.580*
Luminal Area (mm^2)	0.516 \pm 0.170	0.486 \pm 0.130	0.395*
Stromal Area (mm^2)	0.199 \pm 0.067	0.203 \pm 0.069	0.964**
CVI (%)	71.8 \pm 6.2	70.7 \pm 5.3	0.316**
* Student's t-test			
** Mann-Whitney U test			
CT: Choroidal thickness, N1000: 1000 μm nasal of fovea, T1000: 1000 μm temporal of fovea, CVI: Choroidal vascularity index, VAS: Visual analogue scale			

Table 3
Comparison of choroidal parameters in migraine patients with and without aura

	With aura (n=5)	Without aura (n=31)	<i>P</i> value
Total Choroidal Area (mm^2)	0.778 \pm 0.20	0.704 \pm 0.22	0.481*
Luminal Area (mm^2)	0.584 \pm 0.18	0.505 \pm 0.16	0.413*
Stromal Area (mm^2)	0.193 \pm 0.02	0.199 \pm 0.07	0.765*
CVI (%)	73.9 \pm 7.5	71.5 \pm 6	0.101**
* Student's t-test			
** Mann-Whitney U test			
CVI: Choroidal vascularity index			

The mean VAS pain score in the migraine group was 8.17 ± 0.33 (range 1-10). VAS score was not correlated with SFCT or CVI ($r=0$, $p=0.998$ and $r=-0.06$, $p=0.731$, respectively). The mean monthly migraine frequency was 5.60 ± 3.60 and there was also no correlation between migraine frequency and SFCT or CVI ($r=-0.17$, $p=0.328$ and $r=-0.06$, $p=0.731$, respectively) (Table 4).

Table 4
Correlation between VAS score, monthly migraine frequency, choroidal thickness, and CVI in migraine patients

	VAS score (range 1-10)		Attacks per month (range 1-14)	
	r	p	r	p
Subfoveal CT (μm)	0	0.998	-0.17	0.328
N1000 CT (μm)	0.06	0.733	-0.125	0.476
T1000 CT (μm)	-0.039	0.826	-0.128	0.465
Total Choroidal Area (mm^2)	0.183	0.293	-0.046	0.792
Luminal Area (mm^2)	0.152	0.382	0.044	0.800
Stromal Area (mm^2)	0.328	0.054	0.017	0.923
CVI (%)	-0.06	0.731	0.04	0.820

CT: Choroidal thickness, N1000: 1000 μm nasal of fovea, T1000: 1000 μm temporal of fovea, CVI: Choroidal vascularity index, VAS: Visual analogue scale

Discussion

In this study, we observed no significant differences in SFCT and CVI between chronic migraine patients during an attack-free period when compared with the control group. Zengin et al. found that the mean CT was thinner in newly diagnosed (at least 3 months) migraine patients than in the control group ($p=0.001$). In the same study, they determined that CT decreased significantly in 5 patients during a migraine attack. [12] Reggio et al. also reported that CT was thinner in chronic migraine patients with and without aura compared to the control group ($p<0.0001$ for both eyes). [14] Likewise, Karaca et al. found that SFCT was thinner in the attack-free period in migraine patients with and without aura compared to the control group ($p<0.05$). However, CT was similar at all measured points between the migraine subgroups with and without aura ($p>0.05$). [15] Contrary to these studies, Gunes et al. found that CT was thicker in chronic migraine patients compared to the control group ($p<0.001$ for both eyes). [16] In their literature review, Ascado et al. summarized the different CT results obtained in migraine patients. [24] Therefore, a different parameter is needed due to the variable nature of CT. In our study, we determined that there was no statistically significant difference in CT and CVI between migraine patients and the normal healthy group. To the best of our knowledge, this is the first study evaluating the CVI in patients with chronic migraine.

Temel et al. investigated CVI in newly diagnosed migraine patients and reported that CVI was significantly decreased in patients with migraine. [25] However, limitations of their study are that it included a relatively small sample size and they only excluded patients who used ergot alkaloids and triptans within 24 hours before examination. Ergot alkaloids and triptans induce arterial and venous vasoconstriction, [26, 27] and

the long-term effects of these drugs on CT are unknown. Therefore, the decrease in CVI may be related to the use of these drugs.

Many studies have shown that CT is affected by various factors, especially certain drugs such as sildenafil and antihistamines, smoking, age, and the axial length of the eye.[28–31] While choroidal thinning is seen in choroidal dystrophies and AMD, thickening occurs in diseases such as Vogt-Koyanagi-Harada, central serous retinopathy, and polypoidal choroidal vasculopathy.[32–34] In addition, CT measurements may differ due to examiner bias and interobserver variation.[19, 35] On the other hand, CVI gives more reliable information than CT because LA (vascular), SA (interstitial), and TCA are determined from EDI-OCT images by special software using the binarization method.[18, 19] The rich vascular structure and changes in the connective tissue can be examined in more detail, providing more reliable data about the choroidal structure. For this reason, CVI is increasingly used instead of assessing choroidal structure only by its thickness.

EDI-OCT is a non-invasive method that enables detailed visualization of the choroid.[10] Although ICGA is still considered the gold standard imaging modality in choroidal pathologies such as PCV, its clinical use is declining because of its invasiveness.[36] With EDI-OCT, however, the effects of intraocular pressure and perfusion changes on the choroid can be assessed instantly and non-invasively. OCTA imaging has also seen more widespread clinical use in recent years because it allows non-invasive visualization of retinal and choroidal blood flow. This method uses special software to detect the movement of red blood cells in the vasculature and display vascular flow.[11] Guler et al. used OCTA to examine differences in retinal, peripapillary, and choriocapillaris blood flow between 26 patients with migraine without aura and a healthy control group. They observed no significant difference between the two groups in terms of blood flow in the superficial or deep retina, choriocapillaris, or choroid (choroidal flow area was 9.64 ± 0.44 and 9.65 ± 0.21 mm² in the migraine and control groups, respectively, $p=0.495$).[37] Ozcift et al. examined optic disc perfusion, central macular perfusion, and central CT in 38 chronic migraine patients and reported no significant difference in perfusions or CT, although CT was negatively correlated with the duration of migraine disease ($r=-0.46$, $p=0.004$).[38]

OCT is excellent for visualizing the retinal and choroidal anatomy but provides no information about the vasculature or circulation.[39] The fact that we detected no statistically significant difference in CVI values between the two groups in our study is consistent with previous studies indicating no change in choroidal flow on OCTA. The inconsistency between OCTA and CT studies may also be due to the relative subjectivity of CT measurement. According to Guler et al., there was no significant difference in retinal and choroidal blood flow in migraine patients, and retinal blood flow was determined by the dynamics of the vascular microenvironment.[37] Rather than CT, more OCTA and CVI data are needed to explain pathophysiological mechanisms, especially in a disease of unclear pathophysiology such as migraine.

Finally, we determined that mean VAS score (8.17 ± 0.33) and monthly attack frequency in the migraine group were not significantly correlated with CT or CVI. This is consistent with the results reported by Zengin et al., who observed no significant relationship between mean VAS score (5.55 ± 2.93) and CT.[12]

However, Karaca et al. investigated the relationship between CT and VAS score, Migraine Disability Assessment Score, and Wong-Baker faces pain rating scale score and determined that CT moderately correlated with VAS score and Wong-Baker scores in patients with migraine without aura but not in patients with migraines with aura.[15] In this regard, it is clear that more studies are needed to understand the correlation between CT and different pain scores and migraine frequency.

The present study has some limitations. One important limitation of our study is that changes in CVI were not evaluated during migraine attacks. Different results may be obtained during a migraine attack due to the activation of different pathophysiological mechanisms. Another limitation is the small number of patients included in the study. In addition, as in all CT studies, the manual determination of CT in our study is a limitation because manual segmentation remains a potential source of bias. Software-based automatic determination of CT is needed to eliminate this problem.

In conclusion, the results of this study suggest that CT and CVI do not differ significantly in chronic migraine patients during an attack-free period compared to healthy controls. However, considering the complex pathophysiology of migraine disease, more studies are needed to understand the relationship between migraine and CT.

Declarations

Acknowledgments: None

Statement and Declarations

Compliance with ethical standards

Conflict of interest Author Taha Sezer declares that he has no conflict of interest. Author Alper Aziz Hüdai Ayaslı declares that she has no conflict of interest. Author Bayram Meydan declares that he has no conflict of interest.

Ethical approval All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki Declaration and its later amendments or comparable ethical standards. This study was approved by Duzce University Institutional Review Board.

Informed consent Informed consent was obtained from all individual participants included in the study.

References

1. Tepper SJ, Rapoport A, Sheftell F (2001) The pathophysiology of migraine. *Neurologist* 7(5):279–286

2. Rasmussen BK, Olesen J (1992) Migraine with aura and migraine without aura: an epidemiological study. *Cephalalgia: an international journal of headache* 12(4):221–228. discussion 186.
3. Panconesi A, Bartolozzi ML, Guidi L (2009) Migraine pain: reflections against vasodilatation. *J Headache Pain* 10(5):317–325
4. Cole AJ, Aubé M (1990) Migraine with vasospasm and delayed intracerebral hemorrhage. *Arch Neurol* 47(1):53–56
5. De Smet Y, Brucher JM (1991) Migraine with vasospasm and delayed intracerebral hemorrhage. *Arch Neurol* 48(1):17
6. Goadsby PJ, Holland PR, Martins-Oliveira M, Hoffmann J, Schankin C, Akerman S (2017) Pathophysiology of Migraine: A Disorder of Sensory Processing. *Physiol Rev* 97(2):553–622
7. Delaey C, Van De Voorde J (2000) Regulatory mechanisms in the retinal and choroidal circulation. *Ophthalmic Res* 32(6):249–256
8. Brown JS, Flitcroft DI, Ying GS, Francis EL, Schmid GF, Quinn GE et al (2009) In vivo human choroidal thickness measurements: evidence for diurnal fluctuations. *Invest Ophthalmol Vis Sci* 50(1):5–12
9. Freeman WR, Bartsch D-U, Mueller AJ, Banker AS, Weinreb RN (1998) Simultaneous Indocyanine Green and Fluorescein Angiography Using a Confocal Scanning Laser Ophthalmoscope. *Arch Ophthalmol* 116(4):455–463
10. Sezer T, Altınışik M, Koytak İA, Özdemir MH (2016) The Choroid and Optical Coherence Tomography. *Turk J Ophthalmol* 46(1):30–37
11. Spaide RF, Fujimoto JG, Waheed NK, Sadda SR, Staurengi G (2018) Optical coherence tomography angiography. *Prog Retin Eye Res* 64:1–55
12. Zengin MO, Elmas Z, Cinar E, Kucukerdonmez C (2015) Choroidal thickness changes in patients with migraine. *Acta Neurol Belg* 115(1):33–37
13. Karalezli A, Celik G, Koktekir BE, Kucukerdonmez C (2015) Evaluation of choroidal thickness using spectral-domain optical coherence tomography in patients with migraine: a comparative study. *Eur J Ophthalmol* 25(4):338–342
14. Reggio E, Chisari CG, Ferrigno G, Patti F, Donzuso G, Sciacca G et al (2017) Migraine causes retinal and choroidal structural changes: evaluation with ocular coherence tomography. *J Neurol* 264(3):494–502
15. Karaca EE, Koçer EB, Özdek Ş, Akçam HT, Ercan MB (2016) Choroidal thickness measurements in migraine patients during attack-free period. *Neurol Sci* 37(1):81–88
16. Gunes A, Karadag AS, Yazgan S, Celik HU, Simsek A (2018) Evaluation of retinal nerve fibre layer, ganglion cell layer and choroidal thickness with optical coherence tomography in migraine patients: a case-control study. *Clin Exp Optom* 101(1):109–115
17. Dervisogullari MS, Totan Y, Gençler OS (2015) Choroid thickness and ocular pulse amplitude in migraine during attack. *Eye* 29(3):371–375

18. Agrawal R, Gupta P, Tan K-A, Cheung CMG, Wong T-Y, Cheng C-Y (2016) Choroidal vascularity index as a measure of vascular status of the choroid: Measurements in healthy eyes from a population-based study. *Sci Rep* 6:21090
19. Breher K, Terry L, Bower T, Wahl S (2020) Choroidal Biomarkers: A Repeatability and Topographical Comparison of Choroidal Thickness and Choroidal Vascularity Index in Healthy Eyes. *Translational Vision Science & Technology* 9(11):8
20. Velaga SB, Nittala MG, Vupparaboina KK, Jana S, Chhablani J, Haines J et al (2020) Choroidal vascularity index and choroidal thickness in eyes with reticular pseudodrusen. *Retina (Philadelphia, Pa)* 40(4):612–617
21. The International Classification of Headache Disorders: 2nd edition. *Cephalalgia: an international journal of headache* (2004) ;24 Suppl 1:9-160
22. Tan CS, Ouyang Y, Ruiz H, Sadda SR (2012) Diurnal Variation of Choroidal Thickness in Normal, Healthy Subjects Measured by Spectral Domain Optical Coherence Tomography. *Investig Ophthalmol Vis Sci* 53(1):261–266
23. Sonoda S, Sakamoto T, Yamashita T, Uchino E, Kawano H, Yoshihara N et al (2015) Luminal and stromal areas of choroid determined by binarization method of optical coherence tomographic images. *Am J Ophthalmol* 159(6):1123–31e1
24. Ascaso FJ, Marco S, Mateo J, Martínez M, Esteban O, Grzybowski A (2017) Optical Coherence Tomography in Patients with Chronic Migraine: Literature Review and Update. *Frontiers in Neurology.* ; 8(684)
25. Temel E, Aşıkgarip N, Koçak Y, Şahin C, Özcan G, Kocamiş Ö et al (2021) Choroidal vascularity index and retinal nerve fiber layer reflectivity in newly diagnosed migraine patients. *Photodiagnosis Photodyn Ther* 36:102531
26. Schmetterer L, Wolzt M, Krejcy K, Graselli U, Findl O, Eichler HG et al (1996) Cerebral and ocular hemodynamic effects of sumatriptan in the nitroglycerin headache model. *Clin Pharmacol Ther* 60(2):199–205
27. Tepper SJ, Rapoport AM, Sheftell FD (2002) Mechanisms of Action of the 5-HT_{1B/1D} Receptor Agonists. *Arch Neurol* 59(7):1084–1088
28. Vance SK, Imamura Y, Freund KB (2011) The effects of sildenafil citrate on choroidal thickness as determined by enhanced depth imaging optical coherence tomography. *Retina (Philadelphia, Pa)* 31(2):332–335
29. Sizmaz S, Küçükerdönmez C, Pinarci EY, Karalezli A, Canan H, Yilmaz G (2013) The effect of smoking on choroidal thickness measured by optical coherence tomography. *Br J Ophthalmol* 97(5):601–604
30. Ikuno Y, Kawaguchi K, Nouchi T, Yasuno Y (2010) Choroidal thickness in healthy Japanese subjects. *Invest Ophthalmol Vis Sci* 51(4):2173–2176
31. Hirata M, Tsujikawa A, Matsumoto A, Hangai M, Ooto S, Yamashiro K et al (2011) Macular Choroidal Thickness and Volume in Normal Subjects Measured by Swept-Source Optical Coherence

Tomography. *Investig Ophthalmol Vis Sci* 52(8):4971–4978

32. Chung SE, Kang SW, Lee JH, Kim YT (2011) Choroidal thickness in polypoidal choroidal vasculopathy and exudative age-related macular degeneration. *Ophthalmology* 118(5):840–845
33. Maruko I, Iida T, Sugano Y, Oyamada H, Sekiryu T, Fujiwara T et al (2011) Subfoveal choroidal thickness after treatment of Vogt-Koyanagi-Harada disease. *Retina (Philadelphia, Pa)* 31(3):510–517
34. Maruko I, Iida T, Sugano Y, Ojima A, Sekiryu T (2011) Subfoveal choroidal thickness in fellow eyes of patients with central serous chorioretinopathy. *Retina (Philadelphia, Pa)* 31(8):1603–1608
35. Yamashita T, Yamashita T, Shirasawa M, Arimura N, Terasaki H, Sakamoto T (2012) Repeatability and Reproducibility of Subfoveal Choroidal Thickness in Normal Eyes of Japanese Using Different SD-OCT Devices. *Investig Ophthalmol Vis Sci* 53(3):1102–1107
36. Cheung CMG, Lai TYY, Ruamviboonsuk P, Chen SJ, Chen Y, Freund KB et al (2018) Polypoidal Choroidal Vasculopathy: Definition, Pathogenesis, Diagnosis, and Management. *Ophthalmology* 125(5):708–724
37. Güler Ö, Güler M, Tuğan Yıldız CB, Hakkoymaz H (2020) Are Retinal and Peripapillary Blood Flows Affected during Migraine Attack? *Neuro-ophthalmology (Aeolus Press)*. 44:299–3065
38. Gürakar Özçift S, Aydın E, Eriş E (2021) Assessment of the choroidal thickness, central macular vascular and optic disk perfusion in migraine patients with optical coherence tomography angiography. *Photodiagnosis Photodyn Ther* 35:102397
39. Savastano MC, Lumbroso B, Rispoli M (2015) In vivo characterization of retinal vascularization morphology using optical coherence tomography angiography. *Retina (Philadelphia, Pa)* 35(11):2196–2203

Figures




Figure 1

A width of 1.5 mm centered at the fovea selected by manual plotting polygonal tool.




Figure 2

Superimposed binarized image showing segmentation of the choroidal luminal and stromal structures.