Intraoperative mapping of pre-central motor cortex and subcortex: a proposal for supplemental cortical and novel subcortical maps to Penfield's motor homunculus

Prajwal Ghimire (✉ prajwal.ghimire@kcl.ac.uk)

King's College Hospital NHS Foundation Trust <https://orcid.org/0000-0003-2884-956X>

Jose Lavrador

King's College Hospital NHS Foundation Trust

Asfand Mirza

King's College Hospital NHS Foundation Trust

Noemia Pereira

Inomed Neurocare UK Ltd

Hannah Keeble

Inomed Neurocare UK Ltd.

Marco Borri

King's College Hospital NHS Foundation Trust

Luciano Furlanetti

King's College Hospital NHS Foundation Trust

Christian Brogna

King's College Hospital NHS Foundation Trust

Jozef Jarosz

King's College Hospital NHS Foundation Trust

Richard Gullan

King's College Hospital NHS Foundation Trust

Francesco Vergani

King's College Hospital NHS Foundation Trust

Ranj Bhangoo

King Colege London

Keyoumars Ashkan

King's College Hospital NHS Foundation Trust

Keywords: Motor homunculus, subcortical map, craniotomy, corticospinal tract, motor cortex

Posted Date: December 11th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-115348/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Intraoperative mapping of pre-central motor cortex and subcortex: a proposal for supplemental cortical and novel subcortical maps to Penfield's motor homunculus

Prajwal Ghimire^{1*}, Jose Pedro Lavrador¹, Asfand Baig Mirza¹, Noemia Pereira³, Hannah Keeble³, Marco Borri², Luciano Furlanetti¹, Christian Brogna¹, Jozef Jarosz², Richard Gullan¹, Francesco Vergani¹, Ranjeev Bhangoo¹, Keyoumars Ashkan¹

1. Department of Neurosurgery, King's College Hospital NHS Foundation Trust, London, UK
2. Department of Neuroradiology, King's College Hospital NHS Foundation Trust, London, UK
3. Neuromonitoring Team, Inomed Neurocare UK Ltd., UK

Corresponding Author:

Mr Prajwal Ghimire
MBBS MRCSEd. MSc
Department of Neurosurgery
King's College Hospital NHS Foundation Trust
London, UK

Word Count:

Abstract: 250
Manuscript: 2500 (excluding references, figures, tables, acknowledgements)
References: 40

Abstract:

Introduction:

Penfield's motor homunculus describes a caricaturised yet useful representation of the map of various body parts on the pre-central cortex. We propose a supplemental map of the clinically represented areas of human body in pre-central cortex and a novel subcortical corticospinal tract map that are accurate and essential for safe surgery in patients with eloquent brain lesions.

Materials and methods:

A single-institution retrospective cohort study of patients who underwent craniotomy for motor eloquent lesions with intraoperative motor neuromonitoring (cortical and subcortical) between 2015 and 2020 was performed. All positive cortical and subcortical stimulation points were taken into account and cartographic maps were produced to demonstrate cortical and subcortical areas of motor representation and their configuration. A literature review in PubMed was performed.

Results:

180 patients (58.4% male, 41.6% female) were included in the study with 81.6% asleep and 18.4% awake craniotomies for motor eloquent lesions (gliomas 80.7%, metastases 13.8%) with intraoperative cortical and subcortical motor mapping. Based on the data, we propose a supplemental clinical cortical and a novel subcortical motor map to the original Penfield's motor homunculus, including demonstration of localisation of intercostal muscles both in the cortex and subcortex which has not been previously described.

Conclusion:

The supplementary clinical cortical and novel subcortical motor maps of the homunculus presented here have been derived from a large cohort of patients undergoing direct cortical and subcortical brain mapping. The information will have direct relevance for improving the safety and outcome of patients undergoing resection of motor eloquent brain lesions.

Key Words:

Motor homunculus, subcortical map, craniotomy, corticospinal tract, motor cortex

Introduction:

Homunculus as described by Penfield and Boldrey in 1937 after bipolar direct cortical stimulation (DCS) in 126 awake patients has provided the foundation for intraoperative motor mapping in patients undergoing craniotomy for brain lesions¹. Since then, there has been a range of studies on motor mapping of corticospinal tract (CST) using neuro-physiological, viral tracing, microstructural, cadaveric and intraoperative human and animal studies^{2,3,4,5}. More recently, there has been further studies using advanced imaging techniques such as deterministic as well as probabilistic tractography to produce a map of the CST, including U fibre connections, highlighting the complex fibre connections and overlap^{6,7}. Human connectome project has further provided an advanced interface to characterise the configuration of cortical and subcortical CST⁸. Difference in the size of representation of body-parts within the primary motor cortex has been recognized and is thought to be likely related to each regions' functional specialisation and fine motor functions^{1,7,9}.

Advancements in the DCS and subcortical mapping techniques (monopolar stimulation, dynamic continuous monitoring), allow accurate measurement of the distance from CST during intraoperative stimulation to achieve safe lesion resection and avoid inadvertent injury to CST central to preserving the quality of life of the patients^{10,11,12}. Bello et al described subcortical mapping in 57 patients where individual areas of hand, arm and leg were identified during resection of gliomas and accurate identification of CST enhanced surgical performance and safety maintaining a high rate of functional preservation¹³. Increasingly, a combination of pre-operative (functional magnetic resonance imaging(fMRI), diffusion tensor imaging(DTI), navigated transcranial magnetic stimulation(nTMS)) and intra-operative (neuronavigation, intra-operative ultrasound (ioUS), augmented reality microscope, 5-aminolevulinic acid (5-ALA)) tools are combined with direct intra-operative brain mapping to aid surgical planning for maximal safe resection^{10,11,12,13,14,15}. Despite these studies and the data gathered, no updated homunculus nor subcortical map, based on intra-operative direct cortical/ subcortical stimulation, has been proposed since the original 1937 work (**Supplemental Table 1**).

In this study, we propose a supplemental motor cortical map and a novel motor subcortical map of the configuration of the corticospinal tract initially described by Penfield and Boldrey^{1,9} that gives an insight into the cortical and subcortical representation of the body parts; critical for surgery in eloquent brain and for better clinical outcomes.

Materials and Methods:

A single-centre retrospective study was performed between January 2015 and January 2020 to collect intraoperative data on 180 patients who underwent craniotomy for eloquent brain lesions with intraoperative cortical and subcortical motor neuromonitoring at our quaternary referral neurosurgical centre. Patients were consented for intraoperative neuromonitoring including motor mapping. Prior to the surgery, all patients underwent a range of pre-operative brain mapping investigations to include fMRI, DTI and nTMS to evaluate the feasibility of lesion resection, determine the best surgical approach and help consent the patient based on individual's risk profile. At surgery, 3 dimensional (3D) reconstructed structural MRI, tractography from DTI, fMRI hotspots and 3D reconstructed nTMS motor stimulation points were projected onto the brain with augmented reality using ZEISS KINEVO® 900 microscope (CARL ZEISS Meditec AG, Jena, Germany) for the patients. These helped as starting points to define the relationship between the lesions and eloquent motor brain areas, guiding intra-operative mapping with cortical/ subcortical stimulation. Thereafter, all positive documented cortical and subcortical points of stimulation confirmed with intraoperative neuromonitoring were recorded and plotted over cartographical maps by utilizing the intraoperative numerically labelled pictures and correlating with intra-operative neuro-navigation MRI, corrected for intra-operative brain shift using the ioUS, and immediate post-operative anatomical T1 post gadolinium MRI. The overlap of stimulated points were also noted. Demographic, clinical and surgical data were collected from patients' medical records. A PubMed literature review on subcortical motor mapping was performed to provide insights into the current position on the subcortical motor representation of human body in the literature. The demographic and clinical data were analysed with Microsoft Excel 2020.

Functional Magnetic Resonance Imaging (fMRI)

Three different tasks were performed: lip smacking, finger-tapping and foot rocking (**Supplemental Figure 1**). The cortical activated areas were used to constrain the probabilistic tractography.

Diffusion tensor imaging (DTI) and deterministic tractography

DTI sequences were obtained to reconstruct CST. The parameters utilised for obtaining diffusion tensor imaging were b-value: 1,500; diffusion directions: 64; diffusion mode; multi-

directional diffusion weighting; field of view: 32 cm; voxel size, 2.5*2.5*2.5 mm; TR/TE: 9,500/86; scan time: 11:35 minutes. The deterministic tractography was modelled in 3D with StealthViz Medtronic Software (Minneapolis, Minnesota, USA)¹². The dissection of the corticospinal tract was performed according to the regions of interest (ROIs) between the motor cortex and the ipsilateral half of the medulla oblongata below the middle cerebellar peduncle (**Figure 1**). Constrained probabilistic tractography for complex motor function and cortical/subcortical CST was performed with MRTrix3 opensource Software¹⁶ (**Supplemental Figure 2**).

Navigated transcranial magnetic stimulation (nTMS)

Preoperative nTMS was utilised for pre-operative motor mapping, using Nexstim TMS, v.4.3.1 (Nexstim, Helsinki, Finland)¹⁷. nTMS was obtained with single-pulse sequence delivered using a figure-of-eight coil, and the motor mapping was performed at 105% of the determined resting motor threshold¹⁷. The nTMS preoperative positive motor responses were transformed in 3D objects (StealthViz Medtronic software, Minneapolis, Minnesota, USA) superimposed to the 3D tractography model of the CST on the Medtronic Stealth Station S7/S8 neuronavigation software/machine (Minneapolis, Minnesota, USA) (**Supplemental Figure 3**).

Cortical mapping

Cortical motor evoked potentials (cMEPs) were recorded with direct electrical monopolar stimulations of primary motor cortex (**Figure 2**). Train of five pulses, positive pulse form, inter-stimuli interval of 4.0 ms, pulse width of 0.5 ms, 1 Hz and anodal pole were the parameters for stimulations. Continuous cMEPs were monitored using a four-contact strip electrode positioned over the motor cortex recording stable MEPs at the motor threshold. Muscles monitored during intraoperative neuromonitoring were orbicularis oris, masseter, tongue, cricothyroid, deltoid, brachioradialis/flexor carpi ulnaris (BR/FCU), abductor pollicis brevis/abductor digiti minimi (APB/ADM), first dorsal interosseous (FDI), intercostals, quadriceps femoris, tibialis anterior, abductor hallucis.

Subcortical mapping

Subcortical motor evoked potentials (scMEPs) were recorded using modified monopolar suction probe and monopolar probe (Parameters were same as cMEP recording with cathodal pole)^{10,12} (**Figure 3**). Muscles monitored were same as on cortical mapping.

Intraoperative ultrasound

Ultrasound images were obtained intraoperatively prior to corticotomy and at the end of resection with Esaote Ultrasound Machine (Esaote, Genova, Italy) to delineate the margins of the tumour and post resection cavity. It provided the configuration, depth and location of the lesion with respect to the motor cortex and subcortical CST identified with subcortical stimulation and helped to correct for any intra-operative brain shift (**Supplemental Figure 4**).

Results:

Table 1 summarises the demographics. The motor threshold during stimulation represented the estimated distance from the clinically detected anatomical bundle of fibres of CST (1mA=1mm). Cortical motor threshold of DCT of motor cortex/CST was recorded as mean of 7.2 mA (Range:0.8 mA – 25mA, n=180). Cortical and subcortical mean stimulation threshold for leg (c8.4mA, range:2-17mA,n=14; sc5.8mA, range:3-13,n=11), foot (c7mA, range:5-12mA,n=17; sc6.5mA, range:2-14mA,n=12), intercostal muscles (c10mA, range:6-14mA,n=6; sc6mA, range:4-12mA,n=7), arm (c7.6mA, range:0.8-13mA,n=21; sc4.3mA, range:2-6mA,n=6), hand (c8.7mA, range:0.8-20mA,n=47; sc6.6mA, range:5-8.5mA,n=20) and face (c7.4 mA, range:2-11 mA, n=24; sc5.7 mA, range:0.5-8, n=12) were recorded during the procedures (**Table 2,Table 3**). The stimulated cortical and subcortical points were then plotted in the diagrammatic representation of cortex and subcortex with individual area stimulated. The points were then assembled into cartographic maps to demonstrate a clinical cortical and subcortical map of intraoperative corticospinal tract (**Figure 4, Figure 5**).

Cortical and subcortical stimulation of Intercostal Muscles

As a contribution to the previous described homunculus, we present, for the first time in the literature, a detailed representation at both cortical and subcortical level of the intercostal muscles—demonstrated during routine neurophysiological mapping and monitoring (**Supplemental Table 2**). An example of isolated subcortical stimulation recording (scMEP) during intraoperative subcortical mapping with electrode placed in the intercostal muscles is shown in **Supplemental Figure 5**. These findings aided in the illustration of the proposed maps.

Literature Review

A PubMed literature review was performed, in order to investigate if a subcortical motor map was previously described, with the MeSH items ((Subcortical stimulation) OR (subcortical mapping) OR (subcortical intraoperative neuromonitoring) AND Motor AND corticospinal tract)) with filters of “articles with abstracts, adult (19+ years), English language articles, human studies”. It resulted in identification of 141 articles. The articles were reviewed to identify data on subcortical mapping of specific motor areas of patients who underwent craniotomy for brain lesions (**Supplemental Table 1**). There was no article identified suggesting a map for the subcortical motor areas.

Proposed homunculus Maps

Proposal of Supplemental map of Cortical motor Region(Clinical cortical motor Homunculus)

The proposed supplemental cortical map consists of new addition of cortical representation of intercostal muscles in the existing Penfield's motor homunculus (**Figure 4**).

Proposal of map of intraoperative subcortical Corticospinal Tract (isCST) (Clinical Subcortical motor homunculus)

The proposed subcortical novel map of the corticospinal tract along the corona radiata (**Figure 5**) provides an insight into the configuration of CST fibres and highlights the significant differences between the cortical orientation and subcortical orientation of the CST fibres.

Discussion:

Understanding the topography of the motor pathway at both cortical and subcortical levels is crucial for neurosurgeons during surgery for motor eloquent brain lesions if risk of deficits is to be minimised. Penfield's 1937 description of the arrangement of the motor function onto the homunculus, although a simplified map, provides a representative image of the motor arrangement of function onto the precentral cortex¹. Since then, human clinical studies and cadaver anatomical studies have provided further insight into the complexity of the motor cortex and overlap of function^{3,7}. There has been recognition of dynamic, rather than the traditional static, representation of the body parts on the motor cortex, especially for complex motor movement^{18,19} as well as significant overlapping of motor representation in the supplementary motor area (SMA), premotor, primary motor and parietal cortex²⁰. Furthermore, contrary to the traditional orientation of body parts on motor cortex, multiple directions of the representation of have now been described: Lateral-anterior-ventral direction along the precentral gyrus and central sulcus, reflecting a shift from lower- to upper-body muscles²¹. Catani recently described a detailed reappraisal of the original Penfield's findings, suggesting overlaps between the transitional areas and cross over to the somatosensory cortex along with description of role of U fibres⁷. Non-human primate studies, using viral tracers, have added additional dimensions to this complexity by defining phylogenetically new M1 and old M1 cortex⁴. Despite the spectrum of the new knowledge and techniques, from the practical point of view, the original Penfield's motor homunculus remains central as a starting point when planning surgery for lesions in and around motor cortex, and DCS continues as the gold standard for intra-operative refinement. Thus, Duffau and colleagues, through DCS studies, were recently even able to demonstrate the concept of negative motor response whereby movement arrest was induced at specific stimulation points in the pre-central gyrus²². In this context, our supplementary motor homunculus presented here, provides a useful addition by clearly demonstrating the motor cortical area for the intercostal muscles, previously not described in detail. We first reported mapping of intercostal muscles as a technical note²³. This has now been replicated and mapped in detail in this paper (**Supplemental Table 2**). Understanding the cortical topography of intercostal muscles is important to avoid inadvertent damage during the surgery which could lead to paralysis of respiratory muscles and the well-recognised post-operative respiratory complications.

Unlike the motor cortex, our understanding of the detailed functional anatomy of the white matter motor tracts is at an earlier stage. Cadaveric anatomical and imaging studies have shown that corticospinal tract (CST) predominantly begins in the pre-central gyrus (M1), making its way caudally through the corona radiata into the internal capsule, condensing the fibres with a unique formation and thus producing a swirl-like configuration with the twist of the fibres subcortically^{24,25,26,27,28,29}. The CST undergoes a rotation whereby the most medially located fibres at the cortical level (topographically – lower limb) become posterior at the level of the internal capsule and lateral at the level of the cerebral peduncle^{24,29,30,31}. Primate studies have demonstrated the complexity of organisation of the CST fibres corresponding to the functional body areas and its dynamic nature³². Understanding the detailed anatomy of these tracts is therefore essential for safe surgery for lesions in or around the subcortical motor pathways. In the recent years, DTI has proved useful as a starting point to understand the relationship between the CST and the lesions with DTI based maps of subcortex emerging as useful tools for pre-operative planning^{6,29,33,34,35}. The accuracy of the image-based technology, however, remains less than what is required for confident safe surgery and intra-operative direct stimulation of subcortical motor pathways remain the gold standard. Despite the limited number of papers published on subcortical stimulation (**Supplemental Table 1**), accurate intraoperative identification of cortical and subcortical boundaries has been challenging with no clear subcortical map, along the same line as the cortical motor homunculus, has been described. We therefore aimed to address it using the data from our large cohort of patients as demonstrated in our novel subcortical map. In this illustration and based on our intra-operative direct stimulation of CST, the motor areas of leg shifted from medial to the posterior, motor areas of hand/forearm shifted from superior to the centre of the condensed bundle and the face representation shifted from lateral to the anterior aspect of the condensed bundle demonstrating the change in CST configuration during the descent of fibres within the corona radiata. There were further changes in configuration prior to reaching the internal capsule: the leg area shifted from posterior to the midline position with the face area shifting anteriorly and the arm area shifting posteriorly as progressing into the known configuration of fibres in the genu and posterior limb of internal capsule. The subcortical map thus generated, we hope will aid in a better understanding of the functional anatomy of the CST and aid in safe surgery for motor eloquent lesions.

As our understanding of the biology of brain lesions, particularly gliomas, evolve and concepts such as survivorship and preservation of quality of life, quite rightly, gain centre-stage, there

is an ever-increasing need for maximal safe resection. This remains a challenge for lesions in and around the motor pathways. Understanding the cortical and subcortical functional anatomy is therefore crucial and it is best derived from intra-operative direct stimulation of motor pathways. Our findings from a large cohort of patients presented as an updated motor homunculus and a novel subcortical map will aid neurosurgeons in their quest to achieve the best outcome possible for these patients.

Conclusion:

Penfield's motor homunculus represented a landmark in understating the functional anatomy of the motor cortex with real implications for surgery in motor eloquent brain. After an almost a century, however, there is now time for a reappraisal. Our data from a large cohort of patients undergoing modern intra-operative stimulation of motor pathways, the gold standard for brain mapping, generated a supplementary updated motor homunculus and a novel subcortical motor map. Our finding will aid surgical planning for lesions in or around the motor pathways, to reduce the risks and increase the extent of resection.. Further multicentre studies are required to validate the map towards routine utilisation in clinical practice.

Acknowledgements:

Authors would like to acknowledge pioneering work on direct cortical stimulation by Penfield and Boldrey. We are sitting on the shoulders of the giants producing this paper. Authors would like to acknowledge the support provided by the Department of Neurosurgery and Department of Neuroradiology at King's College Hospital, London, UK and the intraoperative neurophysiology Team at Inomed Neurocare UK, London, UK. This paper partly represents independent research on BRCMAP project funded by the NIHR- Wellcome Trust, King's Clinical Research Facility and the National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. We would like to thank PG, Ms Aashika Sharma Lamichhane and Mrs Pilar Camino Alcon for their work in hand painting, illustrating and software processing of Figure 4 and 5.

Author Contributions:

PG drafted the manuscript. ABM carried out the collection of the patient data. PG, JPL and KA participated in the design of the study and in drafting discussion. MB, JPL, JJ participated in obtaining advanced imaging including fMRI, DTI and its analytical modelling. NP, HK participated in performing intraoperative neuromonitoring and its data analysis. PG, JPL, LF, CB, FV, KA participated in obtaining and analysing intraoperative data. RB, RG, FV and KA conceived of the study and participated in its design and coordination and helped to draft the manuscript. All authors read and approved the final manuscript.

Funding: No funding was obtained for the study

Competing interests: The authors report no competing interests.

Supplementary Materials: Supplemental Table 1, 2; Supplemental Figure 1,2,3,4,5

References:

1. Penfield W, Boldrey E. Somatic motor and sensory representation in the cerebral cortex of man as studied by electrical stimulation. *Brain* 1937; 60: 389–443.
2. Foerster O. The motor cortex in man in the light of Hughlings Jackson's doctrines. *Brain* 1936; 59: 135–59.
3. Farrell DF, Burbank N, Lettich E, Ojemann GA. Individual variation in human motor-sensory (rolandic) cortex. *J Clin Neurophysiol* 2007; 24: 286–93.
4. Rathelot, JA, Strick PL. Subdivisions of primary motor cortex based on cortico-motoneuronal cells. *Proceedings of the National Academy of Sciences* 2009; 106(3):918-923.
5. Desmurget M, Sirigu A. Revealing humans' sensorimotor functions with electrical cortical stimulation. *Philos Trans R Soc Lond B Biol Sci* 2015; 370: 20140207.
6. Berman JI, Berger MS, Mukherjee P, Henry RG. Diffusion-tensor imaging-guided tracking of fibers of the pyramidal tract combined with intraoperative cortical stimulation mapping in patients with gliomas. *J Neurosurg* 2004 Jul;101(1):66-72.
7. Catani M. A little man of some importance. *Brain* 2017;140 (11):3055–3061.
8. Human Connectome Project. Available from: <<http://www.humanconnectomeproject.org>> [Accessed 31 July 2020]
9. Penfield W, Rasmussen R. The cerebral cortex of man. New York, NY: The Macmillan Company; 1950.
10. Seidel K, Beck J, Stieglitz L, Schucht P, Raabe A. The warning-sign hierarchy between quantitative subcortical motor mapping and continuous motor evoked potential monitoring during resection of supratentorial brain tumors. *J Neurosurg* 2013;118(2):287-296.
11. Schucht, P, Seidel K, Murek M, Stieglitz L, Urwyler N, Wiest R, et al. Low-threshold monopolar motor mapping for resection of lesions in motor eloquent areas in children and adolescents. *J Neurosurg Pediatr* 2014;13(5):572-578.
12. Lavrador JP, Ghimire P, Brogna C, et al. Pre- and Intraoperative Mapping for Tumors in the Primary Motor Cortex: Decision-Making Process in Surgical Resection. *J Neurol Surg A Cent Eur Neurosurg*. 2020;10.1055/s-0040-1709729. [published online ahead of print, 2020 May 21]

13. Bello L, Gambini A, Castellano A, Carrabba G, Acerbi F, Fava E, et al. Motor and language DTI Fiber Tracking combined with intraoperative subcortical mapping for surgical removal of gliomas. *Neuroimage* 2008 Jan 1;39(1):369-82.
14. Vassal F, Schneider F, Nuti C. Intraoperative use of diffusion tensor imaging-based tractography for resection of gliomas located near the pyramidal tract: comparison with subcortical stimulation mapping and contribution to surgical outcomes. *Br J Neurosurg* 2013 Oct;27(5):668-75.
15. Hervey-Jumper SL, Berger MS. Maximizing safe resection of low- and high-grade glioma. *J Neurooncol* 2016 Nov;130(2):269-282.
16. Tournier JD, Smith R, Raffelt D, Tabbara R, Dhollander T, Pietsch M, et al. MRtrix3: A fast, flexible and open software framework for medical image processing and visualisation. *NeuroImage* 2019;202:116137.
17. Jung J, Lavrador JP, Patel S, et al. First United Kingdom experience of navigated transcranial magnetic stimulation in preoperative mapping of brain tumors. *World Neurosurg* 2019;122:e1578–e1587.
18. Perez MA, Rothwell JC. Distinct influence of hand posture on cortical activity during human grasping. *J Neurosci.* 2015;35(12):4882-4889.
19. Schellekens W, Petridou N, Ramsey NF. Detailed somatotopy in primary motor and somatosensory cortex revealed by Gaussian population receptive fields. *Neuroimage* 2018;179:337-347.
20. Cunningham DA, Machado A, Yue GH, Carey JR, Plow EB. Functional somatotopy revealed across multiple cortical regions using a model of complex motor task. *Brain Res.* 2013;1531:25-36.
21. Verstynen T, Sabes PN. How Each Movement Changes the Next: An Experimental and Theoretical Study of Fast Adaptive Priors in Reaching. *Journal of Neuroscience* 2011;31(27):10050-10059
22. Rech F, Herbet G, Gaudeau Y, et al. A probabilistic map of negative motor areas of the upper limb and face: a brain stimulation study. *Brain* 2019;142(4):952-965.
23. Ghimire P, Bedussa N, Giamouriadis A et al. Somatotopy of corticospinal tract fibres of the intercostal muscles: report of a case and review of literature. *Brain Struct Funct* 2019;224 (2):515-520.
24. Ebeling U, Reulen HJ. Subcortical topography and proportions of the pyramidal tract. *Acta Neurochir (Wien)* 1992;118:164–171.

25. Kim JS, Pope A. Somatotopically located motor fibers in corona radiata: evidence from subcortical small infarcts. *Neurology*. 2005; 64: 1438–1440.
26. Yamada K, Kizu O, Kubota T, Ito H, Matsushima S, Oouchi H, Nishimura T. The pyramidal tract has a predictable course through the centrum semiovale: a diffusion–tensor based tractography study. *J Magn Resonan Imaging* 2007; 26: 519–524.
27. Verstynen T, Jarbo K, Pathak S, Schneider W. In vivo mapping of microstructural somatotopies in the human corticospinal pathways. *J Neurophysiol*. 2011;105(1):336–346.
28. Zolal A, Vachata P, Hejčl A, et al. Anatomy of the supraventricular portion of the pyramidal tract. *Acta Neurochir (Wien)* 2012;154(6):1097–1104.
29. Chenot Q, Tzourio-Mazoyer N, Rheault F, Descoteaux M, Crivello F, Zago L, et al. A population-based atlas of the human pyramidal tract in 410 healthy participants. *Brain Struct Funct* 2019 Mar;224(2):599–612.
30. Englander RN, Netsky MG, Adelman LS. Location of human pyramidal tract in the internal capsule: anatomic evidence. *Neurology* 1975;25:823–826.
31. Catani M, Thiebaut de Schotten M. A diffusion tensor imaging tractography atlas for virtual in vivo dissections. *Cortex* 2008;44:1105–1132.
32. Graziano MS, Taylor CS, Moore T. Complex movements evoked by microstimulation of precentral cortex. *Neuron* 2002; 34: 841–51.
33. Okada T, Mikuni N, Miki Y, Kikuta K, Urayama S, Hanakawa T, et al. Corticospinal tract localization: integration of diffusion-tensor tractography at 3-T MR imaging with intraoperative white matter stimulation mapping--preliminary results. *Radiology* 2006 Sep;240(3):849–57.
34. Rosenstock T, Giampiccolo D, Schneider H, Runge SJ, Bährend I, Vajkoczy P, Picht T. Specific DTI seeding and diffusivity-analysis improve the quality and prognostic value of TMS-based deterministic DTI of the pyramidal tract. *Neuroimage Clin* 2017;16:276–285.
35. Oda K, Yamaguchi F, Enomoto H, Higuchi T, Morita A. Prediction of recovery from supplementary motor area syndrome after brain tumor surgery: preoperative diffusion tensor tractography analysis and postoperative neurological clinical course. *Neurosurg Focus* 2018;44 (6):E3.
36. Cignoni P, Callieri M, Corsini M, Dellepiane M, Ganovelli F, Ranzuglia G. MeshLab: an Open-Source Mesh Processing Tool. *Sixth Eurographics Italian Chapter Conference* 2008:129–136.

37. Fan L, Li H, Zhuo J, Zhang Y, Wang J, Chen L, et al. The Human Brainnetome Atlas: A New Brain Atlas Based on Connectional Architecture. *Cerebral Cortex* 2016. doi 10.1093/cercor/bhw157
38. Duffau H, Capelle L, Denvil D, Sichez N, Gatignol P, Taillandier L, et al. Usefulness of intraoperative electrical subcortical mapping during surgery for low-grade gliomas located within eloquent brain regions: functional results in a consecutive series of 103 patients. *J Neurosurg* 2003;98:764–778.
39. Mikuni N, Okada T, Enatsu R, Miki Y, Hanakawa T, Urayama S, et al. Clinical impact of integrated functional neuronavigation and subcortical electrical stimulation to preserve motor function during resection of brain tumors. *J Neurosurg* 2007 Apr;106(4):593-8.
40. Bello L, Riva M, Fava E, Ferpozzi V, Castellano A, Raneri F, et al. Tailoring neurophysiological strategies with clinical context enhances resection and safety and expands indications in gliomas involving motor pathways. *Neuro Oncol* 2014 Aug;16(8):1110-28.

Captions:

Tables

Table 1: Demographics of the patient data

Table 2: Cortical Stimulation of motor cortex areas with their thresholds

Table 3: Subcortical Stimulation of motor subcortex areas with their thresholds

Supplemental Table 1: Literature Review of Subcortical intraoperative mapping demonstrating each mapped area

Supplemental Table 2: Case series of cortical and subcortical intercostal muscles Stimulation (MEP)

Figures:

Figure 1(a,b,c,d): Pre-operative 3D reconstruction of Corticospinal tract (*blue, green*) with tumour (*red*) utilising the pre-operative DTI and modelling using neuronavigation (Medtronic Stealth S7, StealthViz Software) (*70y female with left precentral metastatic tumour (primary lung small cell carcinoma who underwent gross total resection)*)

Figure 2: Cortical stimulation with monopolar probe stimulating different areas of motor cortex **(a)** 1,2,3: hand and forearm 4,5: foot and leg **(b)** 6,7,8,9,10,11,13,14- hand knob (ADM, FDI, APB, forearm, deltoid) 1,2,5- post central gyrus **(c)** 1,2: intercostal muscles 3,4: deltoid 5: foot **(d)** 1,2: arm, forearm, hand 3,4,5: hand, face 6,7: face

Figure 3: Subcortical stimulation with intraoperative MEPs demonstrating subcortical mapping of the corticospinal tract **(a,b)** with contact numbering corresponding to the mapping/stimulation positive areas with motor thresholds **(1-15)**; **1-** upper limb (8mA), face (5mA); **2-** upper limb (8mA), face (6mA); **3-** upper limb (7mA), face (2mA); **4-** upper limb (5mA), face (2mA); **5-** upper limb (7mA), face (4mA); **6-** face (2mA); **7-** upper limb (6mA), face (5mA); **8-** upper limb (8mA); **9-** face (6mA); **10-** upper limb (7mA), lower limb (10mA), face (6mA); **11-** face (8mA); **12-** upper limb, face (13mA); **13-** upper limb (8mA), face (7mA); **14-** upper limb (12mA), face (11mA); **15-** upper limb (19mA), face (17mA)

Figure 4: Proposed Provisional Supplemental Cortical Motor Homunculus including the cortical somatotopy of intercostal muscles: **(a)** cortical motor representation **(b,c):** Primary motor cortex reconstructed with Meshlab opensource software³⁶ with the data available from The Human Brainnectome Atlas³⁷

Figure 5: Illustration demonstrating intraoperative subcortical Corticospinal Tract (isCST) (the novel “Clinical subcortical motor homunculus”)

Supplementary Figure 1 (a,b,c): fMRI demonstrating complex motor function mapping (**a**-foot rock task, **b**-Finger tapping, **c**-lip smacking) demonstrating cortical complex motor function representation of CST (*36y left-handed female with recurrent right frontal transformed WHO Grade III IDH 1 mutant MGMT methylated anaplastic gemistocytic astrocytoma who underwent subtotal resection*)

Supplementary Figure 2 (a,b,c,d): Pre-operative 3D Reconstruction of Cortical and subcortical corticospinal tract (*foot: blue circle, hand: orange circle, face: yellow circle representation*) utilizing Constrained Probabilistic tractography and modelling (MRTrix opensource software¹⁶) (*36y left-handed female with recurrent right frontal transformed WHO Grade III IDH 1 mutant MGMT methylated anaplastic gemistocytic astrocytoma who underwent subtotal resection*)

Supplemental Figure 3 (a,b,c): Integration of nTMS motor stimulation points with DTI reconstruction with neuronavigation (StealthStation Medtronic S7 machine, StealthViz software) (*White -Tumour, Red – CST (Face + Upper Limb), Green – CST (Lower Limb), Yellow dots – TMS Motor map, Blue Tractography – CST (contralateral to the lesion)*)(*22y male with right cystic-solid parietal WHO grade I IDH wildtype, ATRX wildtype, MGMT unmethylated pilocytic astrocytoma who underwent gross total resection*)

Supplementary Figure 4 (a,b,c): Intraoperative pre-resection ultrasound (ioUS) demonstrating right frontal tumour and its relation to the pre-central cortex at the cortical and subcortical level ; **(d):** Intraoperative image with cortical mapping (Tag 1 – Lower Limb, Tag 2 – Lower Limb and Intercostal Muscles, Tag 3 – Intercostal Muscles, Tag 4 – Forearm, Strip Number 1 – Monitoring the Foot, Strip Number 2 – Monitoring the Hand)(Black arrow: tumour location, blue arrow: cortical motor cortex and subcortical CST)(*36y left-handed female with recurrent right frontal transformed WHO Grade III IDH 1 mutant MGMT methylated anaplastic gemistocytic astrocytoma who underwent subtotal resection*)

Supplementary Figure 5: Isolated Subcortical stimulation of intercostal muscles during intraoperative subcortical mapping

Figure 1(a,b,c,d): Pre-operative 3D reconstruction of Corticospinal tract (*blue, green*) with tumour (*red*) utilising the pre-operative DTI and modelling using neuronavigation (Medtronic Stealth S7, StealthViz Software) (*70y female with left precentral metastatic tumour (primary lung small cell carcinoma who underwent gross total resection)*)

Figure 2: Cortical stimulation with monopolar probe stimulating different areas of motor cortex **(a)** 1,2,3: hand and forearm 4,5: foot and leg **(b)** 6,7,8,9,10,11,13,14- hand knob (ADM, FDI, APB, forearm, deltoid) 1,2,5- post central gyrus **(c)** 1,2: intercostal muscles 3,4: deltoid 5: foot **(d)** 1,2: arm, forearm, hand 3,4,5: hand, face 6,7: face

Figure 3: Subcortical stimulation with intraoperative MEPs demonstrating subcortical mapping of the corticospinal tract (**a,b**) with contact numbering corresponding to the mapping/stimulation positive areas with motor thresholds (**1-15**); **1-** upper limb (8mA), face (5mA); **2-** upper limb (8mA), face (6mA); **3-** upper limb (7mA), face (2mA); **4-** upper limb (5mA), face (2mA); **5-** upper limb (7mA), face (4mA); **6-** face (2mA); **7-** upper limb (6mA), face (5mA); **8-** upper limb (8mA); **9-** face (6mA); **10-** upper limb (7mA), lower limb (10mA), face (6mA); **11-** face (8mA); **12-** upper limb, face (13mA); **13-** upper limb (8mA), face (7mA); **14-** upper limb (12mA), face (11mA); **15-** upper limb (19mA), face (17mA)

675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708

Figure 4: Proposed Provisional Supplemental Cortical Motor Homunculus including the cortical somatotopy of intercostal muscles: **(a)** cortical motor representation **(b,c):** Primary motor cortex reconstructed with Meshlab opensource software³⁶ with the data available from The Human Brainnectome Atlas³⁷

725 **Figure 5:** Illustration demonstrating intraoperative subcortical Corticospinal Tract (isCST) (the
726 novel “Clinical subcortical motor homunculus”)

727

728

Table 1: Demographics of the patient data

Demographics	Data
Mean Age (years) (with Range)	50 (16-79)
Laterality of the lesion	N (%)
Right	97 (53.8%)
Left	83 (46.2%)
Gender	N (%)
Male	105 (58.4%)
Female	75 (41.6%)
Pathology of the lesion	
Glioma	
-High Grade (III,IV)	104 (57.7%)
-Low Grade (I,II)	41 (23%)
Meningioma	8 (4.4%)
Metastasis	25 (13.8%)
Vascular malformations	2 (1.1%)

748 **Table 2: Cortical Stimulation of motor cortex areas with their thresholds**

Cortical Stimulation	MT	Intraoperative Representation (Tag Numbers)
Leg	12mA	<p>1,2,3: hand and forearm 4, 5: foot and leg</p>
Intercostal Muscles	12mA	<p>1,2-intercostals 3,4-deltoid 5 -Foot</p>
Arm	10mA	<p>1,2 – arm, forearm, hand 3,4,5 – hand and face 6,7 – face</p>
Hand	6mA	<p>6,7,8,9,10,11,13,14- hand knob (<i>ADM, FDI, APB, forearm, deltoid</i>) 1,2,5- post central gyrus</p>
Face	10mA	<p>5,6,7- Face</p>

749 *Black Arrow: Tumour causing expansion of motor/pre-motor cortex, ADM: abductor digiti minimi ,FDI: first*
750 *dorsal interossei , APB: abductor pollicis brevis , MT: motor threshold*

751 **Table 3: Subcortical Stimulation of motor subcortex areas with their thresholds**
752

Subcortical Stimulation	MT	Representative Demonstration
Leg	4mA	<p>1,2,3- hand 5- face 6- foot 7,8,9- UL, LL</p> <p>6,7,8- 5-7mA 9- 4mA</p>
Intercostal muscles	4mA	<p>1,2,3,10- hand, forearm (10 mA) 4,5- foot, leg (12 mA) 6- Intercostal Muscles 7,8- foot, leg 9- hand, forearm (5 mA)</p>
Arm	20mA	<p>1,2,3- Intercostal muscles (cortical) 4- Deltoids (arm) 5- foot 6,8- Hand 7- UL, LL, Face, Intercostal muscles 9- Intercostal muscles (subcortical) 12mA 10- Hand (subcortical) Green- Superimposed CST model with operating microscope</p>
Hand		
Face		

753 UL: upper limbs, LL: lower limbs, MT: motor threshold, CST: corticospinal tract
754

Figures

Figure 1

(a,b,c,d): Pre-operative 3D reconstruction of Corticospinal tract (blue, green) with tumour (red) utilising the pre-operative DTI and modelling using neuronavigation (Medtronic Stealth S7, StealthViz Software) (70y female with left precentral metastatic tumour (primary lung small cell carcinoma who underwent gross total resection))

Figure 2

Cortical stimulation with monopolar probe stimulating different areas of motor cortex (a) 1,2,3: hand and forearm 4,5: foot and leg (b) 6,7,8,9,10,11,13,14- hand knob (ADM, FDI, APB, forearm, deltoid) 1,2,5- post central gyrus (c) 1,2: intercostal muscles 3,4: deltoid 5: foot (d) 1,2: arm, forearm, hand 3,4,5: hand, face 6,7: face

Figure 3

Subcortical stimulation with intraoperative MEPs demonstrating subcortical mapping of the corticospinal tract (a,b) with contact numbering corresponding to the mapping/stimulation positive areas with motor thresholds (1-15); 1- upper limb (8mA), face (5mA); 2- upper limb (8mA), face (6mA); 3- upper limb (7mA), face (2mA); 4- upper limb (5mA), face (2mA); 5- upper limb (7mA), face (4mA); 6- face (2mA); 7- upper limb (6mA), face (5mA); 8- upper limb (8mA); 9- face (6mA); 10- upper limb (7mA), lower limb (10mA), face (6mA); 11- face (8mA); 12- upper limb, face (13mA); 13- upper limb (8mA), face (7mA); 14- upper limb (12mA), face (11mA); 15- upper limb (19mA), face (17mA)

Figure 4

Proposed Provisional Supplemental Cortical Motor Homunculus including the cortical somatotopy of intercostal muscles: (a) cortical motor representation (b,c): Primary motor cortex reconstructed with Meshlab opensource software³⁶ with the data available from The Human Brainnectome Atlas³⁷

Figure 5

Illustration demonstrating intraoperative subcortical Corticospinal Tract (isCST) (the novel “Clinical subcortical motor homunculus”)

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementalMaterials.pdf](#)