Large contribution of fossil-derived components to aqueous secondary organic aerosols in China

Buqing Xu

Guangzhou Institute of Geochemistry, Chinese Academy of Sciences

Gan Zhang (✉ Zhanggan@gig.ac.cn)

Guangzhou Institute of Geochemistry, Chinese Academy of Sciences <https://orcid.org/0000-0002-9010-8140>

Örjan Gustafsson

Stockholm University <https://orcid.org/0000-0002-1922-0527>

Kimitaka Kawamura

Chubu University

Jun Lin

Chinese academy of sciences

August Andersson

Stockholm University <https://orcid.org/0000-0002-4659-7055>

Srinivas Bikkina

Chubu University

Bhagawati Kunwar

Chubu University

Ambarish Pokhrel

Chubu University

Guangcai Zhong

Guangzhou Institute of Geochemistry, Chinese Academy of Sciences

Shizhen Zhao

Guangzhou Institute of Geochemistry, Chinese Academy of Sciences

Jing Li

Guangzhou Institute of Geochemistry, Chinese Academy of Sciences

Chen Huang

Guangzhou Institute of Geochemistry, Chinese Academy of Sciences

Zhineng Cheng

Guangzhou Institute of Geochemistry, Chinese Academy of Sciences <https://orcid.org/0000-0002-6598-849X>

Sanyuan Zhu

Chinese Academy of Sciences

Ping'an Peng

Guangzhou Institute of Geochemistry, Chinese Academy of Sciences

Guoying Sheng

Guangzhou Institute of Geochemistry, Chinese Academy of Sciences

Article

Keywords:

Posted Date: December 14th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1155038/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Nature Communications on August 31st, 2022. See the published version at <https://doi.org/10.1038/s41467-022-32863-3>.

Abstract

Incomplete understanding of the sources of secondary organic aerosol (SOA) leads to large uncertainty in both air quality management and in climate change assessment. Chemical reactions occurring in the atmospheric aqueous phase represent an important source of SOA mass, yet, the effects of anthropogenic emissions on the aqueous SOA (aqSOA) are not well constrained. Here we use compound-specific dual-carbon isotopic fingerprints ($\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$) of dominant aqSOA molecules, such as oxalic acid, to track the precursor sources and formation mechanisms of aqSOA. Substantial stable carbon isotope fractionation of aqSOA molecules provides robust evidence for extensive aqueous-phase processing. Contrary to the paradigm that these aqSOA compounds are largely biogenic, radiocarbon-based source apportionments show that fossil precursors produced over one-half of the aqSOA molecules. Large fractions of fossil-derived aqSOA contribute substantially to the total water-soluble organic aerosol load and hence impact projections of both air quality and anthropogenic radiative forcing. Our findings reveal the importance of fossil emissions for aqSOA with effects on climate and air quality.

Introduction

Organic aerosol (OA) is the dominant component of atmospheric fine particulate mass¹. The majority of OA is secondary (SOA), formed in the atmosphere from the oxidation of precursor gases². Most SOA are composed of oxygenated and hygroscopic compounds, thereby, have harmful impacts on respiratory health and large, but uncertain effects on atmospheric radiative forcing³. However, current global models typically underpredict SOA magnitude, distribution, and dynamics, suggesting a limited understanding of their sources and formation processes⁴. An increasing number of modeling and experimental studies point toward aqueous chemical reactions occurring in cloud droplets and wet aerosols as an important missing pathway for SOA formation^{2–8}. Aqueous SOA (aqSOA) formation often involves mixed effects of anthropogenic and biogenic emissions^{2,3}. The extent to which aqSOA is controllable or “natural” is an important subject of ongoing research for both air quality and climate issues^{2,3}. However, the poorly understood aqueous chemical processes and the complexity of anthropogenic-biogenic interactions renders any firm conclusion hard to come by².

In model simulations, aqSOA are put to largely derive from the natural biogenic precursors, such as the oxidation products of isoprene^{3,4,8}. Fossil precursors are noticed to be less polar and less hydrophilic than biogenic precursors, thus diminishing the possibility of partitioning to the aqueous phase, and subsequent aqSOA formation⁹. However, a few recent studies in East Asia suggest that fossil precursors may substantially contribute to the formation of aqSOA. For example, fast aqueous-phase oxidation of polycyclic aromatic hydrocarbons emitted from fossil fuel combustion was observed in Beijing winter haze and may explain the observed SOA¹⁰. Radiocarbon-based estimations attributed ~50% of water-soluble OA to fossil sources in the East Asia outflow^{11,12}, compared to <30% in Europe, U.S., and South Asia^{13–18}. Nevertheless, on aqSOA formation, a quantitative estimation of the fossil contribution remains

challenging, largely due to the lack of specific and reliable technical tools in tracking the sources and formation mechanism of aqSOA in ambient aerosols.

Here we leverage the recent advent of compound-specific dual-carbon isotope fingerprint ($\delta^{13}\text{C}$ - $\Delta^{14}\text{C}$) of aqSOA molecules¹⁹ to quantify and characterize aqSOA sources and atmospheric chemical processes. The molecular-level $\Delta^{14}\text{C}$ contents of aqSOA tracers will provide direct constrain on their origins¹⁹, whereas $\delta^{13}\text{C}$ fingerprints of molecular tracers can differentiate various atmospheric processes/reactions²⁰. The technique was applied to purified oxalic acid and other similarly abundant organic acids (e.g., glyoxylic acid). Oxalic acid, likely represent the highest oxidation states of OA, has frequently been used as a proxy for aqueous processing in both laboratory and field studies^{21–23}. Organic acids that have high O/C ratios (around 1–2), are among the most abundant SOA components and are key end products in the aqueous-phase photochemical oxidation of various volatile organic compounds (VOCs) and intermediates such as glyoxal (Gly) and methylglyoxal (MeGly) in clouds or wet aerosols^{4, 5, 24, 25}. Therefore, these organic acids can serve as signature compounds to trace the aqSOA formation pathways^{21–23, 26}.

We used a set of year-round aerosol samples from a regional receptor site (Heshan Atmospheric Environmental Monitoring Superstation, Supplementary Fig. 1) in South China. We measured the aqSOA compounds and their $\delta^{13}\text{C}$ - $\Delta^{14}\text{C}$ isotope fingerprints during two contrasting phases of the East Asia monsoon system, i.e., continental outflow vis-à-vis South China Sea (SCS) air masses (Fig. 1). By combining the dual-carbon isotopic compositions and other chemical indicators together with meteorological parameters, we elucidate the relative contribution of fossil and biomass aqSOA precursors and their subsequent evolution processes, and provide ambient isotopic observational evidence for a large aqSOA formation from anthropogenic fossil emissions. Finally, the dual $\delta^{13}\text{C}$ - $\Delta^{14}\text{C}$ characterization of aqSOA molecules are extended to wider spatial coverage of China, which further confirm a ubiquitous contribution of aqueous-phase transformation of fossil precursors to organic aerosols.

Results

Meteorological setting and aerosol molecular compositions

The Heshan receptor site features an East Asia Monsoon Climate. The air is controlled by continental monsoon in winter and oceanic monsoon in summer, respectively (Fig. 1). Backward trajectories revealed that clean SCS air mass dominated the airshed during June-August 2017, while continental outflow rendered by enhanced anthropogenic emissions from the Pearl River Delta (PRD) city clusters, dominated during September 2017 to March 2018 (Fig. 1). Therefore, we categorized the 2017-2018 samples into two groups based on their distinctly different air mass transport regimes (Supplementary Table 1), i.e., “coastal background ($n = 11$)” and “continental outflow ($n = 21$)” (see Supplementary Fig. 2 for air mass clusters and Supplementary Fig. 3 for relative cluster contribution).

The chemical composition of aerosols during the campaign varied greatly with the air mass transport regimes. Concentrations of gas-phase pollutants (SO_2 , NO_2 , O_3 , CO) and major components of $\text{PM}_{2.5}$ during the continental outflow were 2-5 times higher than during the coastal background (Supplementary Table 2 and Supplementary Fig. 4). The observed organic matter (OM) and anthropogenic water-soluble inorganic constituents ($\text{WSIC}_{\text{anth}}$) increased in continental outflow samples, along with the aerosol liquid water (ALW) content associated with hygroscopic aerosol components (Supplementary Text 2). The concentration of OM, $\text{WSIC}_{\text{anth}}$, and ALW show a clear increase in continental outflow samples with 3, 5, and 3 times higher values than the coastal background, respectively (Fig. 1).

The molecular composition of diacids and related aqueous processing tracers are compiled in Supplementary Table 3. The malonic to succinic acid ratios (C_3/C_4) are significantly higher in the coastal background samples (1.5 ± 0.4) than the continental outflow (0.9 ± 0.3) (Supplementary Table 4). Since malonic acid are produced by photochemical oxidation of succinic acid in the atmosphere, this indicate that the coastal background aerosols have undergone more extensive photochemical aging^{27, 28}. The ratio of water-soluble organic carbon (WSOC) to organic carbon (WSOC/OC) is another indicator of atmospheric photochemical aging²⁹. For example, the WSOC/OC ratios increased from ~40% in South Asian source region to ~70% in the Indian Ocean receptor site due to the photochemical aging of OC during long-range atmospheric transport^{17, 29}. However, in this study, the WSOC/OC ratios of the continental outflow samples ($44 \pm 17\%$) cannot be distinguished statistically from the coastal background ($39 \pm 10\%$) (Supplementary Table 4), implying that additional mechanisms other than atmospheric aging may exist.

The uptake of water-soluble gaseous organic precursors to wet aerosol particles and fog/clouds, with subsequent aqueous-phase transformation to lower volatile compounds could be a significant pathway for WSOC aerosol formation^{23, 26, 30, 31}. Given the high relative humidity (RH%) ($74 \pm 9\%$; Supplementary Table 2) and the abundant hygroscopic particles (particularly for particulate nitrate as shown in Supplementary Fig. 4) in our continental outflow samples, we expect that the changes in the gas-particle partitioning driven by the increased ALW could have enhanced the formation of aqSOA species, such as oxalic acid^{23, 26, 32}. This can be supported by the sixfold higher aqSOA precursors (Gly + MeGly) mass fractions in OC observed in the continental outflow samples than the coastal background (Supplementary Table 4). With the addition of aqSOA formation, the higher WSOC/OC ratios in the continental outflow samples may thus be explained. Yet more precise and robust evidence/information is needed to give insights into the aqSOA formation.

Stable carbon isotope forensics of aqueous processing of aerosols

The $\delta^{13}\text{C}$ signatures reflect both sources and atmospheric processing. Emissions from different sources typically have different end-member $\delta^{13}\text{C}$ values. Once formed, the $\delta^{13}\text{C}$ signature is then influenced by kinetic isotope effects (KIE) during atmospheric reactions/processing¹⁵. Secondary aerosols stemming

from the oxidation of precursor gases due to the KIE typically lead to lower $\delta^{13}\text{C}$ values in particulate products compared to the gaseous reactants, whereas aerosol aging (i.e., oxidation of organic matter in the aerosol phase) can lead to a release of isotopically lighter carbon such as in CO_2 and CO , leaving residual (aged) aerosols enriched in ^{13}C ^{12, 15, 17, 29}. Figure. 2 shows the temporal variation of the mass concentrations and dual-carbon isotopic signatures of oxalic acid during the year-round observation. The $\delta^{13}\text{C}$ values for oxalic acid are on average $-24.6 \pm 2.7\text{‰}$ in the continental outflow samples and substantially more enriched in ^{13}C in the coastal background samples ($-19.9 \pm 2.3\text{‰}$). This clear contrast can likely be attributable to different emission sources and/or atmospheric processing.

Source mixing is not expected to have a major influence on the $\delta^{13}\text{C}$ composition of oxalic acid in this study. First, the $\delta^{13}\text{C}$ compositions of azelaic acid (C_9 diacids), an oxidation product of biogenic unsaturated fatty acids³³, in the coastal background samples ($-34.1 \pm 2.8\text{‰}$) and continental outflow ($-34.7 \pm 2.6\text{‰}$) are very close (Fig. 3a). These $\delta^{13}\text{C}$ values point to unsaturated fatty acids of C3 plant origin (-38.5‰ to -32.5‰)³⁴, while excluding marine phytoplankton and C4 plant sources that both instead are more enriched in $\delta^{13}\text{C}$ ^{35, 36}. This is consistent with the limited marine-biogenic contributions as indicated by the mass ratios of nss-SO_4^{2-} to SO_4^{2-} ($99 \pm 1\%$) and nss-K^+ to K^+ ($94 \pm 4\%$) in all samples²⁹. Given contributions from marine and C4 plants sources are excluded, the potential precursors of oxalic acid would then be anthropogenic VOCs (AVOCs; e.g., aromatic hydrocarbons: $-27.7 \pm 1.7\text{‰}$ ³⁷) and C3 plant-derived biogenic VOCs (BVOCs; e.g., isoprene: $-27 \pm 2\text{‰}$ ³⁸), of which the $\delta^{13}\text{C}$ values are also similar. Therefore, the isotopic range of various sources of oxalic acid at the supersite is minimal, leaving kinetic isotopic fractionation the most probable processes controlling the $\delta^{13}\text{C}$ composition of oxalic acid.

Oxalic acid is an SOA end product and can be formed by two pathways. The first pathway is a stepwise photochemical breakdown of longer-chain aliphatic diacids^{27, 28}. Recently, significant amounts of oxalic acid followed by succinic acid were also found to produce via ozone oxidation of isoprene under dry condition³⁹. The second pathway is the oxidation of BVOCs and AVOCs to semi-volatile organic compounds (SVOCs, e.g., Gly and MeGly), followed by partitioning of SVOCs to wet aerosols or cloud/fog droplets, with subsequent aqueous-phase transformation to pyruvic acid (Pyr) and glyoxylic acid (ωC_2) and finally oxidized to oxalic acid^{24–26}. Our dataset displays a significant correlation of oxalic acid with ALW ($r^2 = 0.78$, $P < 0.001$, Fig. 3b), suggesting that oxalic acid was dominantly formed *via* gas-to-liquid transfer of precursors and subsequent aqueous-phase reactions. This is consistent with a previous model study revealing that aqSOA formation is limited by ALW rather than by the availability of precursors⁴⁰. As shown in Fig. 3b, the five points with excessive ALW are off the trend, signifying that ALW was no longer the limiting parameter in the specific samples. The oxalic acid and its aqueous-phase intermediate product, glyoxylic acid, also display a significant correlation ($r^2 = 0.65$, $P < 0.001$), which stand for the aqueous-phase pathway.

The compound-specific $\delta^{13}\text{C}$ values of oxalic acids and relevant SOA species can help to explore the formation pathway. First, the higher homologs aliphatic diacids from C_3 to C_9 , in contrast to C_2 , show no significant statistical difference (t -test, $P > 0.05$, Supplementary Table 5) between the two air mass source regimes (Fig. 3a). Instead, ωC_2 and Pyr show significantly lower (t -test, $P < 0.05$, Supplementary Table 5) $\delta^{13}\text{C}$ values of (by 4.4‰ and 3.2‰, respectively, Fig. 3a) during continental outflow than coastal background. As aqSOA formed from VOCs or SVOCs typically leads to lower $\delta^{13}\text{C}$ values relative to the precursors since lighter-isotope-containing precursors are preferentially oxidized to form reaction products. The continental outflow regime was characterized by high RH% and substantial anthropogenic hygroscopic particles, which could facilitate aqSOA formation. Therefore, the vast difference in the $\delta^{13}\text{C}$ composition of oxalic acid between the two air mass source regimes could be mainly resulted from the aqueous-phase reaction pathway, i.e. $\text{AVOCs/BVOCs} \rightarrow \text{SVOCs} \rightarrow \text{Pyr} \rightarrow \omega\text{C}_2 \rightarrow \text{C}_2^{41}$, rather than photochemical breakdown of longer-chain diacids. Note that the source mixing contribution is minimal as mentioned above.

This aqueous reaction effect on isotope fractionation may also apply to water-soluble OA, which is a cocktail of primary OA (e.g., sugars) and SOA species. Figure. 3c reveals a clear decrease in $\delta^{13}\text{C}$ signatures of oxalic acid against an increase in WSOC concentrations ($r^2 = 0.70$, $P < 0.001$). Atmospheric aqueous-phase processing would simultaneously cause a decrease in the $\delta^{13}\text{C}$ signature of reaction products and an increase in the WSOC mass concentrations. This is precisely what was observed during this campaign. Overall, the results indicate that aqSOA formation processes may be a significant route to form water-soluble OA in the atmosphere of South China and similar settings.

Molecular-level radiocarbon isotope forensics of aqSOA sources

Besides $\delta^{13}\text{C}$ signature, which undergoes strong isotope fractionation during atmospheric processing, radiocarbon isotopic fingerprint ($\Delta^{14}\text{C}$) is conservative and highly precise in distinguishing fossil from biogenic/biomass sources¹⁹. Individual water-soluble diacids and related compounds (C_2 , C_3 , C_4 , ωC_2 , and MeGly) were isolated by preparative capillary gas chromatography techniques (*Materials and Methods*). It is possible to harvest sufficient mass of C_2 oxalic acid from aerosol samples, while the other isolated compounds had to each be pooled per month to obtain sufficient carbon mass for the ^{14}C measurement. The determined $\Delta^{14}\text{C}$ compositions reported as the “fraction of modern carbon” (F_m) are shown in Supplementary Table 6 and Supplementary Table 7. It is further possible to translate the F_m signatures into fractional contribution from fossil (f_{fossil}) *versus* non-fossil (including biogenic emissions: *bio* and biomass burning: *bb*; hereafter referred as $f_{\text{bio/bb}}$) sources (Supplementary Text 3).

The excellent sensitivity and precision of molecular-level ^{14}C analysis reveal a clear diversity in ^{14}C signatures of oxalic acid for different samples and regimes (Fig. 2 and Supplementary Table 6). In coastal background air mass regime, $67 \pm 9\%$ of oxalic acid is from biogenic emission and biomass

burning. Although summer monsoon originating from SCS results in efficient precipitation (Fig. 1), there is still a substantial fossil contribution ($f_{\text{fossil}} = 19\% - 48\%; 33 \pm 9\%$) from anthropogenic activities. For the continental outflow samples, the $f_{\text{fossil-C}_2}$ ($f_{\text{fossil}} = 42\% - 78\%; 55 \pm 10\%$) are even higher than biogenic/biomass source contributions. Back trajectory analysis reveals that during the latter period the receptor site was downwind from highly polluted continental source regions, which thus apparently is a source of oxalic acid in the regional aerosol (Fig. 1). The concentrations of C_2 oxalic acid increased with the increment of $f_{\text{fossil-C}_2}$ ($r^2 = 0.52, P < 0.001$; Supplementary Fig. 5), suggesting that the precursors from fossil-fuel emission was the dominant factor in controlling the ambient abundance of oxalic acid. The high ratios of NO_3^- in atmospheric particles ($\sim 10\%$; Supplementary Fig. 4) indicate a significant role of traffic emission-derived precursors⁴² in the formation of oxalic acid. Moreover, terephthalic acid (tPh) was one of the most abundant anthropogenic diacids accounting for 5% of total diacids (Supplementary Table 3). As tPh is a combustion tracer of plastic polymers⁴³, this highlights the importance of precursors from plastic burning to the formation of fossil oxalic acid. Thus, both traffic and plastic burning emissions were important contributors to the fossil source-derived oxalic acid.

The mass ratio of adipic acid to azelaic acid (C_6/C_9) is often used to discriminate the relative significance of biogenic *versus* anthropogenic source contribution to diacids. Here, C_6 adipic acid is the photochemical oxidation product of cyclic hexene emitted from the fossil fuel combustion sources^{27, 44}, whereas C_9 azelaic acid is a photochemical degradation product of biogenic unsaturated fatty acids³³. Lower C_6/C_9 ratios (0.37 ± 0.13) were observed in the coastal background samples than continental outflow (0.76 ± 0.21). Furthermore, C_6/C_9 ratios correlates well with the parameter of $f_{\text{fossil-C}_2}$ ($r^2 = 0.63, P < 0.001$; Supplementary Fig. 6). Hence, C_6/C_9 ratios gives strong and consistent support to the result of diacids ^{14}C -based source apportionment.

The fossil contents of oxalic acid in the continental outflow samples are compared with its presumed precursors of longer-chain diacids (C_3 and C_4 diacids) and its aqueous processing precursors (ωC_2 and MeGly) (Fig. 4c and Supplementary Table 7). The result indicates that fossil sources were the overwhelming contributor to both oxalic acid ($55 \pm 10\%$) and to its aqueous processing precursors, i.e., glyoxylic acid (ωC_2 ; $69 \pm 4\%$) and methylglyoxal (MeGly; $67 \pm 5\%$). Therefore, a dominance from fossil fuel sources to their aqSOA formation in the continental outflow is obvious.

Although malonic C_3 and succinic C_4 acids are believed to be partly produced from aqueous processing^{22, 26}, there is for these C_3 and C_4 a somewhat lower but still large contribution from fossil sources (C_3 : $31 \pm 4\%$; C_4 : $36 \pm 2\%$). This demonstrates that C_3 and C_4 diacids instead were mainly originated from the stepwise photochemical breakdown of biogenic unsaturated fatty acids even in highly polluted continental outflow air mass. The large contrast of fossil fraction among C_2 diacids and its higher homologs (i.e., C_3 and C_4 diacids) further demonstrate that oxalic acid is mainly produced from fossil-fuel derived precursors, rather than formed by the photochemical breakdown of longer-chain diacids of more biogenic origin⁴⁵.

The ^{14}C -derived concentrations of oxalic acid from non-fossil sources ($\text{bio}/\text{bb}\text{-C}_2$) span from 40 to 417 ng m^{-3} ($207 \pm 88 \text{ ng m}^{-3}$), whereas from fossil source ($\text{fossil}\text{-C}_2$) are 21 to 650 ng m^{-3} ($230 \pm 168 \text{ ng m}^{-3}$) (Supplementary Fig. 7). We observed a close similarity in the temporal variability of $\text{bio}/\text{bb}\text{-C}_2$ with nss-K^+ and levoglucosan (Supplementary Fig. 8), especially during fall to winter, when open burning of post-harvest agricultural crop residues is widespread in rural China⁴⁶. For all data from the whole campaign, $\text{bio}/\text{bb}\text{-C}_2$ concentration are positively correlated with levoglucosan ($r^2 = 0.67$, $P < 0.001$; Fig. 4a). The ratios of $\text{bio}/\text{bb}\text{-C}_2$ and nss-K^+ also show significant correlation ($r^2 = 0.58$, $P < 0.001$; Supplementary Fig. 9). In contrast, the coastal background samples record low concentrations of levoglucosan and nss-K^+ (Fig. 4a and Supplementary Fig. 9), signifying an insignificant contribution from biomass burning. Further, a strong linear correlation ($r^2 = 0.81$, $P < 0.001$, Fig. 4b) was found between $\text{bio}/\text{bb}\text{-C}_2$ concentrations and biogenic SOA tracers (i.e., isoprene tracers and α/β -pinene tracers) for the coastal background samples, suggesting that the $\text{bio}/\text{bb}\text{-C}_2$ in low biomass burning scenario was formed mainly from biogenic precursors. These evidences confirm that $\text{bio}\text{-C}_2$ likely represents a background C_2 level resulting from local emissions of C3 plants, whereas $\text{bb}\text{-C}_2$ contribution is an addition of excessive non-fossil C_2 concentrations. We therefore assume that, when there is little biomass burning contribution ($\text{Lev} \approx 0$), the concentration of $\text{bio}/\text{bb}\text{-C}_2$ (128 ng m^{-3} ; Fig. 4a) would approximately represent the background C_2 level, that is, the $\text{bio}\text{-C}_2$. With this estimation, the total concentration of oxalic acid can be quantitatively distributed into three sources, i.e., $\text{fossil}\text{-C}_2$, $\text{bb}\text{-C}_2$ and $\text{bio}\text{-C}_2$.

The anthropogenic C_2 (including $\text{fossil}\text{-C}_2$ and $\text{bb}\text{-C}_2$) accounted for 78% of total C_2 in the continental outflow samples, whereas its contribution is 39% in coastal background (Fig. 4d). A tracer-based estimation suggest that anthropogenic gasSOA, which are formed by photochemical reaction in gas phase followed by thermodynamic partition to the condensed phase⁴, account for 79% of the total gasSOA in the PRD during fall-winter⁴⁷. This anthropogenic content of gasSOA is consistent with our ^{14}C -constrained source apportionment of aqSOA ($f_{\text{anthropogenic}}\text{-C}_2 = 78\%$). By extension, the contribution of anthropogenic emissions in aqSOA and gasSOA are likely to be consistent under ambient conditions with high ALW content. Our observation-based isotope constraints question the paradigm, largely based on model simulations, which instead have held that aqSOA, which may contribute as much mass as gasSOA to SOA budget, are largely formed from biogenic emissions⁴. Previous study considered that fossil precursors may be more hydrophobic than biogenic precursors, and thus are likely form SOA preferentially through gas-phase processing⁹. However, the oxidation products of fossil precursors are largely water-soluble and thus their chemical reaction in the aqueous phase constitutes an unneglected pathway, which may be competitive to SOA formation *via* the gas-phase chemistry.

Figure 5 shows the dual effects of precursor sources and atmospheric processing of aqSOA by combining the ^{13}C and ^{14}C (f_{fossil}) signatures of oxalic acid. Lower oxalic acid concentrations were found in the coastal background samples, with higher non-fossil contributions, and more enriched in ^{13}C . Higher oxalic acid concentrations in the continental outflow samples are characterized by higher fossil

contributions and more depleted in ^{13}C . The hygroscopic particles and precursors in continental outflow air masses facilitate the aqueous-phase chemical processes, which results in lighter $\delta^{13}\text{C}$ values and higher fossil contribution to oxalic acid.

Conversely, such an aqSOA formation process became less important during coastal background period, when there are less significant anthropogenic activities, and hence less precursors and hygroscopic particles in the atmosphere. In this case, the photochemical degradation (aging) of oxalic acid would surpass its aqueous-phase formation, which is evident of the increasing $\delta^{13}\text{C}$ values of oxalic acid (Fig. 5). The interpretation may also apply to several outliers of the continental outflow samples (pink shadow in Fig. 5), of which the OC/EC ratios are much lower than the others, referring to an exhaustive atmospheric aging.

Effects of aqueous-phase processing on organic aerosols

Many studies have investigated the formation of aqSOA from biogenic/biomass precursors (e.g., isoprene)^{3, 7, 8, 22, 48, 49}, while much less attention has been paid to fossil/anthropogenic precursors (e.g., aromatic hydrocarbons). Fossil precursors have been thought to consist mostly of hydrophobic components, and thus fossil-derived SOA have been expected to be less soluble in water⁹. Smog chamber experiments show that water-soluble organic aerosols exhibited more similarities to biogenic SOA than fossil SOA^{50, 51}. In fact, low fossil contributions to WSOC (0–29%) have been reported in Europe, North America, and South Asia across different seasons^{13–18}, and ^{14}C analyses of aerosol OC and its sub-fractions have been interpreted such that the majority of fossil-derived OA was water insoluble^{52, 53}. Hence, the picture from these earlier field and laboratory studies is that water-soluble OA is only to very limited extent stemming from fossil sources. In contrast, the results from the current study, using an extensive set of molecular $^{13}\text{C}/^{14}\text{C}$ data in China, strongly points to anthropogenic-fossil emissions as an important source to water-soluble OA. This is also supported by a large contribution of fossil sources to WSOC (30–60%) reported for both Chinese urban areas^{53–56} and a receptor station in the southeast Yellow Sea intercepting a large regional footprint^{11, 12}. Taken together, these two sets of findings imply a significant missing pathway responsible for the enhancement in fossil fuel-derived water-soluble OA in China.

The enhanced atmospheric aqueous-phase processing ($\delta^{13}\text{C}$ of oxalic acid is used as the proxy for aqSOA formation) simultaneously cause an increasing in the WSOC mass concentration, suggesting that aqSOA constitute a substantial water-soluble OA source (Fig. 3c). A recent study also showed that the oxidation of gas-phase components and subsequent uptake of water-soluble oxidation products triggered by the increase of ALW can explain the increase in SOA mass in Chinese megacities³². This is further corroborated by radiocarbon signatures of individual aqSOA tracers (i.e., C_2 , ωC_2 , and MeGly), which reveals a dominant fossil source to aqSOA (Fig. 4c). The aqSOA even showed a fossil content (55–69%) comparable to the water-insoluble OC in other Chinese urban areas such as Beijing (62–68%) and Guangzhou (53–56%) in winter haze⁵⁷. Our dual carbon isotope data of aqSOA tracers indicate that

various fossil precursors may be oxidized to form aqSOA in the atmosphere and contribute substantially to the WSOC aerosols. We suggest that this atmospheric mechanism is responsible for the enhanced formation of fossil-derived WSOC in China.

To test the spatial extent of the findings, the dual-carbon isotopic compositions of individual aqSOA compounds (oxalic acid) and bulk WSOC are further compared in winter and in summer for the five emission hotspot megacities of China (Fig. 6 and Supplementary Table 8). The average $\delta^{13}\text{C}$ and ^{14}C -derived $f_{\text{bio/bb}}$ values of WSOC show minor difference in winter and summer ($\delta^{13}\text{C}$: $-24.2 \pm 0.4\text{‰}$ vs. $-24.9 \pm 0.7\text{‰}$; $f_{\text{bio/bb}}$: $63 \pm 11\%$ vs. $60 \pm 9\%$). In contrast, the $\delta^{13}\text{C}$ and $f_{\text{bio/bb}}$ values of oxalic acid are significantly lower during winter than summer ($\delta^{13}\text{C}$: $-27.1 \pm 1.8\text{‰}$ vs. $-22.7 \pm 2.5\text{‰}$; $f_{\text{bio/bb}}$: $53 \pm 11\%$ vs. $65 \pm 13\%$). These seasonal variability in dual-carbon isotopic compositions of oxalic acid was consistent with the observation at the supersite (refer to Fig. 2). Although oxalic acid accounts for a large fraction of WSOC ($\sim 5.2\%$), probably the single most abundant compound, we observe significant but opposite difference in the $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ compositions between oxalic acid and WSOC during the two seasons (Fig. 6). Both aqSOA formation and aging can lead to the production of more polar compounds such as oxalic acid. Oxalic acid was more enriched in ^{13}C (by 2.2‰) and ^{14}C ($f_{\text{bio/bb}}$ increased by 5.0%) than WSOC in summer, when aerosols were more influenced by atmospheric aging. This is probably due to that biomass/biogenic component from the WSOC pool is preferentially subjected to oxidative aging to small molecules^{12, 58}, which results in more aged oxalic acid (enriched in $\delta^{13}\text{C}$) more derived from biomass/biogenic precursors (higher $f_{\text{bio/bb}}$).

On the contrary, oxalic acid was more depleted in ^{13}C (by 2.9‰) and ^{14}C (lower $f_{\text{bio/bb}}$ by 10%) when compared to the WSOC pool in winter. This agrees with the dominant role of aqueous-phase formation processes triggered by the increased ALW in winter³². As mentioned above, overwhelming fossil contributions to aqSOA compounds were observed, that were consistent with or even higher than the fossil content in water-insoluble OA. Substantial fossil-derived precursors are likely oxidized to WSOC aerosol through secondary aqueous processing in winter, resulting in products (such as oxalic acid) with more negative $\delta^{13}\text{C}$ values and more fossil contributions than the bulk WSOC pool. The seasonality of $\delta^{13}\text{C}$ - $\Delta^{14}\text{C}$ composition patterns of oxalic acid and WSOC in the five Chinese cities further confirm a ubiquitous contribution of aqueous-phase transformation of fossil precursors to the WSOC aerosol.

Discussion

Aqueous-phase processing of fossil precursors, competitive to gas-phase pathway, could amplify the impact of anthropogenic emissions on water-soluble OA, alter the hygroscopicity of aerosols, influence the cloud condensation nuclei (CCN) activity, and hence affect the cloud-mediated effect of anthropogenic aerosols on the climate system⁵⁹. Condensation of oxalic acid and its salts in the aqueous phase may significantly impact the CCN activity of aerosol particle⁶⁰. The formation of aqSOA can also promote positive feedback, that the dissolution of aqSOA in the particle phase increases ALW, which in

turn promotes further gas-to-liquid transfer of water-soluble precursors^{23, 32}. Oxalic acid is identified as the upper bound of the hygroscopic properties of OA, with a *k*-Köhler of 0.48⁷. Using the Zdanovskii–Stokes–Robinson (ZSR) mixing model⁶¹, the effect of anthropogenic oxalic acid on aerosol water accounts for 3–38% (average: 10%) of the ALW contributed by organic compounds (Supplementary Text 2).

A positive feedback mechanism between hygroscopic particles and ALW has been reported^{31, 62}, in which the enhancement of aqSOA formation triggered by the increase of ALW may play a key role^{6, 32}. Our year-round observation at the regional receptor site witnesses a positive feedback loop among hygroscopic particles, ALW, and aqSOA formation. The hygroscopic particles induce ALW growth, facilitating the partitioning of gas-phase oxidation products into aqueous-phase medium, and promoting the formation of hygroscopic particles and aqSOA. The production of hygroscopic particles and aqSOA in turn enhances the ALW, enforcing the trend. The ALW and hygroscopic particles (mostly contributed by secondary inorganic aerosols) can be considered anthropogenic in most continental regions^{3, 30, 40}. We provide ¹⁴C-based evidence that, besides biogenic precursors, anthropogenic precursors can also contribute extensively to aqSOA formation. Therefore, a broad control of various anthropogenic emissions such as SO₂, NO_x, and VOC precursors is vital for reducing the organic particulate pollution.

Nitrate is the dominant atmospheric hygroscopic particles in China^{30, 31, 42}. A model study predicted that 92% of increased ALW in eastern China by the year of 2100 can be attributed to the increase in anthropogenic nitrate aerosol³⁰. Here, the average nitrate concentration in the continental outflow samples was nearly 20 times higher than coastal background (Supplementary Table 2). Thus, this nitrate enhancement in the particulate phase could be the dominant factor that facilitated the increase of ALW in the continental outflow regime. Furthermore, there has been a rapid transition from coal combustion to natural gas in China. However, natural gas combustion can produce more than three times as much water vapor as coal burning⁶³. The combustion-derived water was interpreted to constitute 6.2% of the atmospheric moisture and added 5.1% of the anthropogenic PM_{2.5} in northwestern China⁶³. Therefore, the enhancing effects of particulate nitrate and combustion-derived water vapor on aqSOA formation should be addressed when changing energy structure in future climate and air quality scenarios.

In conclusion, novel compound-specific dual carbon isotopic evidence from field observations show that fossil anthropogenic precursors contribute to the formation of aqSOA to a much higher extent than expected. Large fractions of fossil source-derived aqSOA significantly change the chemical characterization and hygroscopicity of organic aerosols, which in turn facilitate the gas-to-liquid transfer of water-soluble precursors and promote the subsequent aqSOA production. Atmospheric aqueous-phase processes are expected to increase due to the enhanced evapotranspiration in a warmer world⁶¹. An absence of accounting such aqueous processing incurred by fossil precursors could lead to an underestimation of the anthropogenic contribution to organic aerosols. Understanding the role of anthropogenic emissions in aqSOA formation and the chemical mechanisms involved are of critical concern for future climate and air quality projections, and within the context of energy needs and choices.

Materials And Methods

The regional receptor site was positioned in Heshan Atmospheric Environmental Monitoring Superstation (22.711° N, 112.927° E, 60 m asl), a rural site located at 50 km southwest to the megacity of Guangzhou in the central PRD. The surrounding area of the sampling site is dominated by farmlands and forests and is far from local anthropogenic emissions⁶⁴. During the winter northeast-monsoon season, this specific site well intercepts high anthropogenically dominated outflow air mass from Chinese continental. The strength of anthropogenic impact decreases in the summer southeast/southwest monsoon, with the arrival of clean air mass originating from the Western Pacific/Indian Ocean. Field experiments started in June 2017 and ended in May 2018. PM_{2.5} aerosol samples ($n=32$) were collected onto pre-combusted quartz-fiber filters by using a high-volume sampler at an air-flow rate of 1m³ /min for 48 hours.

PM_{2.5} samples were additionally studied in five emission hot spot megacities of China (Beijing, Shanghai, Guangzhou, Chengdu, Wuhan), which were collected continuously every 24h for 1 week in winter (January 2018) and in summer (July 2018). Continuous sets of 7 filters collected in one week were combined to represent the seasonal characteristic of each urban site. The sample filters were stored in darkness at -20°C prior to analysis. Field blank filters were collected for each campaign.

Backward air mass trajectories and ALW estimation

Data of atmospheric concentration of meteorological parameters and gaseous pollutants were obtained from the hourly monitor by Heshan Atmospheric Environmental Monitoring Superstation. To characterize different geographical sources of air masses encountered at Heshan, 3-day isentropic backward air mass trajectories were computed at an elevation of 100 m for every 12 h using the HYSPLIT-4 model. The back trajectories are grouped into two major transport pathways with vastly different source regions during the sampling campaigns.

The total aerosol liquid water (ALW) is taken as the sum of water associated with individual aerosol chemical components, with the assumption that the particles are internally mixed⁶⁵. ALW from inorganic species was estimated using a thermodynamic equilibrium model, ISORROPIA- β ; <http://nenes.eas.gatech.edu/ISORROPIA/> (Supplementary Text 2) using particle concentration of inorganic ions and meteorological parameters (temperature and RH%). The organic contribution for ALW was calculated by Zdanovskii–Stokes–Robinson (ZSR) mixing rule as discussed in Supplementary Text 2.

Chemical analysis.

The concentration of typical aqSOA species (diacids, oxoacids, and α -dicarbonyls) was measured following earlier protocols.^{19, 27} An aliquot of filter samples was extracted with Milli-Q water. The extracts were concentrated to dryness and derivatized with 10% BF₃ in 1-butanol (Sigma-Aldrich, St Louis, MO) in a hot water bath at 100°C for about 1 h. The derivatives were extracted with n-hexane and finally

quantified using GC-MS (Agilent 7890A gas chromatography and Agilent 5975C mass spectrometry). The concentrations reported here are corrected for field blank, but are not corrected for recovery efficiencies.

Relevant details of the analysis method for aerosol carbonaceous components (organic carbon: OC, elemental carbon: EC; water-soluble organic carbon: WSOC), water-soluble inorganic constituents (WSIC, i.e., Na^+ , NH_4^+ , K^+ , Mg^{2+} , Ca^{2+} , Cl^- , NO_3^- , and SO_4^{2-}), biogenic SOA tracers; and sugar compounds are provided in the Supplementary Text 1.

Compound-specific $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ analysis

The $\delta^{13}\text{C}$ values of diacids derivatives were determined using GC ISOLink2/IRMS (Thermo Fisher Scientific). The $\delta^{13}\text{C}$ values of free diacids and related compounds were then calculated by an isotopic mass balance approach, based on the measured $\delta^{13}\text{C}$ values of derivatives and the derivatizing agent (1-butanol, $\delta^{13}\text{C} = -30.21\text{‰}$). The isotopic fractionation during the derivatization step has been reported to be less than 0.73‰ in our earlier publications¹⁹. Each sample was measured in duplicate and the mean $\delta^{13}\text{C}$ values were reported. The difference in $\delta^{13}\text{C}$ of diacids (i.e., C_2 , C_3 , C_4 , C_5 , C_6 , Ph, tPh, C_9) in replicate analyses was generally $< 1\text{‰}$, though the analytical accuracy was $< 2\text{‰}$ for other compounds (i.e., Pyr, ωC_2).

The analytical method for the isolation of individual diacids to determine $\Delta^{14}\text{C}$ compositions builds on our previously publications¹⁹. In brief, the filter was extracted with the Milli-Q water and derivatized with BF_3 /1-butanol as described above. Microgram quantities of C_2 , C_3 , C_4 , ωC_2 , and MeGly were isolated and collected by preparative capillary gas chromatography (pcGC) through around 50 consecutive runs (5 μl per injection), which was sufficient for off-line natural abundance radiocarbon analysis. The pcGC isolates were combusted at 920°C and the resulting CO_2 was finally converted to graphite via hydrogen reduction method. Measurements of ^{14}C were performed using Accelerator Mass Spectrometry (1.5SDH-1, 0.5MV, NEC, USA)¹⁹. In all cases, ^{14}C results are reported as “fraction of modern carbon” (F_m) relative to NBS Oxalic Acid I standard normalized to a $\delta^{13}\text{C}$ value of -25‰ . The F_m values were further converted into the fraction of non-fossil source ($f_{\text{bio/bb}}$) with a conversion factor of 1.06 to compensate for the excess ^{14}C produced by nuclear bomb testing in the 1950s-1960s. The concentration of diacids in blank filters was $< 0.06\%$ of the real samples, therefore, no filter blank subtraction was performed for the isotope analysis in this study. Sample results were corrected for procedural blanks using a pair of processing standards with modern or fossil radiocarbon composition. An isotopic mass-balance approach was adopted to correct the carbon contribution of butanol groups (1-butanol, $F_m = 0.0029 \pm 0.001$) added in the derivatives of diacids, where appropriate. Calculations for blank correction, F_m results and error propagation are shown in Supplementary Text 3.

$\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ analysis of WSOC

The analytical method for the measurements of $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$ composition of WSOC were described previously⁵⁴ and reported in Supplementary Text 4. Measurements of the $\delta^{13}\text{C}$ composition of WSOC were performed on a Flash 2000 elemental analyzer connected to a Thermo Scientific Delta V isotope ratio mass spectrometer. Measurements of the $\Delta^{14}\text{C}$ composition were performed using accelerator mass spectrometry facility (1.5SDH-1, 0.5MV, NEC, USA) of the Guangzhou Institute of Geochemistry of the Chinese Academy of Sciences (GIGCAS).

Declarations

Data availability

All data analyzed during the current study are included in this article and its Supplementary Information.

Acknowledgements

This study was funded by the Natural Science Foundation of China (42030715), the Alliance of International Science Organizations research and cooperative projects (ANSO-CR-KP-2021-05), the Guangdong Foundation for Program of Science and Technology Research (2017BT01Z134), and the National Key R&D Program of China (2017YFC0212000). We would like to thank Dr Weiwei Hu for helpful discussions and Dr Lili Ming for support with instruments.

Author contributions

B.X. and G.Z. designed the experiment. JUN.L., Z.C. and S.-Z.Z. provided the samples. B.X., A.A., B.K., A.P., JING.L., C.H., Z.C. and S.-Y.Z. processed data and performed analyses. B.X. wrote the paper. G.Z., Ö.G., K.K., A.A., S.B., G.-C.Z., P.P., and G.S. commented on the manuscript.

Additional information

Supplementary Information is available for this paper

Competing financial interests: The authors declare no conflict of interests.

References

1. Jimenez JL, *et al.* Evolution of Organic Aerosols in the Atmosphere. *Science* **326**, 1525–1529 (2009).
2. Carlton AG, Christiansen AE, Flesch MM, Hennigan CJ, Sareen N. Multiphase Atmospheric Chemistry in Liquid Water: Impacts and Controllability of Organic Aerosol. *Acc. Chem. Res.* **53**, 1715–1723 (2020).

3. McNeill VF. Aqueous organic chemistry in the atmosphere: sources and chemical processing of organic aerosols. *Environ. Sci. Technol.* **49**, 1237–1244 (2015).
4. Ervens B, Turpin BJ, Weber RJ. Secondary organic aerosol formation in cloud droplets and aqueous particles (aqSOA): a review of laboratory, field and model studies. *Atmos. Chem. Phys.* **11**, 11069–11102 (2011).
5. Lim YB, Tan Y, Perri MJ, Seitzinger SP, Turpin BJ. Aqueous chemistry and its role in secondary organic aerosol (SOA) formation. *Atmos. Chem. Phys.* **10**, 10521–10539 (2010).
6. Hennigan CJ, Bergin MH, Dibb JE, Weber RJ. Enhanced secondary organic aerosol formation due to water uptake by fine particles. *Geophys. Res. Lett.* **35**, (2008).
7. Gilardoni S, *et al.* Direct observation of aqueous secondary organic aerosol from biomass-burning emissions. *Proc. Natl Acad. Sci. USA* **113**, 10013-10018 (2016).
8. Lamkaddam H, *et al.* Large contribution to secondary organic aerosol from isoprene cloud chemistry. *Sci. Adv.* **7**, eabe2952 (2021).
9. Favez O, Sciare J, Cachier H, Alfaro SC, Abdelwahab MM. Significant formation of water-insoluble secondary organic aerosols in semi-arid urban environment. *Geophys. Res. Lett.* **35**, (2008).
10. Wang J, *et al.* Aqueous production of secondary organic aerosol from fossil-fuel emissions in winter Beijing haze. *Proc. Natl Acad. Sci. USA* **118**, e2022179118 (2021).
11. Fang W, *et al.* Divergent Evolution of Carbonaceous Aerosols during Dispersal of East Asian Haze. *Sci. Rep.* **7**, 10422 (2017).
12. Kirillova EN, Andersson A, Han J, Lee M, Gustafsson Ö. Sources and light absorption of water-soluble organic carbon aerosols in the outflow from northern China. *Atmos. Chem. Phys.* **14**, 1413–1422 (2014).
13. Weber RJ, *et al.* A study of secondary organic aerosol formation in the anthropogenic-influenced southeastern United States. *J. Geophys. Res.* **112**, (2007).
14. Zhang YL, *et al.* Fossil and non-fossil sources of different carbonaceous fractions in fine and coarse particles by radiocarbon measurement. *Radiocarbon* **55**, 1510–1520 (2013).
15. Kirillova EN, *et al.* ¹³C- and ¹⁴C-based study of sources and atmospheric processing of water-soluble organic carbon (WSOC) in South Asian aerosols. *J. Geophys. Res.: Atmos.* **118**, 614–626 (2013).
16. Szidat S, *et al.* Fossil and non-fossil sources of organic carbon (OC) and elemental carbon (EC) in Goteborg, Sweden. *Atmos. Chem. Phys.* **9**, 1521–1535 (2009).
17. Bosch C, *et al.* Source-diagnostic dual-isotope composition and optical properties of water-soluble organic carbon and elemental carbon in the South Asian outflow intercepted over the Indian Ocean. *J. Geophys. Res.: Atmos.* **119**, 11,743-711,759 (2014).
18. Wozniak AS, Bauer JE, Dickhut RM. Characteristics of water-soluble organic carbon associated with aerosol particles in the eastern United States. *Atmos. Environ.* **46**, 181–188 (2012).
19. Xu B, *et al.* Compound-specific radiocarbon analysis of low molecular weight dicarboxylic acids in ambient aerosols using preparative gas chromatography: method development. *Environ. Sci.*

Technol. Lett. **8**, 135–141 (2021).

20. Cao F, *et al.* New insights into the sources and formation of carbonaceous aerosols in China: potential applications of dual-carbon isotopes. *Natl. Sci. Rev.* **4**, 804+ (2017).
21. Sorooshian A, *et al.* Particulate organic acids and overall water-soluble aerosol composition measurements from the 2006 Gulf of Mexico Atmospheric Composition and Climate Study (GoMACCS). *J. Geophys. Res.: Atmos.* **112**, (2007).
22. Chen Y, *et al.* Low-Molecular-Weight Carboxylic Acids in the Southeastern U.S.: Formation, Partitioning, and Implications for Organic Aerosol Aging. *Environ. Sci. Technol.* **55**, 6688–6699 (2021).
23. Wang J, *et al.* Enhanced aqueous-phase formation of secondary organic aerosols due to the regional biomass burning over North China Plain. *Environ. Pollut.* **256**, 113401 (2020).
24. Carlton AG, *et al.* Atmospheric oxalic acid and SOA production from glyoxal: Results of aqueous photooxidation experiments. *Atmos. Environ.* **41**, 7588–7602 (2007).
25. Lim YB, Tan Y, Turpin BJ. Chemical insights, explicit chemistry, and yields of secondary organic aerosol from OH radical oxidation of methylglyoxal and glyoxal in the aqueous phase. *Atmos. Chem. Phys.* **13**, 8651–8667 (2013).
26. Bikkina S, Kawamura K, Sarin M. Secondary organic aerosol formation over coastal ocean: inferences from atmospheric water-soluble low molecular weight organic compounds. *Environ. Sci. Technol.* **51**, 4347–4357 (2017).
27. Kawamura K, Ikushima K. Seasonal changes in the distribution of dicarboxylic acids in the urban atmosphere. *Environ. Sci. Technol.* **27**, 2227–2235 (1993).
28. Kawamura K, Sakaguchi F. Molecular distributions of water soluble dicarboxylic acids in marine aerosols over the Pacific Ocean including tropics. *J. Geophys. Res.: Atmos.* **104**, 3501–3509 (1999).
29. Dasari S, *et al.* Photochemical degradation affects the light absorption of water-soluble brown carbon in the South Asian outflow. *Sci. Adv.* **5**, eaau8066 (2019).
30. Hodas N, *et al.* Aerosol liquid water driven by anthropogenic nitrate: implications for lifetimes of water-soluble organic gases and potential for secondary organic aerosol formation. *Environ. Sci. Technol.* **48**, 11127–11136 (2014).
31. Wu Z, *et al.* Aerosol liquid water driven by anthropogenic inorganic salts: implying its key role in haze formation over the North China Plain. *Environ. Sci. Technol. Lett.* **5**, 160–166 (2018).
32. Gkatzelis GI, *et al.* Uptake of water-soluble gas-phase oxidation products drives organic particulate pollution in Beijing. *Geophys. Res. Lett.* **48**, e2020GL091351 (2021).
33. Kawamura K, Gagosian RB. Implications of ω -oxocarboxylic acids in the remote marine atmosphere for photo-oxidation of unsaturated fatty acids. *Nature* **325**, 330–332 (1987).
34. Ballentine DC, Macko SA, Turekian VC. Variability of stable carbon isotopic compositions in individual fatty acids from combustion of C4 and C3 plants: implications for biomass burning. *Chem. Geol.* **152**, 151–161 (1998).

35. Miyazaki Y, Kawamura K, Jung J, Furutani H, Uematsu M. Latitudinal distributions of organic nitrogen and organic carbon in marine aerosols over the western North Pacific. *Atmos. Chem. Phys.* **11**, 3037–3049 (2011).
36. Martinelli LA, Camargo PB, Lara L, Victoria RL, Artaxo P. Stable carbon and nitrogen isotopic composition of bulk aerosol particles in a C4 plant landscape of southeast Brazil. *Atmos. Environ.* **36**, 2427–2432 (2002).
37. Goldstein AH, Shaw SL. Isotopes of volatile organic compounds: An emerging approach for studying atmospheric budgets and chemistry. *Chem. Rev.* **103**, 5025–5048 (2003).
38. Rudolph J, *et al.* The stable carbon isotope ratio of biogenic emissions of isoprene and the potential use of stable isotope ratio measurements to study photochemical processing of isoprene in the atmosphere. *J. Atmos. Chem.* **44**, 39–55 (2003).
39. Bikkina S, Kawamura K, Sakamoto Y, Hirokawa J. Low molecular weight dicarboxylic acids, oxocarboxylic acids and α -dicarbonyls as ozonolysis products of isoprene: Implication for the gaseous-phase formation of secondary organic aerosols. *Science of The Total Environment* **769**, 144472 (2021).
40. Carlton AG, Turpin BJ. Particle partitioning potential of organic compounds is highest in the Eastern US and driven by anthropogenic water. *Atmos. Chem. Phys.* **13**, 10203–10214 (2013).
41. Zhang Y-L, Kawamura K, Cao F, Lee M. Stable carbon isotopic compositions of low-molecular-weight dicarboxylic acids, oxocarboxylic acids, α -dicarbonyls, and fatty acids: Implications for atmospheric processing of organic aerosols. *J. Geophys. Res.: Atmos.* **121**, 3707–3717 (2016).
42. Zhang Q, *et al.* Drivers of improved PM2.5 air quality in China from 2013 to 2017. *Proc. Natl Acad. Sci. USA* **116**, 24463–24469 (2019).
43. Pavuluri CM, Kawamura K, Swaminathan T. Water-soluble organic carbon, dicarboxylic acids, ketoacids, and α -dicarbonyls in the tropical Indian aerosols. *J. Geophys. Res.* **115**, (2010).
44. Hatakeyama S, *et al.* Ozone-cyclohexene reaction in air: quantitative analysis of particulate products and the reaction mechanism. *Environ. Sci. Technol.* **19**, 935–942 (1985).
45. Kawamura K, Kasukabe H, Barrie LA. Source and reaction pathways of dicarboxylic acids, ketoacids and dicarbonyls in arctic aerosols: One year of observations *Atmos. Environ.* **30**, 1709–1722 (1996).
46. Yuan B, Liu Y, Shao M, Lu S, Streets DG. Biomass burning contributions to ambient VOCs species at a receptor site in the Pearl River Delta (PRD), China. *Environ. Sci. Technol.* **44**, 4577–4582 (2010).
47. Ding X, *et al.* Tracer-based estimation of secondary organic carbon in the Pearl River Delta, south China. *J. Geophys. Res.: Atmos.* **117**, (2012).
48. Lim H-J, Carlton AG, Turpin BJ. Isoprene forms secondary organic aerosol through cloud processing: Model simulations. *Environ. Sci. Technol.* **39**, 4441–4446 (2005).
49. Carlton AG, *et al.* CMAQ model performance enhanced when in-cloud secondary organic aerosol is included: comparisons of organic carbon predictions with measurements. *Environ. Sci. Technol.* **42**, 8798–8802 (2008).

50. Baltensperger U, *et al.* Secondary organic aerosols from anthropogenic and biogenic precursors. *Faraday Discuss.* **130**, 265–278 (2005).
51. Updyke KM, Nguyen TB, Nizkorodov SA. Formation of brown carbon via reactions of ammonia with secondary organic aerosols from biogenic and anthropogenic precursors. *Atmos. Environ.* **63**, 22–31 (2012).
52. Wozniak AS, Bauer JE, Dickhut RM, Xu L, McNichol AP. Isotopic characterization of aerosol organic carbon components over the eastern United States. *J. Geophys. Res. Atmos* **117**, (2012).
53. Zhang Y-L, *et al.* Large contribution of fossil fuel derived secondary organic carbon to water soluble organic aerosols in winter haze in China. *Atmos. Chem. Phys.* **18**, 4005–4017 (2018).
54. Mo Y, *et al.* Dual carbon isotope-based source apportionment and light absorption properties of water-soluble organic carbon in PM_{2.5} over China. *J. Geophys. Res.: Atmos.*, e2020JD033920 (2021).
55. Liu J, *et al.* Source apportionment and dynamic changes of carbonaceous aerosols during the haze bloom-decay process in China based on radiocarbon and organic molecular tracers. *Atmos. Chem. Phys.* **16**, 2985–2996 (2016).
56. Liu J, *et al.* Radiocarbon-derived source apportionment of fine carbonaceous aerosols before, during, and after the 2014 Asia-Pacific Economic Cooperation (APEC) summit in Beijing, China. *J. Geophys. Res.: Atmos.* **121**, 4177–4187 (2016).
57. Zhang Y-I, *et al.* Micro-scale (μg) radiocarbon analysis of water-soluble organic carbon in aerosol samples. *Atmos. Environ.* **97**, 1–5 (2014).
58. Kirillova EN, *et al.* Water-soluble organic carbon aerosols during a full New Delhi winter: Isotope-based source apportionment and optical properties. *J. Geophys. Res.: Atmos.* **119**, 3476–3485 (2014).
59. Glassmeier F, *et al.* Aerosol-cloud-climate cooling overestimated by ship-track data. *Science* **371**, 485–489 (2021).
60. Boreddy SKR, Kawamura K. Investigation on the hygroscopicity of oxalic acid and atmospherically relevant oxalate salts under sub- and supersaturated conditions. *Environ. Sci.: Processes Impacts* **20**, 1069–1080 (2018).
61. Nguyen TKV, Zhang Q, Jimenez JL, Pike M, Carlton AG. Liquid water: ubiquitous contributor to aerosol mass. *Environ. Sci. Technol. Lett.* **3**, 257–263 (2016).
62. Wang Y, *et al.* Mutual promotion between aerosol particle liquid water and particulate nitrate enhancement leads to severe nitrate-dominated particulate matter pollution and low visibility. *Atmos. Chem. Phys.* **20**, 2161–2175 (2020).
63. Xing M, *et al.* Vapor isotopic evidence for the worsening of winter air quality by anthropogenic combustion-derived water. *Proc. Natl Acad. Sci. USA* **117**, 33005-33010 (2020).
64. Kong L, *et al.* Aerosol optical properties under different pollution levels in the Pearl River Delta (PRD) region of China. *J. Environ. Sci. (Beijing, China)* **87**, 49–59 (2020).

65. Guo H, *et al.* Fine-particle water and pH in the southeastern United States. *Atmos. Chem. Phys.* **15**, 5211–5228 (2015).

Figures

Figure 1

The average aerosol optical depth (AOD) at 550 nm during June 2017 to May 2018 over East Asia with indication of the location of the sampling site (Heshan) and the mean air mass back trajectories (BT) based on HYSPLIT cluster analysis. The BT show two dominant air mass transport pathways: continental outflow (62% of the total clusters) and oceanic air masses (38% of total clusters). See Supplementary Fig. 2 for the 72-h BT with a 6-h interval and Supplementary Fig. 3 for relative cluster contribution. The bar charts depict the mass concentrations of total carbonaceous aerosol [TCA = organic matter (OM) + elemental carbon (EC)], anthropogenic WSIC [WSIC_{anth} = non-sea-salt (nss) K⁺ + nss-SO₄²⁻ + NH₄⁺ + NO₃⁻ + nss-Cl⁻], and aerosol liquid water (ALW). The AOD data was obtained from Moderate Resolution Imaging Spectroradiometer (MODIS) observations (<https://giovanni.gsfc.nasa.gov/giovanni/>)

Figure 2

Year-round variability in mass concentration, $\delta^{13}\text{C}$ composition, and ^{14}C -based source apportionment (fraction of non-fossil) for individual oxalic acid in $\text{PM}_{2.5}$ samples collected from June 2017 to May 2018 at regional receptor site (Heshan city). The unshaded periods represent coastal background air mass regime, while the shadowed periods represent continental outflow air mass regime. Red and blue bars represent the concentration of oxalic acid during coastal background air mass regime and continental outflow air mass regime, respectively. The orange and green symbols show the $\delta^{13}\text{C}$ composition and ^{14}C -based fraction of non-fossil for oxalic acid.

Figure 3

Stable carbon isotopic evidence of aqueous secondary organic aerosol (aqSOA) formation. (a) Average $\delta^{13}\text{C}$ values of dicarboxylic acids (C₂-C₉) and oxocarboxylic acids (ωC_2 and Pyr) for the two air mass regimes. Whiskers indicate the 95% confidence interval of $\delta^{13}\text{C}$ values in each air mass regime. Pink shadows represent the compounds for which the $\delta^{13}\text{C}$ values are statistically significantly different ($p < 0.05$) between the two air mass origins. For abbreviations, see Supplementary Table 3. (b) Relationships

between oxalic acid mass concentration and aerosol liquid water (ALW). (c) Relationships between $\delta^{13}\text{C}$ -oxalic acid and mass concentration of water-soluble organic carbon (WSOC) as a proxy for aqSOA abundance.

Figure 4

Radiocarbon-based source apportionment of dicarboxylic acids and related compounds. (a) Relationships between biogenic/biomass burning (bio/bb) source-derived oxalic acid vs. levoglucosan. (b) Relationships between bio/bb source-derived oxalic acid vs. biogenic SOA tracer (include isoprene SOA tracers and α/β -pinene SOA tracers; Supplementary Text 1). (c) Monthly variability of non-fossil contribution to oxalic acid, malonic acid, succinic acid, glyoxylic acid, and methylglyoxal. (d) Relative contribution of biogenic emission, biomass burning, and fossil-fuel emission to oxalic acid in coastal background and continental outflow.

Figure 5

Two-dimensional dual-carbon isotope characterization of individual oxalic acid for coastal background and continental outflow. The color of the symbols represents the oxalic acid concentration. The $\delta^{13}\text{C}$ source signatures of oxalic acid precursor gases from biogenic VOCs (BVOCs; i.e., isoprene) and from anthropogenic VOCs (AVOCs; i.e., anthropogenic emitted non-methane hydrocarbons) are obtained from reported literature values^{37, 38}.

Figure 6

The $\delta^{13}\text{C}$ composition (inserted triangular symbols) and $\Delta^{14}\text{C}$ -based source apportionment (bars) of WSOC and oxalic acid in PM_{2.5} collected during winter (January 2018) and summer (July 2018) at urban sites of Beijing (BJ), Shanghai (SH), Guangzhou (GZ), Chengdu (CD), and Wuhan (WH). The central map presents locations of the five sites and aerosol optical depth (AOD) at 550 nm during the sampling period.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supportingmaterial.docx](#)