Synthesis and Biological Evaluation of Prodrugs For Nitroreductase Based 4- β -Amino-4'-Demethylepipodophyllotoxin As Potential Anticancer Agents

Zhengrong Wu (✉ wuzhengrong@lzu.edu.cn)

Lanzhou University

Wei Deng

Lanzhou Medical College: Lanzhou University

Dian He

Lanzhou Medical College: Lanzhou University

Research Article

Keywords: Podophyllotoxin, O-nitrophenoxycetyl acid, Cell cycle arrest, Apoptosis, Structure-Activity Relationship

Posted Date: December 22nd, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1158885/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.
[Read Full License](#)

Version of Record: A version of this preprint was published at Medicinal Chemistry Research on January 24th, 2022. See the published version at <https://doi.org/10.1007/s00044-022-02847-5>.

Abstract

A series of prodrugs for nitroreductase based 4- β -amino-4'-Demethylepipodophyllotoxin as potential anticancer agents were synthesized, and their antiproliferative activities in vitro showed compounds **2b** (IC_{50} =0.77, 0.83 and 1.19 μ M) and **2d** (IC_{50} =0.98, 0.91 and 1.58 μ M) were greatly selectively toxic to tumor cells A-549, HeLa and HepG2, respectively, and lower damage to normal WI-38 cells in comparison with positive agent Etoposide and Demethylepipodophyllotoxin, and induced cell cycle arrest in the G2/M phase with a concomitant decrease in the population of G1 phase in HeLa cells, which were accompanied by apoptosis. Furthermore, Molecular docking model showed that compounds **2b** and **2d** appeared to form stable bonds with NTR 1DS7. Taken together, these conjugates have the potential to be developed as anti-tumor drugs.

Introduction

With the rapid growth of cancer cases across the globe, a great number of approaches are being investigated to develop an effective cure for this deadly disease [1]. Among various approaches, naturally occurring substances account for the abundant sources to identify novel hits and leads. However, a number of antitumor naturally-occurring substances do not show target specificity, which may even show adverse toxicity to non-carcinoma cells. In order to reduce the cytotoxicity of the natural compounds, appropriate structural modification is the vital way and has been adopted to develop new drug [2].

Podophyllotoxin is a natural occurring aryltetralin lignin with pronounced cytotoxic activity. However, its clinical application for cancer treatment has been blocked due to its poor water solubility and selectivity [3]. For reducing cytotoxicity and increasing water solubility of the bioactive compound, a great number of structural modification have been adopted to address these limitations and obtain better therapeutic agents [4–9], and these efforts have led to the structure–activity relationships (SAR) unambiguously demonstrate that C4 is the major molecular area tolerable to significant structural diversification [10]. Therefore, most of the current derivatization on Podophyllotoxin focuses on C-4 position. Recently, A series of 4 β -N substituted 4'-*O*-demethyl-4-deoxypodophyllotoxins have been synthesized, and exhibited significant anticancer activity [11–15], Among these, the substituent of *p*-nitrobenzylpiperazine markedly affected the activity profiles of this compound class, and may serve as a potential anti-cancer drug [16].

Prodrug strategies based on nitroreductases (NTR) have been extensively studied in recent years. NTR can catalyze nitro-substituted aromatic compounds into aromatic amines and undergoes internal cyclization to a non-toxic derivative, which reduces serious side effects on healthy cells and increases concentration of drugs at cancer cells is remarkable. So the discoveries of new prodrugs/NTR combinations which are of increasing interest as potential cancer therapeutics [17–22].

O-nitrophenoxyacetyl acid have known as potential inhibitors of vascular endothelial growth factor (VEGF) in cancer chemotherapy. NTR can catalyze nitro substituent into aromatic amines and undergoes

internal cyclization to a non-toxic derivative, which reduces serious side effects on healthy cells and increases concentration of drugs at cancer cells is remarkable [23].

Inspired by the above background, herein, we present the design, synthesis, of a novel series of 4 β -N-(4-Nitrophenyl piperazinyl)-4'-*O*-demethyl-4- deoxypodophyllotoxin and four different *O*-nitrophenoxyacetyl acids(substituent of electron-donating groups, electron-withdrawing groups, steric groups, comprehensive groups, respectively) hybrid and evaluation of their cytotoxic activity against several cancer cell lines.

Results And Discussion

Synthesis

The synthetic route to the target compounds is shown in Scheme 1. The *O*-nitrophenoxyacetyl acids were prepared according to our previous report [24]. 4 β -N-(4-Nitrophenyl piperazinyl)-4'-*O*-demethyl-4- deoxypodophyllotoxin was according to the procedure described previous [16]. The chemical structures of the final compounds were fully characterized by ¹H NMR, ¹³C NMR and HRMS spectroscopic techniques.

Cytotoxicity

The in vitro cytotoxicities of conjugates 2a–2t were evaluated against a panel of three human cancer cell lines (lung carcinoma A-549, cervical carcinoma HeLa, hepatocellular carcinoma HepG2, and human embryonic lung fibroblasts WI-38 cells), with Etoposide and Demethylepipodophyllotoxin as reference compounds. In addition, their toxicity evaluation was tested on normal embryonic lung fibroblast WI-38 cells. The screening procedure was based on the standard MTT growth inhibition assay, and the results are summarized in Table 1. As shown, most of the compounds were more potent against three tumor cell, these compounds were most effective in A549 cells and HeLa cells, and had lowest potency in HepG2 cells. The IC₅₀ value of compounds **2b** (0.77, 0.83 and 1.19 μ M) and **2d** (0.98, 0.91 and 1.58 μ M), and which were comparable to positive agent Etoposide and more potent than the positive agent Demethylepipodophyllotoxin (9.64, 15.29 and 12.73 μ M) for HeLa, HepG2 and A549 cells. In comparison, compounds **2b** (IC₅₀=688.54 μ M) and **2d** (IC₅₀=477.96 μ M) showed greater selectivity, and were relatively less active toward normal WI-38 cells in comparison with positive agent Etoposide (IC₅₀<100 μ M) and Demethylepipodophyllotoxin (IC₅₀<100 μ M). Based on these results, it was possible to deduce some preliminary SAR. First of all, *O*-nitrophenoxyacetyl acids with electron-donating group substitution (2a-2e) appeared to be more potent than those with electron-withdrawing (2f-2k), steric (2l-2o), and comprehensive (2p-2t) groups.

Table 1
Cytotoxicity of target compounds to different cells following 24h of exposure.

Compds.	Cytotoxicity (IC ₅₀ , μ M) ^a			
	A-549	Hela	Hep-G2	WI-38
2a	5.7 $\pm$ 0.73	22.92 $\pm$ 2.37	13.25 $\pm$ 2.53	583.78 $\pm$ 61.57
2b	1.19 $\pm$ 0.67	0.77$\pm$0.76	0.83$\pm$0.37	688.54 $\pm$ 70.58
2c	21.99 $\pm$ 2.7	12.93 $\pm$ 1.53	19.31 $\pm$ 0.84	>1000
2d	1.58 $\pm$ 0.81	0.98 $\pm$ 0.74	0.91 $\pm$ 0.45	477.96 $\pm$ 32.84
2e	13.81 $\pm$ 3.93	18.55 $\pm$ 0.71	15.96 $\pm$ 2.23	>1000
2f	16.87 $\pm$ 1.18	18.94 $\pm$ 1.72	20.67 $\pm$ 3.62	>1000
2g	17.64 $\pm$ 2.15	16.82 $\pm$ 0.36	28.63 $\pm$ 2.81	>1000
2h	27.44 $\pm$ 1.55	9.75 $\pm$ 2.42	23.42 $\pm$ 4.07	>1000
2i	8.29 $\pm$ 0.84	10.44 $\pm$ 1.38	18.91 $\pm$ 3.09	744.26 $\pm$ 89.42
2j	16.54 $\pm$ 1.38	19.38 $\pm$ 2.14	22.55 $\pm$ 1.85	>1000
2k	8.81 $\pm$ 1.15	5.49 $\pm$ 0.67	16.12 $\pm$ 2.60	656.67 $\pm$ 85.32
2l	7.08 $\pm$ 0.97	4.8 $\pm$ 0.36	19.87 $\pm$ 1.90	508.54 $\pm$ 42.67
2m	6.32 $\pm$ 1.16	10.02 $\pm$ 0.62	10.21 $\pm$ 0.95	727.23 $\pm$ 67.76
2n	7.26 $\pm$ 0.25	8.84 $\pm$ 0.34	15.19 $\pm$ 3.10	675.45 $\pm$ 37.87
2o	18.72 $\pm$ 1.89	10.14 $\pm$ 1.00	30.68 $\pm$ 2.19	>1000
2p	19.82 $\pm$ 1.03	23.21 $\pm$ 1.33	19.68 $\pm$ 1.45	>1000
2q	30.46 $\pm$ 4.87	20.31 $\pm$ 0.27	14.35 $\pm$ 1.41	>1000
2r	28.29 $\pm$ 2.03	17.99 $\pm$ 1.38	14.64 $\pm$ 1.12	421.42 $\pm$ 2.10
2s	18.32 $\pm$ 1.55	15.44 $\pm$ 1.36	20.37 $\pm$ 2.02	575.11 $\pm$ 36.26
2t	22.21 $\pm$ 2.43	17.28 $\pm$ 1.72	18.88 $\pm$ 1.14	>1000
Etoposide	1.42 $\pm$ 0.11	2.64 $\pm$ 0.45	1.64 $\pm$ 0.32	<100
DEPPT	12.73 $\pm$ 1.19	9.64 $\pm$ 0.58	15.29 $\pm$ 0.17	<100
Date are the mean of three indepdent experiments and recorded as the mean of standard deviation (x $\pm$ SD, n=3).				
DEPPT is Etoposide Demethylepipodophyllotoxin.				

Cell cycle arrest and apoptotic assay

The data above revealed that compounds **2b** and **2d** the most potent and selective were further investigate the effects on cell cycle progression by means of fluorescence-activated cell sorting analysis of HCT-116 cells stained with propidium iodide.

As shown in Fig. 1, treatment with compounds **2b** and **2d** led to a dose-dependent accumulation of cells in the G2/M phase with a concomitant decrease in the population of G1 phase cells respectively, compared with untreated cultures and Etoposide. Similar effects also occurred with the Annexin V/PI apoptosis detection (Fig. 2). These results demonstrated that compounds **2b** and **2d** were interfering with cell proliferation by arresting the cell cycle and induced G2/M arrest accompanied by apoptosis in HeLa cells.

Molecular docking assay

To prove the anticancer mechanism of the action, a molecular docking investigation was undertaken. As shown in Fig. 3, which revealed that compounds **2b** and **2d** were well inserted into the two active pockets of NTR 1DS7, Hydrogen bonds were shown with yellow dash lines and dark green spheres. The hydrophobic interactions which are van der Waals were represented by light green spheres, these results reveal that compounds **2b** and **2d** having hydrophobic property, good membrane permeability and specifically targeting NTR. The binding forms are demonstrated that the para nitro and carbonyl groups in phenyl ring were able to affect the interaction on NTR significantly. These data provide certain theoretical support for experimental results.

Conclusions

In conclusion, we have designed and synthesized a number of prodrugs for nitroreductase based 4- β -amino-4'-Demethylepipodophyllotoxin as potential anticancer agents were synthesized, and their antiproliferative activities in vitro showed compounds **2b** and **2d** were greatly selectively toxic to tumor cells A-549, HeLa and HepG2, respectively, and lower damage to normal WI-38 cells in comparison with positive agent Etoposide and Demethylepipodophyllotoxin, and induced cell cycle arrest in the G2/M phase with a concomitant decrease in the population of G1 phase in HeLa cells, which were accompanied by apoptosis. Furthermore, Molecular docking model showed that compounds **2b** and **2d** appeared to form stable bonds with NTR 1DS7. Taken together, these conjugates have the potential to be developed as anti-tumor drugs. Further studies on the relevant action mechanisms and structural modification of identified hits are on-going.

Materials And Methods

Chemistry

All reactions were performed with commercially available reagents and solvents without further purification. All reactions were monitored by thin-layer chromatography (TLC). ^1H and ^{13}C NMR spectra were recorded on Bruker AVANCE-III HD 400MHz or NEO 500MHz (Bruker Daltonics Inc., Germany) spectrometers using TMS as a reference. Mass spectra were recorded on a Bruker APEXII49e spectrometer (Bruker Daltonics Inc., Germany) with ESI source as ionization.

Data for Compound 2-(3-methyl-2-nitrophenoxy)acetic acid-4 β -N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2a). ^1H NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ : 8.06 (d, $J=9.2\text{Hz}$, 2H), 7.49 (t, $J=8.0\text{Hz}$, 1H), 7.43 (s, 1H), 7.1(d, $J=8.4\text{Hz}$, 1H), 7.06 (d, $J=9.2\text{ Hz}$, 1H), 7.02 (d, $J=9.2\text{ Hz}$, 2H), 6.58 (s, 1H), 6.38 (s, 2H), 6.01 (s, 2H), 5.32 (s, 2H), 4.63 (d, $J=7.2\text{ Hz}$, 1H), 4.14 (dd, $J_1=16.0\text{ Hz}$, $J_2=12.0\text{ Hz}$, 3H), 3.65 (s, 6H), 3.51-3.36 (m, 4H), 3.18 (dd, $J_1=12.0\text{ Hz}$, $J_2=4.0\text{ Hz}$, 1H), 2.63-2.51 (m, 5H), 2.37 (s, 3H); ^{13}C NMR (400 MHz, DMSO- d_6 , TMS, ppm) δ : 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI): 783.2536 for $[\text{M}+\text{H}]^+$ (calculated 783.7597 for $\text{C}_{40}\text{H}_{38}\text{N}_4\text{O}_{13}$).

Data for Compound 2-(4-methyl-2-nitrophenoxy)acetic acid-4 β -N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2b). ^1H NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ : 8.07 (d, $J=9.2\text{Hz}$, 2H), 7.72 (s, 1H), 7.51 (d, $J=8.4\text{ Hz}$, 1H), 7.42 (s, 1H), 7.20 (d, $J=8.4\text{ Hz}$, 1H), 7.04 (d, $J=9.2\text{ Hz}$, 2H), 6.57 (s, 1H), 6.38 (s, 2H), 6.01 (s, 2H), 5.30 (s, 2H), 4.59 (d, $J=7.2\text{ Hz}$, 1H), 4.14 (dd, $J_1=16.0\text{ Hz}$, $J_2=12.0\text{ Hz}$, 3H), 3.65 (s, 6H), 3.51-3.35 (m, 4H), 3.18 (dd, $J_1=12.0\text{ Hz}$, $J_2=4.0\text{ Hz}$, 1H), 2.63-2.50 (m, 5H), 2.31 (s, 3H); ^{13}C NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI): 783.2562 for $[\text{M}+\text{H}]^+$ (calculated 783.7597 for $\text{C}_{40}\text{H}_{38}\text{N}_4\text{O}_{13}$).

Data for Compound 2-(5-methyl-2-nitrophenoxy)acetic acid-4 β -N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2c). ^1H NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ : 8.06(d, $J=9.2\text{Hz}$, 2H), 7.82 (d, $J=8.4\text{ Hz}$, 1H), 7.42 (s, 1H), 7.12 (s, 1H), 7.04 (d, $J=9.2\text{ Hz}$, 2H), 6.99 (d, $J=8.0\text{ Hz}$, 1H), 6.57 (s, 1H), 6.39 (s, 2H), 6.01 (s, 2H), 5.33 (s, 2H), 4.59 (d, $J=7.2\text{ Hz}$, 1H), 4.14 (dd, $J_1=16.0\text{ Hz}$, $J_2=12.0\text{ Hz}$, 3H), 3.66 (s, 6H), 3.51-3.34 (m, 4H), 3.18 (dd, $J_1=12.0\text{ Hz}$, $J_2=4.0\text{ Hz}$, 1H), 2.62-2.50 (m, 5H), 2.35 (s, 3H); ^{13}C NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI): 783.2472 for $[\text{M}+\text{H}]^+$ (calculated 783.7597 for $\text{C}_{40}\text{H}_{38}\text{N}_4\text{O}_{13}$).

Data for Compound 2-(5-methyl-2-nitrophenoxy)acetic acid-4 β -N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2c). ^1H NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ : 8.06(d, $J=9.2\text{Hz}$, 2H), 7.82 (d, $J=8.4\text{ Hz}$, 1H), 7.42 (s, 1H), 7.12 (s, 1H), 7.04 (d, $J=9.2\text{ Hz}$, 2H), 6.99 (d, $J=8.0\text{ Hz}$, 1H), 6.57 (s,

1H), 6.39 (s, 2H), 6.01 (s, 2H), 5.33 (s, 2H), 4.59 (d, $J=7.2$ Hz, 1H), 4.14 (dd, $J_1=16.0$ Hz, $J_2=12.0$ Hz, 3H), 3.66 (s, 6H), 3.51-3.34 (m, 4H), 3.18 (dd, $J_1=12.0$ Hz, $J_2=4.0$ Hz, 1H), 2.62-2.50 (m, 5H), 2.35 (s, 3H); ^{13}C NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI): 783.2472 for $[\text{M}+\text{H}]^+$ (calculated 783.7597 for $\text{C}_{40}\text{H}_{38}\text{N}_4\text{O}_{13}$).

Data for Compound 2-(4-methoxy-2-nitrophenoxy)acetic acid-4 β -N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2d). ^1H NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ : 8.06 (d, $J=9.2$ Hz, 2H), 7.47 (s, 1H), 7.42 (s, 1H), 7.28 (d, $J=8.0$ Hz, 2H), 7.04 (d, $J=8.0$ Hz, 2H), 6.57 (s, 1H), 6.38 (s, 2H), 6.01 (s, 2H), 5.25 (s, 2H), 4.59 (d, $J=4.0$ Hz, 1H), 4.14 (dd, $J_1=16.0$ Hz, $J_2=12.0$ Hz, 3H), 3.79 (s, 3H), 3.64 (s, 6H), 3.51-3.35 (m, 4H), 3.18 (dd, $J_1=12.0$ Hz, $J_2=4.0$ Hz, 1H), 2.63-2.51 (m, 5H); ^{13}C NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI): 799.2507 for $[\text{M}+\text{H}]^+$ (calculated 799.7582 for $\text{C}_{40}\text{H}_{38}\text{N}_4\text{O}_{14}$).

Data for Compound 2-(5-methoxy-2-nitrophenoxy)acetic acid-4 β -N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2e). ^1H NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ : 8.02 (d, $J=8.0$ Hz, 2H), 7.94 (d, $J=8.0$ Hz, 1H), 7.38 (s, 1H), 7.00 (d, $J=8.0$ Hz, 2H), 6.73 (d, $J=9.2$ Hz, 2H), 6.53 (s, 1H), 6.35 (s, 2H), 5.97 (s, 2H), 5.33 (s, 2H), 4.58 (d, $J=4.0$ Hz, 1H), 4.09 (dd, $J_1=8.0$ Hz, $J_2=4.0$ Hz, 3H), 3.85 (s, 3H), 3.60 (s, 6H), 3.47-3.31 (m, 4H), 3.14 (dd, $J_1=12.0$ Hz, $J_2=4.0$ Hz, 1H), 2.61-2.46 (m, 5H); ^{13}C NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI): 799.2469 for $[\text{M}+\text{H}]^+$ (calculated 799.7582 for $\text{C}_{40}\text{H}_{38}\text{N}_4\text{O}_{14}$).

Data for Compound 2-(3-fluoro-2-nitrophenoxy)acetic acid-4 β -N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2f). ^1H NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ : 8.04 (d, $J=9.2$ Hz, 2H), 7.89 (dd, $J_1=8.0$ Hz, $J_2=4.0$ Hz, 1H), 7.66-7.62 (m, 1H), 7.41 (s, 1H), 7.36 (dd, $J_1=12.0$ Hz, $J_2=4.0$ Hz, 1H), 7.02 (d, $J=9.2$ Hz, 2H), 6.55 (s, 1H), 6.37 (s, 2H), 5.99 (s, 2H), 5.32 (s, 2H), 4.57 (d, $J=4.4$ Hz, 1H), 4.12 (dd, $J_1=16.0$ Hz, $J_2=8.0$ Hz, 3H), 3.63 (s, 6H), 3.49-3.35 (m, 4H), 3.16 (dd, $J_1=12.0$ Hz, $J_2=4.0$ Hz, 1H), 2.62-2.49 (m, 5H); ^{13}C NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI): 787.2260 for $[\text{M}+\text{H}]^+$ (calculated 787.7220 for $\text{C}_{39}\text{H}_{35}\text{FN}_4\text{O}_{13}$).

Data for Compound 2-(4-fluoro-2-nitrophenoxy)acetic acid-4 β -N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2g). ^1H NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ : 8.05 (d, $J=8.0$ Hz, 2H), 7.68 (dd, $J_1=16.0$ Hz, $J_2=8.0$ Hz, 1H), 7.41 (s, 1H), 7.19 (dd, $J_1=24.0$ Hz, $J_2=12.0$ Hz, 2H), 7.02 (d, $J=9.2$ Hz, 2H), 6.56

(s, 1H), 6.37 (s, 2H), 5.99 (s, 2H), 5.38 (s, 2H), 4.57 (d, $J=4.4$ Hz, 1H), 4.12 (dd, $J_1=16.0$ Hz, $J_2=8.0$ Hz, 3H), 3.64 (s, 6H), 3.62-3.45 (m, 4H), 3.16 (dd, $J_1=12.0$ Hz, $J_2=4.0$ Hz, 1H), 2.62-2.49 (m, 5H); ^{13}C NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI): 787.2280 for $[\text{M}+\text{H}]^+$ (calculated 787.7220 for $\text{C}_{39}\text{H}_{35}\text{FN}_4\text{O}_{13}$).

Data for Compound 2-(5-fluoro-2-nitrophenoxy)acetic acid-4 β -N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2h). ^1H NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ : 8.02 (d, $J=8.0$ Hz, 2H), 7.76 (dd, $J_1=16.0$ Hz, $J_2=8.0$ Hz, 1H), 7.39 (s, 1H), 7.19 (dd, $J_1=24.0$ Hz, $J_2=12.0$ Hz, 2H), 7.02 (d, $J=9.2$ Hz, 2H), 6.53 (s, 1H), 6.35 (s, 2H), 5.97 (s, 2H), 5.39 (s, 2H), 4.58 (d, $J=4.4$ Hz, 1H), 4.12 (dd, $J_1=16.0$ Hz, $J_2=8.0$ Hz, 3H), 3.61 (s, 6H), 3.48-3.29 (m, 4H), 3.16 (dd, $J_1=12.0$ Hz, $J_2=4.0$ Hz, 1H), 2.62-2.49 (m, 5H); ^{13}C NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI): 787.2286 for $[\text{M}+\text{H}]^+$ (calculated 787.7220 for $\text{C}_{39}\text{H}_{35}\text{FN}_4\text{O}_{13}$).

Data for Compound 2-(4-bromo-2-nitrophenoxy)acetic acid-4 β -N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2i). ^1H NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ : 8.14 (s, 1H), 8.05 (d, $J=8.0$ Hz, 2H), 7.91 (d, $J=8.0$ Hz, 1H), 7.41 (s, 1H), 7.28 (d, $J=12.0$ Hz, 1H), 7.03 (d, $J=12.0$ Hz, 2H), 6.55 (s, 1H), 6.37 (s, 2H), 6.00 (s, 2H), 5.36 (s, 2H), 4.57 (d, $J=4.0$ Hz, 1H), 4.12 (dd, $J_1=16.0$ Hz, $J_2=8.0$ Hz, 3H), 3.64 (s, 6H), 3.50-3.33 (m, 4H), 3.17 (dd, $J_1=16.0$ Hz, $J_2=4.0$ Hz, 1H), 2.62-2.49 (m, 5H); ^{13}C NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI): 847.1464 for $[\text{M}+\text{H}]^+$ (calculated 847.1383 for $\text{C}_{39}\text{H}_{35}\text{BrN}_4\text{O}_{13}$).

Data for Compound 2-(5-bromo-2-nitrophenoxy)acetic acid-4 β -N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2j). ^1H NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ : 8.05 (d, $J=8.0$ Hz, 2H), 7.87 (d, $J=8.4$ Hz, 1H), 7.54 (s, 1H), 7.40 (d, $J=8.0$ Hz, 2H), 7.02 (d, $J=8.0$ Hz, 2H), 6.55 (s, 1H), 6.38 (s, 2H), 5.99 (s, 2H), 5.45 (s, 2H), 4.57 (d, $J=4.0$ Hz, 1H), 4.12 (dd, $J_1=16.0$ Hz, $J_2=8.0$ Hz, 3H), 3.66 (s, 6H), 3.49-3.37 (m, 4H), 3.17 (dd, $J_1=16.0$ Hz, $J_2=4.0$ Hz, 1H), 2.62-2.49 (m, 5H); ^{13}C NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI): 847.1489 for $[\text{M}+\text{H}]^+$ (calculated 847.1383 for $\text{C}_{39}\text{H}_{35}\text{BrN}_4\text{O}_{13}$).

Data for Compound 2-(5-chloro-2-nitrophenoxy)acetic acid-4 β -N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2k). ^1H NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ : 8.07 (d, $J=9.2$ Hz, 2H), 7.97 (d, $J=8.8$ Hz, 1H), 7.43 (d, $J=3.2$ Hz, 2H), 7.38 (d, $J=8.8$ Hz, 1H), 7.04 (d, $J=9.2$ Hz, 2H), 6.57 (s, 1H), 6.39 (s, 2H), 6.01 (s, 2H), 5.46 (s, 2H), 4.59 (d, $J=4.4$ Hz, 1H), 4.14 (dd, $J_1=16.0$ Hz, $J_2=12.0$ Hz, 3H), 3.66 (s, 6H), 3.52-3.36 (m, 4H), 3.18 (dd, $J_1=12.0$ Hz, $J_2=4.0$ Hz, 1H), 2.63-2.50 (m, 5H); ^{13}C NMR (400

MHz, DMSO-*d*₆, TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI): 803.1993 for [M+H]⁺ (calculated 803.1897 for C₃₉H₃₅ClN₄O₁₃).

Data for Compound 2-(2-nitrophenoxy) acetic acid-4β-N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2l). ¹H NMR (400 MHz, DMSO-*d*₆, TMS, ppm): δ: 8.06(d, *J*=8.0 Hz, 1H), 7.49(d, *J*=8.0 Hz, 2H), 7.70 (t, *J*=8.0 Hz, 1H), 7.42 (s, 1H), 7.31(d, *J*=8.0 Hz, 1H), 7.18 (t, *J*=8.0 Hz, 1H), 7.04 (d, *J*=8.0 Hz, 2H), 6.57 (s, 1H), 6.38 (s, 2H), 6.01 (s, 2H), 5.34 (s, 2H), 4.59 (d, *J*=4.0 Hz, 1H), 4.13 (dd, *J*₁=12.0 Hz, *J*₂=8.0 Hz, 3H), 3.65 (s, 6H), 3.51-3.35 (m, 4H), 3.18 (dd, *J*₁=12.0 Hz, *J*₂=4.0 Hz, 1H), 2.63-2.50 (m, 5H); ¹³C NMR (400 MHz, DMSO-*d*₆, TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI): 769.2347 for [M+H]⁺ (calculated 769.2287 for C₃₉H₃₆N₄O₁₃).

Data for Compound 2-(2-nitrophenoxy)propanoic acid-4β-N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2m). ¹H NMR (400 MHz, DMSO-*d*₆, TMS, ppm): δ: 8.04 (d, *J*=4.0 Hz, 2H), 7.85 (d, *J*=8.0 Hz, 1H), 7.66 (t, *J*=4.0 Hz, 1H), 7.38 (s, 1H), 7.24 (d, *J*=8.0 Hz, 1H), 7.14 (t, *J*=8.0 Hz, 1H), 7.01 (d, *J*=4.0 Hz, 2H), 6.53 (s, 1H), 6.33 (s, 2H), 5.96 (s, 2H), 5.46 (dd, *J*₁=12.0 Hz, *J*₂=4.0 Hz, 1H), 4.54 (d, *J*=4.0 Hz, 1H), 4.09 (dd, *J*₁=16.0 Hz, *J*₂=8.0 Hz, 3H), 3.58 (s, 6H), 3.47-3.31 (m, 4H), 3.13 (dd, *J*₁=12.0 Hz, *J*₂=4.0 Hz, 1H), 2.58-2.46 (m, 5H), 1.63 (d, *J*=8.0 Hz, 3H); ¹³C NMR (400 MHz, DMSO-*d*₆, TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI): 783.2518 for [M+H]⁺ (calculated 783.7594 for C₄₀H₃₈N₄O₁₃).

Data for Compound 2-methyl-2-(2-nitrophenoxy)propanoic acid-4β-N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2n). ¹H NMR (400 MHz, DMSO-*d*₆, TMS, ppm): δ: 8.06(d, *J*=8.0 Hz, 2H), 7.98 (d, *J*=8.0 Hz, 1H), 7.66 (t, *J*=4.0 Hz, 1H), 7.42 (s, 1H), 7.24 (d, *J*=8.0 Hz, 1H), 7.18 (t, *J*=4.0 Hz, 1H), 7.01 (d, *J*=4.0 Hz, 2H), 6.57 (s, 1H), 6.39 (s, 2H), 6.01 (s, 2H), 4.58 (d, *J*=4.0 Hz, 1H), 4.12 (dd, *J*₁=8.0 Hz, *J*₂=4.0 Hz, 3H), 3.64(s, 6H), 3.51-3.34 (m, 4H), 3.18 (dd, *J*₁=12.0 Hz, *J*₂=4.0 Hz, 1H), 2.62-2.49 (m, 5H), 1.98 (s, 6H); ¹³C NMR (400 MHz, DMSO-*d*₆, TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI): 797.2677 for [M+H]⁺ (calculated 797.7865 for C₄₁H₄₀N₄O₁₃).

Data for Compound 2-(2-nitrophenoxy)butanoic acid-4β-N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2o). ¹H NMR (400 MHz, DMSO-*d*₆, TMS, ppm) δ: 8.04 (d, *J*=4.0 Hz, 2H), 7.85 (d, *J*=8.0 Hz, 1H), 7.66 (t, *J*=4.0 Hz, 1H), 7.38 (s, 1H), 7.24 (d, *J*=8.0 Hz, 1H), 7.14 (t, *J*=8.0 Hz, 1H), 7.01 (d, *J*=4.0 Hz, 2H), 6.53 (s, 1H), 6.33 (s, 2H), 5.96 (s, 2H), 5.46 (t, *J*=4.0 Hz, 1H), 4.54 (d, *J*=4.0 Hz, 1H), 4.09 (dd, *J*₁=16.0 Hz, *J*₂=8.0 Hz, 3H), 3.58(s, 6H), 3.47-3.31 (m, 4H), 3.13 (dd, *J*₁=12.0 Hz, *J*₂=4.0 Hz, 1H), 2.58-2.46

(m, 5H), 1.94-1.80 (m, 2H), 1.47(t, $J=8.0$ Hz, 3H); ^{13}C NMR (400 MHz, $\text{DMSO}-d_6$, TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI):797.2673 for $[\text{M}+\text{H}]^+$ (calculated 797.7865 for $\text{C}_{41}\text{H}_{40}\text{N}_4\text{O}_{13}$).

Data for Compound 2-methyl-2-(4-methyl-2-nitrophenoxy) prop-anoic acid-4 β -N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2p). ^1H NMR (400 MHz, $\text{DMSO}-d_6$, TMS, ppm) δ : 8.07(d, $J=9.2$ Hz, 2H), 7.72 (s, 1H), 7.50 (d, $J=8.0$ Hz, 1H), 7.43 (s, 1H), 7.20 (d, $J=8.0$ Hz, 1H), 7.03 (d, $J=12.0$ Hz, 2H), 6.57 (s, 1H), 6.39 (s, 2H), 6.01 (s, 2H), 4.59 (d, $J=4.0$ Hz, 1H), 4.14 (dd, $J_1=8.0$ Hz, $J_2=4.0$ Hz, 3H), 3.65(s, 6H), 3.51-3.35 (m, 4H), 3.18 (dd, $J_1=12.0$ Hz, $J_2=4.0$ Hz, 1H), 2.63-2.49 (m, 5H), 2.30 (s, 3H), 1.69 (s, 6H); ^{13}C NMR (400 MHz, $\text{DMSO}-d_6$, TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI):811.2916 for $[\text{M}+\text{H}]^+$ (calculated 811.8137 for $\text{C}_{42}\text{H}_{42}\text{N}_4\text{O}_{13}$).

Data for Compound 2-methyl-2-(5-methyl-2-nitrophenoxy) prop-anoic acid-4 β -N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2q). ^1H NMR (400 MHz, $\text{DMSO}-d_6$, TMS, ppm) δ : 8.06(d, $J=8.0$ Hz, 2H), 7.79 (d, $J=8.0$ Hz, 1H), 7.43 (s, 1H), 7.27 (s, 1H), 7.18 (d, $J=8.0$ Hz, 1H), 7.01 (d, $J=12.0$ Hz, 2H), 6.57 (s, 1H), 6.38 (s, 2H), 6.01 (s, 2H), 4.60 (d, $J=4.0$ Hz, 1H), 4.14 (dd, $J_1=8.0$ Hz, $J_2=4.0$ Hz, 3H), 3.67(s, 6H), 3.51-3.35 (m, 4H), 3.18 (dd, $J_1=12.0$ Hz, $J_2=4.0$ Hz, 1H), 2.62-2.50 (m, 5H), 2.27 (s, 3H), 1.67 (s, 6H); ^{13}C NMR (400 MHz, $\text{DMSO}-d_6$, TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI):811.4158 for $[\text{M}+\text{H}]^+$ (calculated 811.8137 for $\text{C}_{42}\text{H}_{42}\text{N}_4\text{O}_{13}$).

Data for Compound 2-(5-bromo-2-nitrophenoxy)-2-methylprop-anoic acid-4 β -N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2r). ^1H NMR (400 MHz, $\text{DMSO}-d_6$, TMS, ppm) δ : 8.07 (d, $J=8.0$ Hz, 2H), 7.82 (d, $J=8.4$ Hz, 1H), 7.53 (s, 1H), 7.42 (s, 1H), 7.39 (d, $J=8.0$ Hz, 1H), 7.02 (d, $J=8.0$ Hz, 2H), 6.56 (s, 1H), 6.38 (s, 2H), 5.99 (s, 2H), 4.59 (d, $J=4.0$ Hz, 1H), 4.11 (dd, $J_1=16.0$ Hz, $J_2=8.0$ Hz, 3H), 3.66 (s, 6H), 3.49-3.37 (m, 4H), 3.17 (dd, $J_1=16.0$ Hz, $J_2=4.0$ Hz, 1H), 2.62-2.49 (m, 5H), 1.66 (s, 6H); ^{13}C NMR (400 MHz, $\text{DMSO}-d_6$, TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI):876.2146 for $[\text{M}+\text{H}]^+$ (calculated 876.2067 for $\text{C}_{41}\text{H}_{39}\text{BrN}_4\text{O}_{13}$).

Data for Compound 2-(4-methyl-2-nitrophenoxy) propanoic acid-4 β -N-(4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2s). ^1H NMR (400 MHz, $\text{DMSO}-d_6$, TMS, ppm) δ : 8.07 (d, $J=8.0$ Hz, 2H), 7.71 (s, 1H), 7.53 (d, $J=8.0$ Hz, 1H), 7.42 (s, 1H), 7.15 (d, $J=8.0$ Hz, 1H), 7.06 (d, $J=8.0$ Hz, 2H), 6.58 (s, 1H), 6.38 (s, 2H), 6.01 (s, 2H), 5.46 (dd, $J_1=12.0$ Hz, $J_2=4.0$ Hz, 1H), 4.59 (d, $J=4.0$ Hz, 1H), 4.14 (dd, $J_1=16.0$ Hz, $J_2=8.0$ Hz, 3H), 3.63 (s, 6H), 3.49-3.35 (m, 4H), 3.15 (dd, $J_1=12.0$ Hz, $J_2=4.0$ Hz, 1H), 2.61-

2.50(m, 5H), 2.31 (s, 3H), 1.65 (d, $J=8.0$ Hz, 3H); ^{13}C NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI):797.2598 for $[\text{M}+\text{H}]^+$ (calculated 797.7868 for $\text{C}_{41}\text{H}_{40}\text{N}_4\text{O}_{13}$).

Data for Compound 2-(4-methoxy-2-nitro-phenoxy)propanoic acid-4 β -N- (4-Nitrophenyl piperazinyl)-4'-O-demethyl-4-deoxypodophyllotoxin (2t). ^1H NMR (400 MHz, DMSO- d_6 , TMS, ppm) δ : 8.06(d, $J=8.0$ Hz, 2H), 7.49(s, 1H), 7.43 (s, 1H), 7.25 (d, $J=8.0$ Hz, 2H), 7.06 (d, $J=8.0$ Hz, 2H), 6.58 (s, 1H), 6.38 (s, 2H), 6.01 (s, 2H), 5.42 (dd, $J_1=12.0$ Hz, $J_2=4.0$ Hz, 1H), 4.61 (d, $J=4.0$ Hz, 1H), 4.14 (dd, $J_1=16.0$ Hz, $J_2=8.0$ Hz, 3H), 3.79 (s, 3H), 3.64 (s, 6H), 3.51-3.39 (m, 4H), 3.18 (dd, $J_1=12.0$ Hz, $J_2=4.0$ Hz, 1H), 2.63-2.51(m, 5H), 1.67 (d, $J=8.0$ Hz, 3H); ^{13}C NMR (400 MHz, DMSO- d_6 , TMS, ppm): δ 176.26, 171.0, 158.7, 153.86, 153.6, 147.3, 140.4, 139.7, 138.5, 135.1, 132.8, 131.9, 129.7, 129.4, 126.5, 125.6, 115.7, 112.6, 110.5, 110.0, 108.1, 101.7, 70.4, 69.2, 65.3, 56.8, 52.0, 50.5, 45.7, 43.7, 40.5, 19.3; MS(ESI):813.3543 for $[\text{M}+\text{H}]^+$ (calculated 813.7857 for $\text{C}_{41}\text{H}_{40}\text{N}_4\text{O}_{14}$).

Cytotoxicity assay

Cells were incubated at 37 °C in a 5% CO_2 atmosphere. The synthetic compounds 2a-2t and reference compound (Etoposide and Demethylepipodophyllotoxin) were dissolved and the cells were plated in 96-well plates determined according to the reported method [16].

Cell cycle and apoptotic assay

For cell cycle and apoptotic assay were determined according to the reported methods [25, 26]. Detection of apoptosis using Annexin V-FITC/PI staining, after incubation for 24 h at 37 °C in a humidified atmosphere with 5% CO_2 , various concentrations of compound 2b and 2d were added to the cultures, and the plate was incubated for further 72 h. After that, HCT-116 cells were collected, washed, resuspended in 200 μL binding buffer of the Annexin V-FITC kit, stained with propidium iodide and incubated for 30 min at 37 °C prior to flow cytometric analysis.

Molecular docking model assay

Based on the redocking process, molecular docking was applied between the chosen two compounds and NTR model. To clarify the binding forms of the chosen compounds, binding affinities were also predicted by using ligand–enzyme molecular docking calculations with different approaches.

Declarations

Conflict of Interest

Compliance with ethics requirements

Authors have no financial relationship with the organization that sponsored the research.

CRediT authorship contribution statement:

Zheng-Rong Wu: Software, Data curation, Writing-original draft. **Wei Deng:** Supervision. **Dian He:** Funding acquisition, Writing - review. Supervision.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Zheng-Rong Wu

Email: wuzhengrong@lzu.edu.cn

References

1. Shareef MA, D D, Ramasatyaveni G et al (2015) *Eur J Med Chem* 89:128
2. Qu Y, Zhang Ch, Ma X, Gao YH et al (2020) *Bioorg Med Chem Lett* 28(24):115821
3. Thi Q, Le-Nhat-Thuy G, Thi T et al (2021) *Bioorg Med Chem Lett* 37(4):127841
4. Kamal JR, Tamboli MJ, Ramaiah SF, Adil S (2013) *N. C. V. L. P. Bioorg Med Chem* 21:6414
5. Yong Y, Shin SY, Lee YH, Lim Y (2009) *Bioorg Med Chem Lett* 19:4367
6. Chen SW, Gao YY, Zhou NN, Liu J, Huang WT, Hui L, Jin Y, Jin YX (2011) *Bioorg Med Chem Lett* 21:7355
7. V K, Singh.; R, Roy KH (2014) *Eur J Med Chem* 74:552
8. Xu H, Lv M, Tian X (2009) *Curr Med Chem* 16:327
9. Zhao Yu, Ge CWang, Wu ZH, Wang CN, Fang JH, Zhu L (2011) *Eur. J. Med. Chem.* 46,901
10. Wu GR, X B, Yang Y, Q. et al (2018) *Eur. J. Med. Chem.* 155,183
11. Wang L, Yang FY, Yang XC, Guan XH, Hu CQ, Liu T, He QJ, Yang B, Hu Y (2011) *Z Eur J Med Chem* 46:285
12. Huang WT, Liu J, Liu JF, Hui L, Ding YL, Chen SW (2012) *Eur J Med Chem* 49:48
13. Kamal A, Suresh P, Ramaiah MJ et al (2011) *Bioorg Med Chem* 19:4589
14. Liu JF, Sang CY, Xu XH, Zhang LL et al (2013) *Eur J Med Chem* 64:621
15. Kamal A, Suresh P, Ramaiah MJ, Mallareddy A et al (2012) *Bioorg Med Chem* 20:2054
16. Liu JF, Sang CY, Qin WW, Zhao J et al (2013) *Bioorg Med Chem* 21(22):6948
17. W A, Denny (2002) *Curr Pharmaceut Des* 8:1349
18. Patterson S, Wyllie (2014) *Trends Parasitol* 30:289
19. Jiang Q, Zhang. Z, Lu J, Huang Y et al (2013) *Bioorgan Med Chem* 21:7735

20. Ao X, Bright. SA, Taylor NC, Org RBP (2017) *Biomol. Chem.* 15, 6104
21. Zhang X, Zhao. Q, Tang Y (2017) *Anal Chem* 89:5503
22. Zhao Y, Ge CW, Wu ZH, Wang CN, Fang JH, Zhu L (2011) *Eur J Med Chem* 46:901
23. Zhou Y, Bobba KN, Lv XW, Yang D et al (2017) *Analyst* 142:345
24. Liu Q, Zhuo J, Xi H et al (2005) *Guangzhou Chem* 30(4):6
25. Riccardi C, Nicoletti I (2006) *Nat Protoc* 1:1458
26. Chen S, Cheng AC, Wang MS, Peng X (2008) *g. W J G*, 14, 2174

Figures

Figure 1

Effects of compound 2b and 2d on HeLa cells cycle progression (48 h). (A) Control; (B) treated with 2.5μmol/L VP-16; (C) treated with 50 μmol/L VP-16; (D) treated with 2.5μmol/L 2b; (E) treated with 50 μmol/L 2b ; (F) treated with 50 μmol/L 2d; (G) treated with 50 μmol/L 2d.

Figure 2

Apoptosis in HeLa cells were treated with compound 2b and 2d for 48 h, as detected using flow cytometry with the annexin V/PI apoptosis detection kit. (A) Control; (B) treated with 2.5 μ mol/L VP-16; (C) treated with 50 μ mol/L VP-16; (D) treated with 2.5 μ mol/L 2b; (E) treated with 50 μ mol/L 2b; (F) treated with 50 μ mol/L 2d; (G) treated with 50 μ mol/L 2d.

Figure 3

Molecular binding mode of compound 2b and 2d to NTR 1DS7.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [GraphicalAbstract.png](#)
- [Scheme1.png](#)
- [Supplementarymaterial.docx](#)