On Injectivity of an Integral Operator Connected to Riemann Hypothesis

Dumitru Adam (✉ dumitru_adam@yahoo.ca)

Research Article

Keywords: Approximation Subspaces, Integral Operators, Riemann Hypothesis

Posted Date: December 14th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1159792/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

On Injectivity of an Integral Operator Connected to Riemann Hypothesis *

Dumitru Adam [†]

Nov. 30, 2021

Abstract

In this paper we provided methods for investigating the injectivity of linear bounded operators on separable Hilbert spaces using their approximations on dense families of subspaces.

As an application we proved the injectivity of the integral operator involved in the equivalent formulations of Riemann Hypothesis (see [2] and [4]) in the Example 4.2.

subclass: 65R99, 47G10, 45P05

keywords: Approximation Subspaces, Integral Operators, Riemann Hypothesis

Disclaimer: The RH has been introduced by Bernhard Riemann in 1859 and by today, is considered to be one of the most important unsolved problems in pure mathematics. In 1993, Alcantara-Bode showed ([2]) that RH holds if and only if the integral operator in (1) defined on the Hilbert space $L^2(0, 1)$ is injective.

Surprisingly, in almost 30 years the injectivity of the integral operator used in equivalent formulation of RH has not been addressed, nor has been dissociate from RH.

Our goal is to address the injectivity of the linear bounded operators on separable Hilbert spaces.

§1. Introduction.

In the paper of Alcantara-Bode [2], the author proved (Theorem 1, pg. 152) that RH holds if and only if the following Hilbert-Schmidt integral operator T_ρ defined on $L^2(0, 1)$ by:

$$(T_\rho u)(y) := \int_0^1 \left\{ \frac{y}{x} \right\} u(x) dx \quad (1)$$

*Submitted to JNAAT

[†]No Funding, No Affiliations. Email: mikiad77@gmail.com Windsor, On. Ca. & Brasov, Ro

has its null space $N_{T\rho} = \{0\}$.

The linear bounded integral operator has been introduced in [4] (Beurling, 1955) for providing first equivalent formulation of RH in terms of functional analysis. The Beurling formulation of RH has been used in [2] (Alcantara-Bode, 1993) for providing second equivalent formulation of Riemann Hypothesis in terms of injectivity of the operator in (1) on $L^2(0,1)$.

We introduced in this paper criteria for investigating the injectivity of bounded linear operators on generic separable Hilbert spaces providing tools ready to use for practical purposes. The Theorem 2.1 §2, acting on generic separable Hilbert spaces as well its extensions from §3 acting on integral operators on $L^2(0,1)$ (Lemma 3.1 and 3.2), are the criteria using the operator approximations on a dense family of approximation subspaces. We applied the methods in §4 taking two examples of integral operators with discontinue kernels, showing the injectivity of the integral operator used in RH equivalence in the Example 4.2.

§2. Injectivity Criteria.

Let H be a separable Hilbert space. The norm considered on H is the norm induced by its inner product. Because a linear bounded operator T and its associated Hermitian and positive definite operator (T^*T) have the same null space N_T , we will consider its associated Hermitian operator when we do not have enough information about positivity of T on H . Then, we take T or (T^*T) in order to have its positivity $\langle Tu, u \rangle \geq 0$ on H as we need. In the following T is positive definite on H and our aim in this paragraph is to provide methods for investigating its strict positivity on H , in other words, in finding if its null space $N_T = \{0\}$. We will see later that a weaker property than the positivity on H is enough and is mandatory for investigating the injectivity on H , that is the a -positivity of linear operators, introduced in this paragraph.

Let $B := \{u \in H; \|u\| = 1\}$ be the unit sphere in H . Because a not null element u is in N_T if and only if $u/\|u\| \in N_T$, we will consider the normalized elements of N_T for convenience.

A family of finite dimension subspaces $\mathfrak{S} := \{S_n, n \in N\} \subset H$ is a proper family of approximation subspaces if the following properties hold on the subspaces of the family:

- (a) $S_n \subset S_{n+1}, n \in N$
- (b) $\beta_n(u) := \|u - u_n\| \rightarrow 0$, for $n \rightarrow \infty$

for every $u \in B$ and with u_n its orthogonal projection on S_n .

Observation 2.1: A proper family subspaces is dense in H : $\overline{\cup_{n=1}^{\infty} S_n} = H$, property easy to show. Let $\mathfrak{S}^0 = \cup_{n=1}^{\infty} S_n$ and let consider $H_{\mathfrak{S}^0} = H \setminus \mathfrak{S}^0$.

We say that the bounded linear operator T is almost strict positive definite on H or a -positive, if there exists a proper family $\mathfrak{S}$ on which, for every subspace $S_n \in \mathfrak{S}$, there exists $\alpha_n > 0$ such that:

$$\langle Tv, v \rangle \geq \alpha_n \|v\|^2, \forall v \in S_n$$

i.e. T is strict positive definite on each subspace of the family. We name the set $\{\alpha_n, n \geq 1\}$, the set of a -positive parameters associated to T on $\mathfrak{S}$. In facts, a -positive parameters are the smallest eigenvalues of the operator restrictions to the corresponding subspaces. We will use the term a -positive also associated to the family $\mathfrak{S}$ for convenience.

Let $L_{\mathfrak{S}}$ be the set of linear bounded operators on H that are a -positive with respect to the family $\mathfrak{S}$.

If $T \in L_{\mathfrak{S}}$ and there exists $u \in B \cap N_T$ then $u \notin S_n$ for every $n \geq 1$, i.e. $u \notin \mathfrak{S}^0$. For this reason we will consider in our analysis only $u \in H$ that are eligible, in the sense that these elements from H could be the normalised zeros of T , $u \in B_0 := B \cap H_{\mathfrak{S}^0}$.

We will refer in this paper to B_0 as the set of eligible roots of the a -positive operators given a proper family $\mathfrak{S}$.

Observations 2.2: we point out two extreme situations that we will exclude from our analysis:

1.) - If T is not a -positive, it is not injective and the investigations on its injectivity ends here.

That means, a -positivity is mandatory for injectivity.

2.) - If T is a -positive and the approximation subspaces of $\mathfrak{S}$ are invariant subspaces of T , then T is injective and so, no investigations for its injectivity needed.

As a consequence, from now on the linear bounded operators in consideration would be a -positive operators without having the approximation subspaces of the proper family as invariant subspaces.

We introduce an injectivity criteria for linear, bounded operators on a generic separable Hilbert space that are a -positive on a proper family.

Let $\mathfrak{S}$ be a proper family of approximation subspaces on H and $T \in L_{\mathfrak{S}}$. Once T has no invariant subspaces in $\mathfrak{S}$ there exists $w \in S_n^{\perp}$ for which $Tw \notin S_n^{\perp}$. For a such element w and for every $v \in S_n$ both not null, the inner product $\langle Tw, v \rangle$ is well defined in H . Because an strict positive operator on a finite dimension subspace has the same eigenvalues as its adjoint, in our

case $T_{|S_n}^*$ and $T_{|S_n}$ have both the same largest eigenvalue λ_n^{max} on S_n , we obtain the following estimation for the considered inner product:

$$|\langle Tw, v \rangle| = |\langle w, T^*v \rangle| \leq \|w\| \lambda_n^{max} \|v\|$$

If there exists $u \in N_T \cap B_0$, we take $w := (u_n - u) \in S_n^\perp$ and $v = u_n$, the projection of u on S_n . Obviously, $Tw = Tu_n \notin S_n^\perp$ (because of the a-positivity of T $\langle Tu_n, u_n \rangle \neq 0$ and, because there exists an index $n_0(u)$ from which u is not orthogonal to S_n from the density of the proper family). Then:

$\alpha_n \|u_n\|^2 \leq \langle Tu_n, u_n \rangle = |\langle T(u_n - u), u_n \rangle|$ and, together with the previous relations we obtain:

$$\mu_n \|u_n\| \leq \beta_n(u) \quad (2)$$

where $\mu_n := \lambda_n^{min} / \lambda_n^{max}$ is the ratio between the smallest and highest eigenvalues of the restriction of T to the subspace S_n inequality obtained using the a-positivity property and replacing α_n with λ_n^{min} .

Pointing out: if there exists $u \in N_T \cap B_0$ then (2) holds on every $S_n \in \mathfrak{S}$ for which u is not orthogonal on it. In fact, in virtue of the properties of the proper family given $u \in B_0$, there exists $n_0(u)$ finite index from which for every $n \geq n_0(u)$ its orthogonal projections are not null on the subspaces $S_n, n \geq n_0(u)$.

So, the property (2) for an eligible $u \in N_T$, is valid starting from a finite index $n_0(u)$; however, it is validated also for indexes $n < n_0(u)$ because for such indexes $u_n = 0$ and $\beta_n(u) = 1$.

For T a-positive, given $u \in B_0$ we define the expression $\theta_n(u) := \theta_n(T)(u)$ on every subspace of the proper family $S_n, n \geq n_0(u)$ by:

$$\theta_n(u) := \beta_n^2(u) - \mu_n^2 \|u_n\|^2 = 1 - (\mu_n^2 + 1) \|u_n\|^2 \quad (3)$$

with u_n the orthogonal projection of u on S_n . The last member in the equality is obtained using the relationship for every $u \in B$ and its orthogonal projections on S_n and $S_n^\perp$: $\|u_n\|^2 + \|(u - u_n)\|^2 = \|u\|^2 = 1$.

We will split the set of eligible elements B_0 in two disjoint subsets:

$$1) \quad B_0^1 = B_0^1(T) := \{u \in B_0; \quad \theta_n(u) \geq 0, n \geq n_0(u)\};$$

for which, $B_0^1 \supset B_0 \cap N_T$

$$2) \quad B_0^2 = B_0^2(T) := \{u \in B_0; \quad \theta_n(u) < 0, n \geq n_0(u)\};$$

for which, $B_0^2 \cap N_T = \{\emptyset\}$

Note. Because for every normalised $u \in N_T$ the inequality (2) holds globally on the proper family $\mathfrak{S}$, this behaviour will be propagated from now on for each statement involving (2); it is the reason we would omit carrying out every time the index specifications referring to all subspaces of the family, when it is evident from context. We would do it until the end of

the paragraph.

With the next theorem we provide a sufficient condition for an operator in $L_{\mathfrak{S}}$ in order to have its null space containing only the null element in H .

Theorem 2.1. (Injectivity Criteria.) Let $T \in L_{\mathfrak{S}}$.

A.) If the sequence $\{\mu_n\}, n \geq 1$ is inferior bounded by a strict positive constant independent of n , $\mu_n \geq C > 0$, for every $n \geq 1$ then T is injective.

B.) If Q is a linear, bounded and injective operator on H such that Q (or its associate Hermitian) verifies $\mu_n(Q) \leq \mu_n(T), n \geq 1$ then T is injective.

Proof of A.)

Suppose that the sequence $\{\mu_n\}, n \geq 1$ is inferior bounded by a constant C independent of n , strict positive.

Let $u \in B_0$.

1) If there exists an infinite subsequence of subspaces from the family for which $\theta_{n_m}(u) \geq 0, n_m \geq n_0(u)$ we obtain:

$$\|u_{n_m}\|^2 \leq 1/(1 + \mu_{n_m}^2) < 1/(1 + C^2) < 1$$

and $\lim_{m \rightarrow \infty} \|u_{n_m}\|^2 < 1/(1 + C^2) < 1$.

Or, $\{u_{n_m}\}, n_m \geq n_0(u)$ should converge in norm to 1 on the approximation subspaces for every $u \in H$. Thus, the inequality $\theta_{n_m}(u) \geq 0$ could not take place on an infinity of subspaces of the family. We should have instead at most only a finite number of subspaces verifying it.

If there exists a finite number of subspaces on which $\theta_{n_m}(u) \geq 0, n_m \geq n_0(u)$ then we will shrink the family including these subspaces inside of the subspaces already verifying the opposite relationship, obtaining $u \notin B_0^1$ when the sequence $\{\mu_n\}, n \geq 1$ is inferior bounded by a not null constant.

Then, we are in the situation $\theta_n(u) < 0$ for every $n \geq n_0(u)$ on the $\mathfrak{S}$, i.e. $u \in B_0^2$.

2) If $\theta_n(u) < 0$ holds for $n \geq n_0(u)$, then $\beta_n(u) < \mu_n \|u_n\|, n \geq n_0(u)$. But, u satisfying such relationship could not be in N_T because a reverse inequality given in (2) holds for eligible u in N_T . The only restriction we put on $u \in H$, has been its eligibility, $u \in B_0$. Thus, if $\{\mu_n\}, n \geq 1$ is inferior bounded, does not exist $u \in B_0$ that is in N_T . Or, B_0 contains all normalized zeros of T , so $N_T = \{0\}$.

Proof of B.)

Because Q is injective, then Q or (Q^*Q) is strict positive on H and so, is a -positive on $\mathfrak{S}$. Let $\mu_n(Q), n \geq 1$ be the injectivity parameters of Q or of its associate Hermitian.

For every $u \in B_0$, $u \in B_0^2(Q)$ having $\theta_n(Q)(u) < 0$ otherwise, u should be in N_Q that is a null set.

Let $u \in B_0$.

From $\mu_n(Q) \leq \mu_n(T)$ we obtain $\theta_n(T)(u) \leq \theta_n(Q)(u) < 0$, valid for $n \geq 1$ showing that $u \in B_0^2(T)$. Because u is chosen an arbitrary eligible element, $B_0 \subseteq B_0^2(T)$ that means $B_0^1(T) = \emptyset$, showing that T is injective. ■

§3. On Approximations of Integral Operators. On the separable Hilbert space $L^2(0,1)$, we will take in the following paragraphs the proper family $\mathfrak{S}_\chi := \{S_h, nh = 1, n \geq 2\}$, where $S_h := \text{span}\{\chi_{n,k}, k = 1, n, nh = 1\}$ built on simple functions that are linear combinations of indicator functions of disjoint intervals

$$\chi_{n,k} := \chi_{n,k}(t) = \begin{cases} 1 & t \in \Delta_k := ((k-1)h, kh] \\ 0 & \text{otherwise} \end{cases} \quad (4)$$

is dense in $L^2(0,1)$ and meets the requests imposed by the definition of a proper family of approximation subspaces introduced in the previous paragraph. Citing (Buescu & Paixa~o, 2007), the family of functions $\{\chi_{n,k}\}$, $k = 1, n, nh = 1$, is defining a trace class integral operator with the kernel

$$r_h(x, y) = h^{-1} \sum_{k=1}^n \chi_{n,k}(x) \chi_{n,k}(y)$$

that is an orthogonal projection in $L^2(0,1)$ on S_h having the orthogonal eigenfunctions $\{\chi_{n,k}\}, k = 1, n$. In fact, S_h are generated by the families of the orthogonal eigenfunctions of the projection operators $\{P_h := P|_{S_h}\}, nh = 1, n \geq 2$.

Moreover, because the orthogonality of eigenfunctions is dictated by the disjoint interval support, $\chi_{n,k}(x) \chi_{n,j}(y) = \delta_{k,j}$ the entries in the matrix representation (5) of the operator restrictions on approximation subspaces will be zero outside the diagonal and, it explain the form of the discrete approximations of the kernel functions on approximation subspaces defined bellow.

Thus, the linear integral operator (like in (1))

$$T_\rho u := (T_\rho u)(y) = \int_0^1 \rho(y, x) u(x) dx$$

has the discrete approximations of the kernel function ρ like in [5],[6]:

$$\rho_h(y, x) = h^{-1} \sum_{k=1}^n \chi_{n,k}(x) \rho(y, x) \chi_{n,k}(y)$$

Obviously, $(\forall) u \in H$, its projection on S_h take the form of a simple function: $P_{|S_h} u = \sum_{k=1}^n \langle u, \chi_{n,k} \rangle \chi_{n,k}$.

Accordingly, taking any $v_h \in S_h$ not null, of the form $v_h(t) = \sum_{k=1}^n c_k \chi_{n,k}(t)$ the discrete approximation of the integral operator, verifies the following equality:

$$\begin{aligned} \langle T_\rho^h v_h, v_h \rangle &= \int_0^1 \int_0^1 \rho_h(y, x) v_h(x) \bar{v}_h(y) dx dy \\ &= h^{-1} \mathbf{c}_h^T M_h \bar{\mathbf{c}}_h \end{aligned} \quad (5)$$

Here the $(n \times n)$ diagonal matrix $M_h = [d_{kk}^h]_{k=1,n}$ is the matrix representation of the restriction T_ρ^h of the integral operator T_ρ to the subspace S_h and the vector $\mathbf{c}_h$ is the vector of coordinates $\{c_k, k=1,n\}$ of v_h in C^n verifying $\|v_h\|^2 = h \|\mathbf{c}_h\|_{C^n}^2$.

Taking a look of (5) and the matrix representation M_h of the T_ρ^h on S_h , if the matrix is strict positive definite, we have: $\lambda_{\min}(M_h) = \min_{k=1,n} d_{kk}^h$, from where:

$$\langle T_\rho^h v_h, u_h \rangle \geq h^{-2} \lambda_{\min}(M_h) \|v_h\|^2$$

So, the a-positivity parameters associated to T_ρ should have the form like $\alpha_n = h^{-2} \lambda_{\min}(M_h)$, $nh = 1$, $n \geq 1$.

Remark 3.1) The linear bounded integral operator T_ρ is a-positive on $\mathfrak{S}_\chi$ if and only if on approximation subspaces its kernel function is verifying

$$d_{kk}^h := \int_{\Delta_k} \int_{\Delta_k} \rho(y, x) dx dy > 0 \quad (6)$$

for $k = 1, n$, $nh = 1$, i.e. the diagonal entries in the diagonal matrices M_h are strict positive.

Here is the main theorem (Theorem 2.1.A) reformulated for integral operators on $L^2(0,1)$ taking the proper family $\mathfrak{S}_\chi$.

Lemma 3.1. (*Injectivity Criteria for Integral Operators.*) Let $T_\rho \in L_{\mathfrak{S}_\chi}$. If the sequence $\mu_h(T_\rho)$, $nh = 1$, $n \geq 1$

$$\mu_h(T_\rho) = \frac{\min_{(k=1,n)} d_{kk}^h}{\max_{(k=1,n)} d_{kk}^h} \quad (7)$$

is inferior bounded, $\mu_h(T_\rho) > C$ by a strict positive constant C independent of n , $nh = 1$, then T_ρ is injective.

Proof.

In the view of the previous paragraphs, if $T \in L_{\mathfrak{S}_\chi}$ is a linear bounded integral operator its α -positivity parameters are given by $\alpha_n = h^{-2} \min_{k=1, n} d_{kk}^h$, $n \geq 1$, $n \geq 2$. Thus, the injectivity criteria (Theorem 2.1.A) could be used in determining the injectivity of T_ρ observing that in this case $\{\mu_n, n \geq 1\}$ are given as in (7). ■

We will introduce a method for analyse the cases of operators α -positive whose the injectivity parameters sequence is inferior unbounded.

Let define the I -multiplier operator, the operator obtained as a product of identity by a power of x , $\{I_s\}$, $s \in \mathbb{N}$ on $L^2(0,1)$ as follows: for $u \in L^2(0,1)$

$$(I_s u)(x) = x^s u(x) \quad i.e. \quad I_s : u(x) \rightarrow x^s u(x) \quad (8)$$

It is easy to show that I_s for every s , is linear, bounded, positive definite and injective on $L^2(0,1)$ and so, α -positive on $\mathfrak{S}_\chi$. The injectivity and positivity properties on H are coming from $\langle I_s u, u \rangle = \int_0^1 w(x) \overline{w(x)} dx > 0$ for $u \neq 0$ where $w(x) = x^{s/2} u(x)$. Its injectivity parameters applying the formula (7), are for $n \geq 1$

$$\mu_n(I_s) = 1 / \left(\sum_{i=0, s} n^i (n-1)^{s-i} \right) = o(n^{-s})$$

Lemma 3.2 (*I-multipliers Rule.*) Let T α -positive on $\mathfrak{S}_\chi$. If exists an I -multiplier operator I_s for some $s \geq 1$ such that

$$\mu_n(I_s) \leq \mu_n(T)$$

for $n \geq 1$, then T is injective.

Proof

Suppose that the relationship exists. Then, taking $Q = I_s$ in the Theorem 3.1.B, we obtain T injective. ■

Observation: As a consequence of Lemma 3.2, if T is α -positive on $\mathfrak{S}_\chi$ and its injectivity parameters verifies $\mu_n(T) = o(n^{-s})$ for some $s \geq 1$, then T is injective. ■

§4. On integral operators with discontinue kernel functions.

We will investigate the injectivity of two integral operators having the kernel functions discontinue on $H := L^2(0,1)$. We will use the proper family of approximation subspaces introduced in previous paragraph, $\mathfrak{S}_\chi$ on which, for α -positivity of integral operators we will apply Remark 3.1. The I -multiplier Rule is right method to apply for the injectivity of the integral operator in

Example 1 and, Lemma 3.1 for the integral operator in the Example 2.

Example 1. Let the linear bounded integral operator T_σ defined by

$$(T_\sigma u)(y) := \int_0^1 \left\{ \frac{y}{x} \right\} \left\{ \frac{x}{y} \right\} u(x) dx$$

on $L^2(0,1)$ having the kernel function $\sigma(y, x) = \left\{ \frac{y}{x} \right\} \left\{ \frac{x}{y} \right\}$. The brackets denote the fractional part function. The integral operator is linear, bounded having the kernel function discontinue, positive definite and Hermitian on H . The integral operator T_σ has no invariant subspaces in the proper family.

The diagonal entries in the matrix representations of the operator discretisations on the subspaces of approximations are all strict positive (see Remark 3.1 for formula for computing the matrix entries):

$$d_{11}^h = h^2 \left(1 - \frac{\pi^2}{12} \right); \quad d_{kk}^h = h^2 \left(1/2 - k + k^2 \ln \frac{k+1}{k} \right), k = 1, n, nh = 1$$

for $k = 2, n, nh = 1$. Modulo h^2 , $d_{11} \approx 0.7836$ and $d_{22} \approx 0.12186$ showing that $d_{11}^h > d_{22}^h$. The decreasing monotony of diagonal entries comes from the following observation. The function $f(t) = \left(1/2 - t + t^2 \ln \frac{t+1}{t} \right)$ defined on the interval $[2, \infty)$ is monotone decreasing with the limit 0 for t converging to ∞ . Its values for $t = k$, $k \geq 2$ are the values of the diagonal entries in the matrices associated to the operator discretisations on approximation subspaces, modulo h^2 . From the monotony of these entries we have on S_n : $d_{11}^h > d_{22}^h > \dots > d_{nn}^h$, $nh = 1$.

The injectivity parameters sequence are given by (7):

$$\mu_n(T_\sigma) = (1/2 - n + n^2 \ln((n+1)/n)) / (1 - \pi^2/12)$$

$n \geq 2$, $nh = 1$.

Hence T_σ is a-positive but, $\mu_n(T_\sigma)$ is not inferior bounded. Then we will apply the I-multipliers Rule (Lemma 3.2).

Let consider the denominator of $\mu_n(T_\sigma)$ as prototype for our considerations, being $f(t) = (1/2 - t + t^2 \ln((t+1)/t))$, $t \in [1, \infty)$ that is converging to 0 when $t \rightarrow \infty$.

Observing that $(f(t) - 1/t)$ and $(f(t) - 1/t^2)$, both $\rightarrow 0$ for $t \rightarrow \infty$ and,

$1/t^2 < f(t) < 1/t$ for $t \gtrsim 4$ we could consider the following modulo the factor $(1 - \pi^2/12)$, as I-multipliers I_1 and I_2 ($s=1$ and $s=2$ being the powers of x in the definition of the I-multiplier $I_{x^s}u(x) = x^s u(x)$ on $L^2(0,1)$). Then by Lemma 3.2, taking the I-multiplier I_s , $s=2$, we proved that T_σ is injective.

Example 2. *On integral operator used in RH Equivalence.* Let the linear bounded integral operator T_ρ defined by

$$(T_\rho u)(y) := \int_0^1 \left\{ \frac{y}{x} \right\} u(x) dx \quad (9)$$

on $L^2(0,1)$ having the kernel function $\rho(y, x) := \left\{ \frac{y}{x} \right\}$. This kernel function is the fractional part function of the quantity between brackets having jumps on the lines like $y = k \cdot x$, $k \geq 1$ resulting a discontinue function.

Because $(T_\rho \chi_{n,k})(y) \notin \text{span}(\{\chi_{n,k}\}, 1 \leq k \leq n, nh = 1)$, the approximation subspaces are not invariant subspaces of T_ρ . Computing the entries of the diagonal matrices M_h , (see (6)) of the kernel approximations on the subspaces $\{S_h\}, nh = 1, n \geq 2$, we obtain:

$$d_{11}^h = \frac{h^2}{4}(3 - 2\gamma); d_{kk}^h = \frac{h^2}{2}(-1 + \frac{2k-1}{k-1} \ln(\frac{k}{k-1})^{k-1}) \quad (10)$$

for $k = 2, n, nh = 1$ where $\gamma \cong 0.5772156...$ is the Euler-Mascheroni constant rounded here to 7 digits. The diagonal entries of matrix representations of T_ρ on each subspace S_h are strict positive valued and bounded by d_{11}^h and d_{22}^h . Moreover, the sequence $(-1 + \frac{2k-1}{k-1} \ln(\frac{k}{k-1})^{k-1})$ is monotone and converges to 0.5 for $k \rightarrow \infty$. Approximated values modulo the factor h^2 , of the diagonal entries are:

$$d_{1,1} \approx 0.461392, d_{2,2} \approx 0.636625, d_{3,3} \approx 0.506887, \dots$$

Then by Remark 3.1, T_ρ is a-positive on the approximation family having the a-positivity parameters given by $\alpha_n := h^{-2} d_{11}^h = h^{-2} \frac{1}{4}(3 - 2\gamma) \approx 0.461392 h^{-2}, nh = 1$.

The injectivity parameters $\{\mu_n\}$ defined in (7) are $d_{11}^h/d_{22}^h, n \geq 2, nh = 1$ and will be all independent of h because all entries $d_{kk}^h, 1 \leq k \leq n$ are on the same order of h . The evaluations of $\{\mu_n\}, n \geq 1$ are:

$$\mu_n(T_\rho) = \frac{3 - 2\gamma}{-2 + 6\ln(4/3)} \gtrapprox 0.724746, n \geq 2, nh = 1 \quad (11)$$

Hence the sequence $\mu_n(T_\rho), n \geq 1$, is bounded by a constant strict positive on every approximation subspace of the family.

By Lemma 3.1 follows that the integral operator T_ρ is injective, equivalently, $N_{T_\rho} = \{0\}$.

Theorem 4.1 The linear bounded integral operator T_ρ defined by

$$(T_\rho u)(y) := \int_0^1 \left\{ \frac{y}{x} \right\} u(x) dx \quad (12)$$

on $L^2(0,1)$ having the kernel function $\rho(y, x) := \left\{ \frac{y}{x} \right\}$ used for providing equivalent formulations of Riemann Hypothesis is injective.

Proof

Using Lemma 3.1 we just showed that the integral operator T_ρ is injective, so, $N_{T_\rho} = \{0\}$. ■

References

- [1] Adam, D. (1989) "On Mesh Independence by Preconditioning", *Int. J. Of Comp. Math.*, 1989.
- [2] Alcantara-Bode, J. (1993) "An Integral Equation Formulation of the Riemann Hypothesis", *Integr Equat Oper Th*, Vol. 17, 1993.
- [3] Balazard M., Saias E. (2000) "The Nyman–Beurling equivalent form for the Riemann hypothesis", *Expo. Math.* 18, 131–138 (2000)
- [4] Beurling, A.(1955) "A closure problem related to the Riemann zeta function", *Proc. Nat. Acad. Sci.* 41 pg. 312-314, 1955.
- [5] Buescu, J., Paixa~o A. C. (2007) "Eigenvalue distribution of Mercer-like kernels", *Math.Nachr.*280, No.9–10, pg. 984 – 995, 2007.
- [6] Chang, C.H., Ha, C.W. (1999) "On eigenvalues of differentiable positive definite kernels", *Integr. Equ. Oper. Theory* 33 pg. 1-7, 1999.
- [7] Fan, K. (1954) "Inequalities for eigenvalues of Hermitian matrices", *Nat. Bur. Standards Appl. Math. Ser.* 39 pg. 131-139, 1954.
- [8] Mercer, J., (1909) "Functions of positive and negative type and their connection with the theory of integral equations", *Philosophical Transactions of the Royal Society A* 209, 1909
- [9] Reade, J.P., (1983) "Eigenvalues Of Positive Definite Kernels", *SIAM J. Math. Anal.* Vol. 14, No. 1" January 1983.