The spatiotemporal distribution of human brucellosis in mainland China from 2007-2016

Peifeng Liang

People's Hospital of Ningxia Hui Autonomous Region

Yuan Zhao

Ningxia Medical University

Jianhua Zhao

Ningxia Center for Disease Control and Prevention

Dongfeng Pan

People's Hospital of Ningxia Hui Autonomous Region

Zhongqin Guo (✉ guozq_ny@163.com)

Ningxia Medical University

Research article

Keywords: Human brucellosis, mainland China, geographic information system, spatial autocorrelation analysis

Posted Date: February 25th, 2020

DOI: <https://doi.org/10.21203/rs.2.21000/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Infectious Diseases on March 27th, 2020. See the published version at <https://doi.org/10.1186/s12879-020-4946-7>.

Abstract

Background: Despite the considerable efforts made to address the issue of brucellosis worldwide, its prevalence in dairy products continues to be difficult to estimate and represents a key public health issue around the world. The aim of the study was to understand the epidemiology of this disease in mainland China. We set out to investigate the yearly spatial distribution and possible hotspots of the disease.

Methods: Human brucellosis data from mainland China between 2007 and 2016 were collected from the China Information System for Disease Control and Prevention. A geographic information system ArcGIS10.3 (ESRI, Redlands) was used to identify potential changes in the spatial and temporal distribution of human brucellosis in mainland China during the study period.

These distributions were evaluated using three-dimensional trend analysis and spatial autocorrelation analyse. A gravity-center was used to analyse the migration track of human brucellosis. **Results:** A total of 399,578 cases of human brucellosis were reported during the 10-year study period. The monthly incidence of brucellosis in China demonstrates clear seasonality. Spring and summer are the peak seasons, while May is the peak month for brucellosis. Three-dimensional trend analysis suggests that brucellosis is on the rise from south to north, and that the epidemic situation in northern China is more severe. Between 2007 and 2016, the overall migration distance of the brucellosis incidence gravity-center was 906.43 km, and the direction was southwest. However, the overall gravity center of brucellosis was still in the northern part of China. In the global autocorrelation analysis, brucellosis in China demonstrated a non-random distribution between 2013 and 2014, with spatial autocorrelation ($Z > 1.96$, $P < 0.05$) and a clustering trend, while no clustering trend was found from 2007 to 2012 or from 2015 to 2016. In the local autocorrelation analysis, a Low-Low cluster phenomenon was found in the south of China in 2013 and 2014. **Conclusion:** Human brucellosis remains a widespread challenge, particularly in northern China. The hotspots highlight potential high-risk areas which may require special plans and resources for monitoring and controlling the disease. **Keywords :** Human brucellosis; mainland China; geographic information system; spatial autocorrelation analysis

Background

Brucellosis, which is also known as undulant fever, is one of the most common zoonoses in the world [1-4]. It is an infectious disease which can be acute or chronic. It is caused by brucella, which is a natural epidemic disease [5]. Brucellosis is globally distributed. More than 170 countries and regions around the world have recorded brucellosis epidemics, accounting for approximately one-sixth to one-fifth of the world's population [6]. There are between five and six million patients who are living with brucellosis worldwide, the number of cases reported to the WHO each year is 500,000[7-10]. The Mediterranean region as well as Asia, Central and South America are the areas with the highest incidences[11-15].

In China, the first incidence of brucellosis occurred for two foreign patients in Shanghai in 1905 and was recorded as Malta fever[16]. However, several patients in China with similar clinical symptoms had been observed in the 10 years before that report. Following the 1905 report, three more cases were reported in Chongqing in 1906[17]. By the 1950s and 1960s, brucellosis had become more prevalent for Chinese

people and animals. Subsequently, from the 1980s to the early 1990s, activities for the prevention and control of brucellosis have gradually been introduced in mainland China, meaning that the brucellosis epidemic had been basically controlled[18]. During this time, the human infection rate was just 3%, the incidence rate was only 0.02/100,000[19]. However, brucellosis has rebounded since 1993, with sporadic cases increasing at a rate of between 30% and 50%. In 2015, the incidence of human cases reached 4.18/100,000. In the ranking of statutory infectious diseases, it rose from 16th place in 2000 to 6th place in 2014. A strong upward trend such as this is extremely rare among all such reported infectious diseases.

Previous studies have demonstrated that the global epidemiology of brucellosis has changed dramatically over the past decades, particularly in industrialized countries [20]. The spatial distribution characteristics of a brucellosis outbreak also have changed distinctly[21], although whether this change has temporal and spatial covariance requires further investigation.

The aim of our study was to use a spatial distribution model to analyze the spatial distribution characteristics and correlations of human brucellosis in mainland China between 2007 and 2016, and so to provide theoretical support for the prevention and control of human brucellosis in the context of local conditions.

Methods

(See Methods in the Supplementary Files)

Results

1. Epidemic situation of human brucellosis

(1) Time distribution

During the study period of 2007 to 2016, there were 399,578 reported cases of human brucellosis in mainland China. Annual case totals ranged from 19721 in 2007 to 57222 in 2014, with a mean number of approximately 34498 (95% CI: 25610, 43386) cases reported per year. The annual mean incidence of brucellosis was 2.97/100,000, with a low of 1.50/100,000 in 2007 and a high of 4.22/100,000 in 2014. As a whole, annual incidences follow a growth trend, with a linear relationship of $y=0.0211x+1.7003$ from 2007 to 2016 (Figure 1). Furthermore, annual and monthly incidences fluctuated during this 10-year period. Between 2007 and 2009, the annual average incidence of brucellosis in China demonstrated an upward trend. Following that, a short-term decline occurred in 2010, after which incidences of the disease began to rise again, continuing to rise until 2014, and then decline again in 2015 and 2016. A purely temporal analysis between 2007 and 2016 demonstrated that high-incidence seasons of human brucellosis were the time between late spring and early summer, while the incidence was highest in May.

(The histogram shows monthly incidences and the dots show annual incidences of brucellosis. Annual incidences have a $y=0.0211x+1.7003$ relationship. Results of a purely temporal analysis between 2007 and 2016 demonstrated that high-incidence seasons of human brucellosis occurred between late spring and early summer, with incidences highest in May.)

(2) Spatial distribution

Incidences of human brucellosis in each province vary significantly, with an interquartile range of 4.25/100,000 (0.045~4.297/100,000) and a range of 70.10/100,000 (0~70.10/100,000). We defined regions with an incidence greater than 10/100,000 as high incidence areas, while incidences lower than 0.999/100,000 were defined as low incidence areas. Over the total study period, high incidence areas increased from two to five, with the number of high incidence provinces and cities increasing year by year. In 2007, there were six provinces (6/31, 19.35%) with no cases of human brucellosis. These were mainly located in central and southern regions such as Guizhou, Hunan, Jiangxi and Hainan. As time passed, the epidemic area gradually expanded into all provinces and cities in China. The spatial and temporal distribution map (Figure 2) shows that 31 provinces (municipalities and autonomous regions) had incidences between 2007 and 2016. The epidemic mainly occurred in the north of China, such as the Inner Mongolia and Shanxi provinces. After 2009, the high incidence area grandly expanded in other northern areas, such as the Heilongjiang, Xinjiang and Ningxia provinces. In Inner Mongolia, disease incidence was the highest in history, exceeding the 50/100,000 population between 2009 and 2011. Between 2014 and 2016, incidences increased considerably in areas which had been historically low areas such as Gansu, Qinghai and Shandong. This was most marked in Ningxia, where incidences increased sharply from 0.4636/100,000 in 2007 to 43.6557/100,000 in 2015.

2 Three-dimensional trend analysis

The overall three-dimensional trend for 2007-2016 indicated an uneven distribution of human brucellosis in mainland China (Figure 3). The curve on the XZ plane represents the trend of East-West direction, while the curve on the YZ plane represents the change of North-South direction. Each point represents the projection of incidence. It can be seen that incidences of human brucellosis in the northern region were higher than in the southern region. However, in the east-west direction, incidences of brucellosis were evenly distributed in mainland China..

3 Gravity center migration

Most cases of human brucellosis were reported in northern China during 2007 to 2016, while most provinces in northern China experienced a serious epidemic (Figure 4). Between 2007 and 2012, the gravity-center of brucellosis was concentrated in the central area of Inner Mongolia. After 2012, there was a trend of gradual migration to the southwest. However, the distance and direction of annual migration were not obvious. Results showed that brucellosis was still prevalent in the northern part of China. During the study period, the gravity-center of brucellosis migrated 906.43 km southwest. The distance and direction of annual migration can be seen in Table 1.

Table 1. Distance and direction of center of gravity migration of brucellosis between 2007 and 2016.

Time	Distance	Direction	Time	Distance	Direction
2007-2008	69.89km	northeast	2008-2009	53.39 km	northeast
2009-2010	49.95km	northwest	2010-2011	140.28km	southwest
2011-2012	31.91 km	southwest	2012-2013	281.90km	southwest
2013-2014	344.76km	southwest	2014-2015	119.16km	southwest
2015-2016	80.72km	northwest			

4. Spatial autocorrelation analysis

(1) Global spatial autocorrelation

Using provincial units to carry out the global autocorrelation analysis, we obtained Moran's *I* value, variance, *Z* score and *P* value from 2007 to 2016, respectively (see Table 2). The values of Moran's *I* were 0.1179 and 0.1181 respectively for 2013 and 2014, while *Z* values were greater than 1.96 (all *P* < 0.05), indicating that incidences of brucellosis in China between 2013 to 2014 had a non-random distribution, while there was no spatial autocorrelation in either 2007-2012 or 2015-2016 (*Z* < 1.96, *P* > 0.05, meaning there was no clustering trend).

Table 2. Global autocorrelation of Moran's *I* values of brucellosis in mainland China from 2007 to 2016.

Year	Moran's <i>I</i>	Variance	Z Score	PValue	Aggregation
2007	0.041785	0.002682	1.450621	0.146886	NO
2008	0.051605	0.002669	1.644175	0.100140	NO
2009	0.043654	0.002241	1.626341	0.103877	NO
2010	0.040263	0.002109	1.602407	0.109066	NO
2011	0.043491	0.002197	1.638966	0.101220	NO
2012	0.076004	0.003579	1.827598	0.067610	NO
2013	0.117938	0.004963	2.147162	0.031780	YES
2014	0.118131	0.005379	2.065198	0.038904	YES
2015	0.093595	0.005191	1.761692	0.078121	NO
2016	0.092068	0.005165	1.744833	0.081014	NO

(2) Local spatial autocorrelation

While global clustering indices evaluate the distribution of events, they do not specify the location of hotspots. To identify areas characterized by a higher rate of human brucellosis incidences, a local spatial autocorrelation analysis was carried out for the years which had a statistical significance of Moran's *I* value, and a local spatial autocorrelation aggregation distribution map was generated (Figure 5). There were no High-High, High-Low or Low-High cluster areas in China in 2013, while Low-Low cluster areas tended to be concentrated in the southern provinces of China, including the provinces of Yunnan, Guizhou, Guangxi, Hunan, Guangdong, Hubei, Jiangxi, Fujian, Zhejiang and Anhui, with an average incidence of 0.08/100,000. The new High-Low cluster area in 2014 was Xinjiang, while the Low-Low cluster area decreased compared to 2013, now including Yunnan, Guizhou, Guangxi, Hunan, Guangdong, Jiangxi, Fujian, Zhejiang and Anhui provinces. The mean incidence of Low-Low clusters was 0.14/100,000 in 2014, meaning the incidence of Low-Low clusters in 2014 was higher than in 2013. Incidences in the Hubei Province increased rapidly from 0.08/100,000 in 2013 to 0.34/100,000 in 2014, thus reducing Hubei Province in Low-Low cluster areas in 2014. Cluster maps for 2013 and 2014 demonstrate that the cluster area in China was relatively stable, and that compared to the spatial distribution map of incidence, the Low-Low cluster area in each year tends to be the same as the low incidence area, all of which are located in the south of China.

Discussion

Exploring the spatial distribution of a disease is beneficial for guiding the development and implementation of disease surveillance and a control plan. The application of ArcGIS10.3 (ESRI, Redlands) in disease surveillance has yielded good results. Based on surveillance data about human brucellosis between 2007 to 2016, the temporal and spatial distribution of incidences in mainland China suggested that brucellosis in mainland China has unique spatial and temporal clustering characteristics.

In recent years, incidences of brucellosis in mainland China have been relatively high and growing rapidly. Incidences in 2014 are about 2.8 times of that 2007. In addition, brucellosis has obvious seasonality. Incidences were higher in spring and summer and reached the highest in May every year. This is largely consistent with findings from related studies[37-40]. The high incidence may be related to the vigorous development of animal husbandry in mainland China in the past 10 years, with a sharp increase in the number of livestock raised, frequent transactions and frequent transfer of livestock. However, grass-roots animal epidemic prevention departments are weak, and existing resources are not enough to meet the current increasing demand for epidemic prevention, making it difficult to deal with prevention and control measures such as immunization, quarantine and treatment of dead animals. High incidence rates of the disease in spring and summer are mainly due to the fact that it is the calving season, meaning that the chances of exposure to placenta increase. At this time, the contact probability of the relevant occupational personnel with abortion products and secretions of the sick livestock is greatly increased. If there are no effective protective measures for these occupational personnel, incidences will continue to increase.

It was found that brucellosis in China tends to show an upward trend from south to north, all of which began to spread from highly affected areas before gradually spreading outward. The gravity-center of brucellosis in China in recent years gradually moved southwest. However, overall migration distance and direction were small, and the gravity-center was always located in Inner Mongolia. The direction of migration was related to the location of the four major pastoral areas in China, which is similar to previously published research results[41]. High incidences of brucellosis were in the north of mainland China, where some areas showed much higher incidences than in the south of China. This may be because the north is the main gathering area of pastoral areas in China, meaning there are greater numbers of livestock being raised there as well as frequent livestock exchange.

A global spatial autocorrelation study demonstrated that brucellosis in China had a positive spatial correlation and clustering between 2013 and 2014, while there was no spatial autocorrelation between 2007 and 2012 or 2015 and 2016. Further local autocorrelation analysis found that the Low-Low concentration areas of brucellosis in China in 2013 were mainly in the southern part of the country, including the provinces of Yunnan, Guizhou, Guangxi, Hunan, Guangdong, Hubei, Jiangxi, Fujian, Zhejiang and Anhui, which is basically consistent with the low incidence areas. In 2014, the Low-Low cluster areas remained in the south of China. Understanding the spatial distribution of brucellosis in China is of great significance for controlling its occurrence and reducing the burden it causes, as well as

for optimizing the allocation of resources. A prevention policy of key monitoring should be adopted for High-High cluster areas, while favorable factors for disease control should be found in time for Low-Low cluster areas, so as to provide guidance for infectious disease surveillance. In recent years, incidences of brucellosis in Low-Low areas in China have been clustered in fixed places, meaning the influencing factors of this clustering need further investigation.

At the same time, since brucellosis is a natural epidemic disease, it is not infectious between people, meaning the natural and social environment have a great impact on it. Some natural factors in neighbouring areas may make brucellosis spread more easily, leading to a clustering distribution of the affected areas. It was found that the areas with a high incidence of brucellosis tended to be concentrated in the north of China (Figure 4). Previous studies[42-44] have found that besides raising livestock, brucellosis may also be connected to indirect environmental conditions such as altitude, rainfall, temperature, wind speed and other climatic factors. Li found that areas with a high incidence also tended to be located at middle and slightly higher altitudes (1600-1800m), while incidences of brucellosis at other altitudes were relatively low[45]. Low incidence areas tend to be concentrated in the southern part of China. As well as fewer livestock, there may be other influencing factors. These factors require further study.

Our study is not without limitations. Firstly, incidences of human brucellosis were underestimated to some extent, as our data was passively collected by a monitoring system, while surveillance data quality was influenced by comprehensive factors, such as the capacity of local health workers, availability of laboratory diagnostics, levels of awareness about the need to visit doctors and so on, all of which may have affected the study's accuracy. Secondly, our analysis was based on provincial-level data. Smaller spatial units may provide more location-specific information about the design and implementation stages of public health programs. Thirdly, this study only explored the clustering areas of brucellosis and did not analyze factors affecting the clustering distribution, therefore neglecting the impact of human and natural environment factors. In the future, we will therefore further explore the temporal and spatial distribution of brucellosis as well as the factors affecting its distribution.

In summary, given developments in animal husbandry in China, the prevention and detection of brucellosis urgently need to be strengthened. This spatial clustering study of brucellosis is helpful for identifying high-risk areas for brucellosis, and to a certain extent provides a basis for the decision-making of relevant departments. In terms of the distribution characteristics of brucellosis epidemic in this study, we suggest that the detection of brucellosis in northern China should be further strengthened, and that effective methods for controlling brucellosis should be found in southern China, where incidence of brucellosis is low. Governments at all levels should attach importance to the establishment of joint prevention and control mechanism among high-incidence areas. Relevant departments in different regions should strengthen both prevention and control throughout the year. More resources for prevention and control should be appropriately increased in these areas to curb the spread of brucellosis.

Conclusion

It may be concluded that human brucellosis continues to be a widespread challenge in mainland China, especially in the northwest provinces, which are high risk areas. This study utilized ArcGIS10.3 (ESRI, Redlands) to analyze the temporal and spatial distribution of brucellosis in China, thus contributing to the study of high incidence seasons and areas for brucellosis. However, further research should focus on an analysis of environmental, humanistic and socio-economic factors in order to determine risk factors affecting the occurrence and transmission of brucellosis. Such information has the potential to provide critical guidelines for policy makers to initiate prevention measures and control strategies, aimed at susceptible areas that might be high risk, and to prevent or lessen the incidence of human brucellosis in these areas.

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

Data were collected through an internet-based disease-reporting system (the China Information System for Disease Control and Prevention, CISDCP, <http://www.phsciencedata.cn/>).

Competing interests

None of the authors has a conflict of interest.

Fundings

This work was part supported by the Grant from Ningxia Provincial Natural Science Foundation(NO. NZ17187), by the Basic research project of Northwest University for Nationalities(NO. 31920180088).

Authors' contributions

Peifeng Liang participated in the design of the study, Yuan Zhao drafted the manuscript, Jianhua Zhao and Dongfeng Pan revised manuscript critically for important intellectual content. Zhongqin Guo managed the project implementation. All authors read and approved the final version of the manuscript.

Acknowledgement

We are very grateful to referees and editors for their careful reading and valuable comments. We would also like to thank Profs. Yu Zhao, Jiangping Li for their valuable discussion on this work. This study was supported by the Grant from Ningxia Provincial Natural Science Foundation (NO. NZ17187), by the Basic

research project of Northwest University for Nationalities (NO. 31920180088). The authors would like to express their gratitude to EditSprings (<https://www.editsprings.com/>) for the expert linguistic services provided.

Author details

1. Department of medical record and statistics, people's hospital of ningxia hui autonomous region, Yinchuan 750004, China
2. Department of Epidemiology and Biostatistics, School of Public Health and management, Ningxia Medical University, Yinchuan 750001, China
3. Ningxia Center for Diseases Prevention and Control, Yinchuan 750001, China
4. Department of Emergency, people's hospital of ningxia hui autonomous region, Yinchuan 750004, China

References

1. Mcdermott J, Grace D, Zinsstag J. Economics of brucellosis impact and control in low-income countries[J]. *Rev Sci Tech*. 2013; 32(1):249-261.
2. Farina F, Fuser R, Rossi M, [Scotton PG](#). Brucellosis outbreak in Treviso province caused by infected cheese from an endemic area[J]. *Infez Med*. 2008; 16(3):154-157.
3. [Park SH](#), [Lee YH](#), [Chu H](#), [Hwang SD](#), [Hwang KJ](#), [Choi HY](#), et al. Application of the Microagglutination Test for Serologic Diagnosis of Human Brucellosis[J]. *Osong Public Health & Research Perspectives*. 2012; 3(1):19-23.
4. [Luo J](#), [Zeng Z](#), [Song Y](#), [He H](#). Brucellosis in Takins, China[J]. *Emerging Infectious Diseases*. 2012; 18(9):1527-1529.
5. Ahmadkhani M, Alesheikh A. Space-time analysis of human brucellosis considering environmental factors in Iran[J]. *Asian Pacific Journal of Tropical Disease*. 2017; 7(5):257-265.
6. [Haerry TE](#), [Gehring WJ](#). Intron of the Mouse Hoxa-7 Gene Contains Conserved Homeodomain Binding Sites that can Function as an Enhancer Element in Drosophila[J]. *Proceedings of the National Academy of Sciences of the United States of America*. 1996, 93(24):13884-13889.
7. [Al-Ballaa SR](#), [Al-Balla SR](#), [Al-Aska A](#), [Kambal A](#), [Al-Hedaithy MA](#). Seasonal variation of culture positive brucellosis at a major teaching hospital[J]. *Annals of Saudi medicine*. 1994; 14(1):12-15.
8. Zeki Yumuk, David O'Callaghan. Brucellosis in Turkey-an overview[J]. *International Journal of Infectious Diseases*. 2012; 16(4).
9. Pappas Georgios. The changing Brucella ecology: novel reservoirs, new threats[J]. *International Journal of Antimicrobial Agents*. 2010; 36 Suppl 1.
10. [Chen JD](#), [Ke CW](#), [Deng X](#), [Jiang S](#), [Liang W](#), [Ke BX](#), et al. Brucellosis in Guangdong Province, People's Republic of China, 2005-2010[J]. *Emerging Infectious Diseases*. 2013; 19(5).

11. [Park MY, Lee CS, Choi YS, Park SJ, Lee JS, Lee HB](#). A Sporadic Outbreak of Human Brucellosis in Korea[J]. *Journal of Korean Medical Science*. 2005; 20(6):941-946.
12. [Gwida M, Al Dahouk S, Melzer F, Rösler U, Neubauer H, Tomaso H](#). Brucellosis-Regionally Emerging Zoonotic Disease[J]? *Croatian Medical Journal*. 2010; 51(4): p. 289-295.
13. [Hotez PJ, Savioli L, Fenwick A](#). Neglected Tropical Diseases of the Middle East and North Africa: Review of Their Prevalence, Distribution, and Opportunities for Control[J]. *PLoS Neglected Tropical Diseases*. 2012; 6(2):e1475.
14. [Dean AS, Crump L, Greter H, Schelling E, Zinsstag J](#). Global Burden of Human Brucellosis: A Systematic Review of Disease Frequency[J]. *PLoS Neglected Tropical Diseases*. 2012; 6(10):e1865.
15. Sprague L D, Al-Dahouk S, Neubauer H. A review on camel brucellosis: a zoonosis sustained by ignorance and indifference[J]. *Pathogens and Global Health*. 2012; 106(3):144-149.
16. Shengjie L, Hang Z, Weiyi X, Marius G, Zhuojie H, Jianxing Y et al. Changing Epidemiology of Human Brucellosis, China, 1955–2014[J]. *Emerging Infectious Diseases*, 2017, 23(2):184-194.
17. Zhong Z, Yu S, Wang X, Dong S, Xu J, Wang Y, et al. Human brucellosis in the People\'s Republic of China during 2005–2010[J]. *International Journal of Infectious Diseases*, 2013, 17(5):e289-e292.
18. Peng G, Wei W, Desheng H. Trends of reported human brucellosis cases in mainland China from 2007 to 2017: an exponential smoothing time series analysis[J]. *Environmental Health and Preventive Medicine*, 2018, 23(1):23-.
19. [Lai S, Zhou H, Xiong W, Gilbert M, Huang Z, Yu J, et al](#). Changing Epidemiology of Human Brucellosis, China, 1955-2014[J]. *Emerging Infectious Diseases*. 2017; 23(2):184-194.
20. Seleem M N, Boyle S M, Sriranganathan N. Brucellosis: A re-emerging zoonosis[J]. *Veterinary Microbiology*. 2010; 140(3-4):392-398.
21. [Pappas G, Papadimitriou P, Akritidis N, Christou L, Tsianos EV](#). The new global map of human Brucellosis[J]. *The Lancet Infectious Diseases*. 2006; 6(2):91-99.
22. [Cao M, Feng Z, Zhang J, Ma J, Li X](#). Contextual risk factors for regional distribution of Japanese encephalitis in the People’s Republic of China[J]. *Tropical Medicine & International Health*. 2010; 15(8).
23. Zhang W Y, Guo W D, Sun S H, et al. Human Brucellosis, Inner Mongolia, China[J]. *Emerging Infectious Diseases*. 2010; 16(12):2001-2003.
24. Lauren M. Scott, Mark V. Janikas. Spatial statistics in ArcGIS[M]// *Handbook of Applied Spatial Analysis*. 2010.
25. Du PL, Wu KS, Fang JY, Zeng Y, Xu ZX, Tang WR, et al. [Cervical Cancer Mortality Trends in China, 1991-2013, and Predictions for the Future](#)[J]. *Asian Pac J Cancer Prew*, 16(15), 6391-6396
26. [Rao HX, Zhang X, Zhao L, Yu J, Ren W, Zhang XL, et al](#). Spatial transmission and meteorological determinants of tuberculosis incidence in Qinghai Province, China: a spatial clustering panel analysis[J]. *Infectious Diseases of Poverty*. 2016; 5(1):45.

27. Fang J Y, Dong H L, Wu K S, Du P L, Xu Z X. Characteristics and Prediction of Lung Cancer Mortality in China from 1991 to 2013[J]. Asian Pacific journal of cancer prevention: APJCP, 2015, 16(14):5829-5834.
28. Lei, Song, Yingui, Cao, Wei, Zhou, et al. Study on the variation of arable land use and management countermeasures under rapid urbanization: the application of a gravity model in a regional perspective[J]. Environmental Monitoring and Assessment, 2019.
29. Xiaopeng Q, Liang W, Barker L, Lekachvili A, Xingyou Z. Comparison of ArcGIS and SAS Geostatistical Analyst to Estimate Population-Weighted Monthly Temperature for US Counties[J]. Journal of Resources and Ecology, 2012, 3(3):220-229.
30. Tobler W. A computer movie simulating urban growth in the Detroit region[J]. Econ Geogr. 1970; 46:234-40.
31. [Chen Y](#). New Approaches for Calculating Moran's Index of Spatial Autocorrelation[J]. PLoS ONE. 2013; 8(7):e68336-.
32. Anselin L. Local Indicators of Spatial Association-LISA[J]. Geographical Analysis. 1995; 27(2):93-115.
33. Khalid A A, Ali A Z. Spatial Autocorrelation of Cancer Incidence in Saudi Arabia[J]. International Journal of Environmental Research and Public Health. 2013; 10(12):7207-7228.
34. [Zhao X](#), [Huang X](#), [Liu Y](#). Spatial Autocorrelation Analysis of Chinese Inter-Provincial Industrial Chemical Oxygen Demand Discharge[J]. International Journal of Environmental Research and Public Health. 2012; 9(6):2031-2044.
35. [Xia J](#), [Cai S](#), [Zhang H](#), [Lin W](#), [Fan Y](#), [Qiu J](#), et al. Spatial, temporal, and spatiotemporal analysis of malaria in Hubei Province, China from 2004-2011[J]. Malaria Journal. 2015; 14(1):145.
36. [Zhang Y](#), [Shen Z](#), [Ma C](#), [Jiang C](#), [Feng C](#), [Shankar N](#), et al. Cluster of Human Infections with Avian Influenza A (H7N9) Cases: A Temporal and Spatial Analysis[J]. International Journal of Environmental Research & Public Health. 2015; 12(1):816-828.
37. [Park MY](#), [Lee CS](#), [Choi YS](#), [Park SJ](#), [Lee JS](#), [Lee HB](#). A Sporadic Outbreak of Human Brucellosis in Korea[J]. Journal of Korean Medical Science. 2005; 20(6):941-946.
38. [Karagiannis I](#), [Mellou K](#), [Gkolfinopoulou K](#), [Dougas G](#), [Theocharopoulos G](#), [Vourvidis D](#), et al. Outbreak investigation of brucellosis in Thassos, Greece, 2008[J]. [Euro Surveill](#). 2012; Mar 15;17(11). pii: 20116.
39. [De Massis F](#), [Di Girolamo A](#), [Petrini A](#), [Pizzigallo E](#), [Giovannini A](#). Correlation between animal and human brucellosis in Italy during the period 1997-2002[J]. Clinical microbiology and infection: the official publication of the European Society of Clinical Microbiology and Infectious Diseases. 2005; 11(8):632-636.
40. [Golshani M](#), [Buozari S](#). A review of Brucellosis in Iran: Epidemiology, Risk Factors, Diagnosis, Control, and Prevention[J]. Iran Biomed J. 2017; 21(6):349-359.
41. [Zhang J](#), [Yin F](#), [Zhang T](#), [Yang C](#), [Zhang X](#), [Feng Z](#), et al. Spatial analysis on human brucellosis incidence in mainland China: 2004-2010[J]. BMJ Open. 2014; 4(4):e004470.

42. Mai HM, Irons PC, Kabir J, Thompson PN. Herd-level risk factors for *Campylobacter* fetus infection, *Brucella* seropositivity and within-herd seroprevalence of brucellosis in cattle in northern Nigeria[J]. Preventive Veterinary Medicine. 2013; 111(3-4):256-267.
43. Ma JY, Wang H, Zhang XF, Xu LQ, Hu GY, Jiang H, et al. MLVA and MLST typing of *Brucella* from Qinghai, China[J]. Infectious Diseases of Poverty. 2016; 5(1):26.
44. Silva I, Dangolla A, Kulachelvy K. Seroepidemiology of *Brucella abortus* infection in bovids in Sri Lanka[J]. Preventive Veterinary Medicine. 2000; 46(1):51-59.
45. Li YJ, Li XL, Liang S, Fang LQ, Cao WC. Epidemiological features and risk factors associated with the spatial and temporal distribution of human brucellosis in China[J]. BMC Infectious Diseases. 2013; 13(1):547.

Figures

Figure 1

Time distribution of human brucellosis incidences in China from 2007 to 2016.

Figure 2

Spatial distribution of human brucellosis incidences in mainland China from 2007 to 2016 (1/100,000). Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Figure 3

Three-dimensional trend analysis of mean incidences of brucellosis in mainland China between 2007 and 2016. (The curve on the XZ plane represents the East-West direction trend, while the curve on the YZ plane represents change in the North-South direction. Overall, we can see that incidences of brucellosis in northern China are relatively high, although in recent years, incidences in southern China have gradually increased.)

Figure 4

The center of gravity migration track of human brucellosis in China from 2007 to 2016. (The small black dots in the picture show the gravity center of brucellosis between 2007-2016, which were mainly concentrated in northern China.) Note: The designations employed and the presentation of the material

on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Figure 5

Local spatial autocorrelation analysis of incidences of human brucellosis in mainland China between 2013 and 2014. (Bright red indicates a High-Low association, bright blue indicates Low-High, light red indicates High-High and light blue indicates Low-Low. In 2013, there was a Low-Low aggregation area. In 2014, there was a High-Low aggregation area and a Low-Low aggregation area.)Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Methods.pdf](#)