MiR-9-5p promotes rabbit preadipocyte differentiation by suppressing *leptin* gene expression

Gang Luo

Sichuan Agriculture University

Shenqiang Hu

Sichuan Agricultural University

Tianfu Lai

Sichuan Agricultural University

Jie Wang

Sichuan Agricultural University

Li Wang

Sichuan Agricultural University

Songjia Lai (✉ laisj5794@163.com)

Sichuan Agricultural University

Research

Keywords: MiR-9-5p; rabbits; pre-adipocyte; leptin; differentiation

Posted Date: May 12th, 2020

DOI: <https://doi.org/10.21203/rs.2.21007/v3>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Lipids in Health and Disease on June 5th, 2020. See the published version at <https://doi.org/10.1186/s12944-020-01294-8>.

Abstract

Background: MicroRNAs (miRNAs) are a class of small non-coding RNAs, which participate in the regulation of cell differentiation. Previous studies have demonstrated that miR-9-5p plays a key role in cancer cell development, but the mechanisms by which miR-9-5p regulates adipogenesis remain poorly understood. The present study intended to investigate its significance in producing rabbits with high-quality meat by observing the regulatory effect of miR-9-5p in preadipocytes and finding the related targets.

Methods: In this study, a dual-luciferase reporter assay was employed to validate the targeting relationship between miR-9-5p and *leptin* gene. We also utilized quantitative reverse transcription PCR (qRT-PCR), western blot, oil red-O staining assay, and determination of triglyceride content to analyze the regulation of miR-9-5p and *leptin* gene during adipocyte differentiation.

Results: The analysis demonstrated that during preadipocyte differentiation, miR-9-5p was up-regulated and the fat formation related biomarkers, i.e., fatty acid-binding protein 4 (*FABP4*), CCAAT-enhancer binding protein α (*C/EBP α*), and peroxisome proliferator activated receptor γ (*PPAR γ*) were also up-regulated. Meanwhile, the oil red-O staining assay revealed that the accumulation of lipid droplets increased. We also explored the expression pattern and role of miR-9-5p in adipogenesis using white preadipocytes. The results showed that miR-9-5p was up-regulated during preadipocyte differentiation, and overexpression of miR-9-5p enhanced lipid accumulation. Furthermore, we found that the overexpression of miR-9-5p significantly up-regulated the expression of marker genes, *PPAR γ* , *C/EBP α* and *FABP4*, and increased the protein levels of *PPAR γ* and triglyceride content. The results suggest that miR-9-5p might be involved in the regulation of rabbit preadipocyte differentiation. We predicted that *leptin* is the target gene of miR-9-5p, by using bioinformatics tools and the conclusion was validated by a luciferase reporter assay. Finally, we verified that the knock-down of *leptin* by si-*leptin* promoted preadipocyte differentiation in rabbits.

Conclusion: The results of the present study indicate that miR-9-5p regulates white preadipocyte differentiation in rabbits by targeting the *leptin* gene.

1. Introduction

Adipose tissue, which may be brown, beige, or white, not only plays an important role in energy metabolism by generating heat and storing lipids but also serve as an endocrine organ. White adipose tissue is widely dispersed and has the ability to differentiate. Adipose tissue-derived stem cells (ASCs) are stimulated to undergo transformation into preadipocytes, which in turn rapidly mature to adipocytes (Francisco et al., 2016). Adipogenesis, is a complex process in which fibroblast-like preadipocytes differentiate into lipid-laden and insulin-responsive adipocytes (Lefterova and Lazar, 2009). This process requires sequential activation of numerous transcription factors and noncoding RNAs. Several transcription factors such as peroxisome proliferator-activated receptor- γ (*PPAR γ*) and members of the

CCAAT/enhancer-binding family of proteins (*C/EBPα*) play a significant role in the regulation of adipogenesis (Hamam et al., 2014; Prayitno et al., 1997). Expression of *FABP4* is induced during adipocyte differentiation and is controlled by insulin and *PPARγ* agonists [Hunt et al., 1986]. *Leptin* plays a pivotal role in regulating food intake, energy expenditure, and fat deposition in mammals (Matteri, 1998). It can promote lipolysis and thermogenesis by influencing the sympathetic nervous system (Eikelis, 2004). In addition, it can also directly regulate the metabolism of some tissues and cells (McClelland et al., 2004; Sanigorski et al., 2013). In a study conducted by Thierry et al. (1999), *leptin* resulted in a dose and time-dependent increase in type1 collagen, a decrease of adipsin and *leptin* mRNA levels, and a 50% decrease in lipid droplet formation. Zhou et al. (1999) reported fat loss, downregulation of lipogenic enzymes, and suppression of the transcription factor *PPARγ* caused by adenovirus-induced hyperleptinemia in normal rats. The absence of fat in transgenic mice led to a reduction in *leptin* levels and marked metabolic alterations including hyperlipidemia, insulin resistance, and diabetes (Moitra et al., 1998). Recent literature shows that *leptin* enhances the differentiation of osteoblasts and inhibits the differentiation of adipocytes in human bone marrow stromal cells (Thierry et al., 1999). Altogether, these findings indicate that *leptin* may act locally in adipose tissue to influence the formation of mature adipocytes.

Furthermore, microRNAs (miRNAs), a class of short noncoding RNAs, are involved in the inhibition of gene expression by binding to the 3'-UTRs of the target mRNAs and have been demonstrated to regulate lipid metabolism (Bagga et al., 2005; Lagos-Quintana, 2001). miRNAs have also been shown to be involved in cell differentiation (Zhang et al., 2017). In goats, miR-183 promotes preadipocyte differentiation by suppressing the expression of Smad4 (Zhao et al., 2018) and miR-21a-5p induces preadipocyte differentiation by targeting Mitogen-activated protein kinase 2/kinase 3 (*MAPK2K3*) through *MKK3/p38/MAPK* signaling pathway in mice (Xie et al., 2016). MiR-33b downregulates the differentiation of porcine preadipocytes (Taniguchi et al., 2014). MiR-26b promotes adipocyte differentiation in 3T3-L1 cells, through targeting *PTEN* (Li et al., 2017), another study has demonstrated that miR-148a-3p promotes rabbit adipocyte differentiation by targeting *PTEN* in rabbits preadipocytes (He et al., 2018). MiR-9-5p is a highly conserved miRNA that is primarily expressed in the central nervous system (Kapsimali et al., 2007; Wienholds E., 2005). A study conducted by Li et al. (2017) showed that miR-9-5p can promote mesenchymal stem cell (MSC) migration by activating b-catenin signaling pathway. b-catenin plays an important role in the transduction of Wnt signals. Ross et al. (2000) found that the classical Wnt signal can inhibit fat differentiation. It has also been shown that *C/EBPα* or *PPARγ* can inhibit Wnt signaling, which in turn can affect the differentiation of adipocytes by reducing the level of b-catenin (Moldes et al., 2003). Yet, the role of miR-9-5p in differentiation of adipocytes remains unclear.

In the present study, we isolated and cultured rabbit perirenal adipose cells. Then, we studied the effect of overexpression or inhibition of miR-9-5p on adipocyte differentiation. In addition, we also predicted and verified that *leptin* is the target gene of miR-9-5p. Finally, we found that miR-9-5p can influence the differentiation of adipocytes by downregulating the expression of *leptin* gene.

2. Materials And Methods

2.1. Animal and tissue collection

All the experimental procedures using rabbits in this study were conducted under a protocol approved by the Institutional Animal Care and Use Committee, in the College of Animal Science and Technology, Sichuan Agricultural University, China. The rabbits were killed by neck folding. Perirenal adipose tissues were collected from 3 newborn New Zealand rabbits, which were raised under standard conditions at the Sichuan Agricultural University farm (Yaan, Sichuan, China).

2.2. Cell isolation, culture, and induction of adipogenesis

Adipose tissue was rinsed thrice with PBS and then minced and digested with 0.25% collagenase type I (Gibco, Carlsbad, CA, USA) at 37 °C for 1 h. The mixture was then added to the complete medium (CM; DM/F12, 10% fetal bovine serum, 2% penicillin-streptomycin) (Gibco) followed by filtration through 70-nm and 40-nm cell sieves, respectively. The final mixture was centrifuged at 1200×g for 5 min to collect the preadipocytes. The preadipocytes were then seeded into a culture flask with complete medium and incubated at 37 °C in a humidified incubator with 5% CO₂. Culture medium was changed every two days and the cells were frozen for future studies. After the cells reached about 70% confluence, an adipogenic cocktail (0.5 mM 3-isobutyl-1-methylxanthine, 10% FBS, 1 μM dexamethasone, and 1.7 μM insulin) was added into the growth medium to induce differentiation. The cells were further incubated for 72 h, after which the medium was replaced with maintenance medium (growth medium supplemented with 1.7 mM insulin per 50 mL) and incubated for an additional 72 h. Thereafter, the cells were cultured in a growth medium until maturation at ten days.

2.3. Cell transfection

Preadipocyte cells were seeded into 24-well or 6-well plates and transfected using Lipofectamine 3000 (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions after the cells had reached 70% confluency. The final concentrations of the negative control miRNA mimics (NC miR-Mimic), negative control inhibitor (inhibitor NC), and negative control *leptin* siRNA (siRNA NC) were 50 nM, 100 nM, and 100 nM, respectively. Three independent repetitions were performed for each treatment group. The cells were harvested at different time intervals after transfection and used to study adipogenic differentiation.

2.4. RNA isolation and qPCR analysis

Total cellular RNA was extracted using RNAiso Plus reagent (TaKaRa, Shiga, Japan) following manufacturer's instructions. The concentration and quality of the RNA was evaluated using Nano Drop 2000 UV-Vis Spectrophotometer (Thermo Scientific, Waltham, MA, USA). RNA purity was evaluated by determining the ratios of absorbance at 260 nm and 280nm (A260/A280) and at 260 nm and 230 nm (A260/A230). First-strand cDNA of total RNA and small RNA was synthesized using the Prime Script RT

reagent Kit (Takara, Japan) and the SYBR® Prime Script™ miRNA reverse transcription Kit (Takara), respectively, according to the manufacturer's protocol. The corresponding cDNA was stored at –20 °C. qPCR was performed using SYBR premix Ex Taq II (Tli RNase H Plus) (Catalog No. RR820A; Takara) on CFX96 system (Bio-Rad, Hercules, CA, USA). U6 was selected to normalize the expression of miRNA, while the internal reference used for mRNA expression was actin. The RT-qPCR procedure was as follows: Initial denaturation for 3 min at 95 °C, 39 cycles of denaturation at 95 °C for 30 s, annealing at the optimum temperature for 30 s, and final extension for 66 s with temperature increments of 0.5 °C/s from 65 °C to 95 °C. The $2^{-\Delta\Delta Ct}$ method was used to analyze the relative expression of each gene. The sequences of the primers used for qPCR are shown in Table 1.

2.5. Western blotting

Total protein from the cell samples was extracted with the Total Protein Extraction Kit (Sangon, Shanghai, China) following the manufacturer's protocol and protein content was quantified with the BCA Protein Assay Kit. Proteins (40 mg) were resolved on 8%–12% SDS-polyacrylamide gels and transferred to a PVDF membrane (Bio-Rad). The PVDF membrane was rinsed with TBS-Tween20 (TBST) and blocked for 2 h in skimmed milk. Then, the membranes were incubated overnight with anti-PPAR γ (Santa Cruz Biotechnology,) and anti- β -actin (Abs, Beijing, China) according to the respective instructions. The PVDF membranes were then washed thrice and incubated with the secondary goat anti-mouse IgG (H + L) (Transgen, Illkirch-Graffenstaden, France) for 2 h. The protein bands were incubated with chemiluminescence reagents (Beyotime) after being washed four times. The images were obtained with a Bio-Rad GelDoc system equipped with a Universal Hood III (Bio-Rad), and the integrated optical density (IOD) was calculated using Gel-Pro Analyzer 4.0.0.4. Actin was used as an internal control.

2.6. Oil Red-O staining and determination of triglyceride content

Adipocytes were washed thrice with phosphate-buffered saline (PBS) and fixed in 4% paraformaldehyde for 30 min. The fixed cells were stained with 1% filtered oil red-O solution for about 30 min in the dark. The adipocytes were then washed and observed under a phase contrast microscope. Finally, oil red-O was eluted from the stained cells with isopropanol and quantified by measuring the optical density values at 510 nm wavelength. Intracellular triglyceride (TG) content was quantified with the TG Assay Kit (Applygen, Beijing, China) according to the manufacturer's protocol. The TG content (nmol/mg protein) was normalized to the cellular protein concentration, and the protein concentrations were measured with the BCA Protein Assay Kit (Beyotime, Shanghai, China).

2.7. Prediction and verification of the target gene of miR-9-5p

MiRwalk (<http://zmf.umm.uni-heidelberg.de/apps/zmf/mirwalk2/>) was employed to predict the target genes of miR-9-5p. *Leptin* was found to be the target gene of miR-9-5p. The 3'UTR of *leptin* containing the miR-9-5p target site was cloned into the Sac I-Xba I site of the pmirGLO Vector (Promega, Madison, WI) to construct luciferase reporter plasmids. The primers used for plasmid construction are listed in Table 1. Multiple cloning sites were located downstream of firefly luciferase gene. HeLa cells were seeded into 24-

well plates in triplicate. Then, the pmirGLO-*leptin*-3' UTR of wild or mutant type was co-transfected with the synthetic miR-9-5p mimic into HeLa cells after the cell density reached 70–80%. Firefly luciferase (luc2) activity was measured 48 h after transfection and normalized to Renilla luciferase activity according to the TransDetect® Double Luciferase Reporter Assay Kit instructions (Transgen, Beijing, China).

2.8. Statistical analysis

The RT-qPCR results of mRNA expression were normalized to the geometric average of *Actin* identified stability, and the expression level of miRNA was normalized to the expression of U6 snRNA, by an optimized comparative Ct ($2^{-\Delta\Delta Ct}$) value method. The entire data is presented as mean $\pm$ standard error (SEM). The student's t-test in GraphPad Prism7 (GraphPad Software, La Jolla, CA, USA) was used to assess the difference between groups. $P < 0.05$ and $P < 0.01$ were deemed to be significant and highly significant, respectively.

3. Results

3.1. Expression profiling of miR-9-5p during rabbit preadipocyte differentiation

In the process of inducing differentiation, the adipocytes reached full differentiation, depositing large lipid droplets (Fig. 1A). Results of oil red-O staining showed that lipid droplets rapidly increased during preadipocyte cell differentiation (Fig. 1B). Simultaneously, the expression of miR-9-5p increased rapidly after three days of induction and peaked on the ninth day (Fig. 1C). Correspondingly, adipogenic marker gene *FABP4* had the highest mRNA expression on the sixth day whereas *PPAR γ* and *CEBPa* had the highest expression on the third day after induction (Fig. 1D, E, F).

3.2. Upregulation of miR-9-5p promotes rabbit preadipocyte differentiation

In order to investigate the function of miR-9-5p in rabbit preadipocyte cell differentiation, we first detected the expression level of miR-9-5p in the cells transfected with miR-9-5p mimics and NC during differentiation. After induction, miR-9-5p showed a higher expression level in the mimic group than in the NC group ($P < 0.01$) (Fig. 2A). Oil red-O staining results (including quantitative detection; $P < 0.01$) showed that the lipid accumulation was higher in the mimic group as compared with the NC group (Fig. 4B, C). In addition, there was a significant increase in triglycerides owing to the overexpression of miR-9-5p (Fig. 2D). The mRNA levels of adipogenic markers including *PPAR γ* and *C/EBPa* were significantly higher in the mimic group than NC group on the first and third days ($P < 0.01$) (Fig. 2E, F, and G). Western blot analysis further indicated that *PPAR γ* protein levels were significantly higher in the mimic group than NC group after two days of transfection ($P < 0.05$) (Fig. 2H, I).

3.3. Downregulation of miR-9-5p inhibited rabbit preadipocyte differentiation

Next, we examined the differentiation of preadipocytes when endogenous miR-9-5p was effectively inhibited with the miR-9-5p inhibitor. As shown in Figure 3A, the expression levels in the inhibitor groups

were significantly lower than in the NC group on third, 5th, 7th, and 9th day after transfection ($P < 0.01$). We also found that down-regulation of miR-9-5p decreases lipid accumulation in the preadipocytes cells as detected by oil red-O staining (Fig. 3B, C) and TG content (Fig. 3D). The mRNA expression of the common differentiation-related genes (*PPAR γ* and *C/EBP α*) was lower in the inhibition groups on the first and third day after transfection (Fig. 3E, F), whereas the expression of *FABP4* (another common differentiation-related gene) was lower on the fifth and seventh day after transfection in the inhibition groups ($P < 0.01$) (Fig. 3G). In addition, *PPAR γ* expression level also decreased after miR-9-5p inhibition (Fig. 3H, I).

3.4. *Leptin* is a target gene of miR-9-5p

To further determine the function of miR-9-5p, bioinformatics analysis was carried out to predict direct targets of miR-9-5p. MiRWalk2.0 (<http://zmf.umm.uni-heidelberg.de/apps/zmf/mirwalk2/>) software revealed that *leptin* contains the target sites for miR-9-5P (Fig. 4A). To verify the existence of direct binding sites between miR-9-5p and *leptin*, the luciferase reporter assay was carried out and the results showed that the firefly luciferase activity significantly reduces during co-transfection of pmirGLO-LEPTIN-3' UTR and miR-9-5p agomirs (Fig. 4B).

3.5. MiR-9-5p downregulated *leptin* mRNA expression

To better understand the role of miR-9-5p during preadipocyte differentiation, we further validated the effect of miR-9-5p on the expression of *leptin*. As shown in Figure 5A, overexpression of miR-9-5p led to an obvious decrease in the mRNA levels of *leptin*. In contrast, when the endogenous miR-9-5p was inhibited with the miR-9-5p inhibitor, the expression of *leptin* increased as compared with the NC group (Fig. 5B).

3.6. Downregulation of *leptin* expression promoted rabbit preadipocyte differentiation

To further confirm whether *leptin* is involved in the differentiation of rabbit preadipocytes, si-*leptin* was utilized to knockdown endogenous *leptin*. We found that the accumulation lipids and triacylglycerol accelerated in the preadipocytes cells (Fig. 6A, B, C). When *leptin* was knocked down, the mRNA levels of *PPAR γ* and *C/EBP α* rapidly increased on the first and third day after transfection, whereas *FABP4* rapidly increased on the 5th and 7th (Fig. 6D, E, F). Furthermore, *PPAR γ* protein expression level significantly increased in si-*leptin* group than NC group, three days after transfection ($P < 0.05$) (Fig. 6G, H). However, after knocking down endogenous *leptin*, the expression levels declined rapidly (Fig. 6I).

4. Discussion

In recent years, miRNAs have been reported to be expressed in mammalian adipocytes and play an important role in the regulation of adipogenesis (Xu P et al., 2003). Previous studies have discovered a functional interplay between glucose-dependent insulin secretion, mir-9 levels, and *sirt1* protein in pancreatic islets β -cells (Ramachandran et al., 2011). It has also been found that miR-9 and *sirt1* are involved in glucose-dependent insulin secretion in islet β cells, and it has been proved that *sirt1* is a target

gene of miR-9 (Plaisance et al., 2006b). In addition, miR-9-5p could negatively regulate *sirt1* expression and plays an important role in the regulation of cell proliferation and migration (Qi et al., 2015). *Sirt1* can bind to two coenzyme factors of *PPAR γ* , which inhibit the binding of *PPAR γ* to the target gene and the differentiation of adipocytes (Picard et al., 2004). It has also been shown that miR-9-5p promotes the secretion of Granuphilin-a and reduces the secretion and synthesis of insulin by targeting *Onecut2* (Plaisance et al., 2006a). The results showed that *PPAR γ* ligand enhanced glucose uptake in adipose tissue by enhancing the expression of GLUT-4, which can improve insulin sensitivity (Hamm et al., 1999). Furthermore, *PPAR γ* is a marker gene of preadipocyte differentiation. So, miR-9-5p may regulate the differentiation of adipocytes.

The triglyceride content was found to be more and lipid droplets were larger after transfection of miR-9-5p. The upregulation of miR-9-5p increased the expression of adipogenic marker genes *FABP4*, *PPAR γ* , and *CEBPa*, whereas, the downregulation of miR-9-5p using the synthetic inhibitor markedly reduced the formation of neutral lipid droplets and suppressed the expression of marker genes at both the *PPAR γ* mRNA and protein levels. These results suggest that miR-9-5p can promote the differentiation of rabbit preadipocytes. Previous studies demonstrated that NADPH oxidase 4 (*NOX4*) is a target gene of miR-9-5p (Fierro-Fernandez et al., 2015) and *NOX4* is the key enzyme of reactive oxygen species (ROS). It was reported that insulin release is inhibited by increasing ROS in islet β -cells (Anvari et al., 2015; Den Hartigh et al., 2017) and the content of *leptin* is positively correlated with insulin (Anderwald et al., 2002; Papaspyrou-Rao et al., 1997). In the present study, a dual luciferase reporter assay was employed to validate that *leptin* is the target gene of miR-9-5p. The results suggested that miR-9-5p can negatively regulate *leptin* gene expression. Previous studies have shown miR-9-5p can regulate secretion and synthesis of insulin. Insulin can stimulate the secretion of *leptin* by increasing the mRNA expression of *leptin*, and in turn, *leptin* can directly inhibit the secretion of insulin (Emilsson et al., 1997). Studies have shown that insulin can improve the differentiation rate of preadipocytes and promote the accumulation of fat (Sato et al., 1996). It has also been found insulin may affect the expression of *PPAR γ* by affecting sterol regulatory element-binding protein-1c (*SREBP-1c*) (Garcia-Escobar et al., 2011). *Leptin* can also regulate the differentiation of adipocytes and previous studies have demonstrated that *leptin* can inhibit the synthesis of lipids and promote the decomposition of triglycerides in adipocytes (Flier and Spiegelman, 2001). So, miR-9-5p may target *leptin* to promote the differentiation of rabbit preadipocytes.

Leptin is secreted by adipocytes and the mRNA level of *leptin* decreases at a later stage of adipocyte differentiation (Ailhaud et al., 1992). Previous studies have shown that *leptin* is involved in a local ultra short negative feedback loop that regulates the differentiation of preadipocytes (Thierry et al., 1999). Literature also suggests that *leptin* suppresses specific biochemical processes, which contribute to lipid accumulation and adipocyte differentiation (Shimabukuro et al., 1997; Zhang, 1996). Furthermore, *leptin* can accelerate the proliferation of preadipocytes, inhibit the synthesis of lipids and promote the decomposition of triglycerides (Flier and Spiegelman, 2001). In the present study, we found that the suppressed expression of *leptin* dramatically promotes the differentiation of preadipocytes. This finding is in agreement with a previous study which suggested that *leptin* inhibits the differentiation of preadipocytes. Therefore, we speculated that miR-9-5p positively regulates rabbit preadipocyte

differentiation by inhibiting *leptin*. However, the biological basis and mechanism of this phenomenon need further research to get rabbits of high intramuscular fat content. In recent years, numerous studies have been motivated by the desire to get animals (e.g., cattle and pigs) with high intramuscular fat content, but have not yielded satisfactory outcomes. This is largely due to the fact that there is a need to increase the body fat and muscle fat in the same animal at the same time. However, rabbits are animals with particularly low-fat content. So, we can increase the body fat and intramuscular fat at the same time, and this makes it possible to grow rabbits with high intramuscular fat content.

5. Conclusions

In summary, our data demonstrated that miR-9-5p can promote the differentiation of rabbit preadipocytes and *leptin* acts as one of the downstream targets of miR-9-5p. In addition, knockdown of *leptin* promoted adipogenic differentiation. Altogether, we conclude that miR-9-5p promotes rabbit preadipocyte differentiation by suppressing the expression of *leptin* mRNA.

Abbreviations

FABP4: fatty acid-binding protein 4; C/EBP α : CCAAT-enhancer binding protein α ; PPAR γ : peroxisome proliferator activated receptor; ASCs: Adipose tissue-derived stem cells; MAPK2K3: Mitogen-activated protein kinase 2/ kinase 3; 3T3-L1: Mouse Embryonic mouse fibroblast; PBS: phosphate-buffered saline; NOX4: NADPH oxidase 4; ROS: reactive oxygen species; SREBP-1c: Sterol regulatory element-binding protein-1c.

Declarations

Acknowledgements

We thank the staff at our laboratory for their ongoing assistance.

Authors' contributions

GL, JW, SJL designed and directed the study. GL, TFL, LW performed the experiments, data analysis and drafted the manuscript. GL contributed to the analysis and writing of the manuscript. QSH critically reviewed drafts of the manuscript and made comments to improve clarity. All authors approved the final version of this article.

Funding

This work was supported by National rabbit industry technology system (CARS-43-A-2).

Availability of data and materials

All data generated or analyzed during this study are included in this published article.

Animal ethical approval

All experimental operations are carried out in strict accordance with the Experimental Animal Operation Regulations (No.18 [2014] of Sichuan Agricultural University), which formulated by the Sichuan Agricultural University experimental animal operation regulations and welfare management committee. Surgical procedures involving rabbits were performed according to the approved protocols of the Biological Studies Animal Care and Use Committee, Sichuan Province, China. Rabbits had free access to food and water under normal conditions and were humanely sacrificed as necessary to ameliorate suffering.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Animal Genetic Resources Exploration and Innovation Key Laboratory of Sichuan Province, Sichuan Agricultural University, Chengdu 611130, China.

References

- Ailhaud G, Grimaldi P, A, R N (1992) Cellular and Molecular Aspects of Adipose Tissue Development Annual Review of Nutrition 12:207-233
- Anderwald C, Müller G, Koca G, Fürnsinn C, Waldh usl W, Roden M (2002) Short-Term Leptin-Dependent Inhibition of Hepatic Gluconeogenesis Is Mediated by Insulin Receptor Substrate-2 Molecular Endocrinology 16:1612-1628
- Anvari E, Wikstrm P, Walum E, Welsh N (2015) The novel NADPH oxidase 4 inhibitor GLX351322 counteracts glucose intolerance in high-fat diet-treated C57BL/6 mice Free Radical Research 49:1-32
- Bagga S, Bracht J, Hunter S, Massirer K, Holtz J, Eachus R, Pasquinelli AE (2005) Regulation by let-7 and lin-4 miRNAs Results in Target mRNA Degradation Cell 122:553-563
- Brüning JC, Gautam D, Burks DJ, Gillette J, Kahn CR (2000) Role of Brain Insulin Receptor in Control of Body Weight and Reproduction Science 289:2122-2125
- Den Hartigh LJ, Omer M, Goodspeed L, Wang S, Wietecha T, O'Brien KD, Han CY (2017) Adipocyte-specific deficiency of NADPH oxidase 4 delays the onset of insulin resistance and attenuates adipose tissue inflammation in obesity Arteriosclerosis, thrombosis, and vascular biology 37:466-475

- Eikelis, N. (2004) Extra-adipocyte leptin release in human obesity and its relation to sympathoadrenal function *AJP: Endocrinology and Metabolism* 286:E744-E752
- Emilsson V, Liu Y-L, Cawthorne MA, Morton NM, Davenport M (1997) Expression of the Functional Leptin Receptor mRNA in Pancreatic Islets and Direct Inhibitory Action of Leptin on Insulin Secretion *Diabetes* 46:313-316
- Fierro-Fernandez M et al. (2015) miR-9-5p suppresses pro-fibrogenic transformation of fibroblasts and prevents organ fibrosis by targeting NOX4 and TGFBR2 *Embo Reports* 16:1358-1377
- Flier J, Spiegelman GB (2001) Obesity and the regulation of energy balance *Cell* 104:531-543
- Francisco R-O, Azahara R, Carolina G-L, Angel G, Concepción A (2016) Cell Models and Their Application for Studying Adipogenic Differentiation in Relation to Obesity: A Review *International Journal of Molecular Sciences* 17:1040-1066
- Garcia-Escobar E, Rodriguez-Pacheco F, Haro-Mora J, J (2011) Effect of insulin analogues on 3t3-l1 adipogenesis and lipolysis *European Journal of Clinical Investigation* 41:979-986
- Hamam D, Ali D, Kassem M, Aldahmash A, Alajez NM (2014) MicroRNAs as Regulators of Adipogenic Differentiation of Mesenchymal Stem Cells *Stem Cells and Development* 24:417-425
- Hamm J K, el Jack A k, Pilch P F (1999) Role of PPAR gamma in regulating adipocyte differentiation and insulin-responsive glucose uptake *Ann N Y Acad Sci* 892:134-145
- He H et al. (2018) miR-148a-3p promotes rabbit preadipocyte differentiation by targeting PTEN *In Vitro Cellular & Developmental Biology - Animal* 54:241-249
- Hunt CR, Ro JH, Dobson DE, Min HY, Spiegelman BM (1986) Adipocyte P2 gene: developmental expression and homology of 5'-flanking sequences among fat cell-specific genes *Proceedings of the National Academy of Sciences* 83:3786-3790
- Kapsimali M, Kloosterman WP, Bruijn ED, Rosa F, Wilson SW (2007) MicroRNAs show a wide diversity of expression profiles in the developing and mature CNS *Genome Biology* 8:R173
- Lagos-Quintana, M. (2001) Identification of Novel Genes Coding for Small Expressed RNAs *Science* 294:853-858
- Lefterova MI, Lazar MA (2009) New developments in adipogenesis *Trends in Endocrinology & Metabolism* 20:107-114
- Li G, Ning C, Ma Y, Jin L, Liu H (2017) miR-26b Promotes 3T3-L1 Adipocyte Differentiation Through Targeting PTEN *Dna & Cell Biology* 36:672

- Li X, L H, Q Y (2017) MiR-9-5p promotes MSC migration by activating β -catenin signaling pathway American Journal of Physiology-Cell Physiology 313:C80-C93
- Matteri R (1998) THE BIOLOGY OF LEPTIN: A REVIEW 76:1405-1420
- McClelland GB, Kraft CS, Michaud D, Russell JC, Mueller CR, Moyes CD (2004) Leptin and the control of respiratory gene expression in muscle Biochim Biophys Acta 1688:86-93
- Moitra J, Mason MM, Olive M, Krylov D, Vinson C (1998) Life without white fat: a transgenic mouse Genes & Development 12:3168-3181
- MOLDES M, ZUO Y, MORRISON RF, SILVA D, PARK B-H, LIU J, FARMER SR (2003) Peroxisome-proliferator-activated receptor γ suppresses Wnt/ β -catenin signalling during adipogenesis Biochemical Journal 376:607-613
- Papaspyrou-Rao S, Schneider SH, Petersen RN, Fried SK (1997) Dexamethasone Increases Leptin Expression in Humans in Vivo J Clin Endocrinol Metab 82:1635-1637
- Picard F et al. (2004) Sirt1 promotes fat mobilization in white adipocytes by repressing PPAR γ Nature 429:771-776
- Plaisance V, Abderrahmani A, Perret-Menoud V (2006a) MicroRNA-9 controls the expression of Granuphilin/Slp4 and the secretory response of insulin-producing cells Journal of Biological Chemistry 281:26932-26942
- Plaisance V, Abderrahmani A, Perret-Menoud V, Jacquemin P, Lemaigre F, Regazzi R (2006b) MicroRNA-9 Controls the Expression of Granuphilin/Slp4 and the Secretory Response of Insulin-producing Cells Journal of Biological Chemistry 281:26932-26942
- Prayitno DS, Phillips CJC, Omed H (1997) The Effects of Color of Lighting on the Behavior and Production of Meat Chickens Poultry Science 76:452-457
- Qi F, Hu JF, Liu BH, Wu CQ, Yu HY, Yao DK, Zhu L (2015) Mi R-9a-5p regulates proliferation and migration of hepatic stellate cells under pressure through inhibition of Sirt1 World Journal of Gastroenterology 21:82-97
- Ramachandran D, Roy U, Garg S, Ghosh S, Pathak S, Kolthur-Seetharam U (2011) Sirt1 and mir-9 expression is regulated during glucose-stimulated insulin secretion in pancreatic β -islets Febs Journal 278:1167-1174
- Ross SE, Hemati N, Longo Ka, Bennett CN, Lucas PC, Erickson RL, MacDougald Oa (2000) Inhibition of adipogenesis by Wnt signaling Science 289:950-953

- Sanigorski A, Cameron-Smith D, Lewandowski P, Walder K, Silva Ad, Morton G, Collier GR (2013) Impact of obesity and leptin treatment on adipocyte gene expression in *Psammomys obesus* Journal of Endocrinology 164:45-50
- Sato K, NAKANISHI N, MITSUMOTO M (1996) Culture Conditions Supporting Adipose Conversion of Stromal-Vascular Cells from Bovine Intramuscular Adipose Tissues Journal of Veterinary Medical Science 58:1073-1078
- Shimabukuro M et al. (1997) Direct antidiabetic effect of leptin through triglyceride depletion of tissues Proceedings of the National Academy of Sciences of the United States of America 94:4637-4641
- Taniguchi M, Nakajima I, Chikuni K, Kojima M, Mikawa S (2014) MicroRNA-33b downregulates the differentiation and development of porcine preadipocytes Molecular Biology Reports 41:1081-1090
- Thierry T, Francesca G, Sundeep K, D. JM, Bartolome B, Lawrence RB (1999) Leptin Acts on Human Marrow Stromal Cells to Enhance Differentiation to Osteoblasts and to Inhibit Differentiation to Adipocytes* Endocrinology 140 1630-1638
- Wang J, Obici S, Morgan K (2001) Overfeeding Rapidly Induces Leptin and Insulin Resistance Diabetes 50:2786-2791
- Warner RD, Greenwood PL, Pethick DW, Ferguson DM (2010) Genetic and environmental effects on meat quality Meat Science 86:171-183
- Warriss PD (1999) Meat science: an introductory text International Journal of Food Science & Technology 36:449-449
- Wienholds, E (2005). MicroRNA Expression in Zebrafish Embryonic Development Science 309:310-311
- Wood JD et al. (2008) Fat deposition, fatty acid composition and meat quality: A review Meat Science 78:343-358
- Xie X, Song J, Li G (2016) MiR-21a-5p suppresses bisphenol A-induced pre-adipocyte differentiation by targeting map2k3 through MKK3/p38/MAPK Biochemical and Biophysical Research Communications 473:140-146
- Xu P, Y VS, M G (2003) The Drosophila microRNA Mir-14 suppresses cell death and is required for normal fat metabolism Current Biology 13:790-795
- Zhang, S. (1996) Obese Gene Expression Alters the Ability of 30A5 Preadipocytes to Respond to Lipogenic Hormones Journal of Biological Chemistry 271:13939-13942
- Zhang WR et al. (2017) miR-143 regulates proliferation and differentiation of bovine skeletal muscle satellite cells by targeting IGFBP5 In Vitro Cell Dev Biol Anim 53 265-271

Zhao W, Yang H, Li J, Chen Y, Zhang H (2018) MiR-183 promotes preadipocyte differentiation by suppressing Smad4 in goats Gene 666:158-164

Zhou Y-T, Wang Z-W, Higa M, Newgard CB, Unger RH (1999) Reversing Adipocyte Differentiation: Implications for Treatment of Obesity Proceedings of the National Academy of Sciences of the United States of America 96:2391-2395

Table

Table 1 Primers used in this study

Gene name	Primer sequence (5'-3')	Tm/°C	Product size/bp
<i>leptin</i>	CTGTGCCCATGCGGAAAGTC AGTCCAAACCGACGACCCTC	61.4	104
β - <i>actin</i>	GGAGATCGTGCGGGACAT GTTGAAGGTGGTCTCGTGGAT	61.4	318
<i>PPARγ</i>	GAGGACATCCAGGACAACC GTCCGTCTCCGTCTTCTTT	61	168
<i>FABP4</i>	GGCCAGGAATTTGATGAAGTC AGTTTATCGCCCTCCCGTT	61.4	140
<i>C/EBPα</i>	GCGGGAACGAACAACAT GGCGGTCATTGTCACTGGTC	64	172
miR-9-5p mimic	UCUUUGGUUAUCUAGCUGUAUGA AUACAGCUAGAUAAACCAAAGAUA		
Negative Control (NC)	UUCUCCGAACGUGUCACGUTT ACGUGACACGUUCGGAGAATT		
inhibitor-9-5p	UCAUACAGCUAGAUAAACCAAAGA		
inhibitor NC	CAGUACUUUUGUGUAGUACAA		
si- <i>leptin</i> -1	CCACAAUGGACCAGACGUUTT AACGUCUGGUCCAUUGUGGTT		
si- <i>leptin</i> -2	CCAUUGUCACCAGGAUCAGTT CUGAUCCUGGUGACAAUGGTT		
si- <i>leptin</i> -3	CCGAAAUGUGAUCCAAAUATT UAUUUGGAUCACAUUUCGGTT		
Primers for gene 3'UTR cloning <i>leptin</i>	ATGGGTAAGTACATCAAGAG GAGGTCCGAAGACTCATTT		

Figures

A

B

C

D

E

F

Figure 1

Rabbit preadipocyte differentiation model (A) Oil Red-O staining of lipid droplets ; (B) quantitative detection of oil red O staining; (C) MiR-9-5p expression levels during preadipocyte differentiation under normal culture conditions; (D) FABP4 expression levels during preadipocyte differentiation under normal culture conditions; (E) CEBPa mRNA expression levels during preadipocyte differentiation under normal culture conditions; (F) PPAR γ expression levels during preadipocyte differentiation under normal culture conditions ("*", $P \leq 0.05$; "**", $P \leq 0.01$)

Figure 2

Overexpression of miR-9-5p promoted rabbit preadipocyte differentiation (A) MiR-9-5p expression levels during preadipocyte differentiation after transfecting with miR-9-5p mimics and NC, (B) Oil Red-O staining of lipid droplets on day 9; (C) Quantitative detection of oil red O staining on day 9; (D) Accumulation of triacylglycerol on day 9; (E) PPARγ expression levels during preadipocyte differentiation after transfecting with miR-9-5p mimics and NC; (F) CEBPα expression levels during preadipocyte differentiation after transfecting with miR-9-5p mimics and NC; (G) FABP4 expression levels during preadipocyte differentiation after transfecting with miR-9-5p mimics and NC (H, I) PPARγ protein levels during preadipocyte differentiation after transfecting with miR-9-5p mimics and NC (“*”, $P \leq 0.05$; “**”, $P \leq 0.01$)

Figure 3

Inhibition of miR-9-5p inhibited rabbit preadipocyte differentiation (A) MiR-9-5p expression levels during preadipocyte differentiation after transfecting with the miR-9-5p inhibitor and INC, (B) Oil Red-O staining of lipid droplets on day 9; (C) Quantitative detection of oil red O staining on day 9; (D) Accumulation of triacylglycerol on day 9; (E) PPAR γ expression levels during preadipocyte differentiation after transfecting with the miR-9-5p inhibitor and INC; (F) CEBP α expression levels during preadipocyte differentiation after transfecting with the miR-9-5p inhibitor and INC; (G) FABP4 expression levels during preadipocyte differentiation after transfecting with the miR-9-5p inhibitor and INC (H, I) PPAR γ protein levels during preadipocyte differentiation after transfecting with the miR-9-5p inhibitor and INC (“*”, $P \leq 0.05$; “**”, $P \leq 0.01$)

Conserved

	Predicted consequential pairing of target region (top) and miRNA (bottom)	Site type	Context++ score	Context++ score percentile	Weighted context++ score	Conserved branch length	P _{CT}
Position 181-188 of LEP 3' UTR	5' ...CCCCUUUUUUUGUAA-ACCAAGAG... 	8mer	-0.44	99	-0.44	3.798	0.75
hsa-miR-9-5p	5' AGUAGUCGACUACUUGUUCU

Figure 4

Leptin is a target miR-9-5p gene. (A) prediction of target gene (B) PmirGLO and pmirGLO-leptin 3' UTR were cotransfected with the mimic into Hela cells, and normalized firefly luciferase activity was determined (n=3). (** P<0.01).

Figure 5

MiR-9-5p negatively regulated leptin expression: (A) Leptin expression based on quantitative real-time polymerase chain reaction (QRT-PCR) during adipogenesis with miR-9-5p overexpression. (B) Leptin expression based on QRT-PCR during adipogenesis with miR-9-5p knockdown. ("**", $P \leq 0.01$)

Figure 6

Inhibition of leptin gene promoted rabbit preadipocyte differentiation (A) Oil Red-O staining of lipid droplets on day 9; (B) Quantitative detection of oil red O staining on day 9; (C) Accumulation of triacylglycerol on day 9; (D) PPARγ expression levels during preadipocyte differentiation after transfecting with si-leptin and NC; (E) CEBPα expression levels during preadipocyte differentiation after transfecting with si-leptin and NC; (F) FABP4 expression levels during preadipocyte differentiation after transfecting with si-leptin and NC; (G, H) PPARγ protein levels during preadipocyte differentiation after transfecting with si-leptin and NC. (I) Leptin. expression levels during preadipocyte differentiation after transfecting with si-leptin and NC (“*”, $P \leq 0.05$; “**”, $P \leq 0.01$)