Lateral Advection Supports Nitrogen Export in the Oligotrophic Open-Ocean Gulf of Mexico

Thomas Kelly (✉ tbkelly@alaska.edu)

Florida State University <https://orcid.org/0000-0001-6285-1925>

Angela Knapp

Florida State University <https://orcid.org/0000-0001-5575-3382>

Michael Landry

University of California, San Diego

Karen Selph

University of Hawaii at Manoa

Taylor Shropshire

Florida State University

Rachel Thomas

Florida State University <https://orcid.org/0000-0002-1761-686X>

Michael Stukel

Florida State University

Article

Keywords: surface waters, nutrients, chlorophyll concentrations, regional scale dynamics, biogeochemical balances, Gulf of Mexico

Posted Date: December 14th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-116308/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Nature Communications on June 3rd, 2021.
See the published version at <https://doi.org/10.1038/s41467-021-23678-9>.

Lateral Advection Supports Nitrogen Export in the Oligotrophic Open-Ocean Gulf of Mexico

Thomas B. Kelly^{1,2,*}, Angela N. Knapp¹, Michael R. Landry³, Karen E. Selph⁴, Taylor A. Shropshire^{1,2}, Rachel Thomas¹, and Michael R. Stukel^{1,2}

¹ Department of Earth, Ocean and Atmospheric Sciences, Florida State University, Tallahassee, FL, USA

² Center for Ocean-Atmospheric Prediction Studies, Florida State University, Tallahassee, FL, USA

³ Integrative Oceanography Division, Scripps Institution of Oceanography, La Jolla, CA, USA

⁴ Department of Oceanography, University of Hawaii at Manoa, Honolulu, HI, USA

* Correspondence to Thomas B Kelly (tbk14@fsu.edu)

Abstract

In contrast to its productive coastal margins, the open-ocean Gulf of Mexico (GoM) is notable for highly stratified surface waters with extremely low nutrient and chlorophyll concentrations. Field campaigns in 2017 and 2018 identified low rates of turbulent mixing, which combined with oligotrophic nutrient conditions, give very low estimates for diffusive flux of nitrate into the euphotic zone ($< 1 \mu\text{mol N m}^{-2} \text{ d}^{-1}$). Estimates of local N_2 -fixation are similarly low. In comparison, measured export rates of sinking particulate organic nitrogen (PON) from the euphotic zone are 2 – 3 orders of magnitude higher (i.e. $462 - 1144 \mu\text{mol N m}^{-2} \text{ d}^{-1}$). We reconcile these disparate findings with regional scale dynamics inferred independently from remote-sensing products and a regional biogeochemical model and find that laterally-sourced organic matter is sufficient to support $>90\%$ of open-ocean nitrogen export in the GoM. Results show that lateral transport needs to be closely considered in studies of biogeochemical balances, particularly for basins enclosed by productive coasts.

Main Text

Conventionally, primary production is partitioned between new and regenerated production based on the source of inorganic nitrogen¹: new production (NP) is fueled by N (nitrogen) input from external sources (e.g. upwelled nitrate and N₂-fixation) while regenerated production (RP) comes from the internal recycling of nitrogen (e.g. ammonium). While RP often supports the vast majority of total primary production in open-ocean ecosystems, new nitrogen sources (NP) are necessary to support nitrogen and carbon export into the deep sea because the mass fluxes into and out of a system must balance when integrated over sufficiently large spatiotemporal scales. While the NP and RP dichotomy is well established in the simple biogeochemical metric of the *f*-ratio ($=NP/(NP+RP)$), direct measurement of NP is complicated by the intrinsic complexities of nitrogen cycling. For example, the most common NP measurement technique, nitrate uptake incubations, assumes negligible rates of nitrification (i.e. oxidation of ammonium into nitrate) within the euphotic zone. Nitrification, which regenerates nitrate from ammonium, may at times be a dominant source of nitrate in surface waters²⁻⁵, especially in oligotrophic waters where ammonium concentrations can exceed those of nitrate. Additionally, strong gradients in organic matter concentrations between coastal and offshore environments provide significant lateral transport potential. Through lateral transport, nutrients and organic matter can be carried into the offshore region, although this flux is seldom included in NP calculations. Modeling studies have suggested that net offshore fluxes can be an important addition to regional N balances, supporting 24 – 80 % of total export in subtropical gyres^{6,7}. However, lateral transport is difficult to constrain because spatiotemporal scales of advection are usually poorly matched to those of direct field measurements.

While numerous biogeochemical budgets have been reported for the shallow shelves of the GoM⁸⁻¹¹, few studies have focused on the oceanic region that covers $\sim 2/3^{\text{rds}}$ of the GoM surface area. Data presented here were collected during two field campaigns of *Bluefin Larvae in Oligotrophic Ocean Foodwebs: Investigating Nutrients to Zooplankton in the GoM* (BLOOFINZ-GoM) program targeting the central GoM spawning grounds of the endangered Atlantic Bluefin tuna^{12,13}. Major euphotic zone C and N cycling processes were quantified during five Lagrangian studies¹⁴⁻¹⁷, allowing us to compile comprehensive biogeochemical budgets. Here, we combine these *in situ* measurements with independent remote-sensing observations and

a biogeochemical model to test alternate hypotheses that (1) locally upwelled nitrate, (2) N₂-fixation, or (3) lateral transport support N export from oligotrophic oceanic waters of the GoM.

Results and Discussion

Field Observations and Dynamics

Nitrate+nitrite (hereafter “nitrate”) and ammonium were generally <100 nM throughout the euphotic zone, with nitrate concentrations between the detection limit (~10 nM) and 50 nM in the upper euphotic zone (UEZ; 0 – 60 m) (Figure 1b). Nitrate concentrations remained depressed well below the lower euphotic zone (LEZ), with 2.5 – 12 μ M concentrations at 150 m, consistent with prior characterizations of these water masses (Ref. ¹⁸ and references therein). Ammonium averaged 60 nM, with no depth trend in the euphotic zone (Supplemental Figure 1a). UEZ chlorophyll-*a* was consistently below 0.2 mg m⁻³ (Figure 1c), and UEZ integrated nitrogen pools did not vary significantly throughout the Lagrangian experiments (Supplemental Figure 1c-e). Most (62 – 79 %) net primary production (NPP) occurred in the UEZ, with a surface maximum of 270 – 550 μ mol C m⁻³ d⁻¹. In the LEZ, including the deep chlorophyll maximum (DCM), NPP declined to ~100 μ mol C m⁻³ d⁻¹ (Figure 1d). Nitrate uptake rates were not strongly depth-dependent, with maximum rates typically observed between 20 and 60 m and vertically integrated rates of 0.44 – 1.40 mmol N m⁻² d⁻¹ in UEZ and 0.26 – 2.05 mmol N m⁻² d⁻¹ in LEZ. Vertically integrated *f*-ratios were 0.04 – 0.14 (mean: 0.07) in the UEZ and 0.03 – 0.44 (mean: 0.14) for the LEZ. Although nitrification was not directly measured, ammonium-specific rates from other oligotrophic studies (0.02 – 0.5 d⁻¹; Ref. ³ and references within) suggest potential nitrate sources of 1 and 30 nmol N L⁻¹ d⁻¹, which could supply the majority of nitrate utilized by the phytoplankton community.

Gravitational export of organic matter from the UEZ¹⁴ (at 60 m) ranged from 590 – 1530 μ mol N m⁻² d⁻¹ and generally decreased with depth (460 – 1140 μ mol N m⁻² d⁻¹ at the base of the euphotic zone; 190 – 400 μ mol N m⁻² d⁻¹ at 200 m; Figure 1e, 2a). UEZ export was 164% (range: 90 – 330%) that of LEZ, indicating significant consumption of sinking particles within the LEZ. Active export of euphotic zone N due to ammonium excretion of migratory zooplankton at depth ranged from 20 – 172 μ mol N m⁻² d⁻¹ (mean: 71 μ mol N m⁻² d⁻¹; Figure 2b). This underestimates *in situ* active transport since organic excretion and zooplankton mortality at depth are not considered¹⁹.

The abundance of *Trichodesmium*²⁰, a dominant diazotroph in the GoM, ranged from 0 – 19 trichomes L⁻¹, which based on a photo-fixation model²¹, implies maximum N₂-fixation rates of between 0 and 0.38 nmol N L⁻¹ d⁻¹ and vertically integrated fixation rates of 0.4 – 2.8 μmol N m⁻² d⁻¹ in the UEZ (Figure 2d). While lower than in some other regions^{22,23}, such rates are consistent with Northern GoM findings^{15,24} and the low cell densities observed. Furthermore, the isotopic signature of sinking particulate N (i.e. δ¹⁵N) was nearly identical to that of subsurface nitrate¹⁵, leaving little room in the δ¹⁵N budget for N₂-fixation (see Supplemental Discussion).

Thorpe-scale analysis²⁵ of vertical eddy diffusivity in the LEZ ranged from 10⁻⁶ to 10⁻⁴ m² s⁻¹, which, combined with the small observed gradients in NO₃⁻ concentration, yields LEZ flux estimates of 0.01 – 1 μmol N m⁻² d⁻¹ (Figure 2c; Supplemental Figure 2). UEZ nitrate gradients were ~0, constraining vertical nitrate fluxes to << 0.01 μmol N m⁻² d⁻¹ (Supplemental Figure 2). Although vertical diffusivities can vary significantly in space and time, the stratification strength and vertical diffusivities for our field study are consistent with independent estimates from the NEMURO-GOM biogeochemical model and with previously reported values²⁶.

As vertical mixing of nitrate and N₂-fixation individually provide enough N to support <1% of sinking PON flux in our Lagrangian experiments, locally sourced new N appears insufficient to balance the measured rates of PON export. Instead, a significant role of lateral transport of organic material is suggested by the rapid horizontal displacements in our drifter experiments (up to 60 km d⁻¹) and the strong coastal-offshore particulate gradients in the GoM (Supplemental Figure 3-4). We tested this hypothesis with independent approaches of satellite remote-sensing products and a biogeochemical model tuned to the open-ocean GoM.

Lateral Supply of Nitrogen

MODIS Aqua²⁷ and Terra²⁸ remote-sensing products provide spatial POM data to infer lateral gradients in the GoM. Combined with a remote-sensing circulation product, OSCAR²⁹, we calculate lateral fluxes of particulate organic nitrogen (PON) in the UEZ. Independently, results from a 3D, nitrogen-currency biogeochemical model (NEMURO-GOM³⁰) are used to resolve lower trophic level dynamics in the open-ocean GoM.

The interannual median UEZ lateral transport of PON (i.e. organisms and detritus) into the central GoM zone (Figure 1a, 3) is estimated to be equivalent to a flux of 1150 μmol N m⁻²

d⁻¹ (IQR: 610 – 1530 $\mu\text{mol N m}^{-2} \text{ d}^{-1}$) and 1165 $\mu\text{mol N m}^{-2} \text{ d}^{-1}$ (IQR: 700 – 1725 $\mu\text{mol N m}^{-2} \text{ d}^{-1}$) by remote-sensing and NEMURO-GOM, respectively. Both estimates (Figure 3) indicate that lateral transport of PON alone is sufficient to support sediment trap-derived export out of the UEZ (median: 980 $\mu\text{mol N m}^{-2} \text{ d}^{-1}$; range: 587 - 1526 $\mu\text{mol N m}^{-2} \text{ d}^{-1}$) and the LEZ (median: 520 $\mu\text{mol N m}^{-2} \text{ d}^{-1}$; range: 462 – 1144 $\mu\text{mol N m}^{-2} \text{ d}^{-1}$). DON contributed an additional -8 – 46% to total UEZ lateral transport within NEMURO-GOM in any given day (Figure 3). Taken together, these results clearly show that net lateral transport of organic matter is of sufficient magnitude to support open-ocean export in the GoM.

Circulation patterns in NEMURO-GOM suggest that the open-ocean GoM is predominately a downwelling region and thus an organic nitrogen sink. In the absence of significant vertical input of subsurface nitrate into the LEZ or recently fixed nitrogen into the UEZ, particulate N export can only be supported by either lateral transport or a non-steady state drawdown of bioavailable nitrogen. Since the latter was not observed during any of our five, 4-day Lagrangian experiments (Supplemental Figure 1c-e), lateral input is likely to be responsible for ~100 % of export out of the UEZ and >90 % of export out of the LEZ (Figure 4). While episodic transport events such as storms have been shown to be relevant in other regions^{31,32}, the impact of such transient processes on the N balance of open-ocean GoM is likely negligible in late spring and summer due to the very deep nitracline (80 – 125 m). Such second-order processes are likely more important near the shelves where nitracline and mixed layer depths are closer than in the experimental sites themselves. What is clear in the open-ocean GoM is the prevalence of mesoscale eddies formed through instabilities in the Loop Current³³ (Supplemental Figures 3 & 4). Previous studies have found that, through both enhanced surface velocities and long persistence, such mesoscale features can transport large volumes of water onto and off of the shelf^{34–36}.

Global Connections and Implications

Since the oligotrophic GoM has previously been compared to the mid-ocean gyres due to similar biogeochemical properties (e.g. DCM, oligotrophic, low biomass)^{20,30,37}, it may be reasonable to consider the applicability of our results to these oceanic regions. The mid-ocean gyres are both larger than the GoM and may have less average kinetic energy (leading to circulation time-scales of months-years, rather than weeks). Consequently, lateral PON fluxes

are likely weaker in gyres but may be compensated, in part, by greater DON flux contributions⁷. Indeed DON inputs from lateral advection, entrainment, and mixing have been shown to supply gyre interiors^{38,39}, and based on model results may support 40 – 70% of export production in the North Atlantic gyre⁴⁰. Furthermore, export efficiency (i.e. export/NPP) is lower in the subtropical gyres (~5%) than measured in our study (11 - 25%) indicating a reduced lateral input requirement to support gyre export. Taken together, these analyses suggest that lateral transport of organic matter could potentially be a dominant source of external nitrogen in the oligotrophic gyres, as it is in the GoM.

Our results highlight the importance of inter-disciplinary and integrative process studies for constraining biogeochemical fluxes. Among still unresolved issues, it is not clear whether the laterally transported organic matter was originally supported by nitrate from the shelf, upwelling at the shelf break, upwelling associated with mesoscale features, or from some combination of these. Furthermore, extrapolating our results to other time periods and conditions (e.g., winter) in the GoM should be done with caution and appropriate field measurements. Nevertheless, our results clearly highlight the importance of accounting for contributions of lateral transport to carbon and nitrogen budgets in oligotrophic ocean regions and the role that horizontal mesoscale currents have in supporting vulnerable open-ocean GoM ecosystems.

Methods

Sampling Plan and Environmental Sampling

Data were collected during two cruises in the Northern Gulf of Mexico conducted in May 2017 and April/May 2018. Each cruise began with a bongo survey of water parcels that were identified by remote sensing as having favorable SST and vorticity¹² for larval Atlantic Bluefin Tuna (ABT). At selected locations, a sediment trap array was deployed and a Lagrangian experiment was initiated. Over the following 3 – 4 days, the water parcel was repeatedly sampled for biogeochemical properties and incubations were conducted to measure process rates. Throughout an experiment, profiles of temperature and salinity were taken 3 – 4 times per day through the upper 200 m. Additionally, PAR profiles were conducted at least once per Lagrangian experiment around local noon. Ambient nutrient concentrations were determined from samples collected daily from 6 depths throughout the euphotic zone (same depths as incubations were conducted). The concentrations of nitrate+nitrite were measured using a chemiluminescent

method, while concentrations of NH_4^+ were measured using the fluorescent OPA method. See Knapp et al. (Ref¹⁵) for details.

Trichodesmium Abundance and N_2 Fixation— *Trichodesmium* abundance and chlorophyll contents were assessed as reported in Selph et al. (Ref²⁰). Briefly, 6.6 L samples from daily noon casts at 6 depths (2 - 50 m) were collected and filtered through inline 8- μm , 47-mm filters, preserved (2% paraformaldehyde), and frozen (-80°C). *Trichodesmium* were counted with microscopy (10-30X magnification) under 440-460 nm excitation to detect orange-fluorescing trichomes. For chlorophyll *a* contents, separate samples (6.6 L) were collected onto 20 μm , 47-mm filters and frozen (-80°C). Individual trichomes were picked from thawed filters, extracted (90% acetone), and chlorophyll fluorescence measured with 10-AU (Turner Designs) fluorometer⁴¹. Chlorophyll-specific N_2 -fixation rates were calculated based on the photo-fixation model of Breitbarth et al. (Ref.²¹) based on culture work conducted at temperatures (i.e. 26°C) and photoperiods (i.e. 12 h) consistent with *in situ* conditions observed.

Productivity and Nutrient Dynamics

Rates of net primary production (NPP) and nitrate uptake were measured at 6 depths spanning the euphotic zone each day of a Lagrangian experiment. Incubation bottles (2.7 L) were filled by Niskin rosette and then $\text{H}^{13}\text{CO}_3^-$ (final concentration: 154 or 196 $\mu\text{mol L}^{-1}$ on NF1704 and NF1802, respectively) and/or $^{15}\text{NO}_3^-$ (final concentration: 10 or 8 nmol L^{-1} on NF1704 and NF1802, respectively) was added. Bottles were then incubated in mesh bags attached to an *in situ* drift array⁴² for 24 h—drift array was deployed and recovered pre-dawn, typically ~0500 local time. Upon recovery, incubation bottles were immediately filtered onto precombusted GF/F. Filtered samples were kept at -80°C until analysis. In addition to *in situ* incubations, a series of deck-board experiments were conducted for 4 - 6 hours to measure vertical patterns and diel variability in nitrate (same NO_3^- spike as above) and ammonium uptake¹⁷ (final concentration: 9.8 or 4.6 nmol L^{-1} on NF1704 and NF1802, respectively). Samples were analyzed at the UC Davis stable isotope facility. Nutrient uptake rates and associated uncertainty were calculated as in Stukel (Ref.⁴³).

Thorpe Scale Analysis—Profiles of vertical diffusivity (K_z) were calculated by Thorpe scale analysis²⁵ of CTD-derived density profiles. Thorpe scale analysis estimates turbulent mixing parameters (i.e. vertical eddy diffusivity, K_z) based on the number and size of observed density

inversions. Raw Seabird data files were processed based on the recommendations of Ref. ²⁵. Resulting profiles of K_z were averaged across all casts during a Lagrangian experiment using the geometric mean ($n = 14 - 26$, with decreasing number of casts below the euphotic zone). Nitrate fluxes (F_z) were then calculated using the observed vertical nutrient (NO_3^-) gradients and vertical diffusivity (Eq. 1).

$$F_z = -K_z \left(\frac{d[\text{NO}_3^-]}{dz} \right) \quad \text{Eq. 1}$$

Zooplankton Abundance, Biomass, and Migrant Excretion

Zooplankton were collected daily from paired day-night tows with a 1-m diameter ring net (200- μm mesh) towed obliquely through the euphotic zone¹⁶. The tow contents were filtered through stacked Nitex screens of 5, 2, 1, 0.5 and 0.2-mm Nitex mesh to produce 5 size classes, and carbon, nitrogen, and dry mass for each size class were determined. Active nitrogen transport by daytime excretion of diel migrating mesozooplankton at mesopelagic depths was calculated from the equations of Ikeda (Ref. ⁴⁴), the biomass differences between paired day and night tows (size-fractionated migrant carbon), the average C values of individual animals in each migrant size fraction⁴⁵ and the mean temperature in the 300-500 m depth range ($\sim 10\text{-}12^\circ\text{C}$).

Export Production by Sediment Trap

Each sediment trap deployment consisted of three cross-members with 12 Particle Interceptor Tubes⁴⁶ per depth. Trap depths were determined from CTD fluorescence profiles with one cross member below the mixed layer and within the euphotic zone (60 m), one at the base of the euphotic zone (110 – 150 m), and one at 230 m. The sediment trap array included a 3-m x 1-m holey sock drogue at 15 m depth⁴⁷. Sediment trap tubes were filled with a poisoned brine solution. After recovery, the overlying seawater was siphoned off to within 3 cm of the brine interface. The samples were then filtered through a 100 μm filter which was manually inspected under a stereo-microscope (25X) to remove zooplankton “swimmers”. After washing the >100 μm non-swimmer fraction back in to the sample, whole tube contents were filtered onto a pre-combusted GF/F for C and N isotopic analyses.

Analysis

Nitrogen Isotopic Signatures of New and Export Production— Mass balance requires that the nitrogen $\delta^{15}\text{N}$ isotopic signature of export equals the isotopic signature of the source nitrogen. N_2 -fixation supplies nitrogen with a $\delta^{15}\text{N}$ of approximately $-2 - 0\text{‰}$ ^{48,49} and vertical mixing of subsurface nitrate would maintain the subsurface isotopic signature: $2.0 - 3.8\text{‰}$ ¹⁵. Exported nitrogen (sinking + active transport) was observed to have a $\delta^{15}\text{N}$ of $\sim 2.9 - 5\text{‰}$.

Remote-Sensing Estimates of Transport—Eight-day composites of surface POC concentrations estimated from Moderate Resolution Imaging Spectroradiometer (MODIS) Aqua & MODIS Terra were retrieved from the National Aeronautical and Space Administration's (NASA) data repository^{27,28} for each May from 2000 to 2019. We assumed that particles had Redfield stoichiometry (106:16, C:N, mol:mol). Remote sensing fields were binned to 8-km x 8-km resolution prior to analysis to reduce noise. Regional circulation was prescribed from OSCAR (Ref²⁹), a data-assimilative, remote-sensing data product with $1/3^{\text{rd}}$ degree spatial and 5-day temporal resolution. OSCAR circulation fields were mapped onto the remote-sensing grid using bilinear spatial interpolation and linear temporal interpolation. Fluxes between each grid cell were then calculated for each remote-sensing field ($n = 80$). The total net flux into the control volume was divided by the area of the control volume for comparison to vertical N flux measurements. Results are shown for the control volume as defined in Figure 1a.

NEMURO-GOM Estimates of Transport— The biogeochemical model NEMURO-GoM was used to provide a consistent three-dimensional perspective on nutrient uptake dynamics in the study region. The model consists of 29 z-layers with a vertical resolution of 10 m in the upper 150 m and ~ 4 -km horizontal resolution. NEMURO-GOM is run offline within the MIT general circulation model and forced by daily averaged flow fields obtained from a data assimilative HYbrid Coordinate Ocean Model GoM simulation. The model has been extensively validated against nutrient, biomass, and rate measurements and fully resolves a simple nitrogen cycle with 11 state variables consisting of two phytoplankton and three zooplankton functional groups³⁰. Vertical mixing in the model is parameterized based on a nonlocal K-profile parameterization (KPP) mixing scheme from a bulk Richardson number approach that quantifies the importance of stratification and destabilizing shear⁵⁰. Daily model output for May of 1993 to 2012 were analyzed ($n = 571$) with flux integrations performed for the same control volumes as used for the remote sensing estimations (Figure 1a). Mesoscale eddies were identified in daily averaged model flow fields using the algorithm of Laxenaire et al. (Ref. ⁵¹), which utilizes surface

274 velocities along closed contours of SSH to define eddy boundaries. Lateral eddy flux (see
275 Supplemental Discussion) was determined by calculating the net flux into the control column for
276 all model grid cells that were identified as located within an eddy (i.e. enclosed SSH contour
277 with the greatest velocity).
278

Acknowledgements

We thank the captain and crew of the NOAA Ship *Nancy Foster* for providing their assistance in conducting the fieldwork that led to this manuscript. We also thank our collaborators and colleagues at NOAA Southeast Fisheries/RSMAS/University of Miami for helping plan and carry out two successful field campaigns. We would like to thank Remi Laxenaire for his assistance in the identification of eddies. This study acknowledges BLOOFINZ Program support from National Oceanic and Atmospheric Administration awards NA15OAR4320071, NA16NMF4320058, and U.S. National Science Foundation awards OCE-1851347 and OCE-1851558. We also acknowledge support from the National High Magnetic Field Laboratory NSF/DMR-1644779 and the State of Florida.

Author Contributions

TBK and MRS planned and synthesized the data presented here. TBK, MRL, KES, TAS, RT and MRS collected the field data. ANK provided nutrient data for the analysis. TAS designed the hydrodynamic model and provided output data. TBK wrote the manuscript. All authors took part in editing and submission of the final manuscript.

References

1. Harrison, W. G., Platt, T. & Lewis, M. R. f -Ratio and its relationship to ambient nitrate concentration in coastal waters. *J. Plankton Res.* **9**, 235–248 (1987).
2. Wankel, S. D., Kendall, C., Pennington, J. T., Chavez, F. P. & Paytan, A. Nitrification in the euphotic zone as evidenced by nitrate dual isotopic composition: Observations from Monterey Bay, California. *Glob. Biogeochem. Cycles* **21**, (2007).
3. Yool, A., Martin, A. P., Fernández, C. & Clark, D. R. The significance of nitrification for oceanic new production. *Nature* **447**, 999–1002 (2007).
4. Martens-Habbena, W., Berube, P. M., Urakawa, H., de la Torre, J. R. & Stahl, D. A. Ammonia oxidation kinetics determine niche separation of nitrifying Archaea and Bacteria. *Nature* **461**, 976–979 (2009).
5. Dore, J. E. & Karl, D. M. Nitrification in the euphotic zone as a source for nitrite, nitrate, and nitrous oxide at Station ALOHA. *Limnol. Oceanogr.* **41**, 1619–1628 (1996).
6. Williams, R. G. & Follows, M. J. The Ekman transfer of nutrients and maintenance of new production over the North Atlantic. *Deep Sea Res. Part Oceanogr. Res. Pap.* **45**, 461–489 (1998).
7. Letscher, R. T., Primeau, F. & Moore, J. K. Nutrient budgets in the subtropical ocean gyres dominated by lateral transport. *Nat. Geosci.* **9**, 815–819 (2016).
8. Xue, Z. *et al.* Modeling ocean circulation and biogeochemical variability in the Gulf of Mexico. *Biogeosciences* **10**, 7219–7234 (2013).
9. Fennel, K. & Laurent, A. N and P as ultimate and proximate limiting nutrients in the northern Gulf of Mexico: implications for hypoxia reduction strategies. *Biogeosciences* **15**, 3121–3131 (2018).

- 320 10. Laurent, A., Fennel, K., Ko, D. S. & Lehrter, J. Climate Change Projected to Exacerbate
321 Impacts of Coastal Eutrophication in the Northern Gulf of Mexico. *J. Geophys. Res. Oceans*
322 **123**, 3408–3426 (2018).
- 323 11. Fichot, C. G., Lohrenz, S. E. & Benner, R. Pulsed, cross-shelf export of terrigenous
324 dissolved organic carbon to the Gulf of Mexico. *J. Geophys. Res. Oceans* **119**, 1176–1194
325 (2014).
- 326 12. Domingues, R. *et al.* Variability of preferred environmental conditions for Atlantic
327 bluefin tuna (*Thunnus thynnus*) larvae in the Gulf of Mexico during 1993-2011. *Fish.*
328 *Oceanogr.* **25**, 320–336 (2016).
- 329 13. Landry, M. R., Beckley, L. E. & Muhling, B. A. Climate sensitivities and uncertainties in
330 food-web pathways supporting larval bluefin tuna in subtropical oligotrophic oceans. *ICES J.*
331 *Mar. Sci.* **76**, 359–369 (2019).
- 332 14. Stukel, M. R., Kelly, T. B., Landry, M. R., Selph, K. E. & Swalethorp, R. Sinking carbon,
333 nitrogen, and pigment flux within and beneath the euphotic zone in the oligotrophic, open-
334 ocean Gulf of Mexico. *J. Plankton Res.* (submitted).
- 335 15. Knapp, A. N. *et al.* Constraining the sources of nitrogen fueling phytoplankton and food
336 webs in the Gulf of Mexico using nitrogen isotope budgets. *J. Plankton Res.* (submitted).
- 337 16. Landry, M. R. *et al.* Microbial Food Web Dynamics in the Oceanic Gulf of Mexico. *J.*
338 *Plankton Res.* (submitted).
- 339 17. Yingling, N. *et al.* Taxon-Specific Phytoplankton Growth, Nutrient Limitation, and Light
340 Limitation in The Oligotrophic Gulf of Mexico. *J. Plankton Res.* (submitted).

- 341 18. Howe, S., Miranda, C., Hayes, C. T., Letscher, R. T. & Knapp, A. N. The Dual Isotopic
342 Composition of Nitrate in the Gulf of Mexico and Florida Straits. *J. Geophys. Res. Oceans*
343 **125**, (2020).
- 344 19. Stukel, M. R. *et al.* Plankton food webs of the Gulf of Mexico spawning grounds of
345 Atlantic Bluefin tuna. *J. Plankton Res.* (submitted).
- 346 20. Selph, K. E. *et al.* Phytoplankton assemblages in the open ocean water of the Gulf of
347 Mexico during May 2017 and 2018. *J. Plankton Res.* (submitted).
- 348 21. Breitbarth, E., Wohlers, J., Kläs, J., LaRoche, J. & Peeken, I. Nitrogen fixation and
349 growth rates of *Trichodesmium* IMS-101 as a function of light intensity. *Mar. Ecol. Prog. Ser.*
350 **359**, 25–36 (2008).
- 351 22. Mulholland, M. R. *et al.* High Rates of N₂ Fixation in Temperate, Western North
352 Atlantic Coastal Waters Expand the Realm of Marine Diazotrophy. *Glob. Biogeochem. Cycles*
353 **33**, 826–840 (2019).
- 354 23. Tang, W. *et al.* Revisiting the distribution of oceanic N₂ fixation and estimating
355 diazotrophic contribution to marine production. *Nat. Commun.* **10**, 831 (2019).
- 356 24. Holl, C. M. *et al.* *Trichodesmium* in the western Gulf of Mexico: ¹⁵N₂-fixation and
357 natural abundance stable isotopic evidence. *Limnol. Oceanogr.* **52**, 2249–2259 (2007).
- 358 25. Gargett, A. & Garner, T. Determining Thorpe Scales from Ship-Lowered CTD Density
359 Profiles. *J. Atmospheric Ocean. Technol.* **25**, 1657–1670 (2008).
- 360 26. Whitt, D. B., Lévy, M. & Taylor, J. R. Submesoscales Enhance Storm-Driven Vertical
361 Mixing of Nutrients: Insights From a Biogeochemical Large Eddy Simulation. *J. Geophys.*
362 *Res. Oceans* **124**, 8140–8165 (2019).

- 363 27. NASA Ocean Biology Processing Group. MODIS-Aqua Level 3 Mapped Particulate
364 Organic Carbon Data Version R2018.0. (2017) doi:10.5067/AQUA/MODIS/L3M/POC/2018.
- 365 28. NASA Ocean Biology Processing Group. MODIS-TERRA Level 3 Mapped Particulate
366 Organic Carbon Data Version R2018.0. (2018)
367 doi:10.5067/TERRA/MODIS/L3M/POC/2018.
- 368 29. ESR. OSCAR third degree resolution ocean surface currents. Ver. 1. (2009).
- 369 30. Shropshire, T. A. *et al.* Quantifying spatiotemporal variability in zooplankton dynamics
370 in the Gulf of Mexico with a physical–biogeochemical model. *Biogeosciences* **17**, 3385–3407
371 (2020).
- 372 31. Klein, P. & Coste, B. Effects of wind-stress variability on nutrient transport into the
373 mixed layer. *Deep Sea Res. Part Oceanogr. Res. Pap.* **31**, 21–37 (1984).
- 374 32. Rumyantseva, A. *et al.* Ocean nutrient pathways associated with the passage of a storm.
375 *Glob. Biogeochem. Cycles* **29**, 1179–1189 (2015).
- 376 33. Oey, L.-Y., Ezer, T. & Lee, H.-C. Loop Current, Rings and Related Circulation in the
377 Gulf of Mexico: A Review of Numerical Models and Future Challenges. in *Geophysical*
378 *Monograph Series* (eds. Sturges, W. & Lugo-Fernandez, A.) 31–56 (2013).
379 doi:10.1029/161GM04.
- 380 34. Sahl, L. E., Wiesenburg, D. A. & Merrell, W. J. Interactions of mesoscale features with
381 Texas shelf and slope waters. *Cont. Shelf Res.* **17**, 117–136 (1997).
- 382 35. Toner, M. Chlorophyll dispersal by eddy-eddy interactions in the Gulf of Mexico. *J.*
383 *Geophys. Res.* **108**, 3105 (2003).

- 384 36. Barkan, R. *et al.* Submesoscale Dynamics in the Northern Gulf of Mexico. Part II:
385 Temperature–Salinity Relations and Cross-Shelf Transport Processes. *J. Phys. Oceanogr.* **47**,
386 2347–2360 (2017).
- 387 37. Gomez, F. A. *et al.* Seasonal patterns in phytoplankton biomass across the northern and
388 deep Gulf of Mexico: a numerical model study. *Biogeosciences* **15**, 3561–3576 (2018).
- 389 38. Roussenov, V., Williams, R. G., Mahaffey, C. & Wolff, G. A. Does the transport of
390 dissolved organic nutrients affect export production in the Atlantic Ocean? *Glob. Biogeochem.*
391 *Cycles* **20**, (2006).
- 392 39. Oschlies, A. Nutrient supply to the surface waters of the North Atlantic: A model study.
393 *J. Geophys. Res.* **107**, 3046 (2002).
- 394 40. Torres-Valdés, S. *et al.* Distribution of dissolved organic nutrients and their effect on
395 export production over the Atlantic Ocean. *Glob. Biogeochem. Cycles* **23**, (2009).
- 396 41. Strickland, J. D. H. & Parsons, T. R. *A Practical handbook of Seawater Analysis*.
397 (Fisheries Research Board of Canada, 1972).
- 398 42. Landry, M. R., Ohman, M. D., Goericke, R., Stukel, M. R. & Tsyrklevich, K. Lagrangian
399 studies of phytoplankton growth and grazing relationships in a coastal upwelling ecosystem
400 off Southern California. *Prog. Oceanogr.* **83**, 208–216 (2009).
- 401 43. Stukel, M. R. *Investigating equations for measuring dissolved inorganic nutrient uptake*
402 *in oligotrophic conditions*. <http://biorxiv.org/lookup/doi/10.1101/2020.08.30.274449> (2020)
403 doi:10.1101/2020.08.30.274449.
- 404 44. Ikeda, T. Metabolic rates of epipelagic marine zooplankton as a function of body mass
405 and temperature. *11* (1985).

45. Landry, M. R., Al-Mutairi, H., Selph, K. E., Christensen, S. & Nunnery, S. Seasonal patterns of mesozooplankton abundance and biomass at Station ALOHA. *Deep Sea Res. Part II Top. Stud. Oceanogr.* **48**, 2037–2061 (2001).
46. Knauer, G. A., Martin, J. H. & Bruland, K. W. Fluxes of particulate carbon, nitrogen, and phosphorus in the upper water column of the northeast Pacific. *Deep Sea Res. Part Oceanogr. Res. Pap.* **26**, 97–108 (1979).
47. Stukel, M., Ohman, M., Benitez-Nelson, C. & Landry, M. Contributions of mesozooplankton to vertical carbon export in a coastal upwelling system. *Mar. Ecol. Prog. Ser.* **491**, 47–65 (2013).
48. Carpenter, E. J., Harvey, H. R., Fry, B. & Capone, D. G. Biogeochemical tracers of the marine cyanobacterium *Trichodesmium*. *Deep Sea Res. Part Oceanogr. Res. Pap.* **44**, 27–38 (1997).
49. Minagawa, M. & Wada, E. Nitrogen isotope ratios of red tide organisms in the East China Sea: A characterization of biological nitrogen fixation. *Mar. Chem.* **19**, 245–259 (1986).
50. Large, W. G., McWilliams, J. C. & Doney, S. C. Oceanic vertical mixing: A review and a model with a nonlocal boundary layer parameterization. *Rev. Geophys.* **32**, 363 (1994).
51. Laxenaire, R. *et al.* Anticyclonic Eddies Connecting the Western Boundaries of Indian and Atlantic Oceans. *J. Geophys. Res. Oceans* **123**, 7651–7677 (2018).

Figure Legends

Figure 1. Overview of *in situ* data. Study locations in the central GoM (colored drift trajectories) with a white outline indicating the integration volume from surface to 55 m considered as representative of the eastern Central GoM (depths >2000 m, outside the general flow of the Loop Current). (a) Vertical profiles of nitrate concentrations with inset detailing low surface concentrations. (b) Vertical profiles of chlorophyll concentrations as recorded by calibrated fluorometer. (c) Vertical profiles of NPP and (d) vertical profiles of particulate nitrogen export at three depths (UEZ, LEZ, 200 m). Colors correspond to Lagrangian experiments, and error bars indicate ± 1 SD of means for measurements during experiments.

Figure 2. Summary of observed euphotic zone nitrogen fluxes from the oligotrophic GoM. (a) Sediment trap fluxes include export at the base of the UEZ (open squares) and the LEZ (closed circles) (b) Zooplankton excretion below the euphotic zone. (c) Vertical mixing of nitrate into the LEZ. Mixing into UEZ not shown (~ 0). (d) N_2 -fixation rates modeled from observed *Trichodesmium* biomass (UEZ; circles) and from $\delta^{15}N$ -based budgets for the entire euphotic zone (squares) as reported in Knapp et al.¹⁵. Fluxes $< 0.1 \mu\text{mol N m}^{-2} \text{ d}^{-1}$ are shown on left as open symbols.

Figure 3. Comparison of lateral nitrogen supply to observed export production in the UEZ. Satellite lateral PON flux was calculated from 8-day composite MODIS Terra/Aqua imagery during the month of May (2002 – 2019) with flow fields prescribed by OSCAR circulation. NEMURO-GOM fluxes were calculated separately for PON, DON and DIN and summed for total lateral fluxes. NEMURO-GOM vertical fluxes are integrated for the UEZ and include upwelling and turbulent mixing but not export by sinking particles. Positive flux values indicate net input into the integration volume. Flux values are normalized to lateral area.

Figure 4. Schematic of transport mechanisms and fluxes of the NE GoM. (a) Arrows and values and fluxes observed or estimated from model relative to euphotic zone NPP including turbulent mixing (purple), N_2 -fixation (blue), lateral transport (green), gravitational sinking

457 (grey) and primary production (black). (b) Monthly surface snapshot of N inventories for May
458 1993 in NEMURO-GOM.
459
460

461 Figures

462
 463 **Figure 1. Overview of *in situ* data.** Study locations in the central GoM (colored drift
 464 trajectories) with a white outline indicating the integration volume from surface to 55 m
 465 considered as representative of the eastern Central GoM (depths >2000 m, outside the general
 466 flow of the Loop Current). (a) Vertical profiles of nitrate concentrations with inset detailing low
 467 surface concentrations. (b) Vertical profiles of chlorophyll concentrations as recorded by
 468 calibrated fluorometer. (c) Vertical profiles of NPP and (d) vertical profiles of particulate
 469 nitrogen export at three depths (UEZ, LEZ, 200 m). Colors correspond to Lagrangian
 470 experiments, and error bars indicate ± 1 SD of means for measurements during experiments.
 471

Figure 2. Summary of observed euphotic zone nitrogen fluxes from the oligotrophic GoM.

(a) Sediment trap fluxes include export at the base of the UEZ (open squares) and the LEZ (closed circles) (b) Zooplankton excretion below the euphotic zone. (c) Vertical mixing of nitrate into the LEZ. Mixing into UEZ not shown (~ 0). (d) N_2 -fixation rates modeled from observed *Trichodesmium* biomass (UEZ; circles) and from $\delta^{15}\text{N}$ -based budgets for the entire euphotic zone (squares) as reported in Knapp et al.¹⁵. Fluxes $< 0.1 \mu\text{mol N m}^{-2} \text{d}^{-1}$ are shown on left as open symbols.

Figure 3. Comparison of lateral nitrogen supply to observed export production in the UEZ.

Satellite lateral PON flux was calculated from 8-day composite MODIS Terra/Aqua imagery during the month of May (2002 – 2019) with flow fields prescribed by OSCAR circulation. NEMURO-GOM fluxes were calculated separately for PON, DON and DIN and summed for total lateral fluxes. NEMURO-GOM vertical fluxes are integrated for the UEZ and include upwelling and turbulent mixing but not export by sinking particles. Positive flux values indicate net input into the integration volume. Flux values are normalized to lateral area.

Figure 4. Schematic of transport mechanisms and fluxes of the NE GoM. (a) Arrows and values and fluxes observed or estimated from model relative to euphotic zone NPP including turbulent mixing (purple), N_2 -fixation (blue), lateral transport (green), gravitational sinking (grey) and primary production (black). (b) Monthly surface snapshot of N inventories for May 1993 in NEMURO-GOM.

Figures

Figure 1

Overview of in situ data. Study locations in the central GOM (colored drift trajectories) with a white outline indicating the integration volume from surface to 55 m considered as representative of the eastern Central GOM (depths >2000 m, outside the general flow of the Loop Current). (a) Vertical profiles of nitrate concentrations with inset detailing low surface concentrations. (b) Vertical profiles of chlorophyll concentrations as recorded by calibrated fluorometer. (c) Vertical profiles of NPP and (d) vertical profiles of particulate nitrogen export at three depths (UEZ, LEZ, 200 m). Colors correspond to Lagrangian experiments, and error bars indicate +1 SD of means for measurements during experiments.

Figure 2

Summary of observed euphotic zone nitrogen fluxes from the oligotrophic GOM. (a) Sediment trap fluxes include export at the base of the UEZ (open squares) and the LEZ (closed circles) (b) Zooplankton excretion below the euphotic zone. (c) Vertical mixing of nitrate into the LEZ. Mixing into UEZ not shown (-0). (d) N_2 -fixation rates modeled from observed *Trichodesmium* biomass (UEZ; circles) and from 815N -

based budgets for the entire euphotic zone (squares) as reported in Knapp et al.15. Fluxes < 0.1 $\mu\text{mol N m}^2 \text{ d}^{-1}$ are shown on left as open symbols.

Figure 3

Comparison of lateral nitrogen supply to observed export production in the UEZ. Satellite lateral PON flux was calculated from 8-day composite MODIS Terra/Aqua imagery during the month of May (2002 – 2019) with flow fields prescribed by OSCAR circulation. NEMURO-GOM fluxes were calculated separately

for PON, DON and DIN and summed for total lateral fluxes. NEMURO-GOM vertical fluxes are integrated for the UEZ and include upwelling and turbulent mixing but not export by sinking particles. Positive flux values indicate net input into the integration volume. Flux values are normalized to lateral area.

Figure 4

Schematic of transport mechanisms and fluxes of the NE GOM. (a) Arrows and values and fluxes observed or estimated from model relative to euphotic zone NPP including turbulent mixing (purple), N₂-fixation (blue), lateral transport (green), gravitational sinking (grey) and primary production (black). (b) Monthly surface snapshot of N inventories for May 1993 in NEMURO-GOM.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [KellyetalManuscriptSupplementalv4submitted.pdf](#)