Echocardiographic and Electrocardiographic Findings in COVID-19 Patients; A Cross-Sectional Study

Faeze Keihanian (✉ faezekeihanian@yahoo.com)

Mashhad University of Medical Sciences

Hoorak Poorzand

Mashhad University of Medical Sciences

Amin Saeidinia

Mashhad University of Medical Sciences

Ali Eshraghi

Mashhad University of Medical Sciences

Research Article

Keywords: COVID-19, Echocardiography, Electrocardiography, Right Ventricular Dysfunction

Posted Date: January 3rd, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1164425/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: There are still many gaps in our knowledge regarding the direct cardiovascular injuries due to COVID-19 infection. In this study, we tried to find out the effect of SARS-CoV-2 infection on cardiac function in patients without any history of structural heart disease by electrocardiographic and echocardiographic evaluations.

Methods: This was a cross-sectional study on patients with COVID-19 infection admitted to Imam Reza hospital, Mashhad, Iran between 14 April and 21 September 2020. COVID-19 infection was verified by a positive reverse-transcriptase polymerase chain reaction (PCR) assay for SARS-CoV-2 using nasopharyngeal/oropharyngeal samples. We enrolled all patients over 18 years old with definite diagnosis of COVID-19 infection. All patients underwent a comprehensive transthoracic echocardiography at the first week of admission. Clinical and imaging data were collected prospectively.

Results: In total, 142 patients were enrolled in this study. The mean age of participants was 60.69 ± 15.70 years (range: 30-90 years). Most patients were male (82, 57.7%). Multivariate analysis showed that O₂ saturation at admission was independently a predictor of re-hospitalization ($P < 0.001$). RV size ($P < 0.001$), dyslipidemia ($P < 0.001$), ejection fraction (EF) ($P < 0.001$), age ($P = 0.020$), systolic blood pressure ($P = 0.001$), O₂ saturation ($P = 0.018$) and diabetes ($P = 0.025$) independently predicted 30-days mortality.

Conclusion: Echocardiography can be used for risk assessment in patients with COVID-19, especially in those with previous history of diabetes and dyslipidemia. The infection could result in Ventricular dysfunction, even in those without previous history of structural heart disease.

1. Background

Knowledge about short- and long-term complications of COVID-19 infection is growing worldwide¹. COVID-19 is now a serious pandemic infection with different clinical features². It is shown that severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) can involve respiratory tract and patients with previous cardiovascular disorders are at a higher risk³. The underlying mechanisms are yet to be clear⁴. It has been suggested that cardiac injury in COVID-19 infection is possibly related to its clinical outcomes. It was reported that patients with cardiac injury are associated with higher rate of morbidity, more abnormal laboratory and radiological findings⁵. Pathological findings (mononuclear inflammatory infiltration in heart tissue) in patients with COVID-19 infection suggests a direct damage to cardiac tissues⁶. Moreover, it is indicated that previous cardiovascular disorders may also be more predisposing to cardiac injury due to COVID-19 infection⁵. Keeping in mind the cardiovascular presentations, underlying cardiovascular disorders and its adverse events can help the clinician to make better decisions in the management of COVID-19 infection².

Cardiac adverse events related to COVID-19 infection are prevalent (nearly 25%) and can increase the risk of mortality⁷. Most recent studies on COVID-19 cardiac complications have reported clinical or laboratory

findings with no imaging confirmation.

Transthoracic echocardiography is a reliable technique to detect ventricular function or valve disorders and risk assessment, as an additive guide to select better treatment strategies for hemodynamic stabilization⁸. However, due to a risk of viral contamination, echocardiographic evaluation in patients with COVID-19 infection has been limited^{9,10}. To improve our knowledge about the pathophysiology of direct cardiovascular injuries due to COVID-19 infection², we tried to find the effect of SARS-CoV-2 infection on cardiac function in patients without any history of structural heart disease by electrocardiographic and echocardiographic evaluations.

2. Methods

This was a cross-sectional study, performed on admitted patients with COVID-19 infection, in Imam Reza hospital, Mashhad, Iran between 14 April and 21 September 2020.

2.1.1. Eligibility

COVID-19 infection was verified by a positive reverse-transcriptase polymerase chain reaction (PCR) assay for SARS-CoV-2 using nasopharyngeal/oropharyngeal samples. We included all patients over 18 years old and a definite diagnosis of COVID-19 infection. We excluded patients with previous structural heart disease by history taking or evidences of old myocardial infarction in ECG. Figure1 shows the flowchart of study.

2.1.2. Evaluations

At admission, demographic data, comorbid conditions, medications, physical examination, lung CT scan and laboratory findings were all recorded. Cardiac injury was determined as troponin I (TPI) above the 99th percentile at presentation. Clinical and imaging data were collected prospectively and a 12-lead electrocardiography was performed. All patients were followed about three months for re-admission, cardiac events and in-hospital mortality or mortality.

2.1.3. Echocardiography

All patients underwent a comprehensive transthoracic echocardiography (TTE) within two weeks after symptoms presentation, in a standard manner with the same equipment (M7, Mindray Bio-Medical Electronics, Shenzhen, China) and by one cardiologist. All echocardiographic studies were performed bedside at COVID-19 intensive care unit (ICU) or non-ICU. Personal protection included airborne precautions, N-95 respirator masks, fluid-resistant gowns, and 2 sets of gloves, head covers, eye shields, and shoe covers. Most of measurements were performed offline to reduce the exposure time and contamination with two-to-three-seconds clips derived from standard echo views. ventricular size, systolic and diastolic function, pulmonary artery systolic pressure (PASP), presence of pericardial effusion, valvular stenosis or regurgitation were assessed. Abnormal TTE was defined as LV and RV enlargement,

RV dysfunction (by visual assessment), ejection fraction (EF) $\leq 54\%$, pericardial effusion, PASP ≥ 35 mmHg, moderate and severe diastolic dysfunction and valvular regurgitation or stenosis more than mild.

2.1.4. Exclusion criteria

In order to better assessment of direct effects of COVID-19 on the echocardiographic parameters, patients with documented history of previous structural heart diseases (like valvular stenosis, primary valvular regurgitation, previous MI and heart surgery, etc.) were excluded.

2.1.5. Ethics

The ethics committee of Mashhad University of Medical Sciences approved the study (Ethical No IR.MUMS.REC.1399.016). We obtained an informed consent from all patients or their legal guardians. The study was conducted in accordance with the declaration of Helsinki. Informed written consent was given prior to the inclusion of subjects or their families, in the study.

3. Results

In Total, 142 patients with diagnosis of Covid-19 infection, were enrolled who were admitted in Imam Reza Hospital, Mashhad, Iran. The mean age of participants was 60.69 ± 15.70 years (range: 30-90). Most patients were male (82, 57.7%). Known comorbidities were found in 45.1%, with hypertension (24.6%) and diabetes mellitus (23.9%) as the most common ones. The more frequent symptoms were dyspnea on admission (90.1%), cough (54.9%) and fever (49.3%). Baseline demographic characteristics, comorbidities, clinical and laboratory data were presented in Table 1.

LAD was the most common finding (15.5%) in ECG. Seventy-eight patients had sinus tachycardia (55%), one atrial fibrillation (0.7%) and two AVNRT (1.4%). Ten patients had right axis deviation (RAD) and others normal axis.

Bilateral peripheral ground glass opacification (GGO) was the most common specific CT scan manifestation (38%) followed by diffused bilateral GGO (35.2%), unilateral GGO (14.1%), and bilateral pleural effusion (9.9%). Other non-specific presentations were consolidation (63.4%), and lymphadenopathy (31%).

Abnormal TTE was seen in 71 cases (50%). Preserved left ventricle EF (EF=50-54%) was noted in 17 cases and reduced EF (EF<50%) in 8 (5.6%). Seventy-nine patients had mild diastolic dysfunction (56.3%), two of them moderate (1.4%) and others had normal diastolic function. Mild RV dilation was seen in 14 cases (9.9%) and moderate in two cases (1.4%). Mild pulmonary hypertension (PHT) was found in 12 subjects. Representative example of mild RV enlargement and mild PHT in a COVID-19 patient without prior cardiovascular disorder is shown in Figure2. Pericardial effusion was detected in 32 cases (22 minimal PE and 10 mild PE). The echocardiographic indices are shown in Table 2.

RV dysfunction was reported in 49 patients (34.5%) (39 mild RV dysfunction, 8 moderate, 2 severe). Table 3 demonstrates the patients' characteristics stratified by RV function. RV size was correlated with O2

saturation ($r = -0.302$, $P < 0.001$), TPI ($r = 0.337$, $P = 0.019$), ESR ($r = 0.207$, $P = 0.035$) and Platelet count ($r = 0.355$, $P < 0.001$). CPK ($P = 0.033$) and calcium ($P = 0.005$) levels were significantly higher in patients with RV dysfunction.

Our findings demonstrated that SBP (OR: 0.885, $P = 0.006$), DBP (OR: 1.24, $P = 0.003$), male gender (OR: 0.40, $P = 0.004$), higher age (OR: 1.37, $P = 0.002$), smoking (OR: 0.011, $P = 0.035$), addiction (OR: 0.011, $P = 0.011$), hypertension (OR: 2.41, $P = 0.003$), asthma (OR: 0.10, $P = 0.032$), bilateral peripheral GGO (OR: 23.6, $P = 0.019$) and coronary calcification (OR: 5.74, $P = 0.005$) were predictors of in-hospital mortality. Moreover, age (OR: 1.29, $P = 0.002$), hypertension (OR: 2.99, $P = 0.004$) and bilateral peripheral GGO (OR: 26.81, $P = 0.023$) were predictors of 30-day mortality.

In multivariate regression analysis, lower O₂ saturation at the time of admission was independently predictor of re-admission ($P < 0.001$). Furthermore, PASP ($P = 0.026$), dyslipidemia ($P = 0.002$) and RV dilation ($P = 0.037$) were significantly predictors of in-hospital mortality (after adjusting for possible cofounders). RV enlargement ($P < 0.001$), dyslipidemia ($P < 0.001$), lower LVEF ($P < 0.001$), older age ($P = 0.020$), SBP ($P = 0.001$), O₂ saturation ($P = 0.018$), shorter pulmonary acceleration time (PAT) ($P = 0.005$) and diabetes ($P = 0.025$) independently predicted 30-days mortality.

Patients with cardiac injury were mostly females ($P = 0.001$), had hypertension ($P = 0.001$), more comorbidities ($P = 0.026$), bilateral peripheral GGO ($P = 0.001$), RV dysfunction ($P = 0.029$), a higher level of CPK ($P = 0.018$) and coronary calcification ($P = 0.006$), compared to those without. Likewise, in patients with cardiac injury, pericardial effusion was significantly more reported (16 versus one, $P = 0.021$).

LV dysfunction (EF lesser than 55%) was seen in 17.6% of cases; probably most of them due to COVID-19 related cardiac injury.

4. Discussion

Investigations regarding cardiovascular complications in COVID-19 in developing countries are limited. To the best of our knowledge, our study is the first to assess cardiac function in COVID-19 patients using both echocardiography and ECG findings in those without any significant structural heart diseases. Our aim was to evaluate the direct impact of SARS-CoV-2 infection on cardiovascular indices in hospitalized patients.

As above-mentioned, patients with cardiac injury had a higher level of CPK, comorbidities and RV dysfunction. In a similar study, patients with cardiac injury had higher levels of inflammatory biomarkers, underwent mechanical ventilation more probably and had higher mortality rate and RV dysfunction. They showed that two indices of strain echocardiography, i.e. LV global longitudinal strain and RV FWLS were are powerful predictors of higher mortality in patients with COVID-19 infection ¹¹.

RV dysfunction in COVID-19 patients is common and associated with a pro-thrombotic and inflammatory state which is determined by D-dimer and CRP elevation. In our study, RV systolic dysfunction was

reported in 49 cases (34.5%) and LV systolic dysfunction in 17.6%. We showed a significant correlation between RV size and TPI level. In another retrospective study, LV systolic function was hyper-dynamic or normal in most cases (89%), RV was dilated in 41%, and impaired in 27%. It was demonstrated that RV systolic dysfunction (defined by fractional area change) was significantly associated with elevated D-dimer and CRP but not related to hsTnI ⁹.

Similar to our study, in another investigation, all indices of RV hemodynamics and function were worse compared to the LV parameters. They introduced RV dilation and dysfunction as major abnormal echocardiographic patterns among deteriorating patients ¹⁰. Many conditions can increase pulmonary vascular resistance or pulmonary pressure in hospitalized patients and precipitate acute RV failure. These conditions include pulmonary embolism (PE), hypoxic pulmonary vasoconstriction, decrease in lung volume, excessive positive end-expiratory pressure, pneumonia, hyper-carbia, the use of α -agonists or a combination of all these factors ¹⁰. In COVID-19 infection, the probable mechanisms capable for this condition include: 1) direct myocardial injury by virus; 2) increased RV afterload in ARDS; 3) hypoxic pulmonary vasoconstriction; 4) pulmonary micro-thrombosis, and 5) endothelial and microvascular injury ¹². Besides, RV dilation and dysfunction may also affect LV function by ventricular interdependence and paradoxical septal motion which causes reduction in LV volume and cardiac output ¹¹. We showed that RV dilation is a predictor of in-hospital and 30-days mortality in COVID-19 patients without any previous structural heart diseases. Similarly, Argulian et al. showed that RV dilation was predictive for in-hospital mortality in patients with COVID-19 ¹³. Another observation was reported by Szekely et al. ¹⁰, who demonstrated that increased RV end diastolic area was significantly associated with mortality. We demonstrated that dyslipidemia, lower EF, older age, SBP, O2 saturation, PAT and diabetes were also independently predictors of 30-days mortality. In line with our study, the only baseline echocardiographic parameters significantly associated with mortality were low LVEF, elevated E/e' ratio, increased RV end diastolic area, and higher Tei index ¹⁰. Therefore, identifying RV impairment would guide physicians to limit positive end-expiratory pressure and avoid hyper-capnic state, which could adversely affect RV performance by inducing pulmonary arteriolar vasoconstriction and increased RV afterload ⁹.

During acute RV pressure overload and resultant RV dysfunction and reduced cardiac output, systemic blood pressure will decrease, which may result in lowering coronary perfusion and additional reduction in RV contractility ¹⁰.

In Szekely et al study, patients with shorter PAT were older, had more comorbidities, and worse lung disease, lower oxygen saturation, higher LV filling pressure, and higher biomarkers (D-dimer, BNP, Troponin-I, and CRP). They suggested that elevated pulmonary vascular resistance in COVID-19 infection is multifactorial and related to parenchymal lung disease, pulmonary vascular disease, and elevated left atrial pressure, all leading to cardiac injury ¹⁰. Similar to our findings, there is a low prevalence of LV impairment in prior studies ⁹. Previous case reports and publications showed that COVID-19 is a known cause of severe cardiac dysfunction such as fulminant myocarditis, even without common COVID-19

symptoms or pulmonary involvement^{14–16}. Cardiac injury is prevalent in patients admitted with COVID-19 infection and related to high mortality rate² but LV systolic dysfunction is not very common.

This study performed on a larger sample size compared to previous investigations. We enrolled only patients with positive PCR test for COVID-19. Another strength of this study was excluding patients with previous underlying heart diseases, which helps to elucidate the pure effects of COVID-19 on cardiovascular system. Besides, all of evaluations were performed by a single cardiologist, which eliminates inter-observer discrepancies.

Limitation:

Further studies on the reasons of RV dysfunction are needed.

5. Conclusion

RV systolic dysfunction was prevalent in patients with COVID-19 and can be correlated with higher mortality. Moreover, lower LVEF, shorter PAT and lower O₂ saturation, older age, elevated SBP, dyslipidemia and diabetes were independently predictors of 30-days mortality. Using echocardiography for risk assessment in patients with COVID-19, especially with previous history of diabetes, hypertension and dyslipidemia, can be helpful and lead to better management.

Abbreviations

COVID-19

Coronavirus disease 2019

SARS-CoV-2

Severe acute respiratory syndrome coronavirus 2

PCR

polymerase chain reaction

CT

computerized tomography

ECG

Electrocardiography

RV

Right ventricle

PASP

pulmonary artery systolic pressure

EF

ejection fraction

TPI

troponin I

TTE
transthoracic echocardiography
GGO
ground glass opacification
PHT
pulmonary hypertension
ESR
Erythrocyte Sedimentation Rate
CPK
Creatine phosphokinase
SBP
systolic blood pressure
DBP
diastolic blood pressure
OR
Odds ratio
PAT
pulmonary acceleration time
FWLS
Free wall longitudinal strain
CRP
C reactive protein
hsTnI
high sensitive troponin I
PE
pulmonary embolism
ARDS
Acute respiratory distress syndrome
BNP
Brain natriuretic peptide

Declarations

1. Funding

This study was funded by Mashhad University of Medical Sciences.

2. Availability of data and materials

The datasets used or analyzed in the current study are available from the corresponding author on reasonable request.

3. Ethics approval and consent to participate

This study was approved by the ethical committee of the medical faculty of Mashhad University of Medical Sciences. The ethical code of the study was IR.MUMS.REC.1399.016. All patients filled a written informed consent form before entering the study.

4. Competing interests

The authors declare that they have no competing interests.

5. Authors' contributions

Dr. Faeze Keihanian designed the study, collected data and performed echocardiographic assessments, wrote the first draft of the article and approved the final version; Dr. Hoorak Poorzand participated in data analysis and echocardiographic assessments, read the final version and approved it; Dr. Amin Saeidinia participated in data collection, statistical analysis and writing first draft of the article. Dr. Ali Eshrgahi designed the study, wrote the first draft of the article and approved the final version.

6. Acknowledgement

We would like to thank the deputy of research, Mashhad University of Medical Sciences and physicians and nurses working hard in COVID-19 wards in Imam Reza hospital.

7. Consent for publication

All patients took consent to publish their data anonymously.

References

1. NeJhaddadgar N, Ziapour A, Zakkipour G, Abbas J, Abolfathi M, Shabani M Effectiveness of telephone-based screening and triage during COVID-19 outbreak in the promoted primary healthcare system: a case study in Ardabil province, Iran. *Journal of Public Health* 2020:1–6
2. Keihanian F, Bigdelu L (2020) Cardiovascular Considerations in COVID19: A Comprehensive Review. *Ther Clin Risk Manag* 16:1089–1097
3. Force ADT (2012) Acute respiratory distress syndrome. *JAMA* 307:2526–2533
4. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, Xiang J, Wang Y, Song B, Gu X (2020) Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The lancet*

5. Shi S, Qin M, Shen B, Cai Y, Liu T, Yang F, Gong W, Liu X, Liang J, Zhao Q, Huang H, Yang B, Huang C (2020) Association of Cardiac Injury With Mortality in Hospitalized Patients With COVID-19 in Wuhan, China. *JAMA Cardiology* 5:802–810
6. Xu Z, Shi L, Wang Y, Zhang J, Huang L, Zhang C, Liu S, Zhao P, Liu H, Zhu L (2020) Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *The Lancet respiratory medicine* 8:420–422
7. Guo T, Fan Y, Chen M, Wu X, Zhang L, He T, Wang H, Wan J, Wang X, Lu Z (2020) Cardiovascular implications of fatal outcomes of patients with coronavirus disease 2019 (COVID-19). *JAMA cardiology*
8. Mitchell M, Fonarow J, Peterson G, Geraci P, Horwich S (2013) T. 2013 ACCF/AHA Guideline for the Management of Heart Failure. Executive Summary
9. Mahmoud-Elsayed HM, Moody WE, Bradlow WM, Khan-Kheil AM, Senior J, Hudsmith LE, Steeds RP (2020) Echocardiographic Findings in Patients With COVID-19 Pneumonia. *Can J Cardiol* 36:1203–1207
10. Szekely Y, Lichter Y, Taieb P, Banai A, Hochstadt A, Merdler I, Gal Oz A, Rothschild E, Baruch G, Peri Y (2020) The Spectrum of Cardiac Manifestations in Coronavirus Disease 2019 (COVID-19)-a Systematic Echocardiographic Study. *Circulation*
11. Xie Y, Wang L, Li M, Li H, Zhu S, Wang B, He L, Zhang D, Zhang Y, Yuan H (2020) Biventricular Longitudinal Strain Predict Mortality in COVID-19 Patients. *Frontiers in cardiovascular medicine* 7:418
12. Park JF, Banerjee S, Umar S (2020) In the eye of the storm: the right ventricle in COVID-19. *Pulmonary Circulation* 10:2045894020936660
13. Argulian E, Sud K, Vogel B, Bohra C, Garg VP, Talebi S, Lerakis S, Narula J (2020) Right ventricular dilation in hospitalized patients with COVID-19 infection. *Cardiovascular Imaging* 13:2459–2461
14. Paul J-F, Charles P, Richaud C, Caussin C, Diakov C (2020) Myocarditis revealing COVID-19 infection in a young patient. *European Heart Journal - Cardiovascular Imaging* 21:776–776
15. Beşler MS, Arslan H (2020) Acute myocarditis associated with COVID-19 infection. *The American Journal of Emergency Medicine* ;38:2489.e2481-2489.e2482
16. Doyen D, Mocerì P, Ducreux D, Dellamonica J (2020) Myocarditis in a patient with COVID-19: a cause of raised troponin and ECG changes. *The Lancet* 395:1516

Tables

Table 1- Baseline demographic characteristics, comorbidities, clinical and laboratory data

Parameter	
Age (years) (mean±SD)	60.69± 15.70
Sex (N, %)	
Male	82, 57.7
Female	60, 42.3
Comorbidities (N, %)	
Diabetes	34, 23.9
Hypertension	34, 23.9
Dyslipidemia	16, 11.3
History of Stroke	4, 2.8
Chronic kidney disease	6, 4.2
Lung disease	9, 6.3
History of malignancy	3, 2.1
Current smoker	8, 5.6
Addiction	17, 11.9
Vital sign (Mean± S.D)	
Heart rate (beats/minute)	102.43±19.23
Systolic blood pressure (mmHg)	135.46± 20.66
Diastolic blood pressure (mmHg)	84.62± 20.66
O ₂ saturation (%)	88.31± 7.83
Temperature (°C)	37.38± 0.79
Respiratory rate (breaths/minute)	23.92± 5.42
Symptoms (N, %)	
Cough	78, 54.9
Dyspnea	129, 90.8
Fever	70, 49.3
Weakness	57, 40.1
Chest pain	2, 1.4
Gastrointestinal	25, 17.6
Confusion	8, 5.6

Laboratory data (Mean± S.D)	
Blood sugar (mg/dL)	142.12± 22.88
Urea (mg/dL)	43.94± 17.03
Creatinine (mg/dL)	1.34± 0.7
Sodium (meq/L)	136.28± 3.49
Potassium (meq/L)	4.09± 0.60
Total bilirubin (mg/dL)	0.61± 0.43
Direct bilirubin (mg/dL)	0.25± 0.15
Calcium (mg/dL)	8.54± 0.89
Magnesium (mg/dL)	2.06± 0.42
Aspartate transaminase (IU/L)	44.90± 18.23
Alanine aminotransferase (IU/L)	49.06± 17.66
Alkaline phosphatase (IU/L)	212.32± 80.91
Lactate dehydrogenase (IU/L)	606.65± 230.51
Creatine phosphokinase (IU/L)	350.97± 143.38
Cardiac troponin I (ng/ml)	56.22± 25.24
PCO ₂ (mmHg)	42.10± 13.79
HCO ₃ (meq/L)	27.28± 6.57
White blood cell (10 ⁶ /L)	8860.56± 5732.48
PMN (%)	77.88± 9.31
Lymph (%)	16.65± 7.93
Hemoglobin (g/dL)	13.18± 2.11
MCV (fL)	84.54± 7.01
RDW (%)	14.08± 1.74
Platelet (10 ⁹ /L)	222.18±126.55
CRP (mg/L)	88.63±74.66
ESR (mm/h)	42.9±30.26

PCO₂= Partial pressure of carbon dioxide; *HCO₃*= Bicarbonate; *PMN*= Polymorphonuclear leukocytes; *Lymph*= Lymphocyte; *MCV*= Mean corpuscular volume; *RDW*= Red cell distribution width; *CRP*= C-reactive

protein; ESR= Erythrocyte sedimentation rate

Table 2- Echocardiographic findings of patients

Echocardiographic parameters	Mean± S.D
LVEDD	42.08± 4.70
LVEF	54.96± 4.23
E	74.47± 18.82
A	84.67± 22.01
E'	9.16± 2.39
E/ e'	7.95± 2.34
LVOT VTI	18.87± 3.98
TV annulus	29.10± 4.12
RV diameter	27.27± 3.97
RV S'	12.33± 2.36
TAPSE	19.04± 3.39
RVOT VTI	16.65± 3.24
PAT	97.63± 22.32
PASP	24.43± 6.52

LVEDD=Left ventricular end diastolic diameter; LVEF=Left ventricular ejection fraction; LVOT= Left ventricular outflow tract; VTI= Velocity Time Integral; RV= Right ventricle; TAPSE= Tricuspid annular plane systolic excursion; RVOT= Right ventricular outflow tract; PAT= Pulmonary acceleration time; PASP= Pulmonary artery systolic pressure

Table 3- Patients' characteristics Stratified by RV function

Parameter		RV dysfunction (n=50)	Normal RV function (n=92)	P Value
Age (Mean± S.D)		62.63± 16.29	59.26± 15.63	0.240
Gender (N, %)	Male	23, 46.9	59, 63.4	0.044
	Female	26, 53.1	34, 36.6	
Heart Rate (Mean± S.D)		100.63±19.69	103.24± 19.21	0.464
Systolic Blood Pressure (Mean± S.D)		135.40± 19.86	135.39± 21.45	0.999
Diastolic Blood Pressure (Mean± S.D)		83.73± 12.77	85.48± 16.01	0.504
O ₂ saturation (Mean± S.D)		88.10± 5.88	88.42± 8.85	0.808
Laboratory data (Mean± S.D)				
Urea		41.18± 16.77	44.46± 17.37	0.556
Creatinine		1.23± 0.7	1.40± 0.8	0.595
Calcium		8.87± 0.99	8.35± 0.78	0.005
Lactate dehydrogenase		598.51± 211.97	606.82± 241.96	0.835
Creatine phosphokinase		472.00± 181.02	243.50± 146.01	0.033
White blood cell		7302.04± 3021.20	9621.97± 6665.41	0.064
Hemoglobin		13.58± 2.16	12.92± 2.06	0.108
RDW		14.45± 1.96	13.84± 1.57	0.082
PMN		76.03± 8.69	78.76± 9.61	0.985
Lymph		18.16± 7.04	15.87± 8.37	0.090
Platelet				
CRP		80.44±63.8	92.69±79.49	0.370
ESR		47.25±30.33	39.47±30.10	0.217
ECG assessment				
ECG Rhythm (%)	Sinus Tachycardia	48.9	49.4	0.740
	Atrial Fibrillation	0	1.3	
	Normal sinus rhythm	51.1	49.4	
ECG Axis (%)	Normal axis	73.3	74.7	0.543

	LAD	22.2	15.6	
	RAD	4.4	10.1	
ECG abnormality (%)	RBBB	8.2	6.6	0.42
	LAHB	12.2	9.9	
	LBBB	4.1	0	
	LAHB+ RBBB	4.1	0	
Symptoms (%)				
Chest Pain		4.1	0	0.121
Nausea and vomiting		12.2	13.2	0.550
Diarrhea		8.2	2.2	0.112
Dyspnea		87.8	91.2	0.354
Cough		55.1	56	0.528
Fever		51	49.5	0.50
Weakness		26.5	45.1	0.024
Confusion		8.2	4.4	0.289
Comorbidities (%)				
Diabetes		28.6	22	0.253
Addiction		5.6	6.3	0.667
Smoker		2.1	3.5	0.580
Dyslipidemia		12.2	11	0.513
Hypertension		20.4	26.4	0.284
History of stroke		0	4.4	0.174
Asthma		4.1	6.6	0.424
COPD		0	2.2	0.421
Malignancy		2.2	4.1	0.437
ESRD		4.1	3.3	0.429
CT scan findings (%)				
Bilateral peripheral GGO		46.9	34.1	0.095
Bilateral diffused GGO		32.7	34.7	0.358

Unilateral GGO		8.2	17.6	0.10
Bilateral Pleural effusion		8.2	11	0.416
Consolidation		69.4	59.3	0.161
Lymphadenopathy		24.5	33	0.198
Echocardiographic parameters				
LVEDD (Mean± S.D)		41.61± 4.64	42.27± 4.77	0.089
E/ e' (Mean± S.D)		8.05± 3.09	7.90± 1.85	0.755
LVEF (n, %)	≥55	34, 69.4	79, 84.9	0.024
	50-54	9, 18.4	12, 12.9	
	40-49	6, 12.2	2, 2.2	
Pulmonary hypertension (n, %)		6, 12.8	5, 5.6	0.193
RV size (n, %)	Normal	41, 83.7	85, 90.4	0.145
	Mild enlargement	6, 12.2	8, 8.6	
	Moderate enlargement	2, 1.4	0	

RDW= Red cell distribution width; PMN= Polymorphonuclear leukocytes; Lymph= Lymphocyte; CRP= C-reactive protein; ESR= Erythrocyte sedimentation rate; ECG= Electrocardiogram; LAD= Left axis deviation; RAD= Right axis deviation; RBBB= Right bundle branch block; LBBB= Left bundle branch block; LAHB= Left anterior hemi block; COPD= Chronic obstructive pulmonary disease; ESRD= End stage renal disease; GGO= ground glass opacification; LVEDD: Left ventricular end diastolic diameter; LVEF= Left ventricular ejection fraction; RV= Right ventricle

Figures

Figure 1

Flowchart of study. VHD= valvular heart disease; IHD= Ischemic heart disease; SHD= structural heart disease.

Figure 2

Example of two-dimensional transthoracic echocardiography of a patient without previous cardiovascular disorder. A) Mild right ventricular enlargement in RV focused view; B) Off axis four-chamber view of the heart suggesting moderate TR with TR gradient about 41mmHg. Estimated pulmonary arterial pressure in this patient was 44 mmHg (considering normal IVC size and collapsibility).