

# Prevalence of cardiovascular disease risk factors and long-term outcomes in the Saudi population: Results from the Prospective Urban Rural Epidemiology study (PURE-Saudi)

Khalid Alhabib (✉ [khalidalhabib13@hotmail.com](mailto:khalidalhabib13@hotmail.com))

king fahad cardiac center king saud university. <https://orcid.org/0000-0002-6692-3874>

**Mohammed A Batais**

Department of Family and Community medicine, College of Medicine, King Saud University, Riyadh

**Turky H Almigbal**

King's College London School of Medical Education

**Mostafa Q Alshamiri**

Department of Cardiac Sciences, King Fahad Cardiac Center, College of Medicine, King Saud University, Riyadh

**Hani Altaradi**

Department of cardiac Science, king Fahad cardiac Center, College of Medicine, King Saud University, Riyadh

**Sumathy Rangarajan**

Population Health Research Institute, DBCVS REsearch Institute, McMaster University, Hamilton, Canada

**Salim Yusuf**

Population Health Research Institute, DBCVS Researc Institute, McMaster University, Hamilton, Canada

---

## Research article

**Keywords:** PURE-Saudi, diet, stress, CVD, lifestyle behaviors

**Posted Date:** January 16th, 2020

**DOI:** <https://doi.org/10.21203/rs.2.21056/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on August 8th, 2020. See the published version at <https://doi.org/10.1186/s12889-020-09298-w>.

# Abstract

## Objective

We report the prevalence of unhealthy lifestyle behaviors, cardiovascular disease (CVD) risk factors, and long-term outcomes within the Saudi population, stratified by age, sex, and place of residence.

## Methods

The Prospective Urban Rural Epidemiology (PURE) study is a global cohort study including adults of 35–70 years old in 20 countries. PURE-Saudi study participants were recruited from 19 urban and 6 rural communities randomly selected from the Central province (Riyadh and Alkharij) between February 2012 and January 2015. Clinical follow-up of major CVD events and mortality is ongoing.

## Results

The PURE-Saudi study enrolled 2047 participants (mean age,  $46.5 \pm 9.12$  years; 43.1% women; 24.5% rural). Overall, 69.4% had low physical activity, 49.6% obesity, 34.4% unhealthy diet, 32.1% dyslipidemia, 30.3% hypertension, 25.1% diabetes, 12.2% were current smokers, 15.4% self-reported feeling sad, 16.9% had history of stress (several periods), 6.8% had permanent stress, 0.98% had history of stroke, 0.64% had heart failure, and 2.5% had coronary heart disease (CHD). Compared to women, men were more likely to be current smokers (21% vs. 0.45%,  $p < 0.001$ ), have diabetes (28.2% vs. 21.3%,  $p < 0.001$ ), and have history of CHD (3.2% vs. 1.6%,  $p = 0.02$ ); while women were more likely to be obese (58.6% vs. 42.8%,  $p < 0.001$ ), have central obesity (70.7% vs. 32.7%,  $p < 0.001$ ), self-report sadness (22.7% vs. 9.9%  $p < 0.001$ ), experience stress (several periods), feel permanent stress (9.9% vs. 4.5%,  $p < 0.001$ ), and have low education (46.6% vs. 20.2%,  $p < 0.001$ ). Compared to participants in urban areas, those in rural areas had higher rates of diabetes (31.1% vs. 23.3%,  $p < 0.001$ ), obesity (56.6% vs. 47.3%,  $p < 0.001$ ), and hypertension (35.5% vs. 28.6%,  $p = 0.004$ ); and lower rates of unhealthy diet, self-reported sadness, stress (several periods), and permanent stress. Compared to middle- and old-age individuals, younger participants more commonly reported unhealthy diet, permanent stress, and self-reporting of being sad.

## Conclusion

PURE-Saudi, the first population cohort study in Saudi Arabia, revealed a high prevalence of CVD risk factors in the adult Saudi population, with higher rates in rural than urban areas. National public awareness programs and multi-faceted healthcare policy changes are urgently needed to reduce the future burden of CVD risk and mortality.

## Introduction

Cardiovascular disease (CVD) is the leading cause of mortality worldwide, contributing to 31% of all deaths. [1] CVD is also becoming a major health concern in the Gulf Council Countries (GCC) including Saudi Arabia, as it is estimated that the overall deaths from CVD account for over 45% of all deaths.[1, 2]

The most common CVD risk factors were identified in the INTERHEART and INTERSTROKE studies were hypertension, diabetes, dyslipidemia, obesity, smoking, physical activity, poor diet, and alcohol consumption.[3, 4] In the Gulf countries, lifestyle has changed dramatically due to the rapid urbanization with an increasing in the poor diet and the adoption of a sedentary lifestyle. Consequently, the rates of CVD risk factors and the non-communicable diseases among the Gulf population are also likely to have increased; the prevalence sometimes exceed that of developed countries.[5] In Saudi Arabia, there was very limited national representative data on the prevalence of CVD, high prevalence of key risk factors was observed, [2, 6–32] and we lack data on CVD incidence as no large long term cohort study has been completed that compared CVD risk factors and cardiovascular outcomes and mortality among Saudi population. Therefore, well-designed national population-based cohort study focusing on CVD and its associated risk factors is needed.

The Prospective Urban Rural Epidemiology (PURE) study is a prospective cohort study that aimed to collect data on social, environmental, and individual CVD risk factors and chronic diseases in high-, middle-, and low-income countries.[33] Saudi Arabia has been classified as high-income country and joined the global PURE in 2012, and PURE Saudi is the first cohort study that aims to investigate the prevalence of unhealthy lifestyle behavior, CVD risk factors, events and mortality rates among different age groups, men vs women, and urban vs rural.

## Methods

### Study design and participants

the study design, methods (including sampling, information gathered, and follow-up strategy), and participant characteristics of the PURE study have been published previously. [33-36] Briefly, adults aged between 35 years and 70 years from 367 urban and 302 rural communities in 20 countries were included. Households were eligible if one or more of the households' members was aged between 35 and 70 years and the household members intended to stay at that address for a further four years. Risk factors and medical history were documented, and a physical examination was performed on subjects who provided written informed consent. Details of sampling, information gathered, and follow-up strategy have been previously reported in several publications. [34, 35]

### Procedures

Data regarding the demographic factors, socioeconomic status, medical history, health behaviors (e.g., smoking, physical activity, diet, alcohol intake), and subjects' household members were collected by using standardized questionnaires. In addition, we collected information regarding psychosocial factors and other CVD risk factors such as hypertension, diabetes and obesity as described in the INTERHEART study. [3] Anthropometric measures included weight, height, body mass index, waist circumferences and blood pressure. Physical activity was measured using the International Physical Activity Questionnaire (IPAQ) and was categorized according to the metabolic equivalent of task (MET) per min per week into low (3000 MET min per week) activity. Subjects were considered to have diabetes if a physician had

previously diagnosed them with diabetes and/or if they had a fasting plasma glucose level of  $\geq 126$  mg/dl (7.0 mmol/l) or were being treated with glucose-lowering medication. Subjects with history of hypertension, current use of antihypertensive medication and/or a blood pressure  $\geq 140/90$  mm Hg were considered to have hypertension. All subjects were asked whether they had a medical diagnosis of hypertension (awareness) and whether they were receiving antihypertensive medication (treatment). Hypertension control was defined as the proportion of subjects with an average systolic and diastolic blood pressure of  $< 140/90$  mm Hg. A total cholesterol level  $> 5.2$  mmol/l (201 mg/dl) was considered to be an elevated level. Major CVD (myocardial infarction or angina, stroke or heart failure) were the main clinical outcomes included in the analyses based on subjects self-reported responses.

## **PURE Saudi**

Recruitment of Saudi individuals for the PURE study was carried out between February 2012 and January 2015. Nineteen urban and six rural communities were enrolled from the city of Riyadh and Alkharj (Central province). Urban communities involved were defined according to the governmental geographic distribution of the districts, while rural communities were defined as those regions located at least 50 km away from the center of Riyadh. Due to cultural acceptance, the study team met the participants in the primary health care centers (PHCC) in each community. The database of each PHCC was screened and the eligible subjects were invited to participate in the study by calling and/or sending them a short text message service to their mobile phones. A target number of at least 50 subjects were required to be enrolled per each PHCC. All subjects were encouraged to invite their eligible family members who were living within the same household. All blood samples were shipped to the laboratory in the King Khalid University Hospital at King Saud University, Riyadh. Subsequently, the blood results were returned to the treating physicians at the PHCC for further assessment and management.

During the follow-up period, the study team conducted a reminder call at 18 months after the baseline recruitment by telephone to remind the subjects about their upcoming follow-up at 3, 6, and 9 years.

## **Statistical analysis**

Data are summarized as means and standard deviations for continuous variables and as numbers and percentages for categorical variables. Age was categorized into three groups: 35-49 years, 50-59 years and 60-70 years. Education was categorized as high (i.e., trade school, college or university); medium (i.e., secondary school or high school), low (i.e., primary education or no education) or unknown. Proportions and means were compared using Chi-square test and *t* tests, respectively, using 2-sided testing. A *P* value of less than 0.05 was considered statistically significant.

# **Results**

## **Overall cohort**

The PURE-Saudi study enrolled 2047 participants, with mean age of  $46.5 \pm 9.12$  years (Table 1). There were 1165 men (56.9%) and 882 women (43.1%). Around one third of the total cohort had low educational level. The median body mass index (BMI) of the participants was 29.9, where majority of them were either overweight (35.3%) or obese (49.6%). Among obese patients, 30% had BMI between 30-35, and 19.5% had BMI > 35. The prevalence of abdominal obesity defined as waist circumference >102 cm (men) or >88 cm (women) was 49%. This prevalence is increased to 74.3% when the measured waist circumference >90 cm (men) or >85 cm (women). The prevalence of hypertension was 30.3%, out of which only 61.1% were aware of it, 58.9% were treated and 30.7% participants achieved blood pressure control. Approximately 34.4% of the total cohort reported eating unhealthy diet, 69.4% of participants reported low physical activity, 12.2% were current smokers, and 10.6% were former smokers. The prevalence of diabetes was 25.1%, out of which 2.7% were on insulin therapy, 60.7% received oral hypoglycemic agents (OHA), 6.6% received both (insulin and OHA), and 30% received no treatment. About one third (32.1%) had high total cholesterol level, however, only 0.9% were on statin therapy. Moreover, the prevalence of self-reported of being sad or blue was 15.4%, while 16.9% had reported history of several feeling of stress and 6.8% had permanent feeling of stress. History of stroke was found in 0.98%, heart failure in 0.64%, and coronary heart disease in 2.5%. In addition, the median INTERHEART risk score was 11.

### **Young vs Middle vs Old age**

Compared to the younger and middle age groups (35-49 years and 50-59 years), older participants (60-70 years) had higher prevalence of low physical activity (75.2% vs 66.8% and 73.9% ,  $p=0.002$ ), hypertension (65% vs 18.2% and 48.4%,  $p < 0.001$ ), diabetes (57.5% vs 12.9% and 44.9%,  $p < 0.001$ ), low level of education (62.8% vs 21.7% and 44.9%,  $p < 0.001$ ), stroke (3.9% vs 0.4% and 1.1%,  $p < 0.001$ ), history of heart failure (1.8% vs 0.15% and 1.5%,  $p < 0.001$ ), and history of ischemic heart disease (7.5% vs 1.1% and 4.1%,  $p < 0.001$ ) (Table 2 and Figure A). The awareness and treatment of blood pressure were higher among older participants compared with middle and younger individuals (73.5% vs 70.04% and 45.5%, 73.5% vs 69.2% and 40.7%,  $P < 0.0001$  for all). While, middle age participants had more controlled of blood pressure compared to young and older individuals (39.2% vs 19.9% and 35.4%,  $P < 0.0001$ ). However, younger age groups reported more consumption of unhealthy diet (38.9% and 25.2% vs 26.5%,  $p < 0.001$ ), general feeling of permanent stress (9.1% and 2.4% vs 1.9%,  $p < 0.001$ ), and self-reporting of being sad (18.6% and 10% vs 7.1%,  $p < 0.001$ ). The prevalence of obesity with BMI 30-35 and BMI >35 was 34.1% and 21.8% in the middle age group compared to 28.7% and 19.4% in the young age, and 28.8% and 15.5% in older age groups (0.001) respectively. Moreover, the median level of glucose and triglycerides were significantly increased with advanced age (Additional Table 1).

### **Men vs Women**

Compared to women, men have significantly higher proportion of current smoking (21% vs 0.45%,  $p < 0.001$ ) or former smoking (17.9% vs 0.91%,  $p < 0.001$ ), diabetes (28.2% vs 21.3%,  $p < 0.001$ ), obesity with BMI 30-35 (39.9% vs 32.4%,  $p < 0.001$ ), history of ischemic heart disease (3.2% vs 1.6%,  $p = 0.02$ ), and the

median INTERHEART risk score (13% vs 10%,  $p < 0.001$ ). On the other hand, women had higher prevalence of obesity with BMI > 35 (26.1% vs 14.5%,  $p < 0.001$ ), central obesity (70.7% vs 32.7%,  $p < 0.001$ ), self-reporting of being sad (22.7%, vs 9.9%  $p < 0.001$ ), several feeling of stress (23.1% vs 12.4%,  $p < 0.001$ ) or permanent feeling of stress (9.9% vs 4.5%,  $p < 0.001$ ), and low level of education (46.6% vs 20.2%,  $p < 0.001$ ) (Table 1 and Figure B). The awareness, treatment, and control of blood pressure were similar among women and men (61.3% vs 60.9%, 60.9% vs 57.6%, 34.5% vs 28.3%, ( $P = 0.93$ ,  $P = 0.41$ ,  $P = 0.10$ )) respectively. In addition, men were more likely to have lower HDL-cholesterol levels compared with women (Additional Table 2).

## Urban vs Rural

Participants living in urban areas represented 75.48% of the participants, while 24.5% lived in rural areas. Compared to urban areas, participants living in rural areas reported higher prevalence of diabetes (31.1% vs 23.3%,  $p < 0.001$ ), obesity (56.6% vs 47.3%,  $p < 0.001$ ), hypertension (35.5% vs 28.6%,  $p = 0.004$ ), and low level of education (46.4% vs 26.7%,  $p < 0.001$ ). In addition, the median BMI was higher among rural compared to urban participants (30.8% vs 29.7%,  $p < 0.001$ ). On the other hand, urban participants more likely to consume unhealthy diet (36% vs 29.4%,  $p = 0.007$ ), to self-report of being sad (17.6% vs 8.6%,  $p < 0.001$ ), to have several feeling of stress (19.7% vs 8.7%,  $p < 0.001$ ), and permanent feeling of stress (8.03% vs 3.02%  $p < 0.001$ ) (Table 3 and Figure C). The awareness, treatment, and control of blood pressure were similar among the urban and rural communities (62.2% vs 58.4%, 60.1% vs 55.6%, 32.6% vs 25.8%, ( $P = 0.38$ ,  $P = 0.29$ ,  $P = 0.10$ ) respectively. Furthermore, urban participants showed higher median in the glucose level and proportion in the glucose level between 6-7 mmol/L in non-diabetic patients (Additional Table 3).

## Follow-up

There were 6 patients diagnosed with cancer and total of 22 deaths (at rate of 0.07 and 0.26 per 100 person years of follow-up respectively). The overall rates of major cardiovascular events showed similar pattern to that for mortality, where 24 had a myocardial infarction, 6 had a stroke, 4 had heart failure, and 34 had at least one major cardiovascular events with rates of 0.29, 0.07, 0.05, and 0.41 per 100 person years of follow-up respectively.

## Discussion

The PURE-Saudi is the first cohort study in Saudi Arabia that assessed the current prevalence of CVD risk factors, in addition to the long-term clinical outcomes. Being part of the internationally standardized and validated surveys of the global PURE study allows for a valid interpretation and direct comparison of the results in the context of other enrolled countries with variable economic scales and health care systems. Our study showed two major findings. First, the high prevalence of CVD risk factors among the Saudi population: two-thirds had low physical activity, about half had obesity, one third consumed unhealthy diet, about one third had dyslipidemia, about one third had hypertension, and about one quarter had

diabetes. Second, the relative proportion of the individual CVD risk factors varies according to age, sex, and residence place.

The PURE-Saudi study confirms that the prevalence of un-healthy life styles and CAD risk factors remains high in the Saudi population despite over a decade of several previous population cross-sectional surveys (Additional Table 4). The Coronary Artery Disease in the Saudis Study (CADISS) was a national epidemiological health survey between 1995 and 2000, and included 17,395 Saudis aged 30-70 years through a multistage stratified cluster sampling technique. The overall prevalence of hypercholesterolemia was 54%, obesity 35.6%, hypertension 26.1%, diabetes 23.7%, and smoking 12.8%. [11-13, 23-25]. Another national cross-sectional survey by the Saudi Ministry of Health involved 4758 participants aged 15 year to 64 year and aimed to estimate the prevalence of some of the risk factors of non-communicable diseases (NCD) by using the WHO STEPwise approach for NCD surveillance. It indicated that the prevalence of physical inactivity was 67.6%, obesity 36.2%, hypercholesterolemia 19.1%, hypertension 11.6%, diabetes 15.3%, and current daily smoking 10.9%.[14] The Saudi Health Information Survey (SHIS) was done also by the Ministry of Health and enrolled 10,735 Saudis aged 15 years or older in 2013. The prevalence of physical inactivity was 39.8%, obesity 28.7%, daily consumption of < 2 servings of fruits and vegetables 61.6%, hypertension 15.2%, diabetes 13.4%, hypercholesterolemia 8.5%, and smoking 12.1%.[9, 10, 28, 29, 37]. As a result of such high prevalence of the CVD risk factors, patients in Saudi Arabia present almost a decade younger with acute coronary syndromes and acute heart failure compared to those in the developed countries; hence suffer from being at high risk for cardiovascular complications and mortality. [38-40]

Compared to the general population, patients with diabetes are 2 to 4 times more likely to develop CVD. [41]. According to the International Diabetes Federation diabetes Atlas (8th edition), Saudi Arabia is among the top ten countries in the prevalence of diabetes mellitus. [42] It is estimated that the diabetes prevalence will increase by 110% in the Middle East and North Africa (MENA) by 2045. The prevalence of diabetes in PURE Saudi was among the highest levels reported in the global PURE data. Overall, diabetes prevalence was 11%, and the prevalence of diabetes varied between country income groups, where prevalence of diabetes was lowest in HICs 6.6% and highest in low-income countries (LICs) 12.3%. [43]. Moreover, analysis of the global PURE data showed that hypertension prevalence was 40.8%, and that only third of the patients had reached their blood pressure targets.(44) A report of prevalence, awareness, treatment, and control of hypertension from baseline PURE data from four Middle East countries [Iran, Occupied Palestinian Territory (OPT), Saudi Arabia, and United Arab Emirates (UAE)] showed that one-third had hypertension, about half of them were aware and treated, but only one-fifth were controlled.[45] The prevalence of hypertension was highest in UAE (52%) and lowest in Iran (28%), while the awareness, treatment, and control of hypertension were higher in the OPT and Saudi Arabia, compared with UAE and Iran.[45] Other previous national studies reported suboptimal hypertension control in the Saudi population.[7, 24]. Affordability of medications is one of the main reasons for the low rate of hypertension control in the world. [46] However, in Saudi Arabia, healthcare and medications are free of charge, making them largely accessible to the population. Several barriers facing patients, physicians, and healthcare systems pursuing control of hypertension, indicating the need for multifaceted

interventions.[47, 48] Forgetting to take medical therapies and medication side effects were important barriers to medication adherence.[48] Analysis data from a large household survey of 10,735 participants to identify barriers to healthcare in Saudi Arabia found that neither distance to nor type of healthcare clinic were barriers to management of chronic diseases, and highlighted the importance to individual's healthcare seeking practices rather than system based as potential barriers. Possibly that some Saudi population have specific healthcare-seeking practices and they mostly seek healthcare when they are sick and that contradict the concept of an old Arab proverb 'Prevention is better than treatment'.[49] Lack of primary care physicians knowledge and awareness of hypertension guidelines was also found where one-fourth of 322 primary care physicians had deficient knowledge regarding the correct definition of hypertension.[50] Regarding the healthcare systems barriers, inappropriate coordination between medical sectors has been reported as about one third of Saudi hypertensive patients did not have hypertension file at the primary health care centers and they received medical care at different health care sectors that led to missing their regular appointments.[47].

One of the main findings of PURE Saudi is that women are more obese compared to men. Possible explanation of higher obesity prevalence in women in this cohort may be due to some sociocultural factors and governmental bylaws such as – only until recently- the requirement of women to have a car driver for transportation purposes and barriers to practice physical activities in public places that could lead to increasing prevalence of sedentary lifestyle and obesity in Saudi women. Therefore, increasing accessibility of women to exercise facilities and providing safe walking areas are likely to help to reduce the obesity prevalence. Recently, women gymnasia in Saudi Arabia became more accessible and women were allowed to drive by themselves, which could potentially help in improving access to a healthier lifestyle. On the other hand, the lower rate of diabetes in women than men in our study may attributed to the well-recognized greater willingness of women than men to seek medical advice. [51] In addition, women are also more willing than men to adhere to diabetes daily management such as restricted diet, blood glucose monitoring, and medication adherence. [52]

Another important issue to be highlighted is the common belief that risk of developing CVD is higher in individuals who live in urban areas compared to those in rural areas. [53]. Findings from the global PURE cohort from high-income countries reported that similar INTERHEART risk score among population in rural areas in comparison to urban population. [35] However, the PURE Saudi study showed that population living in rural areas had higher CVD risk factors particularly in the prevalence of diabetes, hypertension and obesity compared to the urban population. Possible reasons might be related to 'urbanization of rural life', a term that described by some researchers,[54] where agriculture became mechanized, and cars are used for rural transport, road infrastructure improved, and more consumption of processed carbohydrates and commercially prepared and processed food through national and transnational companies, all of these would contribute to the increase in rural obesity.[55-57] In addition, the limited time and space for cooking healthy meals and possibly perceptions of large weight as a sign of affluence could also exacerbate these effects.[57, 58] Our findings might also reflect less access to and /or low availability of a health care prevention and management facilities in the rural areas. The reason behind these disparities may be due to inconsistent insurance policies, poor healthcare infrastructure and

privatization, and accessibility to healthcare facilitates which largely focus on the urban population, leaving the rural population at disadvantage. [59] The higher prevalence of diabetes in rural rather than urban areas provide support for the link between diabetes and lifestyle risk factors (lifestyle changes are less prominent in rural areas). In addition, the lower educational level among rural population as demonstrated in this study may also partly explain the differences in risk factor levels, resembling what was found in the Vasterbotten Intervention Program study in Sweden.[60] For instance, rural population with only primary education level had consistently higher prevalence of hypertension than urban population with higher educational levels.[60] The data from global PURE that assessed socioeconomic status and risk of CVD in 20 low, middle, and high-income countries, education, rather than wealth, was the socioeconomic indicator most consistently associated with outcomes where the major CVD events and all cause mortality were more common among people with low levels of education in all types of country studied. However, variances in outcomes between educational levels were not explained by variances in risk factors, which decreased as educational level increased in high-income countries, but increased as educational level increased in low-income countries.(61) Furthermore, results from MONICA study suggested that lower education level among rural population could enhance CVD risk, but causality is difficult to prove.[62]

In the present study, individuals living in urban areas were more prevalent in the consumption of unhealthy diet, sadness and stress. Recent global systematic evaluation of dietary consumption patterns across 195 countries found that improvement of diet prevents one in every five deaths globally and suboptimal diet was responsible for more deaths than other risk factors including smoking, highlighting the urgent need to improve people diet.[63] Urbanization is also associated with factors that could potentially influence the mental health and possibly the development of CVD, such as increased life stressors, overcrowding, higher level of violence, and less social support. [64] However, beside stress caused by transition from rural area to urban area, other cultural factors interplay with urban dynamics might contribute to the development psychological-related problems. Therefore, understanding how cultural dynamics interact with adaptation to urban life may help in appropriate management of mental disorders in cities. [65] Awareness of the negative impact of urbanization on mental health is needed across the Saudi society.

Healthcare is one of the main focus areas of the Saudi Vision 2030 where the Saudi government has initiated radical changes in the structure and function of its health-care system through its National Transformation Program (NTP) to achieve quality care and effective service delivery. In addition, the government has already recognized the importance of the primary prevention of CVD diseases and has announced recently four major projects, which aim to improve people lifestyle. [66-71] Furthermore, the World Heart Federation (WHF) has undertaken an initiative to develop a series of Roadmaps to reduce premature deaths from CVD by at least 25% by 2025. These Roadmaps can be used as guidance for countries toward developing or updating their national NCD programs for the prevention and control of NCD. [72]

There are few limitations of our study. First, the sampling framework of the PURE-Saudi Study was not nationally representative; hence caution is needed in generalizing our findings to the whole Saudi population. Second, follow-up rates of CVD events and mortality were low, this might be due to short duration of follow-up and the sample size that was relatively small compared to the large population of Saudi Arabia. There are currently ongoing efforts to expand PURE Saudi to a larger population across all areas in the country in order to have meaningful event rates in the follow up. Lastly, we can not exclude the role of genetic predisposition to such high prevalence of CVD risk factors, which could be related to the high consanguinity in the Saudi population. We have reported recently a high prevalence of familial hypercholesterolemia in Saudi Arabia and the rest of the Arabian Gulf countries. [73]

In conclusion, the PURE-Saudi is the first population cohort study in Saudi Arabia. It demonstrated the continued high prevalence of CVD risk factors in the adult Saudi population despite over a decade of several population surveys, and some of these factors were even more prevalent in the rural than the urban population. National awareness programs and multi-faceted health care policy changes are urgently needed to reduce the future burden of CVD risk and mortality.

## **Declarations**

### **Contributions**

KFA and SY developed the study design. KFA, HA and MQA facilitated data gathering. KFA, MAB, THA and SR performed the data analysis and drafted the manuscript. All authors contributed to the interpretation of results and the revision of the manuscript as well as approved the final manuscript.

### **Ethics approval and consent to participate**

King Saud University Ethics Committee granted ethical approval. Participation in the study was voluntary and all eligible subjects who provided written informed consent were enrolled.

### **Consent for publication**

Not applicable.

### **Competing interests**

The authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript. The authors declare that they have no competing interests.

### **Acknowledgement**

Saudi Heart Association, Saudi Gastroenterology Association, Dr Mohammad Alfagih Hospital, The Deanship of Scientific Research at King Saud University, Riyadh, Saudi Arabia (research group no RG-1436-013). We thank Sumathy Rangarajan for the overall coordination of the PURE study, Hani Altaradi for coordinating the PURE-Saudi study, Anhar Ullah for statistical analyses, and the following research

assistants for data collection and quality control: Monther Abdulrahman, Rehab Nouri, Omar Alnobani, Nour Alkamel, and Mohammad Ali. The data are not publicly available as the participants or the ethics committees have not given permission for sharing the data publicly and the study is ongoing.

## Funding

Saudi Heart Association, Saudi Gastroenterology Association, Dr Mohammad Alfagih Hospital, the Deanship of Scientific Research at King Saud University, Riyadh, Saudi Arabia (research group no RG-1436-013).

## References

1. Organization WH. Cardiovascular diseases (CVDs) fact sheet (2015). Geneva: WHO. Retrieved at: <http://www.who.int/mediacentre/factsheets/fs317/en/>. Accessed 20 January 2016.
2. Aljefree N, Ahmed F. Prevalence of cardiovascular disease and associated risk factors among adult population in the Gulf region: a systematic review. *Advances in Public Health*. 2015;2015.
3. Yusuf S, Hawken S, Ôunpuu S, Dans T, Avezum A, Lanan F, et al. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *The lancet*. 2004;364(9438):937-52.
4. O'Donnell MJ, Xavier D, Liu L, Zhang H, Chin SL, Rao-Melacini P, et al. Risk factors for ischaemic and intracerebral haemorrhagic stroke in 22 countries (the INTERSTROKE study): a case-control study. *The Lancet*. 2010;376(9735):112-23.
5. Mabry R, Reeves MM, Eakin EG, Owen N. Evidence of physical activity participation among men and women in the countries of the Gulf Cooperation Council: a review. *Obesity reviews*. 2010;11(6):457-64.
6. Alotaibi A, Perry L, Gholizadeh L, Al-Ganmi A. Incidence and prevalence rates of diabetes mellitus in Saudi Arabia: An overview. *Journal of epidemiology and global health*. 2017;7(4):211-8.
7. Ahmed AM, Hersi A, Mashhoud W, Arafah MR, Abreu PC, Al Rowaily MA, et al. Cardiovascular risk factors burden in Saudi Arabia: The Africa Middle East Cardiovascular Epidemiological (ACE) study. *Journal of the Saudi Heart Association*. 2017;29(4):235-43.
8. Moradi-Lakeh M, El Bcheraoui C, Tuffaha M, Daoud F, Al Saeedi M, Basulaiman M, et al. Tobacco consumption in the Kingdom of Saudi Arabia, 2013: findings from a national survey. *BMC Public Health*. 2015;15(1):611.
9. Memish ZA, El Bcheraoui C, Tuffaha M, Robinson M, Daoud F, Jaber S, et al. Peer reviewed: Obesity and associated factors—Kingdom of Saudi Arabia, 2013. *Preventing chronic disease*. 2014;11.
10. El Bcheraoui C, Memish ZA, Tuffaha M, Daoud F, Robinson M, Jaber S, et al. Hypertension and its associated risk factors in the kingdom of Saudi Arabia, 2013: a national survey. *International journal of hypertension*. 2014;2014.

11. Al-Nozha MM, Al-Mazrou YY, Arafah MR, Al-Maatouq MA, Khalil MZ, Khan NB, et al. Smoking in Saudi Arabia and its relation to coronary artery disease. *Journal of the Saudi Heart Association*. 2009;21(3):169-76.
12. Al-Nozha MM, Al-Hazzaa HM, Arafah MR, Al-Khadra A, Al-Mazrou YY, Al-Maatouq MA, et al. Prevalence of physical activity and inactivity among Saudis aged 30-70 years. A population-based cross-sectional study. *Saudi medical journal*. 2007;28(4):559-68.
13. Al-Nozha MM, Al-Mazrou YY, Al-Maatouq MA, Arafah MR, Khalil MZ, Khan NB, et al. Obesity in Saudi Arabia. *Saudi medical journal*. 2005;26(5):824-9.
14. Al-Hamdan N, Kutbi A, Choudhry A, Nooh R, Shoukri M, Mujib S. WHO stepwise approach to NCD surveillance country-specific standard report Saudi Arabia. World Health Organization. 2005.
15. Al-Nozha MM, Arafah MR, Al-Mazrou YY, Al-Maatouq MA, Khan NB, Khalil MZ, et al. Coronary artery disease in Saudi Arabia. *Saudi medical journal*. 2004;25(9):1165-71.
16. Al-Nuaim AA, Bamgboye EA, Al-Rubeaan KA, Al-Mazrou Y. Overweight and obesity in Saudi Arabian adult population, role of sociodemographic variables. *Journal of community health*. 1997;22(3):211-23.
17. Al-Nuaim AR. High prevalence of metabolic risk factors for cardiovascular diseases among Saudi population, aged 30–64 years. *International journal of cardiology*. 1997;62(3):227-35.
18. Abalkhail BA, Shawky S, Ghabrah TM, Milaat WA. Hypercholesterolemia and 5-year risk of development of coronary heart disease among university and school workers in Jeddah, Saudi Arabia. *Preventive medicine*. 2000;31(4):390-5.
19. Hasim T. Smoking habits of students in College of Applied Medical Science, Saudi Arabia. *Saudi medical journal*. 2000;21(1):76-80.
20. Kalantan KA, Mohamed AG, Al-Taweel AA, Ghani HA. Hypertension among attendants of primary health care centers in Al-Qassim region, Saudi Arabia. *Saudi medical journal*. 2001;22(11):960-3.
21. Al-Rafae S, Al-Hazzaa HM. Physical activity profile of adult males in Riyadh City. *Saudi medical journal*. 2001;22(9):784-9.
22. Al Haddad N, Al Habeeb T, Abdel Gadir M, Al Ghamdy Y, Qureshi N. Smoking patterns among primary health care attendees, Al-Qassim region, Saudi Arabia. 2003.
23. Al-Nozha MM, Al-Maatouq MA, Al-Mazrou YY, Al-Harthi SS. Diabetes mellitus in Saudi Arabia. 2004.
24. Al-Nozha MM, Abdullah M, Arafah MR, Khalil MZ, Khan NB, Al-Mazrou YY, et al. Hypertension in Saudi Arabia. *Saudi medical journal*. 2007;28(1):77-84.
25. Al-Nozha MM, Arafah MR, Al-Maatouq MA, Khalil MZ, Khan NB, Al-Marzouki K, et al. Hyperlipidemia in Saudi Arabia. *Saudi medical journal*. 2008;29(2):282-7.
26. Al-Baghli NA, Al-Ghamdi AJ, Al-Turki KA, El-Zubaier AG, Al-Ameer MM, Al-Baghli FA. Overweight and obesity in the eastern province of Saudi Arabia. *Saudi medical journal*. 2008;29(9):1319-25.
27. Midhet F, Al Mohaimeed AR, Sharaf F. Dietary practices, physical activity and health education in Qassim region of Saudi Arabia. *International journal of health sciences*. 2010;4(1):3.

28. El Bcheraoui C, Basulaiman M, Tuffaha M, Daoud F, Robinson M, Jaber S, et al. Status of the diabetes epidemic in the Kingdom of Saudi Arabia, 2013. *International journal of public health*. 2014;59(6):1011-21.
29. Basulaiman M, El Bcheraoui C, Tuffaha M, Robinson M, Daoud F, Jaber S, et al. Hypercholesterolemia and its associated risk factors—Kingdom of Saudi Arabia, 2013. *Annals of epidemiology*. 2014;24(11):801-8.
30. Al-Rubeaan K, Al-Manaa HA, Khoja TA, Ahmad NA, Al-Sharqawi AH, Siddiqui K, et al. Epidemiology of abnormal glucose metabolism in a country facing its epidemic: SAUDI-DM study. *Journal of diabetes*. 2015;7(5):622-32.
31. Alzeidan R, Rabiee F, Mandil A, Hersi A, Fayed A. Non-communicable disease risk factors among employees and their families of a Saudi university: an epidemiological study. *PloS one*. 2016;11(11):e0165036.
32. Alharthi FS, Alrahimi JS, Alotaibi AA, Alhamdi DA, Ibrahim BM, Badeeb YA. Prevalence of Undiagnosed Cardiovascular Risk Factors in Adults Aged 20-40: A Cross-Sectional Study in 2016 in Jeddah, Saudi Arabia. *Cardiology research*. 2017;8(3):111.
33. Teo K, Chow CK, Vaz M, Rangarajan S, Yusuf S. The Prospective Urban Rural Epidemiology (PURE) study: examining the impact of societal influences on chronic noncommunicable diseases in low-, middle-, and high-income countries. *American heart journal*. 2009;158(1):1-7. e1.
34. Corsi DJ, Subramanian S, Chow CK, McKee M, Chifamba J, Dagenais G, et al. Prospective Urban Rural Epidemiology (PURE) study: baseline characteristics of the household sample and comparative analyses with national data in 17 countries. *American heart journal*. 2013;166(4):636-46. e4.
35. Yusuf S, Rangarajan S, Teo K, Islam S, Li W, Liu L, et al. Cardiovascular risk and events in 17 low-, middle-, and high-income countries. *New England Journal of Medicine*. 2014;371(9):818-27.
36. Miller V, Mente A, Dehghan M, Rangarajan S, Zhang X, Swaminathan S, et al. Fruit, vegetable, and legume intake, and cardiovascular disease and deaths in 18 countries (PURE): a prospective cohort study. *The Lancet*. 2017;390(10107):2037-49.
37. El Bcheraoui C, Basulaiman M, AlMazroa MA, Tuffaha M, Daoud F, Wilson S, et al. Fruit and vegetable consumption among adults in Saudi Arabia, 2013. *Nutrition and Dietary Supplements*. 2015;2015(1):41-9.
38. AlHabib KF, Elasar AA, Alfaleh H, Kashour T, Hersi A, AlBackr H, et al. Clinical features, management, and short-and long-term outcomes of patients with acute decompensated heart failure: phase I results of the HEARTS database. *European journal of heart failure*. 2014;16(4):461-9.
39. AlHabib KF, Hersi A, Alfaleh H, AlNemer K, AlSaif S, Taraben A, et al. Baseline characteristics, management practices, and in-hospital outcomes of patients with acute coronary syndromes: results of the Saudi project for assessment of coronary events (SPACE) registry. *Journal of the Saudi Heart Association*. 2011;23(4):233-9.
40. Alhabib KF, Kinsara AJ, Alghamdi S, Al-Murayeh M, Hussein GA, AlSaif S, et al. The first survey of the Saudi Acute Myocardial Infarction Registry Program: Main results and long-term outcomes (STARS-1

- Program). PloS one. 2019;14(5):e0216551.
41. Roper NA, Bilous RW, Kelly WF, Unwin NC, Connolly VM. Cause-specific mortality in a population with diabetes: South Tees Diabetes Mortality Study. *Diabetes care*. 2002;25(1):43-8.
  42. Cho N, Kirigia J, Mbanya J, Ogurstonova K, Guariguata L, Rathmann W. IDF diabetes atlas-eighth. The International Diabetes Federation; 2017.
  43. Dagenais GR, Gerstein HC, Zhang X, McQueen M, Lear S, Lopez-Jaramillo P, et al. Variations in diabetes prevalence in low-, middle-, and high-income countries: results from the prospective urban and rural epidemiological study. *Diabetes care*. 2016;39(5):780-7.
  44. Chow CK, Teo KK, Rangarajan S, Islam S, Gupta R, Avezum A, et al. Prevalence, awareness, treatment, and control of hypertension in rural and urban communities in high-, middle-, and low-income countries. *Jama*. 2013;310(9):959-68.
  45. Yusufali AM, Khatib R, Islam S, Alhabib KF, Bahonar A, Swidan HM, et al. Prevalence, awareness, treatment and control of hypertension in four Middle East countries. *Journal of hypertension*. 2017;35(7):1457-64.
  46. Attaei MW, Khatib R, McKee M, Lear S, Dagenais G, Igumbor EU, et al. Availability and affordability of blood pressure-lowering medicines and the effect on blood pressure control in high-income, middle-income, and low-income countries: an analysis of the PURE study data. *The Lancet Public Health*. 2017;2(9):e411-e9.
  47. Al-Saleem SA, Al-Shahrani A, Al-Khalidi YM. Hypertension care in Aseer region, Saudi Arabia: Barriers and solutions. *Saudi Journal of Kidney Diseases and Transplantation*. 2014;25(6):1328.
  48. Khatib R, Schwalm J-D, Yusuf S, Haynes RB, McKee M, Khan M, et al. Patient and healthcare provider barriers to hypertension awareness, treatment and follow up: a systematic review and meta-analysis of qualitative and quantitative studies. *PloS one*. 2014;9(1):e84238.
  49. El Bcheraoui C, Tuffaha M, Daoud F, Kravitz H, AlMazroa MA, Al Saeedi M, et al. Access and barriers to healthcare in the Kingdom of Saudi Arabia, 2013: findings from a national multistage survey. *BMJ open*. 2015;5(6):e007801.
  50. Al-Gelban KS, Khan MY, Al-Khalidi YM, Mahfouz AA, Abdelmoneim I, Daffalla A, et al. Adherence of primary health care physicians to hypertension management guidelines in the Aseer region of Saudi Arabia. *Saudi Journal of Kidney Diseases and Transplantation*. 2011;22(5):941.
  51. Siddiqui MA, Khan MF, Carline TE. Gender differences in living with diabetes mellitus. *Materia socio-medica*. 2013;25(2):140.
  52. Albargawi M, Snethen J, Gannass AA, Kelber S. Perception of persons with type 2 diabetes mellitus in Saudi Arabia. *International journal of nursing sciences*. 2016;3(1):39-44.
  53. Kulshreshtha A, Goyal A, Dabhadkar K, Veledar E, Vaccarino V. Urban-rural differences in coronary heart disease mortality in the United States: 1999–2009. *Public health reports*. 2014;129(1):19-29.
  54. Popkin BM. Global nutrition dynamics: the world is shifting rapidly toward a diet linked with noncommunicable diseases–. *The American journal of clinical nutrition*. 2006;84(2):289-98.

55. Ng SW, Popkin BM. Time use and physical activity: a shift away from movement across the globe. *Obesity reviews*. 2012;13(8):659-80.
56. Ng SW, Norton EC, Popkin BM. Why have physical activity levels declined among Chinese adults? Findings from the 1991–2006 China Health and Nutrition Surveys. *Social science & medicine*. 2009;68(7):1305-14.
57. Collaboration NRF. Rising rural body-mass index is the main driver of the global obesity epidemic in adults. *Nature*. 2019;569(7755):260.
58. Ettarh R, Van de Vijver S, Oti S, Kyobutungi C. Peer reviewed: overweight, obesity, and perception of body image among slum residents in Nairobi, Kenya, 2008–2009. *Preventing chronic disease*. 2013;10.
59. Almalki M, FitzGerald G, Clark M. Health care system in Saudi Arabia: an overview. 2011.
60. Ng N, Johnson O, Lindahl B, Norberg M. A reversal of decreasing trends in population cholesterol levels in Västerbotten County, Sweden. *Global health action*. 2012;5(1):10367.
61. Rosengren A, Smyth A, Rangarajan S, Ramasundarahettige C, Bangdiwala SI, AlHabib KF, et al. Socioeconomic status and risk of cardiovascular disease in 20 low-income, middle-income, and high-income countries: the Prospective Urban Rural Epidemiologic (PURE) study. *The Lancet Global Health*. 2019.
62. Lindroth M, Lundqvist R, Lilja M, Eliasson M. Cardiovascular risk factors differ between rural and urban Sweden: the 2009 Northern Sweden MONICA cohort. *BMC Public Health*. 2014;14(1):825.
63. Afshin A, Sur PJ, Fay KA, Cornaby L, Ferrara G, Salama JS, et al. Health effects of dietary risks in 195 countries, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *The Lancet*. 2019.
64. Srivastava K. Urbanization and mental health. *Industrial psychiatry journal*. 2009;18(2):75.
65. Caracci G, Mezzich JE. Culture and urban mental health. *Psychiatric Clinics of North America*. 2001;24(3):581-93.
66. Arabia. TgoS. National transformation program. . 2016.
67. Alharbi MF. An analysis of the Saudi health-care system's readiness to change in the context of the Saudi National Health-care Plan in Vision 2030. *International journal of health sciences*. 2018;12(3):83.
68. King Salman Park. The King Salman Park Project 2019, March Available from: <http://riyadhksp.sa/en/>.
69. Green Riyadh. The Green Riyadh Project 2019, March Available from: <http://www.riyadhgreen.sa/en/>.
70. Sport Boulevard. The Sport Boulevard Project 2019, March. Available from: <http://www.riyadhalmasar.sa/en/>.
71. Riyadh Art. The Riyadh Art Project 2019, March Available from: <http://www.riyadhart.sa/en/>.
72. WHF. World Heart Federation Roadmaps Geneva. Available at: [http:// www.world-heart-federation.org/what-we-do/whf-roadmaps/](http://www.world-heart-federation.org/what-we-do/whf-roadmaps/). Accessed May 25, 2016. .

73. Al-Rasadi K, Alhabib KF, Al-Allaf F, Al-Waili K, Al-Zakwani I, AlSarraf A, et al. The Gulf Familial Hypercholesterolemia Registry (Gulf FH): Design, Rationale and Preliminary Results. *Current vascular pharmacology*. 2019.

## Tables

Table 1. Characteristics and the prevalence of cardiovascular disease risk factors; in the PURE-Saudi study.

Characteristics	Overall	Men	Women	p-value
N (%)	2047	1165 (56.91)	882 (43.09)	
Age (year), mean $\pm$ SD	46.50 $\pm$ 9.12	47.55 $\pm$ 9.40	45.10 $\pm$ 8.55	< 0.001
Low educational level, n (%) †	646 (31.56)	235 (20.17)	411 (46.60)	< 0.001
Smoking status, n (%) ‡				
Current smoker	249 (12.16)	245 (21.03)	4 (0.45)	< 0.001
Former smoker	217 (10.60)	209 (17.94)	8 (0.91)	< 0.001
Unhealthful diet, n (%) §	702 (34.39)	397 (56.55)	305 (43.45)	0.827
Low physical activity, n (%) ¶	1415 (69.40)	805 (69.34)	610 (69.48)	0.946
Hypertension, n (%) ♦	620 (30.29)	382 (32.79)	238 (26.98)	0.005
Awareness among known hypertensive patients, n (%)	379 (61.13)	233 (60.99)	146 (61.34)	0.931
Treated hypertension among known hypertensive patients, n (%)	365 (58.87)	220 (57.59)	145 (60.92)	0.412
Controlled (ie SBP < 140 mm Hg) among those known to be hypertensive patients, n (%)	190 (30.65)	108 (28.27)	82 (34.45)	0.104
Treated hypertension and SBP $\geq$ 140 mmHg and/or DBP $\geq$ 90mmHg, n (%)	175 (47.95)	112 (50.91)	63 (43.45)	0.163
Treated hypertension and SBP > 120 mmHg and/or DBP > 80 mmHg, n (%)	318 (87.12)	198 (90)	120 (82.76)	0.043
Treated hypertension and SBP > 130 mmHg and/or DBP > 80 mmHg, n (%)	272 (76.19)	173 (79.36)	99 (71.22)	0.078
Diabetes, n (%) **	516 (25.21)	328 (28.15)	188 (21.32)	< 0.001
Among diabetic patients				0.093
On insulin alone?	14 (2.71)	5 (1.52)	9 (4.79)	
On OHA alone?	313 (60.66)	200 (60.98)	113 (60.11)	
on both	34 (6.59)	19 (5.79)	15 (7.98)	
not on drug treatment	155 (30.04)	104 (31.71)	51 (27.13)	
Dyslipidemia				
Total Cholesterol > 5.2 mmol/L and LDL > 3.5 mmol/L, n (%) ¶	569 (32.1)	234 (31.2)	335 (32.68)	0.508
On statins medication, n (%)	5 (0.88)	3 (0.90)	2 (0.85)	0.959

Characteristics	Overall	Men	Women	p-value
BMI, median (25th-75th, IQR)	29.93 (26.68,33.90)	28.97 (26.06,32.65)	31.36 (27.72,35.26)	< 0.001
BMI, n (%) ††				< 0.001
< 25	310 (15.15)	205 (17.60)	105 (11.92)	
25-30	722 (35.29)	462 (39.66)	260 (29.51)	
30-35	613(29.96)	463(39.78)	286(32.43)	
> 35	399 (19.50)	169 (14.52)	230 (26.08)	
Abdominal obesity, n (%)				
Waist circumference > 102 cm (men) or > 88 cm (women)	1005 (49.1)	381 (32.7)	624 (70.75)	< 0.001
Waist circumference > 90 cm (men) or > 85 cm (women)	1521 (74.3)	835 (71.67)	686 (77.78)	0.002
Current alcohol use, n (%)	24 (1.17)	24 (2.06)	0 (0.00)	< 0.001
Self-report of being sad or "blue", n (%)	315 (15.40)	115 (9.88)	200 (22.68)	< 0.001
General feeling of stress, n (%)				
Several periods	339 (16.93)	143 (12.40)	196 (23.09)	< 0.001
Permanent	136 (6.79)	52 (4.51)	84 (9.89)	< 0.001
History of ischemic heart disease (angina, MI or any coronary revascularization), n (%)	51 (2.49)	37 (3.18)	14 (1.59)	0.022
History of stroke, n (%)	20 (0.98)	12 (1.03)	8 (0.91)	0.779
History of heart failure, n (%)	13 (0.64)	5 (0.43)	8 (0.91)	0.178
INTERHEART Risk Score, median (25th-75th, IQR)	11.00 (8.00,16.00)	13.00 (9.00,18.00)	10.00 (6.00,14.00)	< 0.001

† Low educational level was defined as no education, primary education only, or unknown educational level.

‡ Participants who were not current or former smokers are those who had never smoked (data not shown).

§ Diet quality was determined on the basis of the Alternative Healthy Eating Index, which ranges from 6 to 70, with higher scores indicating more healthful eating. An unhealthy diet was considered to be a score of less than 31. ¶ Low physical activity was defined as 600 or fewer metabolic equivalents per minute per week. ♦ Participants were considered to have hypertension if they had a blood pressure greater than 140/90 mm Hg or if they reported a history of hypertension.\*\* Participants were considered to have diabetes if they reported having diabetes or if they had a fasting glucose level higher than 7.0 mmol per liter (126 mg per deciliter).

□ Participants were considered to have dyslipidemia if they reported having

dyslipidemia or if they had a total cholesterol level higher than 5.2 mmol/l (201 mg/dl); data on lipid profile level were available for 1775 participants.†† The body-mass index (BMI) is the weight in kilograms divided by the square of the height in meters.

‡‡ The INTERHEART Risk Score is a validated score for quantifying risk-factor burden without the use of laboratory testing. Scores range from 0 to 48, with higher scores indicating greater risk-factor burden.

Table 2. Prevalence of cardiovascular disease risk factors stratified by age groups in the PURE-Saudi study.

Characteristics	35-49 yrs	50-59 yrs	60-70 yrs	p-value
N (%)	1352 (66.05)	469 (22.91)	226 (11.04)	
Low educational level, n (%)	293 (21.67)	211 (44.99)	142 (62.83)	< 0.001
Smoking status, n (%)				
Current smoker	181 (13.39)	48 (10.23)	20 (8.85)	0.054
Former smoker	124 (9.17)	60 (12.79)	33 (14.60)	0.011
Unhealthful diet, n (%)	524 (38.90)	118 (25.21)	60 (26.55)	< 0.001
Low physical activity, n (%)	899 (66.84)	346 (73.93)	170 (75.22)	0.002
Hypertension, n (%)	246 (18.20)	227 (48.40)	147 (65.04)	< 0.001
Aware among known hypertensive patients, n (%)	112 (45.53)	159 (70.04)	108 (73.47)	< 0.001
Treated hypertension among known hypertensive patients, n (%)	100 (40.65)	157 (69.16)	108 (73.47)	< 0.001
Controlled among known hypertensive patients, n (%)	49 (19.92)	89 (39.21)	52 (35.37)	< 0.001
Treated hypertension and SBP $\geq$ 140 mmHg and/or DBP $\geq$ 90 mmHg, n (%)	51 (51.00)	68 (43.31)	56 (51.85)	0.304
Treated hypertension and SBP > 120mmHg and/or DBP > 80mmHg, n (%)	86 (86.00)	135 (85.99)	97 (89.81)	0.609
Treated hypertension and SBP > 130 mmHg and/or DBP > 80 mmHg, n (%)	80(80.81)	115(73.72)	77(75.49)	0.423
Diabetes, n (%)	175 (12.94)	211 (44.99)	130 (57.52)	< 0.001
Among diabetic patients				0.831
On insulin alone?	5 (2.86)	6 (2.84)	3 (2.31)	
On OHA alone?	105 (60.00)	130 (61.61)	78 (60.00)	
on both	12 (6.86)	10 (4.74)	12 (9.23)	
not on drug Rx	53 (30.29)	65 (30.81)	37 (28.46)	
Dyslipidemia				
CHOL > 5.2 mmol/L and LDL > 3.5 mmol/L, n (%)	380 (32.45%)	124 (29.95%)	65 (34.21%)	0.514
On statins medication, n (%)	2 (0.53)	3(2.42)	0 (0.00)	0.172
BMI, median (25th-75th, IQR)	29.78 (26.59,33.92)	30.65 (27.60,34.29)	29.00 (25.21,32.96)	< 0.001


Characteristics	35-49 yrs	50-59 yrs	60-70 yrs	p-value
BMI, n (%)				< 0.001
< 25	211 (15.62)	46 (9.81)	53 (23.45)	
25-30	488 (36.12)	161 (34.33)	73 (32.30)	
30- 35	388 (28.72)	160 (34.12)	65 (28.76)	
> 35	262(19.39)	102(21.75)	35(15.49)	
Abdominal obesity, n (%)				
Waist circumference > 102 cm (men) or > 88 cm (women)	628 (46.45)	262 (55.86)	115 (50.88)	0.002
Waist circumference > 90 cm (men) or > 85 cm (women)	960 (71.01)	389 (82.94)	172 (76.11)	< 0.001
Current alcohol use, n (%)	14 (1.04)	7 (1.49)	3 (1.33)	0.632
Self-report of being sad or "blue", n (%)	252 (18.65)	47 (10.02)	16 (7.08)	< 0.001
General feeling of stress, n (%)				
Several periods	249 (18.64)	74 (16.30)	16 (7.55)	< 0.001
Permanent	121 (9.06)	11 (2.42)	4 (1.89)	< 0.001
History of ischemic heart disease (angina or MI), n (%)	15 (1.11)	19 (4.05)	17 (7.52)	< 0.001
History of stroke, n (%)	6 (0.44)	5 (1.07)	9 (3.98)	< 0.001
History of heart failure, n (%)	2 (0.15)	7 (1.49)	4 (1.77)	<0.001
INTERHEART Risk Score, median (25th-75th, IQR)	10.00 (7.00,14.00)	14.00 (10.00,19.00)	16.00 (12.00,21.00)	< 0.001

Table 3. Prevalence of cardiovascular disease risk factors stratified by place of residence in the PURE-Saudi study.

Characteristics	Urban	Rural	p-value
N (%)	1545 (75.48)	502 (24.52)	
Age (year), mean $\pm$ SD	46.31 $\pm$ 8.88	47.07 $\pm$ 9.81	0.124
Low educational level, n (%)	413 (26.73)	233 (46.41)	< 0.001
Smoking status, n (%)			
Current smoker	198 (12.82)	51 (10.16)	0.114
Former smoker	175 (11.33)	42 (8.37)	0.061
Unhealthful diet, n (%)	556 (36.01)	146 (29.38)	0.007
Low physical activity, n (%)	1061 (68.72)	354 (71.52)	0.240
Hypertension, n (%)	442 (28.61)	178 (35.46)	0.004
Aware among known hypertensive patients, n (%)	275 (62.22)	104 (58.43)	0.381
Treated hypertension among known hypertensive patients, n (%)	266 (60.18)	99 (55.62)	0.296
Controlled among known hypertensive patients, n (%)	144 (32.58)	46 (25.84)	0.100
Treated hypertension and SBP $\geq$ 140mmgH and/or DBP $\geq$ 90mmgH, n (%)	122 (45.86)	53 (53.54)	0.192
Treated hypertension and SBP > 120mmgH and/or DBP > 80mmgH, n (%)	227 (85.34)	91 (91.92)	0.095
Treated hypertension and SBP > 130 mmgH and/or DBP > 80mmgH, n (%)	191(73.46)	81(83.51)	0.047
Diabetes, n (%)	360 (23.30)	156 (31.08)	< 0.001
Among diabetic patients			< 0.001
on insulin	11 (3.06)	3 (1.92)	
on OHA	249 (69.17)	64 (41.03)	
on both	31 (8.61)	3 (1.92)	
not on Rx	69 (19.17)	86 (55.13)	
Dyslipidemia			
CHOL> 5.2 mmol/L and LDL > 3.5 mmol/L, n (%)	439 (31.49%)	130 (34.12%)	0.330
On statins medication, n (%)	5 (1.14)	0 (0.00)	0.590
BMI, median (25th-75th, IQR)	29.66 (26.33,33.59)	30.84 (27.20,34.92)	< 0.001


Characteristics	Urban	Rural	p-value
BMI, n (%)			< 0.001
< 25	245 (15.87)	65 (12.95)	
25-30	569 (36.85)	153 (30.48)	
30- 35	454 (29.39)	159 (31.74)	
> 35	275(17.80)	124(24.75)	
Abdominal obesity, n (%)			
Waist circumference > 102 cm (men) or > 88 cm (women)	754 (48.80)	251 (50.00)	0.641
Waist circumference > 90 cm (men) or > 85 cm (women)	1156 (74.82)	365 (72.71)	0.347
Current alcohol use, n (%)	18 (1.17)	6 (1.20)	0.956
Self-report of being sad or "blue", n (%)	272 (17.62)	43 (8.57)	< 0.001
General feeling of stress, n (%)			
Several periods	296 (19.65)	43 (8.67)	< 0.001
Permanent	121 (8.03)	15 (3.02)	< 0.001
History of ischemic heart disease (angina or MI), n (%)	36 (2.33)	15 (2.99)	0.411
History of stroke, n (%)	12 (0.78)	8 (1.59)	0.118
History of heart failure, n (%)	8 (0.52)	5 (1.00)	0.327
INTERHEART Risk Score, median (25th-75th, IQR)	11.00 (8.00,16.00)	12.00 (7.00,17.00)	0.833

## Figures




**Figure 1**

Figure A. Cardiovascular risk factors stratified by age groups.



**Figure 2**

Figure B. Prevalence of cardiovascular risk factors in the overall cohort and stratified by sex



**Figure 3**

Figure C. Cardiovascular risk factors stratified by residence place.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [additionaltable1.docx](#)
- [additionaltable4.docx](#)
- [additionaltable3.docx](#)
- [additionaltable2.docx](#)