Nalbuphine Alleviates Inflammation by Down-regulating NF- κ B in an Acute Inflammatory Visceral Pain Rat Model

Di-jiao Ruan

Stomatological Hospital of Chongqing Medical University

Yuan-yuan Wang

Stomatological Hospital of Chongqing Medical University

Si-si Li

Stomatological Hospital of Chongqing Medical University

Chao Zhang

Stomatological Hospital of Chongqing Medical University

Wen-wen Zheng

Stomatological Hospital of Chongqing Medical University

Cong Yu (✉ 500158@hospital.cqmu.edu.cn)

Stomatological Hospital of Chongqing Medical University

Research Article

Keywords: Nalbuphine, NF- κ B, Inflammation, Visceral pain

Posted Date: January 19th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1169482/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Introduction: Nalbuphine can relief patients' inflammation response after surgery compared to other opioid drugs. However, its molecular mechanism has not been clear. Activation of NF- κ B signaling pathway under oxidative stress and inflammation can maintain pain escalation.

Methods: We firstly investigated the effect of nalbuphine on writhing test and mechanical allodynia using a rat model of inflammatory visceral pain (acetic acid (AA) administrated). Cytokines (including tumor necrosis factor (TNF)- α , Interleukin (IL)-1 β , IL-2, and IL-6 in plasma were tested with ELISA technology. Expression levels of TNF- α , I κ B α and p-NF- κ B p65 at the spinal cord (L3-5) were measured by western blot or RT-qPCR.

Results: We found that the values of paw withdrawal threshold (PWT) were reduced and numbers of writhes were increased in the rats of group model, up-regulated IL-1 β , IL-2, IL-6 and TNF- α in both plasma and spinal cord as well as increased protein of p-NF- κ B p65 and mRNA of NF- κ B p65 expression in spinal cord. Subcutaneously injection of nalbuphine (10 μ g/kg) or PDTC (NF- κ B inhibitor) attenuated acetic acid-induced inflammatory pain, and this was associated with reversal of up-regulated IL-1 β , IL-2, IL-6 and TNF- α in both plasma and spinal cord. Furthermore, acetic acid increased p-NF- κ B and TNF- α protein levels in the white matter of the spinal cord, which was attenuated by nalbuphine. These results suggest that nalbuphine can significantly ameliorate inflammatory pain via modulating the expression of NF- κ B p65 as well as inflammation factors level in spinal cord.

Conclusion: In conclusion, nalbuphine inhibits inflammation through down-regulating NF- κ B pathway at spinal cord in a rat model of inflammatory visceral pain.

Introduction

Acute or chronic visceral pain is one of the most happening diseases in our life. Approximately 10% of chronic visceral pain is transmitted from acute postoperative pain that is difficult to control[1]. Opioid drugs play an important role in controlling acute pain during most surgical procedures (e.g., gallbladder, gastrointestinal, and gynecological surgery)[2–5].

Compared with μ -opioid receptor agonists such as remifentanyl, sufentanyl, and fentanyl, nalbuphine is a kappa-opioid receptor agonist and μ -opioid receptor antagonist with fewer side effects and a higher safety profile [6, 7]. Nalbuphine is structurally similar to naloxone, mainly used in clinical practice for intraoperative and postoperative analgesia [8, 9]. Some researchers reported that nalbuphine is primarily acted on kappa receptors to produce analgesia, which can be reversed by naloxone dose-dependently; when acting on μ -receptors, respiratory depression is less than morphine due to the capping effect [10, 11]. There are some studies reported that kappa-opioid drugs are involved in regulating NF- κ B signaling in rats to exert a protective effect after myocardial ischemic and suppressed breast cancer stemlike properties [12, 13]. In addition, there are some studies have shown that nalbuphine can inhibit the inflammatory response in patients to help them recover and reduce the length of hospital stay

perioperatively. It can reduce the release of proinflammatory cytokines interleukin IL-6, tumor necrosis factor TNF- α , and IL-1 β in plasma [14–18]. Moreover, evidence showed that inflammation plays a vital role in visceral pain, it also has a significant impact on short-and long-term postoperative clinical outcomes[19].

Nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) is reported to play a critical role in inflammatory degenerative processes [20, 21]. NF- κ B-mediated inflammatory responses can lead to progressive extracellular matrix damage [22]. Pain is often accompanied by the activation of various inflammatory cytokines such as IL-1 β , IL-2, TNF- α , and NF- κ B pathways. In addition, many studies have reported on the role of NF- κ B and its downstream pro-inflammatory cytokines in mechanisms underlying inflammatory and neuropathic pain processes [23, 24]. NF- κ B is mostly present in the form of a p65-p50 dimer and is associated with inhibitor I κ B binds in an inactive state[25]. Studies have shown that the expression of NF- κ B p65 is significantly enhanced in various inflammatory models[26–29].

In the preliminary study, we have found that nalbuphine can reduce plasma levels of pro-inflammatory factors such as IL-6 and TNF- α in patients 24 h and 48 h after orthognathic surgery compared with sufentanil[17]. Simultaneously, postoperative nausea, vomiting, and other postoperative complications or anesthesia-related adverse reactions were lower in the nalbuphine group. However, the targets through which nalbuphine exerts its anti-inflammatory and analgesic effects have not been determined.

NF- κ B as a key factor in the inflammatory response, there was no report that whether nalbuphine can inhibit the NF- κ B signaling pathway in visceral pain rats. Thus, we aimed to test the effect of nalbuphine in a model of inflammatory visceral pain to further clarify the relationship between NF- κ B and nalbuphine at the spinal cord in the rat.

Materials And Methods

Animals and Ethics

All methods were carried out in accordance with relevant guidelines and regulations in Ethic approval and consent to participation. All procedures were carried in accordance with the ARRIVE (Animal Research: Reporting In Vivo Experiments) guidelines. The Ethics Committee of the College of Stomatology, Chongqing Medical University approved this study (Project No. CQHS-REC-2021 (LSNo.011)). Forty-eight SPF male Sprague-Dawley rats (8 weeks; 230-260 g) were purchased from Hunan SJA Laboratory Animals (Source, SCXK-2019-0004; Number of qualitative qualification, 430727200101046971; Hunan, China). They were housed in the Animal Care Facility within the Chongqing Key Laboratory of Oral Diseases and Biomedicine (Number permit, SYXK-2019-0004) under a 12h light-dark cycle (lights-on at 7:50 am; lights-off at 7:50 pm). Rats had free access to water and food in a room at 22-25°C and 50-60% humidity. All rats were fasted but not water-deprived 12h before experimentation. Rats were divided randomly into four groups: control, AA, AA+ nalbuphine, and AA+PDTC. Efforts were made to minimize the number of animals used and the stress they experienced.

Drugs and agents

Nalbuphine hydrochloride was purchased from Yichang Human Well Pharmaceuticals (State Food and Drug Administration approval number: 989RG; batch number: H20054171; Hunan, China). Antibodies were obtained from primary antibodies against the anti-GAPDH (2118S, CST, USA), anti-TNF- α (GB11188S, ervicebio), anti-IL-6 (ab259341, Abcam, USA), anti-IL-1 β (12703S, CST, USA), anti-I κ B α (4812s, CST, USA), anti-phosphorylated (p)-NF- κ B p65 (ser536) (310013, Zen Bioscience). pyrrolidine dithiocarbamate (PDTC, S1808, Beyotime).

Inflammatory visceral pain

The effects of nalbuphine on acute inflammatory visceral pain were assessed after pain induction using acetic acid, modified from described previously[30]. Briefly, it was induced with an intraperitoneal injection of 1ml/100g of 1% acetic acid solution into the rats' peritoneal under isoflurane anesthesia. Three minutes before it, 500 μ L of physiologic (0.9%) saline or nalbuphine (1mg/100g body weight) was injected into the tail vein. In control group rats, saline was injected into the peritoneum and tail vein. Model and nalbuphine group rats were treated with injection of acetic acid into the peritoneum and saline or nalbuphine into the tail vein respectively. Rats in the PDTC group were injected intraperitoneally with PDTC and 30 minutes later with acetic acid [31].

Behavioral measurements

The writhing test was adapted from the description made by Schmauss and colleagues[32]. Briefly, a positive reaction was characterized by: dorsal-claw flexion to the left or right; body extension and abdomen flat; a contraction in the abdomen followed by extension of the hind limbs. The number of positive responses was recorded every 5 min for 30 min and recorded as the writhing times.

Paw withdrawal threshold (PWT) was measured using a set of Von Frey monofilaments (VFMs; Aesthesio®; DanMic Global, LLC, American). The target force of VFMs was 0.6, 1, 2, 4, 6, 10, 15, and 26g using the up-down method described by Dixon and Mood, and Bradman[33, 34]. Rats were placed in a plastic cover on a wire mesh floor (5mm $\times$ 5mm) and allowed to habituate for 30min before testing. Starting from a target force of 4g, a series of VFMs was tested vertical to the middle (not the pad) of the right lateral plantar surface of the paw for 7s per VFM. A low-level VFM was used if the agile withdrawal of a paw was observed; otherwise, a high-level VFM was applied. When the first positive reaction was followed by a negative reaction or vice versa, four additional stimuli were administered. Each rat received only one treatment, and researchers were blinded to the treatment applied. Experimental data were converted in the way described by Chaplan and colleagues[35].

Enzyme-linked immunosorbent assay (ELISA)

TNF- α , IL-1 β , IL-2, IL-6 in the plasma of rats were detected by Enzyme-linked immunosorbent assay according to manufacturer instructions (Sinovac Biotech. Shanghai, China). Briefly, plasma was extracted from the supernatant of whole blood collected from rat hearts after centrifugation for 15 min at 2.415($\times$

g) at 4°C, and stored at -80°C. Then, all the supernatants were analyzed for quantification of TNF- α , IL-1 β , IL-2, and IL-6 using the ELISA kits.

Western blotting

Total protein in spinal cord (L3-5) was analyzed by western blot. Briefly, the spinal cord was homogenized in RIPA lysis buffer containing phenylmethylsulfonyl fluoride and phosphatase inhibitor cocktail A (50 $\times$) according to specifications using a Polytron™ homogenizer five times, 10s each, 1min apart. After the protein had been lysed fully on ice for 20min, the protein was centrifugated for 5min at 12000($\times$ g) at 4°C and calculated the concentration using a bicinchoninic acid kit (P0012s, Beyotime) following instructions. After mixing with sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) sample loading buffer (5 $\times$; P0015, Beyotime), protein samples were degenerated at 100°C for 10 min by a Mixer-heat/Cool mixer (Thermo Scientific, TCS10, Waltham, MA, USA) and stored at -80°C. Proteins were separated by SDS-PAGE and then transferred to polyvinylidene fluoride (PVDF) membranes (Millipore, Bedford, MA, USA) with an aperture of 0.45mm. PVDF membranes were blocked with 5% non-fat dry milk containing 0.05% Tween 20-TBS for 2 hours at room temperature, and then incubated with primary antibodies against the anti-GAPDH(1:1000), anti-TNF- α (1:200), anti-IL-6 (1:1000), anti-IL-1 β (1:1000), anti-I κ B α (1:1000), anti-phosphorylated (p)-NF- κ B p65 (ser536) (1:1000) overnight at 4°C. Then, immunoblots were incubated with the horseradish peroxidase (HRP) conjugated secondary antibody appropriated to its primary antibody for 2 hours at room temperature and observed by an ECL detection kit (P0018AS, Beyotime). Imaging software (ChemiDoc™ Touch; Bio-Rad Laboratories, Hercules, CA, USA) was used to scan each immunoblot and record its image.

Real-time reverse transcription-quantitative polymerase chain reaction (RT-qPCR)

Spinal cord (L3-5) separated from rats in different groups under isoflurane anesthesia and collected in a frozen storage tube without RNase or DNase (1.8 mL; Corning). This tissue was stored in liquid nitrogen until further processing. RNA was isolated using TRIzol® Reagent (R0016, Beyotime) and purified with DNase I according to the instructions. Complementary-DNA was synthesized using the Reverse Transcription cDNA Archive kit (Bio-Rad, T100, USA). Expression of IL-1 β , IL-6, NF- κ B p65, and TNF- α was measured by the CFX96™ Real-Time System (Bio-Rad, USA) and TaqMan™ Gene Expression kit (RR047A, TaKaRa Biotechnology). A two-step PCR protocol was used to amplify and each step was based on instructions provided by the manufacturers. Data were collected and the quantification (Cq) was utilized to calculate the gene expression of IL-1 β , IL-6, NF- κ B p65, and TNF- α in all groups using GAPDH expression to normalize (2^{- $\Delta\Delta$ CT} method). Primer sequences are shown in Table 1.

Table 1
Primer sequences used for RT-qPCR.

Gene name	Sense primer (5'-3')	Anti-sense primer (5'-3')
IL-1 β	TCCCAAACAATACCCAAAGAAG	ACTATGTCCCGACCATTGCTG
IL-6	GACAAAGCCAGAGTCATTGAGAG	GGATGGTCTTGGTCCTTAGCC
NF-KB p65	ACCTGGAGCAAGCCATTAGC	CCGCATTCAAGTCATAGTCCC
TNF- α	CTTCTCATTCTGCTCGTGG	CCGCTTGGTGGTTTGCTAC

Immunohistochemical (IHC) analyses

IHC experiment was undertaken following behavioral experiments. Under isoflurane anesthesia, the pinal cord was harvested after pre-fixation performed by injecting 100 mL of 0.9% saline and 200 mL of 4% paraformaldehyde into the heart. After fixation, tissue was placed in 4% paraformaldehyde for 72 hours. Then, specimens were dehydrated, embedded in paraffin, and sliced. Slices were dewaxed, hydrated, antigenic repair, blocked, and then incubated overnight at 4°C with primary antibody. Then, they were incubated for 50 min at room temperature with HRP conjugated secondary antibody followed by DAB colored and microscopic examination (Olympus, SLIDEVIEW™ VS200, Tokyo, Japan). The primary antibodies for Anti-TNF- α (1:100) and anti-p-NF-kB p65 (ser536) (1:100) were applied.

Statistical analyses

All data were expressed as mean $\pm$ standard error of the mean (SEM) and analyzed using SPSS26.0 software. Data were submitted to one-way ANOVA, nonparametric tests, or two-tailed Student's t-test according to the results of testing for normality and homogeneity of variance. The post hoc test (Fisher's least significant difference) was conducted to compare multiple groups only if ANOVA yielded a significant main effect. $P < 0.05$ was considered significant. Graphs were made by GraphPad Prism 8 and grouped by AI or PS.

Results

Nalbuphine intervention made behaviors of acute visceral pain model rats changed

To assess the pain threshold, writhing numbers and PWT were measured respectively. The writhing test showed numbers of writhes in the AA group and AA+nalbuphine group were higher than that in the control group ($p < 0.05$), and the analgesic effect of nalbuphine (10 μ g/kg) in rats led to a significantly diminished number of writhes compared with that in the AA group (model group), started at 10 minutes. (Fig. 1a) ($p < 0.05$). In the mechanical withdraw threshold test, the PWT was decreased significantly in the AA group (3.50 ± 0.73 g) compared with that in the control group (5.29 ± 0.61 g), whereas it was increased in

the AA+ nalbuphine ($11.35 \pm 2.59\text{g}$) and AA+ PDTC group ($10.08 \pm 1.63\text{g}$) (Fig. 1b) ($p < 0.05$). The results showed that nalbuphine and NF- κ B inhibitor PDTC both exerted good analgesic effects in visceral pain.

Nalbuphine suppressed the levels of IL-1 β , IL-2, IL-6, and TNF- α in acute visceral pain rats.

ELISA results showed that AA-treated rats secreted higher levels of IL-1 β , IL-2, IL-6, and TNF- α than control rats in plasma (Fig. 2a-d), whereas nalbuphine inhibited AA-stimulated increase in the above inflammatory factor levels. These results fully proved the anti-inflammatory effect of nalbuphine. PDTC as an inhibitor of NF- κ B also played an anti-inflammatory role in acute visceral pain. The levels of IL-1 β , IL-2, IL-6, and TNF- α in the PDTC group were significantly decreased than in the AA group.

Nalbuphine modulated p-NF- κ B p65 and I κ B α in the spinal cord of visceral pain rats

NF- κ B p65 usually binds to inhibitory I κ B α proteins and remains silent in the cells. When phosphorylated I κ B α becomes degraded, NF- κ B p65 is released and translocated into the nucleus and finally leads to regulation of gene transcription[36, 37]. So, we measured both the levels of p-NF- κ B p65 and I κ B α using western blotting and RT-qPCR. The results showed that AA-treated rats decreased the level of I κ B α and increased the protein expression of p-NF- κ B p65 compared with control group rats (Fig. 3a). In addition, Fig. 3b showed that nalbuphine and PDTC also inhibited an AA-stimulated increase in the mRNA level of NF- κ B p65. Therefore, our study indicated that nalbuphine inhibited inflammation by down-regulating the NF- κ B pathway in the spinal cord.

IL-1 β , TNF- α and IL-6 were down-regulated by nalbuphine and PDTC in the spinal cord

To ascertain whether the inflammatory response in the spinal cord was increased during nalbuphine and PDTC treatment, we measured the expressions of IL-1 β , TNF- α , and IL-6 in the spinal lumbar. Western blotting and RT-qPCR results suggested that acetic acid upregulated the proteins of IL-1 β , TNF- α , IL-6, and their mRNA expressions. Nalbuphine treatment decreased the mRNA and protein levels of IL-1 β , TNF- α , and IL-6 compared with rats in the AA group rats (Fig. 4a-d). This indicated that nalbuphine reduced the expression of inflammatory factors not only in the plasma but also in the spinal cord. The NF- κ B inhibitor PDTC treatment also reduced the levels of IL-1 β , IL-6, and TNF- α in the spinal cord compared to the AA group to exert anti-inflammatory effects.

Nalbuphine decreased p-NF- κ B and TNF- α protein levels in the white matter of the spinal cord

In our study, IHC analyses revealed that acetic acid enhanced p-NF- κ B and TNF- α expression in the AA group, whereas nalbuphine could reverse this change. The result of IHC analyses consisted of western blot and RT-qPCR. So, we further confirmed that nalbuphine exerted its anti-inflammatory effect through

down-regulating NF- κ B. In addition, we found p-NF- κ B and TNF- α mainly distributed in the white matter of the spinal cord in all group rats. (Fig. 5).

Discussion

Visceral pain is a common clinical problem, it can be caused by chemical-physical damage to wounds, as well as by psychological, microbial, and endocrine factors, yet its mechanism is not fully clear [38–40]. The development of visceral pain is regulated by the spinal cord and higher centers of the brain. Visceral pain tends to company physical discomfort and special positions like knee flexion position in humans. Similarly, the number of writhes in a certain period reflects the degree of pain in rats and is often used in pain-model experiments[41–43]. Acute visceral pain like acute appendicitis tends to lead to superficial or deep hyperalgesia or allodynia[44]. In this research, the analgesic drug nalbuphine significantly decreased the writhing numbers of rats caused by acetic acid and increased the PWT. These experimental results are consistent with the data in other studies[41, 42].

TNF- α , IL-6, IL-1 β , and IL-2 are involved in many inflammatory visceral pain diseases like visceral pain after surgery, endometriosis [40, 41]. Nalbuphine is applied in some clinical researches to play valid anti-inflammatory and analgesic effects [42, 43]. Excepting in the clinical studies, in ML's study, expressions of pro-inflammatory cytokines (IL-6, IL-1 β , and TNF- α) in the spinal cord tissues were increased in visceral pain rats and inhibited by dexmedetomidine [44]. These studies indicate that inflammatory factors are widely expressed in visceral pain diseases. In our study, TNF- α , IL-6, IL-1 β , and IL-2 significantly enhanced in the visceral pain model in rats induced by acetic acid, and nalbuphine decreased above inflammation factors not only in plasma but also in the spinal cord. It reveals that nalbuphine exerts its anti-inflammatory and analgesic effects through neurohumoral regulation in the central spinal cord system. These results were consistent with our clinic research before, nalbuphine inhibited inflammation after orthognathic surgery [17].

NF- κ B participates in the regulation of cell growth, proliferation, inflammation, immunity, energy metabolism, tumor metastasis, and other aspects in organs[13, 45–51]. Usually, NF- κ B couples with I κ B α in an inactive state in cells, and when the synthesis of I κ B α is inhibited or decomposed, NF- κ B activation increases [52]. I κ B α as a key inhibitory protein of NF- κ B inhibits the activation of NF- κ B (p65) and then reduces expression of downstream pro-inflammatory factors like TNF- α [53, 54]. In the present study, visceral pain caused by acetic acid (i.p) could increase the expression of NF- κ B (p65) and TNF- α , whereas decreased the level of I κ B α in the spinal cord, especially in the macrophage/microglia at white matter of the spinal cord. The white matter is composed of oligodendrocyte precursor cells (OPCs), oligodendrocytes, astrocytes, and microglia [55]. Macrophage/microglia are extensively involved in the process of inflammatory response in the central nervous system[56, 57]. Our results also suggested that spinal macrophage/microglia are also involved in the development of inflammatory visceral pain. Blocking the activation of NF- κ B with NF- κ B inhibitor (PDTC) remarkably suppressed the AA-induced expression of TNF- α . So, we concluded that increased NF- κ B expression in the spinal white matter is a key central regulating mechanism in visceral pain. Analgesic drug nalbuphine significantly inhibited NF-

κB (p65) and TNF-α and increased the IκBα protein expression in the spinal cord, also mRNA and IHC tests showed consistent results in NF-κB (p65) and TNF-α. The results suggested that nalbuphine, as a κ-receptor agonist exerted its anti-inflammatory effects by inhibiting the NF-κB pathway.

So far, our research has explored from clinical phenomenon to mechanism in the spinal cord of rats and finally found that nalbuphine can exert anti-inflammatory and analgesic effects on the inflammatory visceral pain caused by acetic acid through downregulating NF-κB in the spinal cord.

Conclusion

In conclusion, nalbuphine could inhibit the strong response of acute inflammatory visceral pain by inhibiting NF-κB (p65) activation to reduce the release of downstream pro-inflammatory cytokines.

List Of Abbreviations

ELISA	Enzyme-linked immunosorbent assay
TNF	tumor necrosis factor
IL	Interleukin
p-NF-κB p65	Phosphorylated nuclear factor-kappa B
IHC	Immunohistochemical
AA	acetic acid
VFM	Von Frey monofilament
SDS–PAGE	sodium dodecyl sulfate–polyacrylamide gel electrophoresis
PVDF	polyvinylidene fluoride
RT-qPCR	real-time reverse transcription-quantitative polymerase chain reaction
PWT	paw withdrawal threshold
PDTC	pyrrolidine dithiocarbamate

Declarations

Ethics approval and consent to participate

All methods were carried out in accordance with relevant guidelines and regulations in Ethic approval and consent to participation. All procedures were carried in accordance with the ARRIVE (Animal Research: Reporting In Vivo Experiments) guidelines. The Ethics Committee of the College of Stomatology, Chongqing Medical University approved this study (Project No. CQHS-REC-2021 (LSNo.011)).

Consent for publication

Not applicable.

Availability of data and materials

All data generated or analysed during this study are included in this published article.

Competing interests

The authors declare that they have no competing interests.

Funding

This study was supported by the Joint project of Chongqing Health Commission and Science and Technology Bureau (Grant No: 2020MSXM098; 2021MSXM310), and CSA Clinical Research Fund (CSA-A2021-05).

Authors' contributions

Di-jiao Ruan: Methodology, Software, Formal analysis, Investigation, Writing, Original Draft, Conceptualization. Yuan-yuan Wang: Investigation, Conceptualization. Si-si Li: Conceptualization, Writing - Review & Editing. Chao Zhang: Conceptualization, Resources. Wen-wen Zheng: Methodology. Cong Yu: Methodology, Validation, Conceptualization, Writing - Review & Editing, Supervision, Resources.

Acknowledgements

We acknowledge the support from the Technology Bureau (2020MSXM098; 2021MSXM310), and CSA Clinical Research Fund (CSA-A2021-05).

References

1. Glare P, Aubrey KR, Myles PS: **Transition from acute to chronic pain after surgery.** *The Lancet* 2019, **393**(10180):1537-1546.
2. Lovich-Sapola J, Smith CE, Brandt CP: **Postoperative pain control.** *Surg Clin North Am* 2015, **95**(2):301-318.
3. Jiang Z, Zhou G, Song Q, Bao C, Wang H, Chen Z: **Effect of Intravenous Oxycodone in Combination With Different Doses of Dexmedetomidine on Sleep Quality and Visceral Pain in Patients After Abdominal Surgery: A Randomized Study.** *The Clinical journal of pain* 2018, **34**(12):1126-1132.
4. Lu Q, Dong C, Yu J, Sun H, Sun P, Ma X, Luo C: **The dose response of sufentanil as an adjuvant to ropivacaine in cesarean section for relief from somato-visceral pain under epidural anesthesia in parturients with scarred uterus.** *Medicine* 2018, **97**(38):e12404.

5. An Y, Zhao L, Wang T, Huang J, Xiao W, Wang P, Li L, Li Z, Chen X: **Preemptive oxycodone is superior to equal dose of sufentanil to reduce visceral pain and inflammatory markers after surgery: a randomized controlled trail.** *BMC anesthesiology* 2019, **19**(1):96.
6. Haw AJ, Meyer LCR, Fuller A: **Nalbuphine and butorphanol reverse opioid-induced respiratory depression but increase arousal in etorphine-immobilized goats (*Capra hircus*).** *Veterinary Anaesthesia and Analgesia* 2016, **43**(5):539-548.
7. Sun J, Zheng Z, Li YL, Zou LW, Li GH, Wang XG, She BZ, Huang XL, Li YT: **Nalbuphine versus dexmedetomidine for treatment of combined spinal-epidural post-anesthetic shivering in pregnant women undergoing cesarean section.** *J Int Med Res* 2019, **47**(9):4442-4453.
8. Schoppmann S, Spiess D, Muller D, Burch A, Zimmermann R, Simoes-Wust AP: **Nalbuphine: a candidate for treatment of women overwhelmed with sudden, intense labor pain?** *J Matern Fetal Neonatal Med* 2021:1-3.
9. Raghav R, Jain R, Jacob TG, Dhawan A, Roy TS: **Co-administration of nalbuphine attenuates the morphine-induced anxiety and dopaminergic alterations in morphine-withdrawn rats.** *Psychopharmacology (Berl)* 2021, **238**(4):1193-1211.
10. SCHMIDT. WK, TAM. SW, SHOTZBERGER. GS, SMITH. DH, JR., CLARK. R, VERNIER VG: **Nalbuphine.** *Drug and Alcohol Dependence* 1985, **14**:339-362.
11. Yu P, Zhang J, Wang J: **Nalbuphine for spinal anesthesia: A systematic review and meta-analysis.** *Pain Pract* 2021.
12. Lin J, Wang H, Li J, Wang Q, Zhang S, Feng N, Fan R, Pei J: **kappa-Opioid receptor stimulation modulates TLR4/NF-kappaB signaling in the rat heart subjected to ischemia-reperfusion.** *Cytokine* 2013, **61**(3):842-848.
13. Yu J, Luo Y, Wen Q: **Nalbuphine suppresses breast cancer stem-like properties and epithelial-mesenchymal transition via the AKT-NFkappaB signaling pathway.** *J Exp Clin Cancer Res* 2019, **38**(1):197.
14. Zhang MR: **The effects of preemptive analgesia of nalbuphine on serum level of inflammatory factors and postoperative analgesia of the patients with laparoscopic cholecystectomy.** China Science Periodical Database: INNER MONGOLIA MEDICAL UNIVERSITY; 2018.
15. Gong Y, Zhang Y, Tao S: **Nalbuphine for analgesia after fracture surgery and its effect on circulating inflammatory factors.** *Experimental and Therapeutic Medicine* 2017, **15**:859-963.
16. Zhang Y, Jiang Q, Li T: **Nalbuphine analgesic and anti-inflammatory effects on patients undergoing thoracoscopic lobectomy during the perioperative period.** *Exp Ther Med* 2017, **14**(4):3117-3121.
17. Xi MY, Li. SS, Zhang. C, Zhang. L, Wang. T, Yu2. C: **Nalbuphine for Analgesia After Orthognathic Surgery and Its Effect on Postoperative Inflammatory and Oxidative Stress: A Randomized Double-Blind Controlled Trial.** *J Oral Maxillofac Surg* 2020, **78**(4):528-537.
18. Jiang Q, Zhang R, Liu T: **Effect of nalbuphine on patient controlled intravenous analgesia after radical resection of colon cancer.** *Oncol Lett* 2020, **19**(3):2533-2538.

19. Liu Y, Liu W, Wang X, Wan Z, Liu Y, Leng Y: **Dexmedetomidine Relieves Acute Inflammatory Visceral Pain in Rats through the ERK Pathway, Toll-Like Receptor Signaling, and TRPV1 Channel.** *J Mol Neurosci* 2018, **66**(2):279-290.
20. Kim H, Chang E, Kim H, Lee S, Kim H, Su Kim G, Kim H: **Antioxidant alpha-lipoic acid inhibits osteoclast differentiation by reducing nuclear factor-kappaB DNA binding and prevents in vivo bone resorption induced by receptor activator of nuclear factor-kappaB ligand and tumor necrosis factor-alpha.** *Free radical biology & medicine* 2006, **40**(9):1483-1493.
21. Dai S, Hirayama T, Abbas S, Abu-Amer Y: **The IkappaB kinase (IKK) inhibitor, NEMO-binding domain peptide, blocks osteoclastogenesis and bone erosion in inflammatory arthritis.** *J Biol Chem* 2004, **279**(36):37219-37222.
22. Guo Q, Chen X, Chen J, Zheng G, Xie C, Wu H, Miao Z, Lin Y, Wang X, Gao W *et al*: **STING promotes senescence, apoptosis, and extracellular matrix degradation in osteoarthritis via the NF-kappaB signaling pathway.** *Cell Death Dis* 2021, **12**(1):13.
23. Hartung JE, Eskew O, Wong T, Tchivileva IE, Oladosu FA, O'Buckley SC, Nackley AG: **Nuclear factor-kappa B regulates pain and COMT expression in a rodent model of inflammation.** *Brain Behav Immun* 2015, **50**:196-202.
24. Kim HK, Park SK, Zhou JL, Taglialatela G, Chung K, Coggeshall RE, Chung JM: **Reactive oxygen species (ROS) play an important role in a rat model of neuropathic pain.** *Pain* 2004, **111**(1-2):116-124.
25. Biswas*. DK, Dai. S-C, Cruz. A, Weiser. B, Graner. E, Pardee. AB: **The nuclear factor kappa B (NF-kB): A potential therapeutic target for estrogen receptor negative breast cancers.** *PNAS* 2001, **98**(18):10386–10391.
26. Guo. C-A, Ma L, Su. X-L, . Y-ZW, Zhen. L-L, Zhang. B, An. H, Liu H-B: **Esmolol inhibits inflammation and apoptosis in the intestinal tissue via the overexpression of NF-kB-p65 in the early stage sepsis rats.** *TURKISH JOURNAL OF GASTROENTEROLOGY* 2020, **31**(4):331-341.
27. Ren Q, Guo F, Tao S, Huang R, Ma L, Fu P: **Flavonoid fisetin alleviates kidney inflammation and apoptosis via inhibiting Src-mediated NF-kappaB p65 and MAPK signaling pathways in septic AKI mice.** *Biomed Pharmacother* 2020, **122**:109772.
28. Wei. u, Li. J, Li. Y, Song. J: **MicroRNA-451 inhibits inflammation and proliferation of glomerular mesangial cells through down-regulating PSMD11 and NF-kB p65.** *Bioscience Reports* 2019, **39**(10).
29. Zhu MM, Wang L, Yang D, Li C, Pang ST, Li XH, Li R, Yang B, Lian YP, Ma L *et al*: **Wedelolactone alleviates doxorubicin-induced inflammation and oxidative stress damage of podocytes by IkappaK/IkappaB/NF-kappaB pathway.** *Biomed Pharmacother* 2019, **117**:109088.
30. Satyanarayana PSV, Jain NK, Singh A, Kulkarni SK: **Isobolographic analysis of interaction between cyclooxygenase inhibitors and tramadol in acetic acid-induced writhing in mice.** *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 2004, **28**(4):641-649.
31. Wang H, Zhao J, Xue G, Wang J, Wu J, Wang D, Dong L: **Regulatory effect of cytokine-induced neutrophil chemoattractant, epithelial neutrophil-activating peptide 78 and pyrrolidine**

- dithiocarbamate on pulmonary neutrophil aggregation mediated by nuclear factor-kappaB in lipopolysaccharide-induced acute respiratory distress syndrome mice. *Exp Ther Med* 2016, **12**(3):1785-1794.
32. Craig AD: **Interoception: the sense of the physiological condition of the body.** *Current Opinion in Neurobiology* 2003, **13**(4):500-505.
 33. Dixon WJ, Mood AM: **A Method for Obtaining and Analyzing Sensitivity Data.** *Journal of the American Statistical Association* 1948, **43**(241):109-126.
 34. Bradman MJG, Ferrini F, Salio C, Merighi A: **Practical mechanical threshold estimation in rodents using von Frey hairs/Semmes–Weinstein monofilaments: Towards a rational method.** *Journal of Neuroscience Methods* 2015, **255**:92-103.
 35. Jänig W: **Neurobiology of visceral pain.** *Schmerz (Berlin, Germany)* 2014, **28**(3):233-251.
 36. Spange S, Wagner T, Heinzel T, Kramer OH: **Acetylation of non-histone proteins modulates cellular signalling at multiple levels.** *Int J Biochem Cell Biol* 2009, **41**(1):185-198.
 37. Yu X, Yu W, Wu L, Yang W, Lu Y: **Chitotriosidase attenuates brain inflammation via HDAC3/NF-kappaB pathway in D-galactose and aluminum-induced rat model with cognitive impairments.** *Neurosci Res* 2021, **172**:73-79.
 38. Sun LH, Zhang WX, Xu Q, Wu H, Jiao CC, Chen XZ: **Estrogen modulation of visceral pain.** *J Zhejiang Univ Sci B* 2019, **20**(8):628-636.
 39. Meerveld BG, Johnson AC: **Mechanisms of Stress-induced Visceral Pain.** *J Neurogastroenterol Motil* 2018, **24**(1):7-18.
 40. van Thiel IAM, Botschuijver S, de Jonge WJ, Seppen J: **Painful interactions: Microbial compounds and visceral pain.** *Biochim Biophys Acta Mol Basis Dis* 2020, **1866**(1):165534.
 41. Feehan AK, Morgenweck J, Zhang X, Amgott-Kwan AT, Zadina JE: **Novel Endomorphin Analogs Are More Potent and Longer-Lasting Analgesics in Neuropathic, Inflammatory, Postoperative, and Visceral Pain Relative to Morphine.** *J Pain* 2017, **18**(12):1526-1541.
 42. Kilci O, Demir T, Gunbey M, Kara C, Bayazit D, Ornek D, Baydar M: **The analgesic effect of diclofenac sodium administered via the epidural route in an experimental visceral pain model.** *Niger J Clin Pract* 2016, **19**(6):747-752.
 43. De Caro C, Cristiano C, Avagliano C, Cuzzo M, La Rana G, Aviello G, De Sarro G, Calignano A, Russo E, Russo R: **Analgesic and Anti-Inflammatory Effects of Perampanel in Acute and Chronic Pain Models in Mice: Interaction With the Cannabinergic System.** *Front Pharmacol* 2020, **11**:620221.
 44. Shaballout N, Aloumar A, Manuel J, May M, Beissner F: **Lateralization and Bodily Patterns of Segmental Signs and Spontaneous Pain in Acute Visceral Disease: Observational Study.** *J Med Internet Res* 2021, **23**(8):e27247.
 45. Gao J, Wang Y, Zhang W, Zhang J, Lu S, Meng K, Yin X, Sun Z, He QY: **C20orf27 Promotes Cell Growth and Proliferation of Colorectal Cancer via the TGFbetaR-TAK1-NFkB Pathway.** *Cancers (Basel)* 2020, **12**(2).

46. Nisr RB, Shah DS, Ganley IG, Hundal HS: **Proinflammatory NFkB signalling promotes mitochondrial dysfunction in skeletal muscle in response to cellular fuel overloading.** *Cellular and Molecular Life Sciences* 2019, **76**(24):4887-4904.
47. Tang X, Sun L, Wang G, Chen B, Luo F: **RUNX1: A Regulator of NF-κB Signaling in Pulmonary Diseases.** *Current Protein & Peptide Science* 2017, **19**(2).
48. Kaplan E, Binyaminy B, Gafni M, Keren O, Sarne Y: **Membrane-delimited proteolytic regulation of opioid receptors.** *Brain Res* 2011, **1386**:25-34.
49. Guillot X, Tordi N, Laheurte C, Pazart L, Prati C, Saas P, Wendling D: **Local ice cryotherapy decreases synovial interleukin 6, interleukin 1beta, vascular endothelial growth factor, prostaglandin-E2, and nuclear factor kappa B p65 in human knee arthritis: a controlled study.** *Arthritis Res Ther* 2019, **21**(1):180.
50. Wang M, Xu H, Chong Lee Shin OL, Li L, Gao H, Zhao Z, Zhu F, Zhu H, Liang W, Qian K *et al*: **Compound alpha-keto acid tablet supplementation alleviates chronic kidney disease progression via inhibition of the NF-κB and MAPK pathways.** *J Transl Med* 2019, **17**(1):122.
51. van Delft MAM, Huitema LFA, Tas SW: **The contribution of NF-κB signalling to immune regulation and tolerance.** *European Journal of Clinical Investigation* 2015, **45**(5):529-539.
52. Shim. DJ, Yang. L, Reed. JG, Noebels. JL, Chiao. PJ, Zheng. H: **Disruption of the NF-B/IBa Autoinhibitory Loop Improves Cognitive Performance and Promotes Hyperexcitability of Hippocampal Neurons.** *Molecular Neurodegeneration* 2011, **6**(42):2-14.
53. Reiter E, Ahn S, Shukla AK, Lefkowitz RJ: **Molecular Mechanism of β-Arrestin-Biased Agonism at Seven-Transmembrane Receptors.** *Annual Review of Pharmacology and Toxicology* 2012, **52**(1):179-197.
54. Arenas YM, Cabrera-Pastor A, Juciute N, Mora-Navarro E, Felipe V: **Blocking glycine receptors reduces neuroinflammation and restores neurotransmission in cerebellum through ADAM17-TNFR1-NF-kappabeta pathway.** *J Neuroinflammation* 2020, **17**(1):269.
55. Bae HG, Kim TK, Suk HY, Jung S, Jo DG: **White matter and neurological disorders.** *Arch Pharm Res* 2020, **43**(9):920-931.
56. Sabirzhanov B, Li Y, Coll-Miro M, Matyas JJ, He J, Kumar A, Ward N, Yu J, Faden AI, Wu J: **Inhibition of NOX2 signaling limits pain-related behavior and improves motor function in male mice after spinal cord injury: Participation of IL-10/miR-155 pathways.** *Brain Behav Immun* 2019, **80**:73-87.
57. Ballout N, Rochelle T, Brot S, Bonnet ML, Francheteau M, Prestoz L, Zibara K, Gaillard A: **Characterization of Inflammation in Delayed Cortical Transplantation.** *Front Mol Neurosci* 2019, **12**:160.

Figures

Figure 1

Behaviors of acute visceral pain tested with writhing, PWT. (a) The number of writhes for 30 minutes in each group. (b) Mechanical paw withdraw threshold in each group (data were the mean $\pm$ SEM, n = 12). *p<0.05, AA vs control; #p<0.05, AA+nalbuphine vs AA, &p<0.05; AA+PDTC vs AA.

Figure 2

Results of plasma levels of several inflammatory factors. The expressions of IL-1 β , IL-2, IL-6, and TNF- α were compared between groups (data were the mean $\pm$ SEM, n = 8). *p<0.05, AA vs control; #p<0.05, AA+nalbuphine vs AA, &p<0.05; AA+PDTC vs AA.

Figure 3

Levels of I κ B α and p-NF- κ B p65 (Ser536) in four groups. (a) The protein levels of I κ B α and p-NF- κ B p65 (Ser536) in the spinal cord were detected via WB. GAPDH was used as a loading control. (b) The mRNA expression of NF- κ B p65 in the spinal was detected via RT-qPCR. (Data are the mean $\pm$ SEM, n = 4).

*p<0.05, AA vs control; #p<0.05, AA+nalbuphine vs AA, &p<0.05; AA+PDTC vs AA.

Figure 4

The mRNA and protein levels of IL-1 β , TNF- α , and IL-6 in the lumbar spinal cord of rats. (a-c) The mRNA expressions of IL-1 β , TNF- α , and IL-6 in the spinal cord were detected via RT-qPCR. (d) The protein levels of IL-1 β , TNF- α , and IL-6 in the spinal cord were detected via western blotting. GAPDH was used as a loading control. (Data were the mean $\pm$ SEM, n = 4). *p<0.05, AA vs control; #p<0.05, AA+nalbuphine vs AA, &p<0.05; AA+PDTC vs AA.

Figure 5

The expressions of p-NF- κ B and TNF- α in the spinal were observed by IHC. (a) The number of positive cells at representative locations. (b) Representative images of p-NF- κ B and TNF- α after immunohistochemical staining (SLIDEVIEW VS200; $\times 20$, $\times 200$). (Data are the mean $\pm$ SEM, n = 4).

*p<0.05, AA vs control; #p<0.05, AA+nalbuphine vs AA, &p<0.05; AA+PDTC vs AA.