JAC1 Targets YY1-Mediated JWA/p38 MAPK Signaling to Inhibit Proliferation and Induce Apoptosis in TNBC

Zurong Zhai

Nanjing Medical University

Yanlin Ren

Nanjing Medical University

Chuanjun Shu

Nanjing Medical University

Dongyin Chen

Nanjing Medical University

Xia Liu

Nanjing Medical University

Yan Liang

The First Affiliated Hospital of Nanjing Medical University

Aiping Li

Nanjing Medical University

Jianwei Zhou (✉ jwzhou@njmu.edu.cn)

Nanjing Medical University

Research Article

Keywords: JAC1, JWA, YY1, proliferation, apoptosis, TNBC

Posted Date: January 12th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1170583/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Cell Death Discovery on April 5th, 2022. See the published version at <https://doi.org/10.1038/s41420-022-00992-9>.

Abstract

Background: Triple negative breast cancer (TNBC) is a type of breast cancer with poor prognosis, and still has no adequate therapeutic target and ideal medicine. The public database and the relative studies have shown that low expression of JWA is closely related to the poor overall survival in many cancers including breast cancer. However, the precise biological functions and behind mechanisms of JWA in TNBC are still unclear.

Methods: Both TCGA and GEO databases were used to confirm the relationship between expression levels of JWA and overall survival in TNBC cases. JAC1, an agonistic small compound of JWA gene, was used in TNBC models *in vitro* and *in vivo*. The routine cellular and molecular assays include CCK-8, colony formation, EdU incorporation, the flow cytometry, Western blot, immunohistochemistry, immunofluorescence microscopy and reporter gene assays were conducted in this study.

Results: Low expression of JWA was associated with poor prognosis in TNBC patients. JAC1 treatment inhibited TNBC cells proliferation and promoted apoptosis *in vitro* and *in vivo*. JAC1 specifically combined and targeted YY1 to eliminate its transcriptional inhibition on JWA gene. At the same time, JAC1 promoted ubiquitination and degradation of YY1. The rescued JWA induced G1 phase arrest and apoptosis in TNBC cells through the p38 MAPK signaling pathway. In addition, JAC1 disrupted the interaction between YY1 and HSF1, and suppressed the oncogenic role of HSF1 in TNBC through p-Akt signaling pathway.

Conclusions: We discovered for the first time that JAC1 is a YY1 targeting compound and maybe a potential therapeutic agent for TNBC.

Introduction

Breast cancer is the most common cancer and the leading cause of the cancer-related death for women worldwide [1, 2]. Despite decades of clinical research and corresponding progress, the incidence of breast cancer is still rising [3], affecting one in 20 globally and as many as one in eight in high-income countries. According to the latest data in 2020, the number of new cases of female breast cancer reach 2.26 million worldwide. Breast cancer, replacing lung cancer, becomes the world's largest cancer (<https://www.iarc.fr/faq/latest-global-cancer-data-2020-qa/>).

Triple negative breast cancer (TNBC) is a subtype of breast cancer accounting for 15% ~20% of all breast cancers [4], which is defined as lack of estrogen receptor (ER), progesterone receptor (PR) and human epidermal growth factor receptor 2 (HER2, also known as ERBB2) [5]. The outstanding clinical features of TNBC are the high rate of cell mitosis and strong metastasis [6], high recurrence and mortality [7], and low survival rate [8]. The expression status of hormone receptor (HR) and HER2 in tumor tissues are usually used as important evidence for the choice of therapeutic drugs. Unlike ER-positive (tamoxifen and other ER modulators as preferred regimen) and HER2-positive (Herceptin and other HER2 inhibitors as preferred regimen) breast cancers, there are no currently available first-line regimens for TNBC patients [9, 10]. Because the related receptors' expression is negative, targeted therapy for TNBC has not been

approved by Food and Drug Administration (FDA)[11]. TNBC patients are still mainly rely on conventional chemotherapy such as Paclitaxel and cisplatin [12, 13]. Usually, TNBC patients receive a good initial response to these chemotherapy regimens, but chemoresistance often occurs gradually[14], which greatly decreasing the clinical efficacy of the drugs[15]. So far, the molecular mechanism of chemotherapy resistance of TNBC patients has not been fully elucidated. TNBC accounts for 90% of breast cancer-associated mortality [16], therefore, it is an urgent unmet need to find new drugs to overcome the current challenges in TNBC therapy.

Yin Yang 1 (YY1), a zinc finger protein, is first identified as a member of the YY family in 1991 by several independent research groups[17, 18]. YY1 is shown to regulate lots of downstream molecules including tumor suppressors and oncogenes therefore involves in various homeostatic processes and diseases including cancer [19, 20]. YY1 works as a transcription factor and is associated with cell differentiation [21], apoptosis[22] and tumorigenesis[23]. It is reported that the expression level or function of YY1 has two sides in different cancers[24] ; most researches indicate YY1 is highly expressed in a range of cancer types including breast [25, 26], gastric, brain, liver, lung and melanoma[27]. However, YY1 is also reported to play a tumor suppressor role particularly in the case of pancreatic cancer [28].

JWA, also named ARL6IP5, is an effective environmental response gene. JWA actively participates in the signal pathways of cell response, such as oxidative stress[29], DNA repair[30], apoptosis[31] and differentiation[29]. In addition, JWA is identified as a tumor suppressor gene in gastric cancer[32], liver cancer [33], melanoma[34] and breast cancer[35-37]. JWA is a novel microtubule-binding protein, which regulates cancer cell migration via mitogen-activated protein kinase cascades and inhibits cell adhesion, migration, invasion and the metastasis of melanoma cells by suppressing integrin- $\alpha\beta$ 3 signaling [34, 38]. Moreover, JWA exerts pro-apoptotic role in cisplatin-resistant human gastric cancer cells through the CK2 α /XRCC1 axis [39]. As a multifunctional molecule, JWA could regulate TRAIL-induced apoptosis via MARCH8-mediated DR4 ubiquitination in cisplatin-resistant gastric cancer cells [40]. The mechanistic evidences indicated that most of anticancer functions of JWA in anti-proliferation and pro-apoptosis are mediated by activating MEK-ERK signaling [41]. Interestingly, JWA promotes As₂O₃-induced apoptosis in MCF-7 cells through p38 MAPK signaling[31]. JWA also inhibits invasion of TNBC cells by down-regulating the expression of CXCR4[35]. In addition, JWA down-regulates CDK12 to inhibit the proliferation of trastuzumab-resistant breast cancer[37]. These reports demonstrate that JWA plays an important role in suppressing tumors initiation and development.

Down regulation of JWA is related to poor prognosis in many cancers including breast cancer. Recently, we have reported anti-proliferation role of JWA gene agonist JAC1 in HER2 positive breast cancer cells[36]. Herein, we further investigated the effects of JAC1 in TNBC. We identified at the first time that JWA contributes to maintain the homeostasis of cell proliferation and apoptosis and its deficiency might be a new checkpoint in TNBC. More importantly, we determined JAC1 works as a novel effective inhibitor of YY1, it binds to overexpressed YY1 in TNBC cells and removes its transcriptional inhibitory roles on JWA gene, therefore reverses the dysregulation of proliferation/apoptosis signaling in TNBC cells.

Materials And Methods

Chemicals and reagents

JAC1 and Biotin-JAC1 were synthesized by the laboratory-self, with the purity of >98%. Primary antibodies information is listed in Supplementary Table 1. ECL advance reagent, CCK-8 Kit (C0038), Nucleoplasmic protein isolation Kit (P0027), Luciferase reporter gene Kit (RG027), Protein A/G Plus-Agarose (P2055) were purchased from Beyotime (Shanghai, China). Annexin V-FITC/PI apoptosis detection kit (A211-02), HiScript III RT SuperMix for qPCR (+gDNA wiper) (R323-01), AceQ qPCR SYBR Green Master Mix (without ROX) were obtained from Vazyme Biotech Co (Nanjing, China). CHX (761982), MG132 (C2211) were obtained from Sigma (St. Louis, MO).

Cell culture

Human breast cancer cells (SUM1315, MDA-MB-231 and BT474) and human embryonic kidney cells HEK293T were purchased from the American Type Culture Collection (ATCC; Manassas, VA, USA) and all the cells were maintained in Dulbecco's modified Eagle's medium (DMEM) containing 10% or 20% fetal bovine serum (FBS; Gibco, Grand Island, NY, USA) and 1% penicillin-streptomycin (HyClone, Logan, UT, USA). The human normal breast epithelial cells MCF-10A was purchased from Zhong Qiao Xin Zhou and cultured in a special medium (5% horse serum, 20 ng/mL epidermal growth factor, 0.5 µg/mL hydrocortisone and antibiotic-antimycotics, 0.1% penicillin, streptomycin). All the cells were incubated at 37 °C in a 5% CO₂ humidified atmosphere.

Western blot assays

Cell proteins were lysed with RIPA buffer (50 mM Tris, pH 7.4; 150 mM NaCl; 1% NP-40; 0.5% sodium deoxycholate; 0.1% SDS; and the 1% protease inhibitor) for 30 min, 4°C. The xenograft tumor tissue proteins were prepared in tissue protein extraction reagent (Thermo Fisher Scientific). The protein (40mg/20mL) was separated equal amounts of by SDS-PAGE and transferred to PVDF membrane. Then the membrane was blocked for 1h using 5% skim milk at room temperature, and the primary antibodies were used and incubated at 4°C overnight. After washing 5 times with PBST (10 min/time), the membranes were incubated with peroxidase-conjugated anti-rabbit IgG or anti-mouse IgG (1:1000 diluted, Beyotime, Shanghai, China) for 60 min. The blot images were obtained from multifunctional gel imaging system (Bio-Rad, USA).

Colony formation assay

In total of 800 cells for both SUM1315 or MDA-MB-231 were made into single-cell suspension and seeded in six-well plates and then cultured in DMEM containing 10% FBS for 10-14 days. The fresh medium containing the corresponding dose of JAC1 (or p38 inhibitor) was updated every 72 h. About two weeks later, the cell culture dishes with colonies were fixed with 4% paraformaldehyde for 30 min, stained with 0.1% crystal violet for 2h at room temperature and then photographed.

CCK-8 assay

The SUM1315, MDA-MB-231 and MCF-10A cells were seeded at the density of 3×10^3 cells/well/100mL into 96-well plates and were maintained in an incubator at 37 °C and 5% CO₂. Then CCK8 reagent (10μL) was added to each well and incubated for 2h. The absorbance at 450nm(A450) was measured on a microplate reader and calculated the cell survival ratios. This study conducted at least three replicate holes under each condition.

Cell apoptosis assay

The cells were seeded at the density of 1×10^5 cells/well/2mL in six-well plate and then collected and washed with cold PBS for 3 times at the endpoint of culture. After centrifugation, the cell pellet was suspended with 100 mL of binding buffer; 5 mL Annexin V and 5 mL PI were added and incubated for 15 min in dark at room temperature according to the manufacturer's protocol. Then, flow cytometry (BD Biosciences, NJ) was used to detect apoptotic cells within 1 h. Each experiment had 3 replicates.

Cell cycle analysis

The cells were seeded at the density of 1×10^6 cells/well/2mL in six-well plate, and collected and washed 3 times with PBS at the endpoint of culture; and then fixed with ice-cold 70% ethanol for at least 18h. Centrifugation and washing with PBS, then PI staining solution was added and incubated for 15 min in dark at room temperature. The cell cycle was analyzed using flow cytometry (BD Biosciences, NJ). Each experiment had 3 replicates.

Co-immunoprecipitation assay (CO-IP)

The cells were firstly washed twice with PBS at the end point of culture, then collected and lysed with IP lysis buffer at 4°C for 30 min, and centrifuged for 15 min at 4°C, 12,000×g. The supernatant was transferred to a new centrifuge tube, added the anti-biotin primary antibody to 500 mg total protein and incubated overnight at 4°C. The negative control group was added with the IgG (mouse IgG or rabbit IgG). 20mL of protein A/G was added in all samples and co-incubated overnight at 4°C. The samples were washed with ice-cold PBS for 4 times, then add 60mL 2×SDS loading buffer to agarose magnetic beads precipitation and boiled at 100°C for 5min. Then the protein expressions were detected by Western blot.

Immunofluorescence (IF) assay

The cultured cells were removed medium at first and fixed with methanol for 30 min, further added in 10% normal goat serum for 1h to block non-specific binding sites at room temperature. YY1 antibody was then used to incubate cells for overnight at 4°C; the CY3 labeled fluorescent secondary antibody was incubated the cells for 2 h at room temperature, and then the nucleophilic dye DAPI was stained the nucleus for 15 min. Zeiss AIM software was used to capture confocal images of the cells on a Zeiss LSM 700 confocal microscope system (Carl Zeiss Jena, Oberkochen, Germany).

Quantitative real-time PCR analysis

Total RNA was isolated with Trizol reagent (Invitrogen, Carlsbad, California, USA). cDNA was synthesized from 1 mg of total RNA using the Hiscript Q RT Super Mix reagent according to the manufacturer's protocol. The quantitative RT-qPCR was performed using ABI Prism 7900 Sequence detection system (Applied Biosystems, Canada) and SYBR Green PCR Master Mix (TaKaRa Bio, Japan). Thermal cycling conditions of the PCR was 5 min at 94°C, 36 cycles of 35 sec at 94°C, 30 sec at 56°C and 35 sec at 72°C. The expression levels of mRNA were defined based on the threshold cycle (Ct), and relative expression levels were calculated using the $2^{-\Delta\Delta C_t}$ method. The expression level of GAPDH mRNA was used as reference genes.

The respective primer sequences were:

JWA: 5'-TCTGAGGTCTTCTCTCTGAAACATC-3' and 5'-CTTCGGAACCTCAAGAACAAAC-3'

HSF1: 5'-AACACAGCCCCTATGGACA-3' and 5'-GTCTGCAGGTTGTCCAGGTT-3'

YY1: 5'-TACCTGGCATTGACCTCTC-3' and 5'-GGCCGAGTTATCCCTGA-3'

GAPDH: 5'-AATGAAGGGGTCATTGATGG-3' and 5'-AAGGTGAAGGTCGGAGTCAA-3'

Plasmids and small interfering RNA (siRNA) transfection

Human specific siRNA was synthesized by GENERAY (shanghai, China), and the plasmid pcDNA3.0-Flag-YY1 was synthesized by You Bao (Shanghai, China). The transfection assay was performed using Lipofectamine™ 3000 Reagent in Opti-MEM medium (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's protocols. After 48 h of transfection, the cells were harvested for RT-PCR or Western blot assays.

The siRNA sequences were:

si-YY1:

1. 5'-CCUCCUGAUUAUUCAGAAUTT-3' and 5'-AUUCUGAAUAAUCAGGAGGTT-3';

2. 5'-CCAAACAACUGGCAGAAUUTT-3' and 5'-AAUUCUGCCAGUUGUUUGGTT-3';

3. 5'-GCUCCAAGAACAAUAGCUUTT-3' and 5'-AAGCUAUUGUUCUUGGAGCTT-3';

si-CSDE1:

1. 5'-GGACAGAAAUGGUAAGAATT-3' and 5'-UUCUUUACCAUUUCUGUCCTT-3';

2. 5'-GGGCACGGUUUCAUUUCAUTT-3' and 5'-AUGAAAUGAAACCGUGCCCTT-3';

3.5'-GCCAAGGAUGUGGAAGGAUTT-3'and 5'-AUCCUUCCACAUCCUUGGCTT-3';

sh-p53:

1. GTACCACCATCCACTACAA;
2. AGAGAATCTCCGCAAGAAA;
3. GGAGTATTTGGATGACAGA;

sh-LARP7:

1. GGAGAAAGTTAATGCAACA;
2. GCTGGAAACTCGAGATCCT;
3. GCGAATGGATGGATTTGAA;

sh-ZFX:

1. GCAAATGGATGACAATGAA;
2. GTCGGAAATTGATCCTTGT;
3. TGCTGAAATCGCTGACGAA;

sh-STAT1:

1. CTGGATATATCAAGACTGA;
2. GCACGCTGCCAATGATGTT;
3. CATGCGGTTGAACCCTACA;

Luciferase reporter gene assay

HEK 293T cells, 5×10^4 well/500mL, were seeded and cultured overnight in 24-well plate. Then the cells were transfected with either wild-type or mutant reporter gene plasmid using Lipo3000 reagent according to the manufacturer's instructions. After 48 h, the cell lysates were collected and the activities of firefly and Renilla luciferases were measured by a Dual-Luciferase Reporter System (Promega). The dual luciferase kit (Beyotime, RG027) was used to measure luciferase activity. Relative luciferase activity was standardized by the Renilla luciferase. Each experiment was repeated three times.

Ubiquitination assay

Both SUM1315 and MDA-MB-231 cells transiently transfected with Ubiquitin (Ub) plasmid for 48 h were incubated with 5 μ M JAC1 for 24 h and then treated with MG132 (10 μ M) for another 6 h. Extracted whole cell lysates protein was incubated with anti-YY1 antibody for 24 h at 4 °C. 20 μ L Protein A/G Plus-Agarose beads were added and shaken overnight. And then the beads were washed with pre-cooled PBS for 4 times, and centrifuged at $1000 \times g$ for 5 min at 4°C; the IP product was then detected by Western blot.

Protein structure analysis

Three-dimensional structures of JWA and p38 or JAC1 and Transcription factors, such as YY1, CSDE1, ZFX, LARP7, STAT1, P53 were predicted using the AlphaFold protein structure database (<https://alphafold.ebi.ac.uk/>). The largest possible interaction region of proteins was then obtained by Discovery Studio 3.0, respectively. These predicted binding pockets were utilized to construct an initial coarse model of the complex. Then, the complex was optimized by Rosetta software. The optimization model for complex was then obtained based on energy scores. Binding sites between proteins in complex were then obtained by RING (residue interaction network generator) software. PyMol software was utilized to show High-quality 3-D images of structures.

Xenograft tumor mice model

The mice model was pre-approved by the Ethics Committee of Nanjing Medical University (IACUC-1811067). 5-week-old female BALB/c mice were purchased from Model Animal Research Center of Nanjing University (Nanjing, China) and maintained in SPF facilities. MDA-MB-231 cells (5×10^6) were inoculated subcutaneously into the right axilla of each nude mouse, respectively. When the tumor volume reached in average 60-100 mm³ (tumor volume $V = 0.5 \times \text{length} \times \text{width}^2$), the mice were randomly divided into different groups for further experimental manipulations ($n = 8$ per group). JAC1 was administered daily by oral gavage according to the mouse body weight (100 mg/kg/mouse; the solvent composition contained 50% saline, 47.5% PEG40 and 2.5% absolute ethanol). The mice body weight and tumor volume were measured every two days. At the end of the experiment, the mice were euthanized, and blood was collected to measure the serum biochemical indicators. The tumor tissues were collected either by formaldehyde fixation or -80 °C cryopreservation for further analysis.

Statistical analysis

All data were expressed either as mean $\pm$ standard deviation (SD) or standard error of the mean (SEM), and analyzed by SPSS 23.0 (SPSS Inc., Chicago, IL, USA) and GraphPad Prism 8.0 software. Use Student's t-test to determine the statistical difference between the two groups, while use one-way analysis of variance to determine the statistical difference between multiple groups. $P < 0.05$ was considered to be statistically significant. (* $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$). All data were the results of three parallels.

Result

1. Downregulation of JWA is associated with poor prognosis in TNBC

To identify if JWA expression was associated with the prognosis in breast cancer patients, 1210 cancerous and non-cancerous tissues in The Cancer Genome Atlas (TCGA) database were analyzed (<https://xenabrowser.net/datapages/>). The results showed that JWA was significantly down-regulated in breast cancer tissues ($n=1097$) compared with adjacent normal breast tissues ($n=113$) (Fig. 1A). The Overall Survival (OS) of breast cancer patients with high JWA expression is significantly longer than that

patients with low JWA expression(<https://xenabrowser.net/>)(Fig. 1B). Besides, the mRNA expression of JWA was lower in all subtypes of breast cancer than in adjacent normal breast tissues; the triple negative breast cancer (TNBC) tissues had the lowest JWA expression among the subtypes (Fig. 1C). The RNA-seq datasets (GSE2603) from GEO database further confirmed that the mRNA expression of JWA in TNBC tissues was lower than that in non-TNBC tissues (Fig. 1D). In protein level, the expression of JWA was also down-regulated in both TNBC and non-TNBC tissues compared with its adjacent normal tissues ($P < 0.05$, Fig. 1E). The results from TCGA database also showed that Progression-Free-Survival (PFS) of TNBC patients was positively correlated with the mRNA expression levels of JWA (<https://kmplot.com/analysis/>) ($P < 0.05$, Fig. 1F). For further mechanistic investigation of how JWA involves in TNBC development, we detected both the mRNA and protein expressions of JWA in normal breast epithelial cells (MCF10A) and TNBC cells (MDA-MB-231 and SUM1315). RT-PCR (Fig. 1G) and Western blot (Fig. 1H) results showed that both the mRNA and protein expression levels of JWA were lower in MDA-MB-231 and SUM1315 cells than that in MCF10A cells.

2. JAC1 inhibits the proliferation of TNBC cells

Given that overexpression of JWA (by transfection of Flag-JWA) suppresses migration and invasion in TNBC cells *in vitro* [35], and JAC1 (JWA gene activating compound 1) inhibits HER2 positive breast cancer cells proliferation by JWA-triggered E3 ubiquitin ligase SMURF1 [36]. To determine whether increase JWA expression improves the malignant phenotypes in TNBC cells, both MDA-MB-231 and SUM1315 cells were treated by different doses of JAC1 for 24 h. The CCK-8, EdU incorporating and colony formation assays were conducted to investigate the effects of JAC1 on cell viability and proliferation, respectively. The CCK-8 results showed that JAC1 treatment inhibited the cell viability dose-dependently in both MDA-MB-231 and SUM1315 cells, however, did not show significant effect on MCF-10A cells; the IC_{50} values were 21.83 mM for SUM1315 and 14.16 mM for MDA-MB-231 cells, respectively (Fig. 2A). The inhibitory effects of JAC1 on two TNBC cells were also indicated time-response characteristics although the growth curves in two cell lines were shown little bit differences (Fig. 2B, C). The EdU incorporating assays showed that the number of EdU-positive cells were reduced dose-dependently after JAC1 treatment for 24 h in both MDA-MB-231 and SUM1315 cells (Fig. 2D-G). Western blot assays indicated that JAC1 dose-dependently decreased the expression of PCNA, however, increased JWA expression in both MDA-MB-231 and SUM1315 cells (Fig. 2H, I). Similarly, the colony formation assays showed that JAC1 treatment resulted in inhibition of proliferation dose-dependently in both MDA-MB-231 and SUM1315 cells (Fig. 2J-L). These data suggested that JAC1 via JWA inhibited the proliferation in TNBC cells.

3. JAC1 triggers apoptosis and G1 phase arrest of TNBC cells

The apoptosis and cell cycle arrest are known key events to cell proliferation [42]. Here we observed the effects of JAC1 on cell cycle and apoptosis in the TNBC cell lines. The flow cytometric analysis with Annexin V/PI double staining showed that after treatment of SUM1315 and MDA-MB-231 cells with JAC1 for 24 h, the proportion of apoptotic cells were increased in a dose-dependent manner (Fig. 3A-D). To confirm the proapoptotic effects of JAC1 on TNBC cells, we determined apoptosis-related biomarkers by

Western blot in JAC1 treated TNBC cells. The data showed the expression level of Bcl2 was down regulated dose-dependently, at the same time, Bax and cleaved-caspase3 were upregulated after treatment of JAC1 for 24 h in both MDA-MB-231 and SUM1315 cells (Fig. 3E). The cell cycle assay showed that after treating two TNBC cells with 5 mM JAC1 for 24 h, G1 phase cells were increased, however, S phase cells were decreased significantly. In addition, the effects of JAC1 on cell cycle arrest were more sensitive in SUM1315 than in MDA-MB-231 cells (Fig. 3F-I). When TNBC cells were treated by JAC1 for 24 h, the expressions of p21 and CDK6 (not CDK4) were increased but cyclin D1 were inhibited significantly (Fig. 3J). The data also showed the effects of JAC1 was more sensitive on p21 than on cyclin D1. Taken together, these results suggested that JAC1 exerted double roles of pro-apoptosis and cell cycle arrest *in vitro* in TNBC cells.

Interestingly, there were no obvious proapoptotic effects in JAC1 treated normal MCF10A mammary epithelial cells, even the cells exposed to JAC1 at the dose up to 25 mM for 24 h (Fig. 3K,L). These results suggest that JAC1 exerted selective inhibitory roles on TNBC cells.

4. JAC1 suppresses tumor growth in TNBC xenografted mice

In order to evaluate the *in vivo* anti-cancer effects of JAC1 on TNBC progress, a xenograft model in nude mouse was constructed. MDA-MB-231 cells (5×10^6 cells) were subcutaneously injected into 5-week-old female BALB/c nude mice. The growth curve of xenografted TNBC tumor in mice was shown in Fig. 4A, the average volumes of tumor were increased more slowly in JAC1 (100 mg/kg/d¹⁴d) treated mice than those in solvent control mice. At the endpoint of mice model, data confirmed that both tumor mass weight and the ratio of tumor/body weight were less in JAC1 treated group than in solvent control group; and JAC1 indicated a tumor inhibition rate of 37.4% to TNBC (Fig. 4B-D). Western blot results showed that the expressions of PCNA, Bcl2 and Cyclin D1 were reduced, however, JWA, p21, Bax and cleaved-caspase3 were obviously increased in JAC1 treated mice tumor tissues compared to the control group tumors (Fig. 4E).

To understand the potential toxic side effects of JAC1 on organs in MDA-MB-231 cells xenografted mice, we also determined in total of 21 serum biochemical biomarkers. As shown in Fig. 4F-K, JAC1 treatment obviously reduced serum ALT, AST, CK, CK-MB and LDH, however, increased SOD levels in MDA-MB-231 xenografted mice. The other biochemical biomarkers did not show obvious changes by JAC1 treatment (Supplementary Table 2). In addition, H&E staining showed that JAC1 treatment resulted in obvious tissue necrosis in the central area of xenografted tumor mass compared to that of control group (Fig. 4L); the IHC assay showed Ki67 positive cells were less in JAC1 treated tumor tissue than in control group (Fig. 4M). However, the expressions of cleaved-caspase 3 and JWA were higher in JAC1 treated tumor tissue than in that of control (Fig. 4M). By the way, no differences were observed in body weight growth curve (Additional file 1: Figure S1A) and by H&E staining in liver, heart, spleen, lung and kidney tissues (Additional file 1: Figure S1B) between the two group mice. Moreover, no significant difference was observed between the control group and JAC1-treated group in terms of behavior, feeding

pattern and overall activities. In summary, these results showed that JAC1 effectively inhibited the proliferation of TNBC without significant toxicity both *in vivo* and *in vitro* experimental models.

5. JAC1 inhibits TNBC progression through JWA mediated activation of p38 MAPK signaling

Previous studies have shown that overexpression of JWA induces apoptosis in several cancer cells including MCF-7 breast cancer cells by activating p38 phosphorylation[43]. Herein, we determined if the proapoptotic and G1 cell cycle arrest effects of JAC1 were mediated by p38 signal pathway in TNBC. As we predicted, the TNBC cells treated by JAC1 for 24 h induced phosphorylation of p38 and with a dose-dependent manner (Fig. 5A). To confirm the role of p38 to JAC1 anti-proliferation in TNBC cells, p38 inhibitor (SB203580) was used to repeat the colony formation assay in both SUM1315 and MDA-MB-231 cells. Although SB203580 alone treatment did not show effects on colony formation in both cell lines, it reversed inhibitory effects of JAC1 on cell proliferation (Fig. 5B-D). In addition, CCK8 assay also showed that the p38i reversed the effect of JAC1 on cell viability in MDA-MB-231 cells (Fig. 5E). Flow cytometric analysis indicated that JAC1 induced G1 phase arrest was reversed by p38i when co-treatment of MDA-MB-231 cells with JAC1 and p38i (Fig. 5F, G). The molecule docking assay further showed the interaction between JWA and p38, and the 180th -182nd amino acids in p38 was one of interaction regions to JWA (Fig. 5H). Western blot results also confirmed that JAC1 induced activations of p-p38, Bax, p21 and inhibitions of Bcl2, Cyclin D1 were reversed by SB203580 in both MDA-MB-231 and SUM1315 TNBC cells (Fig. 5I). Since the previous studies have shown that some of JWA anticancer functions due by activation of p-ERK MAPK signal pathway, we also determined if the both p-ERK and p-JNK signaling pathways involved in JAC1 triggered anti-proliferation and pro-apoptosis events. Interestingly, U0126 (the p-ERK inhibitor) intervention only blocked the effects of JAC1 on p-ERK, and failed to reverse the effects of JAC1 on Bax, p21, Bcl2 and Cyclin D1 in the both cell lines (Additional file 2: Figure S2A, B). In addition, JAC1 did not show obvious effects on the expression of p-JNK in the both TNBC cell lines (Additional file 2: Figure S2C, D). Taken together, JAC1 triggered G1 phase arrest and pro-apoptotic effects was only dependent on p38 MAPK signaling in TNBC cells.

6. JAC1 targets YY1 and removes the transcriptional inhibition of YY1 on JWA in TNBC cells

Given that we recently have identified JAC1 via JWA/p38 signaling mediating Her2 degradation by SMURF1 in BT474 and SKBR3 human breast cancer cells[36]. Herein, the antagonist role of JAC1 on TNBC cells were also mediated by activation of JWA/p38 signaling. To identify how JAC1 differentially activated JWA in these breast cancer cells, we completed immunoprecipitation assay and followed protein sequencing by using biotin-JAC1 in BT474 breast cancer cells. The data showed biotin-JAC1 enhanced JWA expression (Additional file 3: Figure S3A), and six potential transcription factors including YY1, STAT1, p53, CSDE1, LARP7 and ZFX were identified may involve in JAC1 induced transcriptional upregulation of JWA (Additional file 3: Figure S3B). To determine which factor(s) worked in target cells, the small interfering RNAs of these six genes were constructed and transfected into cells, separately. As a result, only knocking down YY1 increased the expression of JWA (Additional file 3: Figure S3C, D), and the

other five factors CSDE1, LARP7, STAT1, p53 and ZFX did not show obvious effects on JWA expression (Additional file 3: Figure S3E-I).

The computer molecule docking assay also showed an interaction existed between JAC1 and YY1 and with a Lib Dock score at 113.619. The other five candidates were showed lower Lib Dock scores than that of YY1 (Additional file 3: Figure S3J). In TNBC cells, we confirmed the si-YY1 inhibited YY1 but enhanced JWA expressions; in contract, Flag-YY1 obviously increased YY1 but inhibited JWA protein expressions in both MDA-MB-231 and SUM1315 cells (Fig. 6A, B). To verify JAC1 via negative regulation of YY1 to activate JWA expression, Flag-YY1 was used to treat TNBC cells alone or combined with JAC1; the data indicated that JAC1 increased JWA protein expression could be blocked by Flag-YY1 (Fig. 6C, D). The negative transcriptional regulations of YY1 on JWA were also confirmed (Fig. 6E-H). Especially, the MDA-MB-231 cells transfected with si-YY1 increased the mRNA expression level of JWA (Fig. 6E), while transfected with Flag-YY1 decreased the expression level of JWA (Fig. 6F). Luciferase reporter assays showed over-expression of YY1 also reduced JWA luciferase activity (Fig. 6G); however, down-expression of YY1 increased JWA luciferase activity (Fig. 6H). The dose-response relationships of JAC1 on JWA transcription were also determined in both MDA-MB-231 and SUM1315 TNBC cells (Fig. 6I-J). To determine the YY1 binding motif sequence in JWA gene promoter region, we first predicted the potential YY1 binding sites by the promoter analysis tools PROMO (http://alggen.lsi.upc.es/cgi-bin/promo_v3/promo/promoinit.cgi?dirDB=TF_8.3) and JASPAR (<http://jaspar.genereg.net/>). The results showed three YY1 binding motif sequences may involve in this event (Fig. 6K). Then, we constructed several luciferase reporter gene wild type or mutant plasmids for the responsive JWA promoter region, and conducted assays in HEK293T cells. The results showed only the 3rd motif (-572~-560 bp; 5'-GAGAATGGCATA-3') mutant in JWA promoter region lost response to YY1 (Fig. 6L). These results provided evidence that YY1 worked as a negative regulator to the transcription of JWA.

7. JAC1 activates JWA expression by inhibiting YY1 through the ubiquitin–proteasome pathway

To determine the interaction related detail events between JAC1 and YY1 in TNBC cells, we first observed the intracellular localizations of YY1 after JAC1 treatment. Data showed that JAC1 treatment reduced nuclear YY1 levels in a dose-dependent manner, however, YY1 expression levels were no obvious changes in cytoplasm (Fig. 7A and Additional file 4: Figure S4A). The fluorescence staining assay also confirmed that compared to the control cells, JAC1 treatment significantly reduced nuclear YY1 in SUM1315 cells (Fig. 7D). To elucidate if the physical binding between JAC1 and YY1 resulted in any changes of YY1 stability, we performed protein stability assays. The results showed that JAC1 obviously accelerated degradation of YY1 in MDA-MB-231 cells (Fig. 7B, C). Interestingly, the similar effects of JAC1 on YY1 was relatively weak in SUM1315 (Additional file 4: Figure S4B). We also checked if JAC1 treatment affected expression of YY1, the data showed there was no obvious effects of JAC1 on transcription of YY1 in both TNBC cells (Additional file 4: Figure S4C, D). To confirm if JAC1 triggered YY1 degradation was mediated by ubiquitin modification, we detected ubiquitin-modified YY1 in TNBC cells which were transfected with His-ub. As shown in (Fig. 7E, F), after MG132 blockage, the ubiquitin levels in JAC1 treated cells were obviously increased compared to the control cells. As a results of ubiquitin

modification on YY1, the Western blot assay showed JAC1 treatment induced JWA but reduced YY1 expression in TNBC cells. That is to say, JAC1 treatment caused ubiquitination and degradation of YY1 protein in nucleus, removed its negative transcriptional regulation on JWA promoter, thereby increased JWA expression levels.

8. JAC1 regulates the expression of YY1 downstream genes

The YY1 has been identified extensively overexpressed in malignant tumors including breast cancer. The expression of YY1 was higher in TNBC cells than in normal cells (Fig. 7G). TCGA database also showed that breast cancer patients (Additional file 5: Figure S5A) or TNBC patients (Fig. 7H) with high YY1 expression had shorter survival than those with low YY1 expression ones.

Given that YY1 activates the expression of HSF1 and promotes the proliferation of TNBC through the Akt signaling pathway[44]. We also verified this signaling in JAC1 treated TNBC cells. Data showed HSF1 mRNA expression was increased by transfection of YY1 in both MDA-MB-231 and SUM1315 cells (Additional file 5: Figure S5B, C). Furthermore, YY1-HSF1-p-Akt signaling was also confirmed in JAC1 treated TNBC cells (Additional file 5: Figure S5D). These results suggested that as a small molecule compound targeted to YY1, the tumor suppressive effects of JAC1 on TNBC includes at least the following two paths: on the one hand, it released the effect of downstream tumor suppressive target genes such as JWA; On the other hand, it reduced the role of downstream oncogenes such as HSF1. Based on these above evidences, here we summarized the mode of action how JAC1 worked as a potential TNBC inhibitory agent (Fig. 7I).

Discussion

TNBC is the subtype of breast cancer with the highest recurrence, metastasis, and mortality[45]. For decades, TNBC has been always a clinical problem because there is no targeted therapy drug. This study was designed to address this unmet clinical need, and we evaluated the anti-TNBC biological effects and behind mechanisms of JAC1, an agonist of JWA gene. Our data clearly indicated that the anti-proliferation and pro-apoptosis effects of JAC1 in TNBC were triggered through both released tumor suppressor JWA and disrupted the interactions between YY1 and HSF1. YY1 was determined as the direct target of JAC1, and its overexpression in TNBC cells was determined closely involved in the malignant proliferation and anti-apoptosis. The mechanistic evidences showed that both G1 phase arrest and the re-balance of Bcl2/Bax in TNBC were the action mode of JAC1 driving JWA/p38 MAPK signaling. In addition, JAC1 targeting YY1 also blocked the YY1-HSF1 interaction and consequent events induced by YY1/HSF1/Akt pathway in TNBC cells.

The p38 MAPK is a stress-activated protein kinase, which is mainly activated by multiple extracellular stresses and inflammatory cytokines (such as TNF α and interleukins)[46, 47]. Previous studies have shown that activated p38 MAPK leads to cell cycle arrest and apoptosis[48]. Consistent with this, the phosphorylation of p38 induced by JAC1 was associated with the increased levels of cell cycle inhibitor

p21 and pro-apoptotic protein Bax, at same time reduced expression of Bcl2, activated caspase-3 and triggered apoptosis.

Cell proliferation is primarily controlled by several checkpoints of cell cycle [49]. G1 phase arrest allows the cells stopped its DNA synthesis and proliferation, which is a routine and useful therapeutic strategy to postpone tumor growth. In the present study, JAC1 arrested the cell cycle at G1 phase in a dose-dependent manner by increasing p21 while decreasing the expression of CyclinD1 through p38 MAPK signaling pathway.

As a tumor suppressor gene, JWA is found down-regulated in many cancer tissues including breast cancer when compared with the adjacent normal tissues. The dysregulation of intracellular post-translational modifications, such as overexpression of RNF185, was identified as the E3 ubiquitin ligase to mediate the degradation of JWA in gastric cancer cells [32]. In this study, JAC1 targeted to YY1 and blocked its oncogenic functions in TNBC cells. YY1 is a ubiquitously distributed transcription factor belonging to the GLI-Kruppel class of zinc finger proteins; and it is involved in repressing and activating a diverse number of promoters. YY1 involves in various cellular processes including cellular differentiation [21], DNA repair [50], cell survival [51] vs. apoptosis [22], etc. However, as a two sides transcriptional factor, the mechanism of YY1 is not completely determined [52]. Besides, even in breast cancer, YY1 has been reported with two-side effects (cancer suppression/cancer progression) [25, 53]. YY1 is widely expressed in mammalian cells [54], most studies show that YY1 is overexpressed in most cancers, and elevated levels of YY1 are associated with poor clinical outcomes [27, 55]. The DNA binding zinc fingers partially overlaps with sequences involved in transcriptional repression [54]. In the N-terminal region, a transcriptional activator domain is identified [17]. In this study, YY1 inhibited the transcription of JWA, which may be related to the part of YY1's DNA binding to zinc fingers, and the detailed evidences are need to be further clarified.

Ubiquitination is one of the mechanisms by which cells maintain homeostasis of related protein expression levels [36]. Instigated by E3 ligases, ubiquitylation controls many processes that are fundamental for development, such as cell division, fate specification and migration [56, 57]. In this study, JAC1-YY1 interaction accelerated YY1 ubiquitin modification and degradation; however, the behind mechanisms how JAC1 induced YY1 ubiquitination still needs to clarify. Although the E3 ubiquitin ligase Smurf2 is reported to target and degrade YY1 protein [58] and involves in progression of several cancers, we did not obtain positive evidence of Smurf2 on YY1 in TNBC.

In summary, in this study we determined at the first time that JAC1 is a YY1 targeting compound in TNBC cells. The tumor inhibitory effects of JAC1 in TNBC was due to its binding of YY1 therefore rescued anti-cancer functions of JWA via both G1 phase arrest and pro-apoptosis; in addition, the binding of JAC1 to YY1 also promoted ubiquitination mediated degradation of YY1 therefore removed the oncogenic functions of HSF1 in TNBC. Further investigations are warranted to clarify how the JAC1-YY1 interaction triggers YY1 ubiquitination and degradation; and whether the anti-proliferation and pro-apoptotic effects of JAC1 in TNBC are suitable for other types of cancer.

Conclusion

We discovered for the first time that JAC1 is a YY1 targeting compound and maybe a potential therapeutic agent for TNBC.

Abbreviations

ATCC: American Type Culture Collection;BC: Breast cancer; CCK8: Cell Counting Kit-8; DMEM: Dulbecco's modified Eagle's medium;ER: Estrogen receptor;FDA: Food and Drug Administration;FPS: Free Progression Survival;HER2: Human epidermal growth factor receptor 2;HR: Hormone receptor;IHC: Immunohistochemistry;MAPK: Mitogen-activated protein kinase; OS:Over Survival;PR: Progesterone receptor;RT-PCR: Reverse Transcription Polymerase Chain Reaction; siRNA: small interfering Ribonucleic Acid;TCGA: The Cancer Genome Atlas;TMA: Tissue microarray;TNBC: Triple negative breast cancer;YY1: Yin Yang 1.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

All data generated or analyzed during this study are included in this published article [and its supplementary information files].

Competing interests

The authors declare that they have no competing interests.

Authors' information

Not applicable.

Funding

This study was supported by the National Natural Science Foundation of China (No. 81520108027, 81521004, 81973156).

Authors' contributions

Conception and design: Z. Zhai, J. Zhou

Development of methodology: Z. Zhai, Y. Ren, C. Shu, D. Chen, X. Liu, A. Li, Y. Liang

Acquisition of data: Z. Zhai, C. Shu, X. Liu

Analysis and interpretation of data: Z. Zhai, Y. Ren

Writing, review, and/or revision of the manuscript: Z. Zhai and J. Zhou

Administrative, technical, or material support: Z. Zhai, Y. Ren, C. Shu, D. Chen, X. Liu, A. Li

Study supervision: J. Zhou

Acknowledgements

Not applicable

References

1. Zheng X, Huang M, Xing L, Yang R, Wang X, Jiang R, Zhang L, Chen J: The circRNA circSEPT9 mediated by E2F1 and EIF4A3 facilitates the carcinogenesis and development of triple-negative breast cancer. *Molecular cancer* 2020, 19(1):73.
2. Siegel RL, Miller KD, Jemal A: Cancer statistics, 2020. *CA Cancer J Clin* 2020, 70(1):7-30.
3. Global Burden of Disease Cancer C, Fitzmaurice C, Akinyemiju TF, Al Lami FH, Alam T, Alizadeh-Navaei R, Allen C, Alsharif U, Alvis-Guzman N, Amini E *et al*: Global, Regional, and National Cancer Incidence, Mortality, Years of Life Lost, Years Lived With Disability, and Disability-Adjusted Life-Years for 29 Cancer Groups, 1990 to 2016: A Systematic Analysis for the Global Burden of Disease Study. *JAMA Oncol* 2018, 4(11):1553-1568.
4. Wilcock P, Webster RM: The breast cancer drug market. *Nat Rev Drug Discov* 2021, 20(5):339-340.
5. Ademuyiwa FO, Tao Y, Luo J, Weilbaecher K, Ma CX: Differences in the mutational landscape of triple-negative breast cancer in African Americans and Caucasians. *Breast Cancer Res Treat* 2017, 161(3):491-499.
6. Bianchini G, Balko JM, Mayer IA, Sanders ME, Gianni L: Triple-negative breast cancer: challenges and opportunities of a heterogeneous disease. *Nature reviews Clinical oncology* 2016, 13(11):674-690.
7. Li J, Ma M, Yang X, Zhang M, Luo J, Zhou H, Huang N, Xiao F, Lai B, Lv W *et al*: Circular HER2 RNA positive triple negative breast cancer is sensitive to Pertuzumab. *Molecular cancer* 2020, 19(1):142.
8. O'Reilly EA, Gubbins L, Sharma S, Tully R, Guang MH, Weiner-Gorzel K, McCaffrey J, Harrison M, Furlong F, Kell M *et al*: The fate of chemoresistance in triple negative breast cancer (TNBC). *BBA clinical* 2015, 3:257-275.
9. Waks AG, Winer EP: Breast Cancer Treatment: A Review. *JAMA* 2019, 321(3):288-300.

10. Lebert JM, Lester R, Powell E, Seal M, McCarthy J: Advances in the systemic treatment of triple-negative breast cancer. *Curr Oncol* 2018, 25(Suppl 1):S142-S150.
11. Hwang SY, Park S, Kwon Y: Recent therapeutic trends and promising targets in triple negative breast cancer. *Pharmacol Ther* 2019, 199:30-57.
12. Early Breast Cancer Trialists' Collaborative G, Peto R, Davies C, Godwin J, Gray R, Pan HC, Clarke M, Cutter D, Darby S, McGale P *et al*: Comparisons between different polychemotherapy regimens for early breast cancer: meta-analyses of long-term outcome among 100,000 women in 123 randomised trials. *Lancet* 2012, 379(9814):432-444.
13. Garrido-Castro AC, Lin NU, Polyak K: Insights into Molecular Classifications of Triple-Negative Breast Cancer: Improving Patient Selection for Treatment. *Cancer Discov* 2019, 9(2):176-198.
14. Crown J, O'Shaughnessy J, Gullo G: Emerging targeted therapies in triple-negative breast cancer. *Ann Oncol* 2012, 23 Suppl 6:vi56-65.
15. Denkert C, Liedtke C, Tutt A, von Minckwitz G: Molecular alterations in triple-negative breast cancer-the road to new treatment strategies. *Lancet* 2017, 389(10087):2430-2442.
16. Zhang J, Zhang X, Li Z, Wang Q, Shi Y, Jiang X, Sun X: The miR-124-3p/Neuropilin-1 Axis Contributes to the Proliferation and Metastasis of Triple-Negative Breast Cancer Cells and Co-Activates the TGF-beta Pathway. *Frontiers in oncology* 2021, 11:654672.
17. Shi Y, Seto E, Chang LS, Shenk T: Transcriptional repression by YY1, a human GLI-Kruppel-related protein, and relief of repression by adenovirus E1A protein. *Cell* 1991, 67(2):377-388.
18. Seto E, Shi Y, Shenk T: YY1 is an initiator sequence-binding protein that directs and activates transcription in vitro. *Nature* 1991, 354(6350):241-245.
19. Schug J, Schuller WP, Kappen C, Salbaum JM, Bucan M, Stoeckert CJ, Jr.: Promoter features related to tissue specificity as measured by Shannon entropy. *Genome Biol* 2005, 6(4):R33.
20. Kim JD, Yu S, Kim J: YY1 is autoregulated through its own DNA-binding sites. *BMC Mol Biol* 2009, 10:85.
21. Fu CY, Wang PC, Tsai HJ: Competitive binding between Seryl-tRNA synthetase/YY1 complex and NFKB1 at the distal segment results in differential regulation of human vegfa promoter activity during angiogenesis. *Nucleic acids research* 2017, 45(5):2423-2437.
22. Song H, Liu Y, Li X, Chen S, Xie R, Chen D, Gao H, Wang G, Cai B, Yang X: Long noncoding RNA CASC11 promotes hepatocarcinogenesis and HCC progression through EIF4A3-mediated E2F1 activation. *Clinical and translational medicine* 2020, 10(7):e220.
23. Zhang C, Shen L, Zhu Y, Xu R, Deng Z, Liu X, Ding Y, Wang C, Shi Y, Bei L *et al*: KDM6A promotes imatinib resistance through YY1-mediated transcriptional upregulation of TRKA independently of its demethylase activity in chronic myelogenous leukemia. *Theranostics* 2021, 11(6):2691-2705.
24. Martinez-Ruiz GU, Morales-Sanchez A, Pacheco-Hernandez AF: Roles Played by YY1 in Embryonic, Adult and Cancer Stem Cells. *Stem Cell Rev Rep* 2021.

25. Allouche A, Nolens G, Tancredi A, Delacroix L, Mardaga J, Fridman V, Winkler R, Boniver J, Delvenne P, Begon DY: The combined immunodetection of AP-2alpha and YY1 transcription factors is associated with ERBB2 gene overexpression in primary breast tumors. *Breast cancer research : BCR* 2008, 10(1):R9.
26. Powe DG, Akhtar G, Habashy HO, Abdel-Fatah T, Rakha EA, Green AR, Ellis IO: Investigating AP-2 and YY1 protein expression as a cause of high HER2 gene transcription in breast cancers with discordant HER2 gene amplification. *Breast cancer research : BCR* 2009, 11(6):R90.
27. Khachigian LM: The Yin and Yang of YY1 in tumor growth and suppression. *International journal of cancer* 2018, 143(3):460-465.
28. Zhang JJ, Zhu Y, Xie KL, Peng YP, Tao JQ, Tang J, Li Z, Xu ZK, Dai CC, Qian ZY *et al*: Yin Yang-1 suppresses invasion and metastasis of pancreatic ductal adenocarcinoma by downregulating MMP10 in a MUC4/ErbB2/p38/MEF2C-dependent mechanism. *Molecular cancer* 2014, 13:130.
29. Zhu T, Chen R, Li A, Liu J, Gu D, Liu Q, H CC, Zhou J: JWA as a novel molecule involved in oxidative stress-associated signal pathway in myelogenous leukemia cells. *J Toxicol Environ Health A* 2006, 69(15):1399-1411.
30. Wang S, Wu X, Chen Y, Zhang J, Ding J, Zhou Y, He S, Tan Y, Qiang F, Bai J *et al*: Prognostic and predictive role of JWA and XRCC1 expressions in gastric cancer. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2012, 18(10):2987-2996.
31. Zhou J, Ye J, Zhao X, Li A, Zhou J: JWA is required for arsenic trioxide induced apoptosis in HeLa and MCF-7 cells via reactive oxygen species and mitochondria linked signal pathway. *Toxicol Appl Pharmacol* 2008, 230(1):33-40.
32. Qiu D, Wang Q, Wang Z, Chen J, Yan D, Zhou Y, Li A, Zhang R, Wang S, Zhou J: RNF185 modulates JWA ubiquitination and promotes gastric cancer metastasis. *Biochimica et biophysica acta Molecular basis of disease* 2018, 1864(5 Pt A):1552-1561.
33. Wu X, Chen H, Gao Q, Bai J, Wang X, Zhou J, Qiu S, Xu Y, Shi Y, Wang X *et al*: Downregulation of JWA promotes tumor invasion and predicts poor prognosis in human hepatocellular carcinoma. *Molecular carcinogenesis* 2014, 53(4):325-336.
34. Bai J, Zhang J, Wu J, Shen L, Zeng J, Ding J, Wu Y, Gong Z, Li A, Xu S *et al*: JWA regulates melanoma metastasis by integrin alphaVbeta3 signaling. *Oncogene* 2010, 29(8):1227-1237.
35. Xu L, Cheng L, Yang F, Pei B, Liu X, Zhou J, Zhu Y, Wang S: JWA suppresses the invasion of human breast carcinoma cells by downregulating the expression of CXCR4. *Molecular medicine reports* 2018, 17(6):8137-8144.
36. Ren Y, Chen D, Zhai Z, Chen J, Li A, Liang Y, Zhou J: JAC1 suppresses proliferation of breast cancer through the JWA/p38/SMURF1/HER2 signaling. *Cell death discovery* 2021, 7(1):85.
37. Liang Y, Qian C, Xie Y, Huang X, Chen J, Ren Y, Fu Z, Li Y, Zeng T, Yang F *et al*: JWA suppresses proliferation in trastuzumab-resistant breast cancer by downregulating CDK12. *Cell death discovery* 2021, 7(1):306.

38. Cui J, Shu C, Xu J, Chen D, Li J, Ding K, Chen M, Li A, He J, Shu Y *et al*: JP1 suppresses proliferation and metastasis of melanoma through MEK1/2 mediated NEDD4L-SP1-Integrin alphavbeta3 signaling. *Theranostics* 2020, 10(18):8036-8050.
39. Zhang Y, Chen J, Che Z, Shu C, Chen D, Ding K, Li A, Zhou J: JP3 enhances the toxicity of cisplatin on drug-resistant gastric cancer cells while reducing the damage to normal cells. *J Cancer* 2021, 12(7):1894-1906.
40. Wang Q, Chen Q, Zhu L, Chen M, Xu W, Panday S, Wang Z, Li A, Roe OD, Chen R *et al*: JWA regulates TRAIL-induced apoptosis via MARCH8-mediated DR4 ubiquitination in cisplatin-resistant gastric cancer cells. *Oncogenesis* 2017, 6(7):e353.
41. Wu YY, Ma TL, Ge ZJ, Lin J, Ding WL, Feng JK, Zhou SJ, Chen GC, Tan YF, Cui GX: JWA gene regulates PANC-1 pancreatic cancer cell behaviors through MEK-ERK1/2 of the MAPK signaling pathway. *Oncol Lett* 2014, 8(4):1859-1863.
42. Qi G, Liu Z, Fan R, Yin Z, Mi Y, Ren B, Liu X: Athyrium multidentatum (Doll.) Ching extract induce apoptosis via mitochondrial dysfunction and oxidative stress in HepG2 cells. *Scientific reports* 2017, 7(1):2275.
43. Shen L, Xu W, Li A, Ye J, Zhou J: JWA enhances As(2)O(3)-induced tubulin polymerization and apoptosis via p38 in HeLa and MCF-7 cells. *Apoptosis* 2011, 16(11):1177-1193.
44. Yang W, Feng B, Meng Y, Wang J, Geng B, Cui Q, Zhang H, Yang Y, Yang J: FAM3C-YY1 axis is essential for TGFbeta-promoted proliferation and migration of human breast cancer MDA-MB-231 cells via the activation of HSF1. *Journal of cellular and molecular medicine* 2019, 23(5):3464-3475.
45. Guo Z, He H, Zhang Y, Rao J, Yang T, Li T, Wang L, Shi M, Wang M, Qiu S *et al*: Heavy-Atom-Modulated Supramolecular Assembly Increases Antitumor Potency against Malignant Breast Tumors via Tunable Cooperativity. *Advanced materials* 2021, 33(2):e2004225.
46. Bellahcene M, Jacquet S, Cao XB, Tanno M, Haworth RS, Layland J, Kabir AM, Gaestel M, Davis RJ, Flavell RA *et al*: Activation of p38 mitogen-activated protein kinase contributes to the early cardiodepressant action of tumor necrosis factor. *Journal of the American College of Cardiology* 2006, 48(3):545-555.
47. Thirunavukkarasu C, Watkins SC, Gandhi CR: Mechanisms of endotoxin-induced NO, IL-6, and TNF-alpha production in activated rat hepatic stellate cells: role of p38 MAPK. *Hepatology* 2006, 44(2):389-398.
48. Romanov V, Whyard TC, Waltzer WC, Grollman AP, Rosenquist T: Aristolochic acid-induced apoptosis and G2 cell cycle arrest depends on ROS generation and MAP kinases activation. *Archives of toxicology* 2015, 89(1):47-56.
49. Wang B, Li D, Kovalchuk I, Apel IJ, Chinnaiyan AM, Woycicki RK, Cantor CR, Kovalchuk O: miR-34a directly targets tRNAi(Met) precursors and affects cellular proliferation, cell cycle, and apoptosis. *Proceedings of the National Academy of Sciences of the United States of America* 2018, 115(28):7392-7397.

50. Lu X, Liu R, Wang M, Kumar AK, Pan F, He L, Hu Z, Guo Z: MicroRNA-140 impedes DNA repair by targeting FEN1 and enhances chemotherapeutic response in breast cancer. *Oncogene* 2020, 39(1):234-247.
51. Zurkirchen L, Varum S, Giger S, Klug A, Hausel J, Bossart R, Zemke M, Cantu C, Atak ZK, Zamboni N *et al*: Yin Yang 1 sustains biosynthetic demands during brain development in a stage-specific manner. *Nature communications* 2019, 10(1):2192.
52. Sarvagalla S, Kolapalli SP, Vallabhapurapu S: The Two Sides of YY1 in Cancer: A Friend and a Foe. *Frontiers in oncology* 2019, 9:1230.
53. Shen X, Zhong J, Yu P, Zhao Q, Huang T: YY1-regulated LINC00152 promotes triple negative breast cancer progression by affecting on stability of PTEN protein. *Biochemical and biophysical research communications* 2019, 509(2):448-454.
54. Verheul TCJ, van Hijfte L, Perenthaler E, Barakat TS: The Why of YY1: Mechanisms of Transcriptional Regulation by Yin Yang 1. *Frontiers in cell and developmental biology* 2020, 8:592164.
55. Meliala ITS, Hosea R, Kasim V, Wu S: The biological implications of Yin Yang 1 in the hallmarks of cancer. *Theranostics* 2020, 10(9):4183-4200.
56. Rape M: Ubiquitylation at the crossroads of development and disease. *Nature reviews Molecular cell biology* 2018, 19(1):59-70.
57. Senft D, Qi J, Ronai ZA: Ubiquitin ligases in oncogenic transformation and cancer therapy. *Nature reviews Cancer* 2018, 18(2):69-88.
58. Jeong HM, Lee SH, Yum J, Yeo CY, Lee KY: Smurf2 regulates the degradation of YY1. *Biochimica et biophysica acta* 2014, 1843(9):2005-2011.

Figures

Fig. 1

Figure 1

Downregulation of JWA is associated with poor prognosis in TNBC.

A JWA mRNA expression was interrogated in unpaired cohorts (TCGA database, 1097 cancer samples, 113 non-cancerous samples). **B** Kaplan–Meier curves depicted Overall Survival according to the JWA expression of breast cancer cohort (n=1089). **C** The JWA expression level was analyzed in subtypes of breast cancers. (TCGA database, 114 normal; 566 Luminal; 37 HER2 positive; 116 Triple negative). **D** GSE2603 database was analyzed for JWA expression (24 TNBC; 71 non-TNBC). JWA mRNA level was compared between TNBC and non-TNBC. **E** The protein expression of JWA was compared between normal (n=10), TNBC (n=6) and non-TNBC (n=84) tissues. **F** Kaplan–Meier curves depicted Progression Free Survival according to the expression of JWA in TNBC cohort. *P* values were calculated with the log-rank test. **G** The mRNA expression of JWA in normal breast cell (MCF10A) and TNBC (MDA-MB-231 and

SUM 1315) cells (n=3). **H** The protein expression of JWA in normal breast cell (MCF10A) and TNBC (MDA-MB-231 and SUM1315) cells. $Z\text{-score} = \log_2 \frac{\text{FPKM} + 1}{\text{FPKM} + 1}$. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$.

Fig. 2

Figure 2

JAC1 inhibits the proliferation of TNBC cells.

A Viabilities of MDA-MB-231, SUM1315 and MCF10A cells were analyzed by CCK8 after treated by different doses of JAC1 for 24 h (n=5). **B-C** Viability of SUM1315 cells (**B**) and MDA-MB-231 cells (**C**) were analyzed by CCK8 in 6 days (n=5). **D-E** EdU assay of MDA-MB-231 cells was performed to evaluate cell proliferation after treated by different doses of JAC1 for 24 h (magnification, × 50, Scale bar, 50 μm), (n=3). **F-G** EdU assay of SUM1315 cells was performed to evaluate cell proliferation after treated by

different doses of JAC1 for 24 h (magnification, $\times 50$, Scale bar, 50 μm), ($n=3$). **H-I** The expressions of PCNA protein in MDA-MB-231 (**H**) and SUM1315 (**I**) cells were measured by Western blot after treated with 0, 2, 5 μM JAC1 for 24 h. **J** Colony formation assays were completed for MDA-MB-231 and SUM1315 cells treated by JAC1 (0, 2, 5 μM). **K-L** The quantitative data of colony numbers of SUM1315 (**K**) and MDA-MB-231 (**L**) cells ($n=3$). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$. N.S. no significant differences.

Fig. 3

Figure 3

JAC1 triggers apoptosis and cell cycle arrest of TNBC cells.

A-B Apoptosis rate was tested using flow cytometry in MDA-MB-231 (**A**) and SUM1315 (**B**) cells after treated with 0, 2, 5 μ M JAC1 for 24 h. **C-D** The statistics of apoptosis rate in MDA-MB-231 (**C**) and SUM1315 (**D**) cells (n=3). **E** The expressions of apoptosis-related proteins were determined by western blot After TNBC cells were treated with the JAC1 (0, 2, 5 μ M) for 24 h. **F-G** Cell cycle analysis was performed using flow cytometry in MDA-MB-231 (**F**) and SUM1315 (**G**) cells after treated with 0, 2, 5 μ M JAC1 for 24 h. **H-I** The statistics of cell population in MDA-MB-231 (**H**) and SUM1315 (**I**) cells (n=3). **J** The expressions of cell cycle-related proteins were determined by western blot in MDA-MB-231 and SUM1315 cells after treated with 0, 2, 5 μ M JAC1 for 24 h. **K** Apoptosis rate was tested in MCF10A cells using flow cytometry after treated with 0, 2, 5, 25 μ M JAC1 for 24 h. **L** The statistics of apoptosis rate in MCF10A cells (n=3). * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$. N.S. no significant differences.

Fig. 4

Figure 4

JAC1 suppresses tumor growth in TNBC xenografted mice.

A The tumor growth curves of MDA-MB-231 cells injection in Control and JAC1 treated groups (n=8). **B** The tumor weight in Control and JAC1-treated groups (n=8). **C** The ratio of tumor weight/body weight in Control and JAC1-treated groups (n=8). **D** The representative images of subcutaneous tumors in Control and JAC1 treated groups (n=8). **E** Analysis of the expression levels of indicated molecules in tumor tissues of MDA-MB-231 cells tumor-bearing mouse models by Immunoblotting. **F-K** The serum biochemical indexes (ALT, AST, CK, CK-MB, SOD, LDH) in control and JAC1-treated mice (n=8). **L** H&E staining of tumor tissues in Control and JAC1 treated groups. **M** IHC staining of Ki-67, Cleaved caspase3 and JWA in Control and JAC1-treated tumor tissues. * $P < 0.05$; ** $P < 0.01$. N.S. no significant differences.

Fig. 5

Figure 5

JAC1 inhibits TNBC progression through JWA mediated activation of p38 MAPK signaling.

A The expressions of p-p38 were determined by Western blot in both MDA-MB-231 and SUM1315 cells after JAC1 (0, 2, 5 μM) treated 24 h. **B** The inhibition of JAC1 on colony formation via the p38 MAPK pathway was determined in both SUM1315 and MDA-MB-231 cells. **C-D** The quantitative data of colony numbers in both SUM1315 (**C**) and MDA-MB-231 (**D**) cells (n=3). **E** The proliferation inhibition of JAC1 via the p38 MAPK pathway was determined by CCK-8 assay in MDA-MB-231 cells (n=5). **F** The cell cycle arrest via the p38 MAPK pathway was determined by flow cytometry in MDA-MB-231 cells after treated by 5 μM JAC1 for 24 h. **G** The statistics of relative cell population of MDA-MB-231 cells (n=3). **H** Molecular

docking showed the interaction between JWA and p38. The expression levels of indicated molecules (p-p38, Bax, Bcl2, p21, and CyclinD1) were analyzed by western blot in both MDA-MB-231 and SUM1315 cells after treated with JAC1 or SB203580 for 24 h. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$; **** $P < 0.0001$. N.S. no significant differences.

Fig. 6

Figure 6

JAC1 targets YY1 and removes the transcriptional inhibition of YY1 on JWA in TNBC cells.

A-B The relative expressions of JWA were detected by Western blot after transfected with si-YY1(**A**) or Flag-YY1(**B**) in MDA-MB-231 and SUM1315 cells. **C-D** JWA expressions were analyzed by Western blot in SUM1315 (**C**) and MDA-MB-231(**D**) cells after treated with JAC1 or Flag-YY1 or co-treated. **E-F** The JWA mRNA expression was detected by RT-PCR after transfected by si-YY1(**E**) or Flag-YY1(**F**) for 48 h in MDA-MB-231 cells. **G-H** JWA promoter luciferase activities were analyzed in HEK293T cells after transfected by si-YY1(**G**) or Flag-YY1(**H**) plasmid. **I-J** The relative JWA mRNA expression was detected by RT-PCR after treated with indicated doses of JAC1 in MDA-MB-231(**I**) and SUM1315(**J**) cells. **K** Three putative YY1 binding sites in the JWA promoter regions. **L** JWA promoter Luciferase activities were detected after transfected by luciferase reporter gene plasmid which mutated in putative YY1 binding sites. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$. N.S. no significant differences.

Fig. 7

Figure 7

JAC1 activates JWA expression by inhibiting YY1 through the ubiquitin–proteasome pathway.

A Western blot assay analyzed the YY1 expression in nucleoplasm and cytoplasm treated with increasing doses of JAC1 in both MDA-MB-231 and SUM1315 cells. **B**. YY1 stability assay. MDA-MB-231 cells were treated by JAC1 for 24 h and followed by exposed to CHX for indicated time; expression of YY1 was determined by Western blot. **C** The time-course intensities of the YY1 protein after treated by JAC1 and followed by indicated time of CHX exposed. **D** Immunofluorescence analyzed the expressions and location of YY1 in SUM1315 cells after treated with JAC1 at indicated doses (5 μ M). **E-F** JAC1 increases ubiquitination of YY1. The MDA-MB-231(**E**) and SUM1315(**F**) cells were treated with JAC1, followed by MG132; ubiquitination and expression of YY1 were determined by Co-IP and Western blot, respectively. **G** The expression levels of YY1 in MCF10A, MDA-MB-231 and SUM1315 cells. **H** Kaplan–Meier curves depicted survival according to the expression of YY1 in the TNBC cohort from TCGA database. *P* values were calculated with the log-rank test. **I** Schematic diagram of the molecular mechanism of JAC1 inhibiting the proliferation of TNBC.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryTable.docx](#)
- [SupplementaryFig.docx](#)