Polymerase Chain Reaction Confirmed Mycobacterium Tuberculosis Intermediate Uveitis – Analysis of 22 Eyes of 14 Cases From a Tertiary Care Centre in South India.

Dr. Janani Sreenivasan

Sankara Nethralaya

Dr. Anupreeti Jain

Sankara Nethralaya

Dr. P. Neha Kamalini

Sankara Nethralaya

M. K. Janani

Sankara Nethralaya

Dr. Jyotirmay Biswas (✉ drjb@snmail.org)

Sankara Nethralaya

Research Article

Keywords: Intermediate uveitis, Polymerase Chain Reaction, Mycobacterium tuberculosis, real time PCR, nested PCR

Posted Date: December 29th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1177212/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: To report the role of Polymerase Chain Reaction in confirming the diagnosis of presumed mycobacterium Tuberculous Intermediate Uveitis.

Method: Retrospective analysis of 22 eyes of 14 cases of presumed tubercular intermediate uveitis wherein intraocular fluid was tested for MTB DNA by Nested & Real time PCR, based on clinical suspicion of tubercular aetiology. QuantiFERON TB gold test and High Resolution CT Chest were done. Patients were treated with anti-tubercular therapy with oral steroids & immunomodulators. In the study, eleven were male (79%) and three female (21%). The median age was 34 years. Nested PCR for both IS 6110 & MPB 64 was positive in 64% of the cases, IS 6110 positive in 23% and MPB 64 positive in 15%. Real time PCR was positive in 48% of the cases. Vision improved in 33% cases, maintained in 57% cases and worsened in 10 % of cases.

Conclusion: Presumed Tubercular intermediate uveitis can be confirmed by PCR of intraocular fluids. Anti-tubercular therapy with immunosuppression can improve vision and prevent recurrences in such cases.

Background

Tuberculosis is a major health problem in India. It is the second leading cause of death by a single infectious agent worldwide.^[1] Ocular TB is a recognized form of extrapulmonary TB associated with significant morbidity.^[2] Ocular tuberculosis can manifest as anterior, intermediate, posterior or pan uveitis. It often poses diagnostic dilemma due to its protean manifestations. If it is not treated adequately, it can lead to permanent loss of vision.^[1] In recent years, use of molecular biologic techniques based on DNA amplification to detect small amounts of genomic sequences from fluids or tissues has allowed us to diagnose or confirm infections that were previously difficult to detect.^[1] Polymerase chain reaction (PCR) is one such technique for evaluation of very small amounts of DNA or RNA by enzymatic amplification of nucleic acid sequences.^[3,4] The first use of PCR in detecting *Mycobacterium tuberculosis* (MTB) from the eye was reported from aqueous samples of two patients with active retinal vasculitis.^[3,5] Quantitative (qPCR) or real-time PCR is a variant of PCR which allows the quantification of nucleic acid in the sample and facilitates the monitoring of the progress of a PCR reaction in real time.^[3,6] Although PCR positivity for MTB has been reported before in other tubercular uveitis entities (anterior, posterior) and retinal vasculitis, there are very few studies which has reported PCR positivity for the same in intermediate uveitis. We studied intraocular fluid by PCR in intermediate uveitis for mycobacterial tuberculosis DNA by both nested and real time PCR. The objective of the study is to report the clinical profile, management and visual outcome of 14 cases of intermediate uveitis positive for Mycobacterium tuberculosis DNA by PCR of intraocular fluid (aqueous and vitreous). According to COTS-2 guidelines either one or both immunologic tests (purified protein derivative PPD, interferon gamma release assay IGRA) along with high resolution computed tomography (HRCT) of chest suggestive of healed TB aetiology is required to start anti tubercular treatment.

Methodology

It was a retrospective case study wherein 22 eyes of 14 patients diagnosed as unilateral / bilateral intermediate uveitis with intraocular fluid was positive for *Mycobacterium tuberculosis* (MTB) DNA by PCR – Nested (both IS 6110 & MPB 64 gene) and/or real time with minimum 1 year follow up were analysed.

The diagnosis of intermediate uveitis was made based on Standardization of Uveitis Nomenclature (SUN) working groups international workshop for reporting clinical data which stated that it is subset of uveitis where the vitreous is the major site of the inflammation, with or without peripheral vascular sheathing and macular oedema. ^[7] All patients had a detailed history including systemic illness recorded in the case records. The patients underwent complete ophthalmic examination including best corrected visual acuity, intraocular pressure by applanation tonometer, slit lamp bio microscopy, and posterior segment examination with both slit lamp bio microscopy and indirect ophthalmoscopy at baseline and subsequent follow up visits. Ancillary tests including fundus fluorescein angiography, optical coherence tomography with or without angiography, ultrasound bio microscopy, etc., were done as and when indicated. Systemic assessment of all patients was done by an inhouse internist. All patients underwent laboratory investigations like complete blood count (CBC), erythrocyte sedimentation rate (ESR), C-reactive protein (CRP), syphilitic screening by Venereal Disease Research Laboratory (VDRL), and treponemal pallidum hemagglutination test, high resolution computed tomography (HRCT) chest, Mantoux test, tuberculosis (TB) interferon gamma release assay (Quanti-FERON TB gold test) and serum angiotensin-converting enzyme (ACE) level were done. Also magnetic resonance imaging (MRI) scan of the brain and immune markers consisting of rheumatoid factor (RF), anti- double-stranded deoxyribonucleic acid (anti-dsDNA), antineutrophil cytoplasmic antibodies (ANCA) were done whenever required. After ruling out other causes of intermediate uveitis, the decision to test MTB PCR in ocular fluids was taken if patients show positive results in one of the corroborative investigations suggesting tubercular aetiology which include positive PPD test, IGRA assay, HRCT chest. PCR was done in aqueous sample of 7 patients and vitreous sample of 7 patients. The patients who were tested with vitreous samples had undergone diagnostic as well as therapeutic vitrectomy for non-resolving or recurring inflammation, despite maximal tolerated anti-inflammatory therapy.

DNA was extracted from the specimens (aqueous aspirate) using Qiagen DNA extraction kit (Hilden, Germany) following the manufacturer's instructions. The extracted DNA was stored at -20 degree C till further use. Nested PCR with 2 sets of primers targeting MPB 64 gene and IS6110 region were carried out. ^[8] Real time PCR for the detection of the Mycobacterial load was estimated in the DNA extracts of test sample using a commercial kit - Geno Sen's® MTb Complex RG quantitative and the assay was performed on Rotor Gene (Hilden, Germany) real time PCR equipment based on Taqman principle.

All patients received treatment with oral steroids and four drug anti-tubercular therapy. Those not responding to corticosteroids immunomodulators was added. Decision regarding treatment regimen in

terms of initiation and duration of ATT as well as addition of immunomodulator therapy was taken in consensus with attending physicians in collaboration with respiratory disease physicians.

The primary outcome measure was change in visual acuity, whether improved or maintained or deteriorated at the 1 year follow up. The secondary outcome measured was the proportion of patients with recurrence of inflammation after 6 months of initiation of treatment. The study was approved by the institutional review board and adhered to the tenets of Helsinki.

Results

The study included fourteen patients, out of which 11 were males and 3 were females; The median age was 34 years (Range: 10-61years). The disease was bilateral in 8 patients (2%). Defective vision (93%) was the most common presenting complaint, followed by floaters (70%). Twenty three percent of patients were previously treated with oral steroids and another 23% with posterior sub tenon steroid injections. The mean best corrected visual acuity at presentation and at last follow up were 6/24 and 6/18 respectively. Anterior segment was quiet with broad based posterior synechiae (Figure 4) in twenty nine percent of patients. Vitritis was the most common finding, present in all the eyes and snowballs exudates were present in 33% of the eyes. Fundus examination (pseudocolor fundus picture- Optos 200Tx imaging system, Optos PLC, Dunfemline, Scotland, UK) of both eyes (Figure 1A&1B) of a patient whose aqueous sample was tested positive for PCR MTB showed relatively clear media with snowball vitreous exudates (black arrowheads) and attached retina and the right eye(Figure 1A) also had peripheral vitreous membranes (blue arrow figure 1B). PPD was noted in 46%, positive HRCT in 38% and Quanti-FERON TB gold was positive in 46% cases. All the three were positive in 1% of the patients. PCR was done in aqueous sample of 7 patients and vitreous sample of 7 patients. PCR results are shown in Table 1. Nested PCR for both IS 6110 & MPB 64 was positive in 64% of the cases. IS 6110 positive in 23% and MPB 64 positive in 15%. Real time PCR was positive in 48% of the cases. Nested and Real time PCR positivity in aqueous sample of 7 patients (Figure 2) and that of a vitreous sample of 7 patients (Figure 3). All patients were treated with oral steroids with Anti-tubercular treatment for 9 months after being evaluated by attending inhouse physician and a pulmonologist. Posterior sub tenon steroid injections were given in 15 % of the patients. Immunomodulator therapy was needed in 46% of the patients and most common drug used was Mycophenolate Mofetil. Ocular complications encountered were epiretinal membranes (29% of eyes), cystoid macular oedema (14%), posterior sub capsular cataract (0.1%), hypotony (0.1%), choroidal neo vascular membrane (CNVM) (0.1%). No systemic complications encountered. Vision improved in 33% of the cases, maintained in 57% and worsened in 10 % of cases. Factors associated with poor visual acuity include cystoid macular oedema, epiretinal membrane and cataract. Recurrences were observed in 23% of the eyes at 6 months after initiation of treatment.

Table 1: PCR for MTB results of intraocular fluids of patients included in the study

PCR	PERCENTAGE OF CASES POSITIVE
NESTED PCR (MPB 64 & IS 6110)	64%
PCR IS 6110	23%
PCR MPB 64	15%
REAL TIME PCR	48%

Discussion

Mycobacterium tuberculosis is one of the important infectious agent responsible for intermediate uveitis. It was thought that eye shows hypersensitivity to the non-viable or tubercular bacilli DNA causing such uveitis. For definite diagnosis of aetiology, isolation of the MTB bacilli from vitreous sample, aqueous paracentesis or retinal biopsy is required.

Our study provided the clinical spectrum and visual outcome of patients with intermediate uveitis with positive PCR of intraocular fluids for MTB. Pubmed search did not show any study on Real Time/nested Quantitative PCR positivity for MTB in intermediate uveitis. PCR has emerged as a powerful tool for rapid detection of the mycobacterial genome, with a high specificity and a variable sensitivity. The utility of PCR analysis lies in important facts like ocular samples usually are tested negative for the acid fast bacilli (AFB) with direct smear and culture due to paucibacillary nature of the disease and the systemic involvement may not be present at the time of ocular disease, causing delay in diagnosis and management.^[9]

PCR has a very low false-positive rate on intraocular fluids. External contamination leads to false-positive and polymorphism, specimen degradation, or failure to sample in the acute stages of disease causes false-negative results.¹⁹ Arora et al. evaluated the role of PCR for detection of Mycobacterium tuberculosis in aqueous humour samples obtained from eyes with active uveitis and showed that it can be effectively used for the diagnosis of intraocular tuberculosis which was supported by the review report by Gupta et al.²⁰ In our study we performed PCR testing for presumed tubercular intermediate uveitis. As culture is difficult and time consuming, presently, the use of real-time PCR can be done to establish tubercular aetiology, our study also proved PCR to be an important tool for rapid detection of the mycobacterial genome in suspected tubercular intermediate uveitis cases.

According to COTS-2 guidelines either one or both immunologic tests (purified protein derivative) PPD, interferon gamma release assay IGRA) along with high resolution computed tomography (HRCT) of chest suggestive of healed TB infections were candidates for ATT treatment.¹⁶

Parchand et al discussed the clinical profile of intermediate uveitis in Indian population and concluded that addition of anti-tuberculosis therapy in cases of intermediate uveitis of presumed tubercular origin can reduce the recurrences.¹⁰ According to them, patients were diagnosed as intermediate uveitis of

presumed tubercular aetiology if there was a documented positive tuberculin skin test (10 mm of induration or more) at 48–72 hours with evidence of vitritis, snowballs, or snow banking; and all known causes of infectious uveitis except TB and known non-infectious uveitis syndromes were ruled out. Madhavan et al. proved the presence of tuberculosis genome in the vitreous chamber fluid and the epiretinal membrane in Eales disease. They found that 5 out of 14 vitreous (20.8%) fluid samples were positive for tuberculosis genome. In our study we found that about 64% of the patients showed positivity for TB genome in nested PCR and 48 percent cases showed positivity in real time PCR in intermediate uveitis. The PCR results showcased circulating antigens of tuberculosis leading to an inflammatory response. In our study, fundus evaluation of patients with positive PCR (AC tap or Vitreous tap) demonstrated features snow ball opacities and vitritis. Our study provides clinical profile of cases of intermediate uveitis with documented evidence of MTB DNA in ocular fluids. Of note is that 48 % of patients in the study had actively multiplying copies of MTB DNA by Real-time PCR. In cases with signs and symptoms of intermediate tubercular uveitis but not fitting into COTS 2 criteria, ocular fluids PCR testing can also be considered as diagnostic modality for appropriate diagnosis and treatment. Figueira et al. described the role of anti- tuberculosis treatment in inflammatory diseases suggestive of tubercular aetiology.²¹ The consensus was that the positively screened patients should be treated for active tuberculosis with 4 drugs (isoniazid, rifampicin, pyrazinamide and ethambutol) for 6–9 months. Patients should be reviewed at end of initiation phase which is 2 months and the end of treatment which is 9 months. Sixteen percent of our patients had recurrences in one year period.

The major limitations of the study include small sample size, retrospective nature, lack of comparison with other causes of intermediate uveitis with intraocular fluids negative for PCR and relatively short follow up, lack of standardization of PCR and low sensitivity. Further studies are needed to find out the role of PCR in intermediate uveitis of tubercular origin.

Conclusion

Tuberculosis should be considered as an important aetiology of intermediate uveitis in endemic countries like India. Intermediate uveitis of tubercular aetiology can be diagnosed by polymerase Chain Reaction of intraocular fluids. It can also be considered as a diagnostic modality in cases strong clinical suspicion of tubercular origin. Although anti-inflammatory therapy in the form of oral steroids with or without immunosuppression remain the mainstay of treatment of intermediate uveitis, addition of anti-tubercular therapy to treatment regime in PCR MTB positive cases can cause resolution of uveitis and reduce the recurrences.

Abbreviations

PCR: Polymerase Chain Reaction

MTB: Mycobacterium Tuberculosis

HRCT: High Resolution Coherence Tomography

qPCR: quantitative Polymerase Chain Reaction

COTS: Collaborative Ocular Tuberculosis Study

IGRA: Interferon Gamma Release Assay

SUN: Standardization of Uveitis Nomenclature

PPD: Purified Protein Derivative

ATT: Anti Tubercular Treatment

Declarations

All manuscripts must contain the following sections under the heading 'Declarations':

- Ethics approval and consent to participate: taken
- Consent for publication Yes
- Availability of data and material: available with corresponding author
- Competing interests: The authors declare that they have no competing interests
- Funding Not applicable
- Authors' contributions: Dr JS: manuscript preparation

Dr AJ: Manuscript preparation

Dr NK: Manuscript preparation

Dr JB: Manuscript preparation and supervision

MKJ: PCR related work with manuscript preparation

- Acknowledgements: Not applicable

References

1. Ortega-Larrocea, G., Bobadilla-del-Valle, M., Ponce-de-León, A., & Sifuentes-Osornio, J. (2003). Nested Polymerase Chain Reaction for Mycobacterium tuberculosis DNA Detection in Aqueous and Vitreous of Patients with Uveitis. Archives of Medical Research, 34(2), 116–119. doi:10.1016/s0188-4409(02)00467-8
2. Shakarchi FI. Ocular tuberculosis: current perspectives. Clin Ophthalmol. 2015;9:2223-2227

3. Sharma P, Bansal R, Gupta V, Gupta A. Diagnosis of tubercular uveitis by quantitative polymerase chain reaction. *J Ophthalmic Inflamm Infect*. 2010 Nov 11;1(1):23-7. doi: 10.1007/s12348-010-0004-8. PMID: 21475653; PMCID: PMC3062759.
4. Chi-Chao C, DeFen C, Tuo J. Polymerase chain reaction in the diagnosis of uveitis. *Int Ophthalmol Clin*. 2005;45:41–55. doi: 10.1097/01.iio.0000155936.98330.3d.
5. Kotake S, Kimura K, Yoshikawa K, et al. Polymerase chain reaction for detection of *Mycobacterium tuberculosis* in ocular tuberculosis [letter] *Am J Ophthalmol*. 1994;117:805–806.
6. Heid CA, Stevens J, Livak KJ, Williams PM. Real time quantitative PCR. *Genome Res*. 1996;6:986–994. doi: 10.1101/gr.6.10.986.
7. Wakabayashi T, Morimura Y, Miyamoto Y, Okada AA. (2003) Changing patterns of intraocular inflammatory diseases. *Ocul Immunol Inflamm* 11: 277– 286
8. Therese KL, Gayathri R, Dhanurekha L et al (2013) Diagnostic appraisal of simultaneous application of two nested PCRs targeting MPB64 gene and IS6110 region for rapid detection of *M. tuberculosis* genome in culture proven clinical specimens. *Indian J Med Microbiol*. 31(4):366–369.
9. Verma, Aditya & Biswas, Jyotirmay & Lakshmipathy, Dhanurekha & Ramasubban, Gayathri & Therese, Lily. (2015). Detection of *Mycobacterium tuberculosis* with nested polymerase chain reaction analysis in enucleated eye ball in Eales' disease. *International ophthalmology*. 36. 10.1007/s10792-015-0144-9.
10. Parchand, S., Tandan, M., Gupta, V. et al. Intermediate uveitis in Indian population. *J Ophthal Inflamm Infect* 1, 65–70 (2011). <https://doi.org/10.1007/s12348-011-0020-3>
11. Agrawal R, Gunasekeran DV, Grant R, et al. Clinical Features and Outcomes of Patients With Tubercular Uveitis Treated With Antitubercular Therapy in the Collaborative Ocular Tuberculosis Study (COTS)–1. *JAMA Ophthalmol*. 2017;135(12):1318–1327. doi:10.1001/jamaophthalmol.2017.4485
12. Sagar Sudhakar Bhuibhar & Jyotirmay Biswas (2012) Nested PCR-positive Tubercular Ampiginous Choroiditis: A Case Report, *Ocular Immunology and Inflammation*, 20:4, 303-305, DOI: 10.3109/09273948.2012.685684
13. Agarwal M, Shrivastav A (2017) An Update on Tubercular Uveitis. *J Clin Exp Ophthalmol* 8: 676. doi:10.4172/2155-9570.1000676
14. Biswas J, Kazi MS, Agarwal VA, Alam MS, Therese KL. Polymerase chain reaction for *Mycobacterium tuberculosis* DNA detection from ocular fluids in patients with various types of choroiditis in a referral eye center in India. *Indian J Ophthalmol*. 2016 Dec;64(12):904-907. doi: 10.4103/0301-4738.198857. PMID: 28112131; PMCID: PMC5322705.
15. Gupta A, Gupta V, Arora S et al (2001) PCR-positive tubercular retinal vasculitis. *Retina* 21(5):435–444
16. Agrawal R, Testi I, Bodaghi B, Barisani-Asenbauer T, McCluskey P, Agarwal A, Kempen JH, Gupta A, Smith JR, de Smet MD, Yuen YS, Mahajan S, Kon OM, Nguyen QD, Pavesio C, Gupta V; Collaborative Ocular Tuberculosis Study Consensus Group. Collaborative Ocular Tuberculosis Study Consensus

Guidelines on the Management of Tubercular Uveitis-Report 2: Guidelines for Initiating Antitubercular Therapy in Anterior Uveitis, Intermediate Uveitis, Panuveitis, and Retinal Vasculitis. *Ophthalmology*. 2021 Feb;128(2):277-287. doi: 10.1016/j.ophtha.2020.06.052. Epub 2020 Jun 27. PMID: 32603726.

17. Madhavan HN, Therese KL, Gunisha P, Jayanthi U, Biswas J (2000) Polymerase chain reaction for detection of *Mycobacterium tuberculosis* in epiretinal membrane in Eales' disease. *Invest Ophthalmol Vis Sci* 41(3):822–859
18. Kharel (Sitaula), R., Janani, M.K., Madhavan, H.N. et al. Outcome of polymerase chain reaction (PCR) analysis in 100 suspected cases of infectious uveitis. *J Ophthal Inflamm Infect* 8, 2 (2018). <https://doi.org/10.1186/s12348-017-0144-1>
19. Van Gelder RN (2003) CME review: polymerase chain reaction diagnostics for posterior segment disease. *Retina* 23(4):445–452
20. Gupta V, Gupta A, Rao NA (2007) Intraocular tuberculosis—an update. *Surv Ophthalmol* 52(6):561–587
21. Figueira L, Fonseca S, Ladeira I, Duarte R (2017) Ocular tuberculosis: Position paper on diagnosis and treatment management. *Rev Port Pneumol* 23(1):31–38

Figures

Figure 1

Wide field fundus photo (pseudocolor fundus picture- Optos 200Tx imaging system, Optos PLC, Dunfermline, Scotland, UK) of right eye (Figure 1A) and left eye (Figure 1B) of a patient whose aqueous sample was tested positive for PCR MTB. It showed relatively clear media with snowball vitreous

exudates (black arrowheads) and attached retina. The right eye also had peripheral vitreous membranes (blue arrow).

Figure 2

Aqueous sample of a patient which showed positive result in Nested and Real time PCR for MTB

Figure 3

Vitreous sample of a patient which showed positive results in Nested and Real time PCR for MTB

Figure 4

Slit lamp photograph of right eye spill over anterior uveitis with broad based posterior synechiae