COVID-19 and Cognitive Impairment: A Cross-Sectional Clinic Based Study

Amira Siddig

AlNeelain University, Faculty of Medicine

Khabab Abbasher Hussien Mohamed Ahmed (✉ Khabab9722@gmail.com)

University of Khartoum, Faculty of Medicine <https://orcid.org/0000-0003-4608-5321>

Ahmed Abdelrahman

Daoud Research Group Charity Clinic

AlHussien Abbasher

AlYarmouk College

Abubaker Alsedig Abbasher

Zamzam University College

Mohammed Abbasher

AlNeelain University, Faculty of Medicine

Abbasher Hussien

University of Khartoum, Faculty of Medicine <https://orcid.org/0000-0002-9027-0420>

Research Article

Keywords: COVID-19, Cognitive impairment, Dementia, Sudan

Posted Date: December 17th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1178928/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Brain and Behavior on February 23rd, 2022.
See the published version at <https://doi.org/10.1002/brb3.2538>.

Abstract

Background: Patients with dementia are more prone to acquire COVID-19 infection. Patients with COVID-19 showed a tendency to develop cognitive impairment.

Objectives: We aimed to study the clinical manifestations of COVID-19 infection among adult Sudanese demented patients and the prevalence of cognitive impairment among adult Sudanese non demented patients.

Methodology: This is a descriptive cross-sectional study which took place in Sudan, Khartoum state in the period (September-December 2021) in a private neurology/psychiatry clinic. 135 adult Sudanese patients were included in this study and were divided into two groups. The first group consists of 100 patients with a known history of dementia that got infected recently with COVID-19, while the second group consists of 35 patients who developed some sort of cognitive impairment after recovering from COVID-19 infection. Regarding the second group, cognitive functions were assessed by senior consultant neurologist and senior consultant psychiatrist using a well validated neuropsychological measure.

Results: Out of 100 patients in the first group, females were 60 and males were 40. Age distribution is between 63 -98. The common presenting symptoms of COVID-19 among this group were: Cough and fever (90 patients), diarrhea and vomiting (5 patients), breathlessness (4 patients), coughing of blood (5 patients), convulsions (1 patient), paraplegia (1 patient) and hemiplegia (1 patient). Regarding the second group, Age distribution varied from 30 to 80 years. Cognitive functions impairment was noticed as follows: Memory recall (22%), memory recognition (23%), memory encoding (24%), processing speed (16%), executive functioning (19%), phonemic fluency (17%) and category fluency (17%).

Conclusion: Patients with dementia are more susceptible to develop COVID-19 infection. Patients with COVID-19 Infection are at risk of developing some sort of cognitive impairment after recovery.

Introduction

The SARS-CoV-2 virus causes Corona virus Disease (COVID-19). The majority of those infected with the virus will have mild to moderate respiratory symptoms and will recover without the need for specific treatment. Some, on the other hand, will become gravely unwell and require medical attention. People over the age of 65, as well as those with underlying medical disorders such as cardiovascular disease, diabetes, chronic respiratory disease, or cancer, are more likely to develop serious illnesses. COVID-19 can make anyone sick, and they can get seriously ill or die at any age [1].

Dementia is a syndrome that causes cognitive function (i.e. the ability to process thoughts) to deteriorate beyond what would be expected from biological aging. Memory, thought, orientation, comprehension, computation, learning capacity, language, and judgment are all affected. The ability to think is unaffected. Changes in mood, emotional control, behavior, or motivation are frequently accompanied by, and sometimes preceded by, impairment in cognitive performance [2].

Dementia can be caused by a number of diseases and injuries that damage the brain either directly or indirectly, such as Alzheimer's disease or stroke [2].

Dementia is the sixth greatest cause of death worldwide, and one of the top causes of impairment and dependency among the elderly. Dementia has medical, psychological, social, and economic consequences for people with dementia, as well as their caregivers, families, and society in general. Dementia is widely misunderstood and stigmatized, leading in delays in diagnosis and treatment [2].

Dementia has become much more common in recent decades around the world. Over 50 million individuals worldwide suffer from dementia, with that number expected to rise to 152 million by 2050 [3]. People with dementia have become the most vulnerable population due to aging, weakness, reduced autonomy, chronic immunological disorders [4], and other chronic diseases, particularly comorbidities of vascular diseases [5]. Patients with dementia are sensitive to viral infections such as respiratory syncytial virus disease, severe acute respiratory syndrome coronavirus, and influenza A, according to epidemiological research [6]. Their post-infection clinical result is worse than the normal population [7,8].

The incidence of seizures, psychosis, and memory issues among Covid-19 patients is yet unknown. According to some research, up to 84 percent of individuals with acute Covid-19 recover with mental disorientation and abrupt mood fluctuations. Even if the true rate of long-term brain damage is lower, because so many people have been infected, the number of people with long-term cognitive impairments will be significant [9]. In this study we aimed to study the clinical manifestations of COVID-19 infection among Sudanese demented patients and the prevalence of cognitive impairment among Sudanese non demented patients.

Methods And Materials

This is descriptive cross-sectional study which was conducted in Sudan, Khartoum state in a private neurology/psychiatry clinic. Total coverage of participants who fulfilled the inclusion criteria (n= 135) was done in the study period (from September-December 2021). All adult Sudanese patients who are well-known cases of dementia and developed COVID-19 infection were put in the first study group. All adult Sudanese patients with no past medical history of cognitive impairment or dementia and who developed cognitive impairment after recovering from COVID-19 infection were included in the second study group.

The first group consisted of 100 adult Sudanese patients who are already know cases of dementia and had COVID-19 infection. A well-validated semi-structured questionnaire was used. Full detailed history and proper clinical examination were done for each patient. The history includes the following:

1. Personal data like (name, age, sex, residence, occupation).
2. Duration of dementia.
3. History of diabetes mellitus, hypertension, ischemic heart disease, renal diseases and malignancy.

4. Symptoms in favor of involvement of:

- Respiratory system; cough, breathlessness.
- Gastrointestinal: vomiting and diarrhea.
- CNS: convulsion, weakness of upper and lower limbs.
- Cardiovascular system: symptoms of heart failure and symptoms of ischemic heart disease.

The clinical examination includes systemic examination looking for signs in favor of involvement of respiratory, cardiac, gastrointestinal or central nervous system (CNS). Cognitive impairment was also assessed. Oxygen saturation was observed. Overall mortality was reported.

The second group includes 35 patients who survived COVID-19 infection and they had no past medical history of dementia or cognitive impairment. All patients were adult Sudanese and they tended to speak Arabic. The same history and clinical examination were done and we stress on assessment of cognitive functions impairment by senior consultant neurologist and senior consultant psychiatrist.

Cognitive functions impairment was assessed using a well validated neuropsychological measures including:

1. Memory recall, memory recognition and memory encoding.
2. Phonemic fluency and category fluency.
3. Processing speed and executive functioning.

Results

Out of 100 patients with dementia, females were 60 and males were 40. Age distribution was between 63-98. Distribution of chronic diseases was found as follow:

Diabetes mellitus (40 patients), hypertension (20 patients), ischemic heart disease (5 patients), renal disease (3 patients) and cancer (2 patients). The common presenting symptoms were:

- Cough and fever (90 patients).
- Diarrhea and vomiting (5 patients).
- Breathlessness (4 patients)
- Coughing of blood (5 patients).
- Convulsion (1 patient).
- Paraplegia (1 patient).

-Hemiplegia (1 patient).

-Brain MRI showed evidence of brain atrophy in 76% of the patients, while 30% had small vessel disease and 5% had multi infarcts.

Vascular dementia was observed in 30% of the patients, Alzheimer in 20%, senile dementia in 10%, non-senile dementia in 10%, post traumatic dementia in 2% while 28% had dementia of unknown origin. Low oxygen in the blood was reported in 80% of the patients. Out of the 100 patients 49 died.

The second group consists of 35 patients who survived COVID-19 infection and they had no past history of dementia. Age distribution varies from 30 to 80 years.

Diabetes mellitus was observed in 4 patients, hypertension in 3 patients, 3 patients have ischemic heart disease, 2 patients have renal diseases and one patient have cancer. Respiratory symptoms were the commonest presentation (30 patients), followed by cardiovascular symptoms (9 patients), gastrointestinal symptoms (8 patients) and CNS (4 patients). Brain MRI was normal in 30 patients and showed evidence of small vessel disease in 5 patients. Low blood oxygen was noticed in 30 patients.

Cognitive functions impairment was noticed as follow:

1-Memory recall (22%)

2-Memory recognition (23%)

3-Memory encoding (24%)

4-Processing speed (16%)

5-Executive functioning (19%)

6-Phonemic fluency (17%)

7-Category fluency (17%)

Discussion

Dementia is a clinical syndrome characterized by memory disturbance and cognitive impairment. Vascular dementia is the commonest cause, followed by Alzheimer's disease, senile dementia, presenile dementia and post traumatic dementia. Rare causes include Huntington chorea, lewy body dementia, frontotemporal dementia and advanced stage of Parkinson's disease [10]. Dementia is associated also with advanced age, hypertension, diabetes and Ischemic heart disease. Patients with dementia are more prone to develop COVID-19 infection twice more than dementia patients [11]. There are many hypotheses that can explain the increased incidence of COVID-19 infection among dementia patients, one of these is that patients with dementia cannot stick to the preventive measures that can minimize spread of COVID-

19 infection like wearing masks, frequent washing of hands and social distancing. The second hypothesis is that patients with dementia, especially vascular dementia, tend to have damage to the blood brain barrier so they become more vulnerable to infection (ex. bacteria, fungal and viral infections) that can spread to the brain [12,13]. Patients with dementia who have COVID-19 infection will have a chance to experience a severe form of the disease more than non-demented patients this due to the fact that dementia is associated with diabetes, hypertension and ischemic heart disease. All these factors increase morbidity and mortality among patients with COVID-19 Infection [14]. Hypoxia, superadded brain infection, drugs used with anesthesia and damage to the blood brain barrier all these factors can increase mortality among demented patients who developed COVID-19 infection [15].

Regarding the second group, cognitive function was assessed using well-validated semi-structured neuropsychiatric measures. We tend to see the patients who were referred from the isolation centers monthly for six months. All of them were adult Sudanese patients, they had no history of dementia, they were speaking Arabic fluently and the test for COVID-19 antibodies was positive. The common cognitive function impairments which were noticed were as follows: memory recall, memory recognition, memory encoding, followed by executive functioning and processing speed and a considerable number of patients had category fluency and phonetic fluency impairment. At the end of this study questions were raised regarding the second group: Are these cognitive impairments transient or will they be permanent? Will those patients end with a full bloom picture of dementia in the future? Does the ongoing COVID-19 pandemic cause a significant increase in the number of dementia patients in the long term? Does COVID-19 pandemic increase numbers of patients with dementia and Alzheimer's disease worldwide?

So we need more studies to know the long-term impact of COVID-19 on the brain's health.

Conclusion

COVID-19 infection most likely causes massive acceleration in cognitive decline, especially in patients with dementia.

Declarations

Conflicts of interest:

All authors declare that there are no conflicts of interest.

Funding:

There was no fund.

Informed consents:

Both verbal and written consents were obtained from each patient and/or care giver before starting the study.

Ethical approval;

Ethical approval was obtained from the Sudan State Ministry of Health.

Authors' contributions:

All authors participated in planning the study, data collection, analysis and writing draft.

Acknowledgement:

Not applicable.

References

1. https://www.who.int/health-topics/coronavirus#tab=tab_1
2. <https://www.who.int/news-room/fact-sheets/detail/dementia>
3. (2019) Alzheimer's Association Report: 2019 Alzheimer's disease facts and figures. *Alzheimers Dement* 3, 321–387.
4. Rogers NT , Steptoe A , Cadar D (2017) Frailty is an independent predictor of incident dementia: Evidence from the English Longitudinal Study of Ageing. *Sci Rep* 1, 15746.
5. Nikolich-Zugich J , Knox KS , Rios CT , Natt B , Bhattacharya D , Fain MJ (2020) SARS-CoV-2 and COVID-19 in older adults: what we may expect regarding pathogenesis, immune responses, and outcomes. *Geroscience* 2, 505–514.
6. Azarpazhooh MR , Amiri A , Morovatdar N , Steinwender S , Rezaei Ardani A , Yassi N , Biller J , Stranges S , Tokazebani Belasi M , Neya SK , Khorram B , Sheikh Andalibi MS , Arsang-Jang S , Mokhber N , Di Napoli M (2020) Correlations between COVID-19 and burden of dementia: An ecological study and review of literature. *J Neurol Sci* 416, 117013.
7. Killen A , Olsen K , Mckeith IG , Thomas AJ , O'Brien JT , Donaghy P , Taylor J (2020) The challenges of COVID-19 for people with dementia with Lewy bodies and family caregivers. *Int J Geriatr Psychiatry* 10, 1002/gps.5393.
8. Iaboni A , Cockburn A , Marcil M , Rodrigues K , Marshall C , Garcia MA , Quirt H , Reynolds KB , Keren R , Flint AJ (2020) Achieving safe, effective, and compassionate quarantine or isolation of older

- adults with dementia in nursing homes. *Am J Geriatr Psychiatry* 28, 835–838.
9. Covid-19 may result in brain damage and increase the risk of dementia <https://www.clinicaltrialsarena.com/comment/covid-19-brain-damage-dementia-risk/>
 10. Hugo J, Ganguli M. Dementia and Cognitive Impairment. *Clinics in Geriatric Medicine*. 2014;30(3):421-442.
 11. Livingston G, Huntley J, Sommerlad A, Ames D, Ballard C, Banerjee S et al. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *The Lancet*. 2020;396(10248):413-446.
 12. Ryoo N, Pyun J, Baek M, Suh J, Kang M, Wang M et al. Coping with Dementia in the Middle of the COVID-19 Pandemic. *Journal of Korean Medical Science*. 2020;35(42).
 13. Hardan L, Filtchev D, Kassem R, Bourgi R, Lukomska-Szymanska M, Tarhini H et al. COVID-19 and Alzheimer's Disease: A Literature Review. *Medicina*. 2021;57(11):1159.
 14. Matias-Guiu J, Pytel V, Matías-Guiu J. Death Rate Due to COVID-19 in Alzheimer's Disease and Frontotemporal Dementia. *Journal of Alzheimer's Disease*. 2020;78(2):537-541.
 15. Fuh J, Yang C. Exposure to general anesthesia and the risk of dementia. *Journal of Pain Research*. 2015;711.