Parental Stress around Ophthalmological Health Conditions: A Systematic Review of Literature Protocol

JULIO CESAR SOUZA-SILVA (✉ juliocesarsouzasilva@hotmail.com)

Universidade Federal de Goiás <https://orcid.org/0000-0002-4545-0019>

Cleusa Alves Martins

Universidade Federal de Goiás Faculdade de Medicina

Marco Túlio Antonio Garciazapata

Universidade Federal de Goiás Faculdade de Medicina

Maria Alves Barbosa

Universidade Federal de Goiás Faculdade de Medicina

Protocol

Keywords: Parenting Stress, Ophthalmology, Ocular Diseases, Systematic Review, Eye Diseases

Posted Date: December 1st, 2020

DOI: <https://doi.org/10.21203/rs.2.21203/v2>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Systematic Reviews on August 13th, 2021.
See the published version at <https://doi.org/10.1186/s13643-021-01773-8>.

Abstract

Background

Parents can be psychologically impacted when children have eye diseases such as blindness, strabismus, and eye cancer. Stress can reduce the quality of parental care, and may be linked to the deterioration of parents' and children's mental and physical health, and family dynamics. No systematic literature review on parental stress in ophthalmology has been found to provide evidence synthesis capable of stimulating and defining new studies promoting research in this field. To address this important gap, the present review aims to synthesize evidence about approaches, methods, instruments, and results from research regarding ophthalmology-related parental stress.

Methods

Primary epidemiological observational studies should be original in addressing parental stress caused by ophthalmological health conditions in children, and their originality should be explained in both the summary and full text. They should present the characteristics of the study population, and the clinical and ophthalmic characterizations of children.

MEDLINE (via Ovid), EMBASE, PsycINFO, Google Scholar, and Gray Literature (PsycEXTRA, NTIS and OpenSINGLE) will be searched. Controlled vocabulary, Boolean operators, and defined search strategies will be used. There will be no restrictions on the studies' publication language, which will be selected in two screening stages. Two reviewers will independently retrieve full-text studies, assess methodological quality, and extract data. Data available through December 2021 will be considered for inclusion.

Discussion

The socioeconomic characterization of the participants, the identification of which ophthalmological diseases have been studied in relation to parental stress, and the knowledge of each instrument and methodology peculiarities potentially contribute to this study. The results may promote the development or enhancement of public policies focused on this specific theme, thereby providing the means for potential improvement of physical and mental health for parents and children with eye diseases.

Systematic review registration

The review protocol was registered in PROSPERO: number CRD42018094972.

Background

Description of the disease

The World Health Organization estimates that 1.4 million children—defined as individuals between 0 and 15 years old—are blind, with a prevalence of 0.3/1000 children in developed countries and 1.5/1000 in

poor/very poor communities. These children face a lifetime of blindness, which equates to a combined estimated 75 million blind years (blind individuals x life expectancy) [1].

Each year, five hundred thousand children (approximately one per minute) are born blind or become blind before their fifth birthday. The causes of blindness, which vary according to geographic region and socioeconomic status, include corneal scarring, cataracts, glaucoma, retinopathy of prematurity, and refractive errors. Although eye disease may not always lead to low vision or blindness, it can still cause significant parental stress stemming from the processes of diagnosis, treatment, and rehabilitation of vision. In all such cases, parental involvement and understanding are of paramount importance [1].

Being the parent of a child with any chronic disease can cause stress[2, 3], which is defined by Hans Selye as “a non-specific body response to any demand made upon it.” [4] Parental stress refers to a series of processes that lead to psychological and physiological reactions when attempting to adapt to parenting activities [3]. Early identification of increased stress in the parent-child system and suitable interventions can help reduce stress and diminish the frequency and intensity of the child’s emotional and behavioral disorders [5]. The reduction of parental stress improves parental health [6] and positively impacts the quality of care provided to sick children [7].

Conversely, increased and chronic levels of parental stress may put parents, children, and other family members at risk of adverse physical and psychological effects, such as anxiety and depression [8]. Among children, prolonged stress can lead to social incompetence, maladaptive behaviors, and cognitive impairment [2]. Stressful events often influence the pathogenesis of physical illness, causing negative affective states. In turn, these directly impact biological processes or care patterns, increasing the risk of becoming sick [9].

Changes in care may occur due to caregivers’ adaptations or coping responses to stress, such as increased smoking, decreased physical activity, insomnia, disinterest, and poor adherence to medical prescriptions and healthy habits. Furthermore, stress has been linked to major depression and poor adherence to treatments [9].

There are records of interference in the parent-child system due to parental stress around a child’s ophthalmological issues, but we did not find any systemic literature reviews on the subject. The current published studies on psychological/psychiatric diseases in the ophthalmological field [10-17] mainly focus on depression, anxiety, and parental burden and do not address the parental stress and child with visual disorders. A systematic literature review aimed at generating evidence and synthesis of what has been studied and published is needed to support the conception and planning of new studies. The evidence synthesis will guide new research and promote further studies to improve care for children with eye diseases and the physical and mental health of their parents.

Eye disorders in children and psychological implications for parents and family

During pregnancy, parents idealize the birth of a perfect child [18]. The birth of a blind child with strabismus, glaucoma, or congenital eye diseases can create a discrepancy between the idealized and the real child. One aspect of motherhood necessary to establish a healthy mother-child relationship is being able to deal with such discrepancies. Failure in overcoming expectation-reality discrepancies can lead mothers to become depressed, distance themselves from the child, and become unable to provide the warmth and love needed to promote the child's healthy development [10].

The establishment of an official diagnosis of a disabled child (blindness, for example) marks the occurrence of a family crisis, where members start expressing feelings of sadness, anger, guilt, helplessness, and isolation. Stress can be linked to the disruption of what was idealized versus the reality and the breakdown of family routine, and parents with very high levels of stress need to be guided to enable them to effectively provide higher levels of care [19].

The possibility of parental stress becoming an agent of change in the relationship between a parent and child raises questions as to what levels of resilience and coping strategies must be adopted by children's families to alleviate the possible psychological distress caused by children's eye diseases. It is unclear whether stress associated with the basal parental role of daily life is sufficiently robust to lead to clinical disturbances or whether the parental stress experienced by the parents of children with eye diseases correlates with psychological conditions that are potentially harmful to the parents, child, or family health.

More research about parental stress related to ophthalmological disorders in children is necessary because (1) blindness affects over a million children worldwide, (2) stress is a known contributor to many serious health issues, and (3) quality of life and appropriate allocation of healthcare resources are high-priority issues. Systematic reviews will help translate knowledge into action and promote more relevant studies. The first step to progress research in this field is to create a protocol for the synthesized evidence on the methodology used by researchers to evaluate parental stress related to children's eye diseases; the research instruments; the psychometric characteristics of these conditions; and study design advantages and disadvantages, limitations, and peculiarities.

Review questions

The review questions for this systematic review of literature are as follows:

- i. What approaches have been used to research parental stress resulting from pediatric eye disease?
- ii. What methodologies are employed in studies of parental stress resulting from pediatric eye diseases?
- iii. What parental stress assessment tools are used in pediatric eye disease research?
- iv. What peculiarities and psychometric characteristics of parental stress assessment instruments are employed in pediatric eye disease research?

v. What are the main results of research conducted on parental stress resulting from pediatric eye diseases?

Objectives

To identify studies and their respective authors, as well as the bibliographical references related to parental stress in ophthalmology.

To characterize the sociodemographic aspects of participants in studies on parental stress in ophthalmology, as well as the clinical and ophthalmic conditions of their children.

To identify the methodological trajectory of selected studies on parental stress in ophthalmology.

To describe the methodology of the instrument—if present—used to measure parental stress, with emphasis on its psychometric characteristics (internal reliability, test and retest reliability, and validation criteria).

To highlight the main results and conclusions of authors of published research on parental stress around ophthalmology.

To detect the peculiarities of each study presented in the research on parental stress around ophthalmology.

Inclusion criteria

Population, exposure, comparator, outcome, and study design components (PECOS) to be analyzed in the studies:

i. Population/participants: father, mother, or both

ii. Exposure of interest: eye disease in the child resulting from severe reduction in visual acuity, dysfunction of extrinsic eye motility, cancer, congenital eye diseases, deforming / stigmatizing eye, and periocular changes or any eye disease with the potential to increase parental stress

iii. Comparator: no specific comparator

iv. Outcome: parental stress

v. Study design: primary epidemiological observational studies.

The study's originality must be explained in its summary and full text. Studies should be original in addressing parental stress (only father, mother, or both parents) caused by eye disease in children. The terms "parental stress" or "parenting stress" must be included in title, summary, or in the full text. Characterization of the study population and clinical and ophthalmic characterization of the children should be present. Regarding the type of research, experimental studies, reviews, editorials, comments,

mathematical models, methodological articles, expert opinions, and other methodological modalities will be excluded due to the methodological differences that may hinder making comparisons. Studies on parental stress in caregivers or other family members will also not be eligible. Similar studies that do not meet all the inclusion criteria, but that have relevant information, will be analyzed separately in the discussion section of the systematic review.

Methods

The proposed literature review will be conducted in accordance with the Joanna Briggs Institute methodology for systematic reviews of observational epidemiological studies that report prevalence and cumulative incidence data [20].

Study record

This systematic literature review protocol is an integral part of a doctoral dissertation, entitled “Parental Stress in Mothers and Quality of Life of Infants and Blind Children.” It has been registered in the international database PROSPERO (under registration number CRD42018094972) and is being reported according to the *Preferred Reporting Items for Systematic Reviews and Meta-Analyses Protocols* (PRISMA-P) [21, 22] (see checklist in Additional File 1).

Information Sources

The research strategy will be developed using *Medical Subject Headings* (MeSH) and *Elsevier Life Science Thesaurus* (Emtree) descriptors. Reviewers (JCSS and CAM) will systematically search in the following databases: MEDLINE (via Ovid), EMBASE, PsycINFO, and Google Scholar.

Other data sources, such as ProQuest Dissertations and Theses, databases, and gray literature (i.e. PsycEXTRA, NTIS and OpenSINGLE), will also be consulted. Data will be considered for inclusion until December 2021.

Search strategy

Searches will use a controlled vocabulary defined by initial search terms. Examples of search terms and search strategies are presented in Additional_File_2. The snowballing method [23] will be used to identify other studies from the references of selected articles. Search terms in Emtree will be matched with equivalent terms in MEDLINE, in order to be able to conduct a search in EMBASE. Searches will be designed to be conducted in MEDLINE and EMBASE and will be adapted to other electronic databases and gray literature, aiming always at achieving search equivalence. There is no time limit filter for the inclusion of articles in all databases. The search strategy will be developed by an information science expert, aiming at conducting a high specificity and sensitivity search for the review.

Study selection

Studies must present their originality in the summary and full text. Studies should be original in addressing parental stress (only father, mother, or both parents) caused by eye disease in children. Parental Stress (or Parenting Stress) must be written in the title, summary, or in the full text. Additionally, the characterization of the study population and clinical and ophthalmic characterization of the children should be present. The study search will be conducted by two independent researchers [24] in a two-stage selection process.

There will be no blind strategy for reviewers regarding authors' names, institutional affiliations, or country of origin of the analyzed studies.

The raw data obtained from the databases will be processed and duplicate studies eliminated using the ENDNOTE[®] (ENDNOTE X7, Thomson Reuters, USA) reference manager program.

Selection Phase I

In the first phase, the articles will undergo the Relevance Test 1 (RT-1); reviewers will analyze the title and abstract of the article, determining whether it fulfills the eligibility criteria listed in Table 1.

Selection Phase II

After meeting the RT-1 criteria, studies will be subjected to the Relevance Test 2 (RT-2). This verifies, more strictly, if there is presence of full text, characterization of the study population, and clinical and ophthalmic characterization of the children (see RT-2 in Table 2).

Excluded studies will be recorded in a separate table along with the reason for exclusion. "Results" will include a comment section containing the characteristics of each excluded study and the reason that led to their exclusion.

In phases I and II, articles having items evaluated as "not clear" will be the object of a joint analysis among reviewers. In cases where disagreement persists, a third reviewer will provide his/her opinion essential to the resolution of the case. Studies that meet all the inclusion criteria will be accepted for research. The agreement achieved among the reviewers of this systematic review will be assessed using Cohen's (k) kappa agreement index [25].

The results of the search will be reported in full in the final report and presented in a Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) flow diagram (see Figure 1) [26].

Assessment of Methodological Quality

The quality of the included articles will be assessed using the Joanna Briggs Institute assessment tool for methodological guidance for systematic reviews of observational epidemiological studies reporting prevalence and cumulative incidence data (see Additional_File_3) [20].

The methodological quality of the included articles will be assessed by two reviewers. In case of disagreement regarding the quality of a particular study, a third reviewer will be the tiebreaker, providing the final assessment. There is no intention of conducting meta-analysis of the data.

Data Extraction and Management

Data Extraction

Data will be extracted from papers included in the scoping review by the two independent reviewers using a data extraction tool developed by the reviewers. In this phase, the text of the article will be analyzed in full, elaborating a standardized cataloging form to record product information – see form in Additional_File_4.

Additional relevant information not foreseen in the fields above and the peculiarities of each study will be recorded in the “observations” field.

If some information or detail is missing from the article, the study will be included, although the absence of data will be noted.

In case of lack/loss/uncertainty of information, the author of the study will be contacted by email three times. If the problem persists, the study will be excluded.

Data Synthesis

Because our study focuses on methodological analysis by assessing approaches, methodologies, instrument characteristics, and psychometric characteristics, there will be no combination of individual outcomes—that is, no statistical calculations or meta-analysis of results will be performed. This will generate a measurement capable of summarizing the outcomes. We will critically analyze the data and quality of the studies included, providing descriptive tabulations and summaries. The results extracted from the cataloging form will be categorized and compared between the different studies. If any included studies show important methodological differences, a sensitivity analysis will be carried out. Gaps and limitations in the description or methodology will be noted and discussed.

Protocol changes

Any substantive change to this protocol will be recorded upon its occurrence, in PROSPERO, and documented in the final publication.

Discussion

To our knowledge, this will be the first systematic literature review related to a comparative assessment of approaches, methods, and results in the research on parental stress around ophthalmology.

Systematic analysis and synthesis of peculiarities, time of administration, cost, reliability, and stability can offer researchers support for instrument analysis for choosing the best match for their research. Psycho-ophthalmology is a field linking ophthalmology and psychiatry [27]. Few studies related to parental stress have been conducted on this subject. The socioeconomic characterization of the participants, the identification of which ophthalmological diseases have been studied in relation to parental stress, and the knowledge of each instrument and methodology peculiarities potentially contribute to this study. The results may promote the development or enhancement of public policies focused on this specific theme, thereby providing the means for potential improvement of physical and mental health for parents and children with eye diseases.

This review protocol may have potential limitations if attempted to predict the evaluation of psychometric characteristics, such as equivalence and construct analysis [28]; however, as it does not attempt to do so, it keeps the specific evaluation as the object of study of future research.

Abbreviations

PS: Parental stress; NHLBI: National Heart, Lung, and Blood Institute; CAPES: *Coordenação de Aperfeiçoamento de Pessoal de Nível Superior* (Coordination for the Improvement of Higher Education Personnel); FAPEG: *Fundação de Amparo à Pesquisa do Estado de Goiás* (Research Support Foundation of the State of Goiás); ROP: Retinopathy of prematurity; RP: Retinitis pigmentosa.

Declarations

Ethical Approval and Consent for Participation

This systematic review is an integral part of a doctoral dissertation and the research project was approved by the Research Ethics Committee of Clinical Hospital of the Federal University of Goiás and published in *Plataforma Brasil* CAAE 90833718.2.0000.5078 under opinion number 2.746.572.

Consent for publication

Not applicable

Availability of data and materials

Not applicable.

Competing Interests

The authors declare no conflicts of interest.

Funding

This systematic literature review was funded by the authors themselves.

Author Contributions

JCSS is the review guarantor and first reviewer, working on all phases of the systematic literature review. CAM is the second reviewer. MAB and CAM contributed to the elaboration of the protocol, research question, research strategy and final revision of the protocol text. MTAG contributed to the process of evaluation and synthesis of evidence quality. MAB and MTAG will compose the arbitration panel and complete the revision of the protocol text.

Acknowledgments

This study was financed by the Coordination for the Improvement of Higher Education Personnel (CAPES) and the Research Support Foundation of the State of Goiás (FAPEG), providing a Ph.D. scholarship to the first author (reference number 88887.295574/2018-00).

This study received financial support for publication from the Federal University of Goiás (UFG).

We are grateful to Pollyanna Moreira, who acted as the research assistant and was responsible for collecting data from mothers.

Authors' Information

Not applicable

References

1. World Health Organization. Global Initiative for the Elimination of Avoidable Blindness: Action Plan 2006-2011. Geneva: World Health Organization; 2007.
2. Golfenshtein N, Srulovici E, Medoff-Cooper B. Investigating parenting stress across pediatric health conditions - A systematic review. *Issues Compr Pediatr Nurs*. 2015;1–49.
3. Deater-Deckard K, Panneton R. Parental stress and early child development: Adaptive and maladaptive outcomes. Cham: Springer; 2017.
4. Selye H. The evolution of the stress concept. *Am Sci*. 1973; 61: 692–9.
5. Abidin RR. Parenting Stress Index™, (PSI™-4) Professional Manual. 4 ed ed. Lutz, Florida: Par Inc.; 2012.
6. Golfenshtein N, Srulovici E, Deatrck JA. Interventions for reducing parenting stress in families with pediatric conditions: An integrative review. *J Fam Nurs*. 2016; 22:460–92.
7. Chaplin TM, Turpyn CC, Fischer S, Martelli AM, Ross CE, Leichtweis RN, et al. Parenting-focused mindfulness intervention reduces stress and improves parenting in highly stressed mothers of adolescents. *Mindfulness*. 2018;1-13.
8. Vismara L, Rolle L, Agostini F, Sechi C, Fenaroli V, Molgora S, et al. Perinatal parenting stress, anxiety, and depression outcomes in first-time mothers and fathers: A 3- to 6-months postpartum follow-up

- study. *Front Psychol.* 2016; 7:938.
9. Cohen S, Janicki-Deverts D, Miller GE. Psychological stress and disease. *Jama.* 2007;298(14):1685-7.
 10. Tolchin JG, Lederman ME. Congenital (infantile) esotropia: psychiatric aspects. *J Pediat Ophth Strab.* 1978;15(3):160-3.
 11. Bambara JK, Owsley C, Wadley V, Martin R, Porter C, Dreer LE. Family caregiver social problem-solving abilities and adjustment to caring for a relative with vision loss. *Invest Ophthalmol Vis Sci.* 2009;50(4):1585-92.
 12. Duman NS, Gökten ES, Duman R, Duman R, Çevik SG. Evaluation of depression and anxiety levels in mothers of babies' following due to premature Rrtinopathy. *Arch Psychiat Nurs.* 2018;32(3):439-43.
 13. Hahm B-J, Shin Y-W, Shim E-J, Jeon HJ, Seo J-M, Chung H, et al. Depression and vision-related quality of life in patients with retinitis pigmentosa. *Brit J Ophthalmol.* 2008.
 14. Zhang X, Bullard KM, Cotch MF, Wilson MR, Rovner BW, McGwin G, Jr., et al. Association between depression and functional vision loss in persons 20 years of age or older in the United States, NHANES 2005-2008. *JAMA Ophthalmol.* 2013; 131:573–81.
 15. Drews C, Celano M, Plager DA, Lambert SR. Parenting stress among caregivers of children with congenital cataracts. *J Am Assoc Pediat Ophthalmol Strab.* 2003;7(4):244-50.
 16. Celano M, Hartmann EE, Drews-Botsch CD. Parenting stress in the infant aphakia treatment study. *J Pediatr Psychol.* 2013 Jun;38(5):484-93.
 17. Wang LJ, Zhong WX, Ji XD, Chen J. Depression, caregiver burden and social support among caregivers of retinoblastoma patients in China. *Int J Nurs Pract.* 2016; 22:478–85.
 18. Solnit AJ, Stark MH. Mourning and the birth of a defective child. *Psychoanal Study Child.* 1961; 16:523–37.
 19. Fortier LM, Wanlass RL. Family crisis following the diagnosis of a handicapped child. *Fam Relat.* 1984:13–24.
 20. Munn Z, Moola S, Lisy K, Riitano D, Tufanaru C. Methodological guidance for systematic reviews of observational epidemiological studies reporting prevalence and cumulative incidence data. *Int J Evid Based Healthc.* 2015;13:147–53.
 21. Moher D, Shamseer L, Clarke M, Gherzi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev.* 2015 Jan 1; 4:1.
 22. Shamseer L, Moher D, Clarke M, Gherzi D, Liberati A, Petticrew M, et al. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ (Clinical research ed).* 2015;350:g7647.
 23. Tsafnat G, Glasziou P, Choong MK, Dunn A, Galgani F, Coiera E. Systematic review automation technologies. *Syst Rev.* 2014; 3:74.
 24. Higgins J. Green S. *Cochrane handbook for systematic reviews of interventions Version 5.1. 0.* The Cochrane Collaboration. Confidence intervals. 2011.

25. Cohen J. Weighted kappa: Nominal scale agreement provision for scaled disagreement or partial credit. *Psychol Bull.* 1968; 70:213–20.
26. Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: explanation and elaboration. *PLoS Med.* 2009;6(7):e1000100.
27. Rajsekar K, Rajsekar YL, Chaturvedi SK. Psycho ophthalmology: The interface between psychiatry and ophthalmology. *Indian J Psychiatry.* 1999; 41:186–96.
28. Souza ACd, Alexandre NMC, Guirardello EdB. Psychometric properties in instruments evaluation of reliability and validity. *Epidemiol Serv Saude.* 2017; 26:649–59.

Tables

Table 1. Relevance Test 1 (RT-1) of studies included in the systematic review

	Yes	Not clear	No
Is the research original?			
Does the population include only father, only mother, or both parents?			
Does the child have an eye disease/condition?			
Does the child's condition result in parental stress?			

Table 2. Relevance test 2 (RT-2) for the systematic literature review

	Yes	Not clear	No
Is the full text of the article available in databases?			
Is there a characterization of the study population?			
Is there a clinical and ophthalmic characterization of the children?			

Figures

Figure 1

Preferred Reporting Items for Systematic Reviews and Meta-analyses (PRISMA) flow diagram

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AdditionalFile1.docx](#)
- [AdditionalFile2.docx](#)
- [AdditionalFile3.docx](#)
- [AdditionalFile4.docx](#)