A Study of (2+1)-Dimensional Konopelchenko-Dubrovsky (KD) System: Closed-Form Solutions, Solitary Waves, Bifurcation Analysis and Quasi-Periodic Solution

Sachin Kumar (✉ sachinambariya@gmail.com)

University of Delhi <https://orcid.org/0000-0003-4451-3206>

Nikita Mann

University of Delhi

Harsha Kharbanda

University of Delhi

Research Article

Keywords: Konopelchenko-Dubrovsky equations, generalized exponential rational function method, soliton solutions, bifurcation analysis, quasi-periodic solution.

Posted Date: January 5th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1184920/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

A study of (2+1)-dimensional Konopelchenko-Dubrovsky (KD) system: Closed-form solutions, Solitary waves, Bifurcation analysis and Quasi-periodic solution

Sachin Kumar^{a*}, Nikita Mann^{a†} and Harsha Kharbanda^{a‡}

^aDepartment of Mathematics, Faculty of Mathematical Sciences, University of Delhi, Delhi-110007, India

Abstract

Nonlinear evolution equations (NLEEs) are extensively used to establish the elementary propositions of natural circumstances. In this work, we study the Konopelchenko-Dubrovsky (KD) equation which depicts non-linear waves in mathematical physics with weak dispersion. The considered model is investigated using the combination of generalized exponential rational function (GERF) method and dynamical system method. The GERF method is utilized to generate closed-form invariant solutions to the (2+1)-dimensional KD model in terms of trigonometric, hyperbolic, and exponential forms with the assistance of symbolic computations. Moreover, three-dimensional graphics are displayed to depict the behavior of obtained solitary wave solutions. The model is observed to have single and multiple soliton profiles, kink-wave profiles, and periodic oscillating nonlinear waves. These generated solutions have never been published in the literature. All the newly generated soliton solutions are checked by putting them back into the associated system with the soft computation via Wolfram Mathematica. Moreover, the system is converted into a planar dynamical system using a certain transformation and the analysis of bifurcation is examined. Furthermore, the quasi-periodic solution is investigated numerically for the perturbed system by inserting definite periodic forces into the considered model. With regard to the parameter of the perturbed model, two-dimensional and three-dimensional phase portraits are plotted.

Keywords: Konopelchenko-Dubrovsky equations; generalized exponential rational function method; soliton solutions; bifurcation analysis; quasi-periodic solution.

1 Introduction

Nonlinear evolution equations (NLEEs) have extensive significance in the area of applied mathematics and physics. Finding the exact solutions for NLEE is an essential task as NLEE describes numerous phenomenon in nonlinear dynamics, engineering, optical fibre, plasma physics, fluid mechanics, natural sciences, etc. A large number of researchers and mathematicians have developed various effective techniques for computing exact solutions of NLPDEs (nonlinear partial differential equations), for instance, tanh function method [1], Hirota's bilinear method [2,3], the Jacobi elliptic function expansion method [4], Kudryashov method [5], the $\frac{G'}{G}$ -expansion method [6], Darboux transformation method [7], the Backlund transformation method [8], the inverse scattering method [9], Lie-symmetry analysis [10], multiple exp-function method, and many others. Among these techniques, GERF method [11–14] is

*sachinambariya@gmail.com (Corresponding author)

†nikitamann.21@gmail.com

‡ssdn0112@gmail.com

very effective, robust and straightforward approach for finding the abundant exact soliton-form solutions of various NLPDEs.

The dynamics of NLPDEs grants us to understand and predict the acceptable structures of the associated complex nonlinear systems. A soliton or solitary wave is the particle-like object with the finite energy and amplitude, which save its form during propagation and restore it after the collision with another solitons. Nowadays, as a consequence, it is a very hot subject matter to derive the exact closed form solutions of NLPDEs. The soliton-form solutions of such type of NLPDEs are extensively favorable in the various areas such as nonlinear sciences, mathematical physics, plasma physics, applied mathematics, engineering, applied sciences and nonlinear dynamics.

In the recent study, we will analyze the following (2+1)-dimensional Konopelchenko-Dubrovsky (KD) model

$$\begin{aligned} u_y &= v_x, \\ u_t - u_{xxx} - 6buu_x + \frac{3}{2}a^2u^2u_x - 3v_y + 3au_xv &= 0, \end{aligned} \quad (1)$$

where u and v are the differentiable functions with respect to x , y and t variables. Here, a and b are arbitrary constants.

Many reseachers have used some productive techniques to investigate the exact analytical solutions of the KD system. Konopelchenko and Dubrovsky [15] derived (2+1)-dimensional KD (Konopelchenko-Dubrovsky) equation in 1984. In 2014 Kumar et. al [16] used Lie symmetry analysis as well as travelling wave hypothesis to extract solutions to KD equations. In 2019 Alfalqi et al. [17] applied the modified simplest equation method and B-spline method to KD-equation. Recently, Khater et al. [18] implement modified auxiliary equation technique to this system to find analytical traveling wave solutions. Ren et al. [19] in 2016 obtained the non-local symmetries for the K-D equation with the truncated Painleve method and the Mobius (conformal invariant form.

Recently, the investigation of differential equations through the bifurcation analysis has become an important topic in the field of research. Bifurcation is a rapid quantitative shift in the model with a gentle change in the values of parameters. The exploration of the dynamics of nonlinear periodic forms is a significant part for the investigation of the physical propositions in detail. For example, the occurrence of homoclinic orbits, smooth heteroclinic orbits and periodic orbits for travelling wave models describes the periodic wave solutions, oscillatory travelling wave solutions, smooth kink wave solutions and smooth solitary solutions for considered PDEs, respectively. One can refer to [20–24] for the study of recent work in this field.

In this research article, the study of $(2 + 1)$ -dimensional Konopelchenko-Dubrovsky (KD) model (1) is investigated using GERF (generalized exponential rational function) technique. The primary goal of this research study is to construct abundant exact analytical closed-form solutions for the system (1). Moreover, we exhibit the dynamics of solitary wave profiles of some soliton solutions in three dimensional and two dimensional graphics in numerical simulations, and hence we believe that the evolutionary profile dynamics of generated exact closed-form solutions are very impressive and advantageous for physical phenomena. The obtained soliton solutions are entirely new, novel and have never been reported in the previous findings. Also, we have investigated the dynamical system (1) using the bifurcation analysis. Numerically, the quasi-periodic solution is investigated for the perturbed system by including definite forces to the considered model, and two-dimensional and three-dimensional phase portraits are also plotted for the perturbed system.

The strategy of this article is organised as follows: Section 2 deals with the introduction and methodology of GERF approach is presented. In section 3, we find exact travelling wave solution of KD equations. This section also includes some particular three-dimensional shapes which provide more explanation to the behavior of these generated solutions. Graphically, multi-soliton structures,

kink-wave structures, and the interaction of multi-soliton and kink wave solution have been observed for some soliton solutions. Section 4 deals with the bifurcation analysis of the dynamical system of eq. (1), and relative phase portraits are plotted for the considered system. Section 5 is related to the investigation of quasi-periodic solution for the perturbed system by inserting perturbation term to the associated model (1). Finally, the conclusion is given at the end.

2 Methodology of GEF method

This GEF method was introduced by Ghanbari and Inc [11]. We will provide an explanation of GEF technique step-wise in this section:

- Let us consider the system of two nonlinear PDEs including three variables x , y and t given as

$$\begin{aligned}\mathcal{M}(u, v, u_x, u_y, u_t, v_x, v_y, v_t, u_{xx}, u_{xt}v_x, u_{yx}\dots) &= 0. \\ \mathcal{N}(u, v, u_x, u_y, u_t, v_x, v_y, v_t, u_{xx}, u_{xt}v_x, u_{yx}\dots) &= 0.\end{aligned}\quad (2)$$

Here, $u(x, y, t) = U(\xi)$ and $v(x, y, t) = V(\xi)$ are the wave transformations where $\xi = \alpha x + \beta y - \mu t$ is used to obtain a system of ODEs (ordinary differential equations) which further provides an ordinary differential as

$$\mathcal{W}(U, U', U'', \dots) = 0, \quad (3)$$

where $U = U(\xi)$, $U' = \frac{dU}{d\xi}$ and α , β and μ are constants, to be calculated later.

- Assume that the solitary wave solution of (3) can be represented as

$$U(\xi) = R_0 + \sum_{k=1}^M R_k \Psi(\xi)^k + \sum_{k=1}^M S_k \Psi(\xi)^{-k}, \quad (4)$$

where

$$\Psi(\xi) = \frac{r_1 e^{s_1 \xi} + r_2 e^{s_2 \xi}}{r_3 e^{s_3 \xi} + r_4 e^{s_4 \xi}}, \quad (5)$$

and r_i , s_i ($1 \leq i \leq 4$), R_0 , R_k and S_k ($1 \leq k \leq M$) are constants to be calculated and M (a positive integer) is evaluated by homogeneous balancing method.

- Substituting (4) into (3) plugging with (5) and arranging all the terms yields

$$\mathfrak{A}(e^{s_1 \xi}, e^{s_2 \xi}, e^{s_3 \xi}, e^{s_4 \xi}, \dots) = 0. \quad (6)$$

Equalizing the coefficients of $\mathfrak{A}$ to zero, yields a set of algebraic equations.

- With the assistance of Mathematica, we find the values of the coefficients R_0 , R_k , S_k , α , β and μ by solving above mentioned algebraic equations.
- Inserting all these values into (4) we get the solution of determining equation (3). Accordingly we can get other function and hence we achieve some new solitary wave solutions of (2).

3 GEF method - Applications

In this section, we apply the GEF technique to construct the exact closed-form solutions of (2+1)-dimensional Konopelchenko-Dubrovsky (KD) model (1).

3.1 GEF method for the (2+1)-D Konopelchenko-Dubrovsky(KD) model

Making use of wave transformation $u(x, y, t) = U(\xi)$ and $v(x, y, t) = V(\xi)$ where $\xi = \alpha x + \beta y - \mu t$ in eq.(1), we obtain the following system of ODEs

$$\begin{aligned} \beta U'(\xi) &= \alpha V'(\xi) \\ \frac{3}{2}a^2\alpha U(\xi)^2 U'(\xi) + 3a\alpha V(\xi)U'(\xi) - 6\alpha b U(\xi)U'(\xi) - \mu U'(\xi) - \alpha^3 U^{(3)}(\xi) - 3\beta V'(\xi) &= 0. \end{aligned} \quad (7)$$

Integrating the first equation we find,

$$\beta U(\xi) = \alpha V(\xi) \quad (8)$$

Putting it into the second equation of (7) and integrating the resulting equation, and neglecting the constant of integration, the following ODE is obtained as

$$\alpha^3 U''(\xi) - \frac{a^2}{2}\alpha U(\xi)^3 + \frac{3}{2}(2\alpha b - a\beta)U(\xi)^2 + \left(\frac{3\beta^2}{\alpha} + \mu\right)U(\xi) = 0. \quad (9)$$

Using balancing principle on terms $U^{(3)}$ and U'' of the equation (9), we get $M + 2 = 3M$ which yields $M = 1$. Employing $M = 1$ in (4), we obtain the trial solution given as

$$U(\xi) = R_0 + R_1 \Psi(\xi) + \frac{S_1}{\Psi(\xi)}. \quad (10)$$

By using equation (10) into (9) and according to the step of GEF method, the following cases are considered for finding the exact solitary wave solutions of (1) with the assistance of Mathematica.

Family 1: For $[r_1, r_2, r_3, r_4] = [1, -3, -1, 1]$ and $[s_1, s_2, s_3, s_4] = [1, -1, 1, -1]$, eq. (5) yields

$$\Psi(\xi) = \coth(\xi) - 2. \quad (11)$$

With the assistance of soft computation via Mathematica, we solve algebraic equation for obtaining the values of parameters and hence following set of solutions can be achieved.

Solution set 1.1:

$$R_0 = \mp \frac{6\alpha}{a}, \quad R_1 = \mp \frac{2\alpha}{a}, \quad S_1 = 0, \quad \beta = \frac{2(b\alpha \pm a\alpha^2)}{a}, \quad \mu = -\frac{4(4a^2\alpha^3 \pm 6a\alpha^2b + 3\alpha b^2)}{a^2}.$$

Substituting the values of above known-constant parameters into eq.(10) and plugging it with eq. (9), we obtain the expression for U as

$$U(\xi) = \mp \frac{6\alpha}{a} \mp \frac{2\alpha(\coth(\xi) - 2)}{a}. \quad (12)$$

Hence, from eq.(8), we obtain the expression for V as

$$V(\xi) = \mp \frac{6\beta}{a} \mp \frac{2\beta(\coth(\xi) - 2)}{a}. \quad (13)$$

Accordingly, the solution of (1) is obtained as

$$\begin{aligned} u(x, y, t) &= \mp \frac{6\alpha}{a} \mp \frac{2\alpha \left(\coth \left(\frac{4t(4a^2\alpha^3 \pm 6a\alpha^2b + 3\alpha b^2)}{a^2} + \frac{2y(\alpha b \pm a\alpha^2)}{a} + \alpha x \right) - 2 \right)}{a} \\ v(x, y, t) &= \mp \frac{12(\alpha b \pm a\alpha^2)}{a^2} \mp \frac{4(\alpha b \pm a\alpha^2) \left(\coth \left(\frac{4t(4a^2\alpha^3 \pm 6a\alpha^2b + 3\alpha b^2)}{a^2} + \frac{2y(\alpha b \pm a\alpha^2)}{a} + \alpha x \right) - 2 \right)}{a^2}. \end{aligned} \quad (14)$$

(a) 3D

(b) 2D

(c) 3D

(d) 2D

Figure 1: Interacting multisoliton and kink wave profile structure for solution (14) at $\alpha = 0.459$, $a = 2$, $b = -0.019$ under the range space $-20 \leq x \leq 20$, $-90 \leq y \leq 90$ and $t = 0.1$.

Solution set 1.2:

$$R_0 = \pm \frac{4\alpha}{a}, \quad R_1 = \pm \frac{2\alpha}{a}, \quad S_1 = 0, \quad \beta = \frac{2\alpha b}{a}, \quad \mu = \frac{2(a^2\alpha^3 - 6\alpha b^2)}{a^2}.$$

Substituting the values of above known-constants into eq.(10) and plugging it with eq. (9), we obtain the expression for U as

$$U(\xi) = \pm \frac{4\alpha}{a} \pm \frac{2\alpha(\coth(\xi) - 2)}{a}. \quad (15)$$

Hence, from eq.(8), we obtain the expression for V as

$$V(\xi) = \pm \frac{4\beta}{a} \pm \frac{2\beta(\coth(\xi) - 2)}{a}. \quad (16)$$

Accordingly, the solution of (1) is obtained as

$$\begin{aligned} u(x, y, t) &= \pm \frac{4\alpha}{a} \pm \frac{2\alpha \left(\coth \left(-\frac{2t(a^2\alpha^3 - 6\alpha b^2)}{a^2} + \frac{2\alpha by}{a} + \alpha x \right) - 2 \right)}{a} \\ v(x, y, t) &= \pm \frac{8\alpha b}{a^2} \pm \frac{4\alpha b \left(\coth \left(-\frac{2t(a^2\alpha^3 - 6\alpha b^2)}{a^2} + \frac{2\alpha by}{a} + \alpha x \right) - 2 \right)}{a^2}. \end{aligned} \quad (17)$$

Solution set 1.3:

$$R_0 = -\frac{2\alpha}{a}, \quad R_1 = -\frac{2\alpha}{a}, \quad S_1 = 0, \quad \beta = -\frac{2(a\alpha^2 - \alpha b)}{a}, \quad \mu = -\frac{4(4a^2\alpha^3 - 6a\alpha^2b + 3\alpha b^2)}{a^2}.$$

Substituting the values of known-constants mentioned above into eq. (10) and plugging it with eq. (9), we obtain the value for U as

$$U(\xi) = -\frac{2\alpha(\coth(\xi) - 2)}{a} - \frac{2\alpha}{a}. \quad (18)$$

Hence, from eq.(8), we obtain the expression for V as

$$V(\xi) = -\frac{2\beta(\coth(\xi) - 2)}{a} - \frac{2\beta}{a}. \quad (19)$$

Accordingly, the solution of (1) is obtained as

$$\begin{aligned} u(x, y, t) &= -\frac{2\alpha}{a} - \frac{2\alpha \left(\coth \left(\frac{4t(4a^2\alpha^3 - 6a\alpha^2b + 3\alpha b^2)}{a^2} - \frac{2y(a\alpha^2 - \alpha b)}{a} + \alpha x \right) - 2 \right)}{a} \\ v(x, y, t) &= \frac{4(a\alpha^2 - \alpha b)}{a^2} + \frac{4(a\alpha^2 - \alpha b) \left(\coth \left(\frac{4t(4a^2\alpha^3 - 6a\alpha^2b + 3\alpha b^2)}{a^2} - \frac{2y(a\alpha^2 - \alpha b)}{a} + \alpha x \right) - 2 \right)}{a^2}. \end{aligned} \quad (20)$$

Family 2: For $[r_1, r_2, r_3, r_4] = [5 + i, 5 - i, 1, 1]$ and $[s_1, s_2, s_3, s_4] = [-i, i, -i, i]$, eq. (5) yields

$$\Psi(\xi) = \tan(\xi) + 5. \quad (21)$$

With the assistance of soft computation via Mathematica, we solve algebraic equation for obtaining the values of parameters and hence following set of solutions can be achieved.

Solution set 2.1:

$$R_0 = -\frac{10\alpha}{a}, \quad R_1 = \frac{2\alpha}{a}, \quad S_1 = 0, \quad \beta = \frac{2\alpha b}{a}, \quad \mu = -\frac{2(a^2\alpha^3 + 6\alpha b^2)}{a^2}.$$

We substitute the values of known-constants mentioned above and expression (21) into eq.(10), and plugging it with eq. (9), we obtain the value for U as

$$U(\xi) = \frac{2\alpha(\tan(\xi) + 5)}{a} - \frac{10\alpha}{a}. \quad (22)$$

Hence, from eq.(8), we obtain the expression for V as

$$V(\xi) = \frac{2\beta(\tan(\xi) + 5)}{a} - \frac{10\beta}{a}. \quad (23)$$

Accordingly, the solution of eq. (1) is obtained as

$$\begin{aligned} u(x, y, t) &= \frac{2\alpha \left(\tan \left(\frac{2t(a^2\alpha^3 + 6\alpha b^2)}{a^2} + \frac{2\alpha by}{a} + \alpha x \right) + 5 \right)}{a} - \frac{10\alpha}{a} \\ v(x, y, t) &= \frac{4\alpha b \left(\tan \left(\frac{2t(a^2\alpha^3 + 6\alpha b^2)}{a^2} + \frac{2\alpha by}{a} + \alpha x \right) + 5 \right)}{a^2} - \frac{20\alpha b}{a^2}. \end{aligned} \quad (24)$$

(a) 3D

(b) 2D

(c) 3D

(d) 2D

Figure 2: Evolutionary profile dynamics of closed form solution (24) at $\alpha = 2$, $a = 3$, $b = 1$ under the range space $-20 \leq x \leq 20$, $-20 \leq y \leq 20$ and $t = 0.01$.

Solution set 2.2:

$$R_0 = -\frac{(10 + 2i)\alpha}{a}, \quad R_1 = \frac{2\alpha}{a}, \quad S_1 = 0, \quad \beta = \frac{2i(a\alpha^2 - iab)}{a}, \quad \mu = \frac{4(4a^2\alpha^3 - 6ia\alpha^2b - 3ab^2)}{a^2}.$$

We substitute the values of known-constants mentioned above and expression (21) into eq.(10), and plugging it with eq. (9), we obtain the value for U as

$$U(\xi) = \frac{2\alpha(\tan(\xi) + 5)}{a} - \frac{(10 + 2i)\alpha}{a}. \quad (25)$$

Hence, from eq.(8), we obtain the expression for V as

$$V(\xi) = \frac{2\beta(\tan(\xi) + 5)}{a} - \frac{(10 + 2i)\beta}{a}. \quad (26)$$

Accordingly, the solution of (1) is obtained as

$$\begin{aligned} u(x, y, t) &= \frac{2\alpha \left(5 + \tan \left(-\frac{4t(4a^2\alpha^3 - 6ia\alpha^2b - 3\alpha b^2)}{a^2} + \frac{2iy(a\alpha^2 - i\alpha b)}{a} + \alpha x \right) \right)}{a} - \frac{(10 + 2i)\alpha}{a} \\ v(x, y, t) &= \frac{4i(a\alpha^2 - i\alpha b) \left(5 + \tan \left(-\frac{4t(4a^2\alpha^3 - 6ia\alpha^2b - 3\alpha b^2)}{a^2} + \frac{2iy(a\alpha^2 - i\alpha b)}{a} + \alpha x \right) \right)}{a^2} + \frac{(4 - 20i)(a\alpha^2 - i\alpha b)}{a^2}. \end{aligned} \quad (27)$$

Solution set 2.3:

$$R_0 = \frac{(10 + 2i)\alpha}{a}, \quad R_1 = -\frac{2\alpha}{a}, \quad S_1 = 0, \quad \beta = -\frac{2i(a\alpha^2 + i\alpha b)}{a}, \quad \mu = \frac{4(4a^2\alpha^3 + 6ia\alpha^2b - 3\alpha b^2)}{a^2}.$$

We substitute the values of known-constants described above and expression (21) into eq.(10), and plugging it with eq. (9), we obtain the expression for U as

$$U(\xi) = -\frac{2\alpha(\tan(\xi) + 5)}{a} + \frac{(10 + 2i)\alpha}{a}. \quad (28)$$

Hence, from eq.(8), we obtain the expression for V as

$$V(\xi) = -\frac{2\beta(\tan(\xi) + 5)}{a} + \frac{(10 + 2i)\beta}{a}. \quad (29)$$

Accordingly, the solution of (1) is obtained as

$$\begin{aligned} u(x, y, t) &= \frac{(10 + 2i)\alpha}{a} - \frac{2\alpha \left(5 + \tan \left(-\frac{4t(4a^2\alpha^3 + 6ia\alpha^2b - 3\alpha b^2)}{a^2} - \frac{2iy(a\alpha^2 + i\alpha b)}{a} + \alpha x \right) \right)}{a} \\ v(x, y, t) &= \frac{4i(a\alpha^2 + i\alpha b) \left(5 + \tan \left(-\frac{4t(4a^2\alpha^3 + 6ia\alpha^2b - 3\alpha b^2)}{a^2} - \frac{2iy(a\alpha^2 + i\alpha b)}{a} + \alpha x \right) \right)}{a^2} + \frac{(4 - 20i)(a\alpha^2 + i\alpha b)}{a^2}. \end{aligned} \quad (30)$$

Family 3: For $[r_1, r_2, r_3, r_4] = [1, 0, 1, 1]$ and $[s_1, s_2, s_3, s_4] = [1, 0, 1, 0]$, eq. (5) yields

$$\Psi(\xi) = \frac{1}{1 + e^{-\xi}}. \quad (31)$$

With the assistance of soft computation via Mathematica, we solve algebraic equation for obtaining the values of parameters and hence following set of solutions can be achieved.

Solution set 3.1:

$$R_0 = \frac{\alpha}{a}, \quad R_1 = -\frac{2\alpha}{a}, \quad S_1 = 0, \quad \beta = \frac{2\alpha b}{a}, \quad \mu = \frac{\alpha(a^2\alpha^2 - 24b^2)}{2a^2}.$$

We substitute the values of known-constants described above and expression (31) into eq.(10), and plugging it with eq. (9), we obtain the expression for U as

$$U(\xi) = \frac{\alpha}{a} - \frac{2\alpha e^\xi}{a(e^\xi + 1)}. \quad (32)$$

Hence, from eq.(8), we obtain the expression for V as

$$V(\xi) = \frac{\beta}{a} - \frac{2\beta e^\xi}{a(e^\xi + 1)}. \quad (33)$$

Accordingly, the solution of eq. (1) is obtained as

$$\begin{aligned} u(x, y, t) &= \frac{\alpha}{a} - \frac{2\alpha \exp\left(-\frac{\alpha t(a^2\alpha^2 - 24b^2)}{2a^2} + \frac{2\alpha by}{a} + \alpha x\right)}{a\left(\exp\left(-\frac{\alpha t(a^2\alpha^2 - 24b^2)}{2a^2} + \frac{2\alpha by}{a} + \alpha x\right) + 1\right)} \\ v(x, y, t) &= \frac{2\alpha b}{a^2} - \frac{4\alpha b \exp\left(-\frac{\alpha t(a^2\alpha^2 - 24b^2)}{2a^2} + \frac{2\alpha by}{a} + \alpha x\right)}{a^2\left(\exp\left(-\frac{\alpha t(a^2\alpha^2 - 24b^2)}{2a^2} + \frac{2\alpha by}{a} + \alpha x\right) + 1\right)}. \end{aligned} \quad (34)$$

Solution set 3.2:

$$R_0 = \frac{2\alpha}{a}, \quad R_1 = -\frac{2\alpha}{a}, \quad S_1 = 0, \quad \beta = \frac{2\alpha b - a\alpha^2}{a}, \quad \mu = -\frac{4(a^2\alpha^3 - 3a\alpha^2b + 3\alpha b^2)}{a^2}.$$

We substitute the values of known-constants mentioned above and expression (31) into eq.(10), and plugging it with eq. (9), we obtain the expression for U as

$$U(\xi) = \frac{2\alpha}{a} - \frac{2\alpha e^\xi}{a(e^\xi + 1)}. \quad (35)$$

Hence, from eq.(8), we obtain the expression for V as

$$V(\xi) = \frac{2\beta}{a} - \frac{2\beta e^\xi}{a(e^\xi + 1)}. \quad (36)$$

Accordingly, the solution of (1) is obtained as

$$\begin{aligned} u(x, y, t) &= \frac{2\alpha}{a} - \frac{2\alpha \exp\left(\frac{4t(a^2\alpha^3 - 3a\alpha^2b + 3\alpha b^2)}{a^2} + \frac{y(2\alpha b - a\alpha^2)}{a} + \alpha x\right)}{a\left(\exp\left(\frac{4t(a^2\alpha^3 - 3a\alpha^2b + 3\alpha b^2)}{a^2} + \frac{y(2\alpha b - a\alpha^2)}{a} + \alpha x\right) + 1\right)} \\ v(x, y, t) &= \frac{2(2\alpha b - a\alpha^2)}{a^2} - \frac{2(2\alpha b - a\alpha^2) \exp\left(\frac{4t(a^2\alpha^3 - 3a\alpha^2b + 3\alpha b^2)}{a^2} + \frac{y(2\alpha b - a\alpha^2)}{a} + \alpha x\right)}{a^2\left(\exp\left(\frac{4t(a^2\alpha^3 - 3a\alpha^2b + 3\alpha b^2)}{a^2} + \frac{y(2\alpha b - a\alpha^2)}{a} + \alpha x\right) + 1\right)}. \end{aligned} \quad (37)$$

Solution set 3.3:

$$R_0 = 0, \quad R_1 = -\frac{2\alpha}{a}, \quad S_1 = 0, \quad \beta = \frac{\alpha(a\alpha + 2b)}{a}, \quad \mu = -\frac{4(a^2\alpha^3 + 3a\alpha^2b + 3\alpha b^2)}{a^2}.$$

(a) 3D

(b) 2D

(c) 3D

(d) 2D

Figure 3: Kink wave structures for solution (37) at $\alpha = 2.5$, $a = 1$, $b = 4.59$ under the range space $-20 \leq x \leq 20$, $-30 \leq y \leq 30$ and $t = 0.01$.

We substitute the values of known-constants mentioned above and expression (31) into eq.(10), and plugging it with eq. (9), we obtain the expression for U as

$$U(\xi) = -\frac{2\alpha e^\xi}{a(e^\xi + 1)}. \quad (38)$$

Hence, from eq.(8), we obtain the expression for V as

$$V(\xi) = -\frac{2\beta e^\xi}{a(e^\xi + 1)}. \quad (39)$$

Accordingly, the solution of (1) is obtained as

$$\begin{aligned} u(x, y, t) &= -\frac{2\alpha \exp\left(\frac{4t(a^2\alpha^3+3a\alpha^2b+3\alpha b^2)}{a^2} + \frac{\alpha y(a\alpha+2b)}{a} + \alpha x\right)}{a \left(\exp\left(\frac{4t(a^2\alpha^3+3a\alpha^2b+3\alpha b^2)}{a^2} + \frac{\alpha y(a\alpha+2b)}{a} + \alpha x\right) + 1\right)} \\ v(x, y, t) &= -\frac{2\alpha(a\alpha+2b) \exp\left(\frac{4t(a^2\alpha^3+3a\alpha^2b+3\alpha b^2)}{a^2} + \frac{\alpha y(a\alpha+2b)}{a} + \alpha x\right)}{a^2 \left(\exp\left(\frac{4t(a^2\alpha^3+3a\alpha^2b+3\alpha b^2)}{a^2} + \frac{\alpha y(a\alpha+2b)}{a} + \alpha x\right) + 1\right)}. \end{aligned} \quad (40)$$

Family 4: For $[r_1, r_2, r_3, r_4] = [-2i, -2i, 5, -5]$ and $[s_1, s_2, s_3, s_4] = [3i, -3i, 3i, -3i]$, eq. (5) yields

$$\Psi(\xi) = \frac{2}{5} \cot(3\xi). \quad (41)$$

With the assistance of soft computation via Mathematica, we solve algebraic equation for obtaining the values of parameters and hence following set of solutions can be achieved.

Solution set 4.1:

$$R_0 = 0, \quad R_1 = -\frac{15\alpha}{a}, \quad S_1 = \frac{12\alpha}{5a}, \quad \beta = \frac{2\alpha b}{a}, \quad \mu = -\frac{12(6a^2\alpha^3 + \alpha b^2)}{a^2}.$$

We substitute the values of known-constants described above and expression (41) into eq.(10), and plugging it with eq. (9), we obtain the expression for U as

$$U(\xi) = \frac{6\alpha \tan(3\xi)}{a} - \frac{6\alpha \cot(3\xi)}{a}. \quad (42)$$

Hence, from eq.(8), we obtain the expression for V as

$$V(\xi) = \frac{6\beta \tan(3\xi)}{a} - \frac{6\beta \cot(3\xi)}{a}. \quad (43)$$

Accordingly, the solution of (1) is obtained as

$$\begin{aligned} u(x, y, t) &= \frac{6\alpha \tan\left(3\left(\frac{12t(6a^2\alpha^3+\alpha b^2)}{a^2} + \frac{2\alpha by}{a} + \alpha x\right)\right)}{a} - \frac{6\alpha \cot\left(3\left(\frac{12t(6a^2\alpha^3+\alpha b^2)}{a^2} + \frac{2\alpha by}{a} + \alpha x\right)\right)}{a} \\ v(x, y, t) &= \frac{12\alpha b \tan\left(3\left(\frac{12t(6a^2\alpha^3+\alpha b^2)}{a^2} + \frac{2\alpha by}{a} + \alpha x\right)\right)}{a^2} - \frac{12\alpha b \cot\left(3\left(\frac{12t(6a^2\alpha^3+\alpha b^2)}{a^2} + \frac{2\alpha by}{a} + \alpha x\right)\right)}{a^2}. \end{aligned} \quad (44)$$

Solution set 4.2:

$$R_0 = -\frac{6i\alpha}{a}, \quad R_1 = 0, \quad S_1 = -\frac{12\alpha}{5a}, \quad \beta = \frac{2(\alpha b + 3ia\alpha^2)}{a}, \quad \mu = \frac{12(12a^2\alpha^3 - 6ia\alpha^2b - \alpha b^2)}{a^2}.$$

We substitute the values of known-constants mentioned above and expression (41) into eq.(10), and plugging it with eq. (9), we obtain the expression for U as

$$U(\xi) = -\frac{6\alpha \tan(3\xi)}{a} - \frac{6i\alpha}{a}. \quad (45)$$

Figure 4: Multisoliton structures for solution (44) at $\alpha = 1$, $a = 1$, $b = 0.3$ under the range space $-20 \leq x \leq 20$, $-300 \leq y \leq 300$ and $t = 0.01$.

Hence, from eq.(8), we obtain the expression for V as

$$V(\xi) = -\frac{6\beta \tan(3\xi)}{a} - \frac{6i\beta}{a}. \quad (46)$$

Accordingly, the solution of (1) is obtained as

$$u(x, y, t) = -\frac{6\alpha \tan \left(3 \left(-\frac{12t(12a^2\alpha^3 - 6ia\alpha^2b - \alpha b^2)}{a^2} + \frac{2y(\alpha b + 3ia\alpha^2)}{a} + \alpha x \right) \right)}{a} - \frac{6i\alpha}{a}$$

$$v(x, y, t) = -\frac{12(\alpha b + 3ia\alpha^2) \tan \left(3 \left(-\frac{12t(12a^2\alpha^3 - 6ia\alpha^2b - \alpha b^2)}{a^2} + \frac{2y(\alpha b + 3ia\alpha^2)}{a} + \alpha x \right) \right)}{a^2} - \frac{12i(\alpha b + 3ia\alpha^2)}{a^2}. \quad (47)$$

Solution set 4.3:

$$R_0 = \frac{6i\alpha}{a}, \quad R_1 = \frac{15\alpha}{a}, \quad S_1 = 0, \quad \beta = \frac{2(\alpha b - 3ia\alpha^2)}{a}, \quad \mu = \frac{12(12a^2\alpha^3 + 6ia\alpha^2b - \alpha b^2)}{a^2}.$$

We substitute the values of known-constants described above and expression (41) into eq.(10), and plugging it with eq. (9), we obtain the expression for U as

$$U(\xi) = \frac{6\alpha \cot(3\xi)}{a} + \frac{6i\alpha}{a}. \quad (48)$$

Hence, from eq.(8), we obtain the expression for V as

$$V(\xi) = \frac{6\beta \cot(3\xi)}{a} + \frac{6i\beta}{a}. \quad (49)$$

Accordingly, the solution of (1) is obtained as

$$\begin{aligned} u(x, y, t) &= \frac{6\alpha \cot \left(3 \left(-\frac{12t(12a^2\alpha^3+6ia\alpha^2b-\alpha b^2)}{a^2} + \frac{2y(\alpha b-3ia\alpha^2)}{a} + \alpha x \right) \right)}{a} + \frac{6i\alpha}{a} \\ v(x, y, t) &= \frac{12(\alpha b - 3ia\alpha^2) \cot \left(3 \left(-\frac{12t(12a^2\alpha^3+6ia\alpha^2b-\alpha b^2)}{a^2} + \frac{2y(\alpha b-3ia\alpha^2)}{a} + \alpha x \right) \right)}{a^2} + \frac{12i(\alpha b - 3ia\alpha^2)}{a^2}. \end{aligned} \quad (50)$$

Family 5: For $[r_1, r_2, r_3, r_4] = [4, 0, 1, -1]$ and $[s_1, s_2, s_3, s_4] = [5, -5, 5, -5]$, eq. (5) yields

$$\Psi(\xi) = 2(1 + \coth(5\xi)). \quad (51)$$

With the assistance of soft computation via Mathematica, we solve algebraic equation for obtaining the values of parameters and hence following set of solutions can be achieved.

Solution set 5.1:

$$R_0 = \frac{10\alpha}{a}, \quad R_1 = -\frac{5\alpha}{a}, \quad S_1 = 0, \quad \beta = \frac{2\alpha b}{a}, \quad \mu = \frac{2(25a^2\alpha^3 - 6\alpha b^2)}{a^2}.$$

We substitute the values of known-constants mentioned above and expression (51) into eq.(10), and plugging it with eq. (9), we obtain the expression for U as

$$U(\xi) = \frac{10\alpha}{a} - \frac{10\alpha(\coth(5\xi) + 1)}{a}. \quad (52)$$

Hence, from eq.(8), we obtain the expression for V as

$$V(\xi) = \frac{10\beta}{a} - \frac{10\beta(\coth(5\xi) + 1)}{a}. \quad (53)$$

Accordingly, the solution of (1) is obtained as

$$\begin{aligned} u(x, y, t) &= \frac{10\alpha}{a} - \frac{10\alpha \left(\coth \left(5 \left(-\frac{2t(25a^2\alpha^3-6\alpha b^2)}{a^2} + \frac{2\alpha by}{a} + \alpha x \right) \right) + 1 \right)}{a} \\ v(x, y, t) &= \frac{20\alpha b}{a^2} - \frac{20\alpha b \left(\coth \left(5 \left(-\frac{2t(25a^2\alpha^3-6\alpha b^2)}{a^2} + \frac{2\alpha by}{a} + \alpha x \right) \right) + 1 \right)}{a^2}. \end{aligned} \quad (54)$$

Solution set 5.2:

$$R_0 = -\frac{20\alpha}{a}, \quad R_1 = \frac{5\alpha}{a}, \quad S_1 = 0, \quad \beta = \frac{2(5a\alpha^2 + \alpha b)}{a}, \quad \mu = -\frac{4(100a^2\alpha^3 + 30a\alpha^2b + 3\alpha b^2)}{a^2}.$$

We substitute the values of known-constants described above and expression (51) into eq.(10), and plugging it with eq. (9), we obtain the expression for U as

$$U(\xi) = \frac{10\alpha(\coth(5\xi) + 1)}{a} - \frac{20\alpha}{a}. \quad (55)$$

Hence, from eq.(8), we obtain the expression for V as

$$V(\xi) = \frac{10\beta(\coth(5\xi) + 1)}{a} - \frac{20\beta}{a}. \quad (56)$$

Accordingly, the solution of (1) is obtained as

$$\begin{aligned} u(x, y, t) &= \frac{10\alpha \left(\coth \left(5 \left(\frac{4t(100a^2\alpha^3 + 30a\alpha^2b + 3\alpha b^2)}{a^2} + \frac{2y(5a\alpha^2 + \alpha b)}{a} + \alpha x \right) \right) + 1 \right)}{a} - \frac{20\alpha}{a} \\ v(x, y, t) &= \frac{20(5a\alpha^2 + \alpha b) \left(\coth \left(5 \left(\frac{4t(100a^2\alpha^3 + 30a\alpha^2b + 3\alpha b^2)}{a^2} + \frac{2y(5a\alpha^2 + \alpha b)}{a} + \alpha x \right) \right) + 1 \right)}{a^2} \\ &\quad - \frac{40(5a\alpha^2 + \alpha b)}{a^2}. \end{aligned} \quad (57)$$

Solution set 5.3:

$$R_0 = 0, \quad R_1 = \frac{5\alpha}{a}, \quad S_1 = 0, \quad \beta = -\frac{2(5a\alpha^2 - \alpha b)}{a}, \quad \mu = -\frac{4(100a^2\alpha^3 - 30a\alpha^2b + 3\alpha b^2)}{a^2}.$$

We substitute the values of known-constants mentioned above and expression (51) into eq. (10), and plugging it with eq. (9), we obtain the expression for U as

$$U(\xi) = \frac{10\alpha(\coth(5\xi) + 1)}{a}. \quad (58)$$

Hence, from eq.(8), we obtain the expression for V as

$$V(\xi) = \frac{10\beta(\coth(5\xi) + 1)}{a}. \quad (59)$$

Accordingly, the solution of (1) is obtained as

$$\begin{aligned} u(x, y, t) &= \frac{10\alpha \left(\coth \left(5 \left(\frac{4t(100a^2\alpha^3 - 30a\alpha^2b + 3\alpha b^2)}{a^2} - \frac{2y(5a\alpha^2 - \alpha b)}{a} + \alpha x \right) \right) + 1 \right)}{a} \\ v(x, y, t) &= -\frac{20(5a\alpha^2 - \alpha b) \left(\coth \left(5 \left(\frac{4t(100a^2\alpha^3 - 30a\alpha^2b + 3\alpha b^2)}{a^2} - \frac{2y(5a\alpha^2 - \alpha b)}{a} + \alpha x \right) \right) + 1 \right)}{a^2}. \end{aligned} \quad (60)$$

4 Bifurcation analysis

We explore the new dynamics of KD system (1), in this section, by utilizing the concepts of bifurcation theory.

After simplification, equation (9) can be rewritten in the following form of planar dynamical system as

$$\begin{aligned}\frac{dU}{d\xi} &= P, \\ \frac{dP}{d\xi} &= A_1 U^3 - A_2 U^2 - A_3 U,\end{aligned}\tag{61}$$

where, $A_1 = \frac{a^2}{2\alpha^2}$, $A_2 = \frac{3}{2} \left(\frac{2b}{\alpha^2} - \frac{a\beta}{\alpha^3} \right)$ and $A_3 = \frac{3\beta^2}{\alpha^4} + \frac{\mu}{\alpha^3}$.

The three equilibrium points for the above system of differential equations are computed as $(0, 0)$, $(U_1, 0)$ and $(U_2, 0)$ on U-axis, where

$$U_1 = \frac{1}{2A_1} \left(A_2^2 + \sqrt{\delta} \right), \quad U_2 = \frac{1}{2A_1} \left(A_2^2 - \sqrt{\delta} \right).$$

and $\delta = A_2^2 + 4A_1A_3$.

Let $M(U_i, 0)$ is the coefficient matrix of the linearized system of (61) at equilibrium point $(U_i, 0)$; J and T be the determinant and trace of the matrix M , respectively.

Here,

$$\begin{aligned}J &= -(3A_1U^2 - 2A_2U - A_3) \text{ and} \\ T &= 0.\end{aligned}$$

By using the theory of planar dynamical system [25], we can discuss following definitions for the critical points $(U_i, 0)$.

- (i) When $J < 0$, then $(U_i, 0)$ is a saddle point.
- (ii) When $J > 0$ and $T^2 - 4J \geq 0$, then $(U_i, 0)$ is a node; which is stable if $T < 0$ and unstable if $T > 0$.
- (iii) When $J > 0$, $T^2 - 4J < 0$ and $T \neq 0$, then $(U_i, 0)$ is a focus; which is stable if $T < 0$ and unstable if $T > 0$.
- (iv) When $J > 0$ and $T = 0$, then $(U_i, 0)$ is a center.
- (v) When $J = 0$ and Poincare index of $(U_i, 0)$ is zero, then it is called the zero point.

For different choices of parameters A_1, A_2 and A_3 , various cases in detail are explained as:

Case 1: $A_1 > 0$, $A_2 > 0$, and $A_3 > 0$: Figure 5(a) exhibits the phase portrait for the values of parameters considered as $A_1 = 1$, $A_2 = 1$, $A_3 = 1$. For this case, we have three equilibrium points where $(0, 0)$ is center point whereas $(U_1, 0)$ and $(U_2, 0)$ are saddle points. Here, presence of nonlinear periodic trajectory and nonlinear homoclinic trajectory ensure the occurrence of closed-form solutions of the KD equation (1).

Case 2: $A_1 > 0$, $A_2 > 0$, and $A_3 < 0$: Figure 5(b) explains the phase portrait for the values of parameters given as $A_1 = 1$, $A_2 = 1$, $A_3 = -1$. For this case, we have only one real equilibrium point, trivial equilibrium point $(0, 0)$, which is a saddle point. Here, it has been observed that the closed form trajectories are not obtained.

(a) $A_1 > 0$, $A_2 > 0$, and $A_3 > 0$

(b) $A_1 > 0$, $A_2 > 0$, and $A_3 < 0$

(c) $A_1 > 0$, $A_2 < 0$, and $A_3 > 0$

(d) $A_1 > 0$, $A_2 < 0$, and $A_3 < 0$

Figure 5: Phase Portraits

Case 3: $A_1 > 0$, $A_2 < 0$, and $A_3 > 0$: Figure 5(c) describes the phase portrait for the values of parameters given by $A_1 = 1$, $A_2 = -1$, $A_3 = 1$. For this case, we have three equilibrium points, where $(0,0)$ is a center point whereas $(U_1, 0)$ and $(U_2, 0)$ are saddle points. In this case also, the presence of nonlinear homoclinic trajectory and nonlinear periodic trajectory ensure the occurrence of closed form solutions of the KD equation (1).

Case 4: $A_1 > 0$, $A_2 < 0$, and $A_3 < 0$: Figure 5(d) exhibits the phase portrait for the parameter values $A_1 = 1$, $A_2 = -1$, $A_3 = -1$. For this case, we have only one equilibrium points namely $(0,0)$ which is a saddle point. Here, closed form trajectories are not obtained.

Case 5: $A_1 < 0$, $A_2 > 0$, and $A_3 > 0$: Figure 6(a) describes the phase portrait for the values of parameters $A_1 = -1$, $A_2 = 1$, $A_3 = 1$. For this case, we have obtained only one real equilibrium point, namely $(0,0)$ which is a center point. For this case, presence of non-linear periodic trajectories

(a) $A_1 < 0$, $A_2 > 0$, and $A_3 > 0$

(b) $A_1 < 0$, $A_2 > 0$, and $A_3 < 0$

(c) $A_1 < 0$, $A_2 < 0$, and $A_3 > 0$

(d) $A_1 < 0$, $A_2 < 0$, and $A_3 < 0$

Figure 6: Phase Portraits

ensures the existence of closed form solutions.

Case 6: $A_1 < 0$, $A_2 > 0$, and $A_3 < 0$: Figure 6(b) exhibits the phase portrait for the parameter values $A_1 = -1$, $A_2 = 1$, $A_3 = -1$. For this case, we have three equilibrium points, where $(0,0)$ is a saddle point whereas $(U_1, 0)$ and $(U_2, 0)$ are center points. Here, the presence of nonlinear periodic trajectory and nonlinear homoclinic trajectory ensure the occurrence of closed form solutions of the KD equation (1).

Case 7: $A_1 < 0$, $A_2 < 0$, and $A_3 > 0$: Figure 6(c) describes the phase portrait for the values of parameters $A_1 = -1$, $A_2 = -1$, $A_3 = 1$. For this case, we have obtained only one real equilibrium point namely $(0,0)$ which is a center point. For this case also, presence of non-linear periodic trajectories ensures the existence of closed for solutions.

Case 8: $A_1 < 0$, $A_2 < 0$, and $A_3 < 0$: Figure 6(d) explains the phase portrait for the parameter values $A_1 = -1$, $A_2 = -1$, $A_3 = -1$. For this case, we have three equilibrium points where $(0, 0)$ is a saddle point whereas $(U_1, 0)$ and $(U_2, 0)$ are center points. Here, the presence of nonlinear periodic trajectory and nonlinear homoclinic trajectory insure the occurrence of closed form solutions of the KD equation (1).

5 Quasi-periodic solution

The exploration of different dynamics of the perturbed system of the main system (1) by adding perturbation term $\rho_1 \cos(\sigma\eta)$ is investigated in this section. Thus, equation (7) with the addition of perturbation term becomes

$$\begin{aligned} \frac{dU}{d\xi} &= P, \\ \frac{dP}{d\xi} &= A_1 U^3 - A_2 U^2 - A_3 U + \rho_1 \cos(\sigma\eta), \end{aligned} \quad (62)$$

where ρ_1 is constant representing the intensity of the perturbed term and σ represents the frequency. To analyze the chaotic and periodic behavior of system (7) with the existence of a perturbation term, we will fix the influence of force and frequency of disruption and retain the physical parameters of the system under observation.

(a) 2D–Phase portraits with perturbation term

(b) 3D–Phase portraits with perturbation term

Figure 7: For $A_1 = -0.1$, $A_2 = 1.056$, $A_3 = 8.12$, $\rho_1 = -0.8$ and $\sigma = \pi$, 2-D and 3-D phase portraits of the perturbed system (62).

Figure 7 exhibits the 2-D and 3-D phase portraits of the perturbed model (62) for $A_1 = -0.1$, $A_2 = 1.056$, $A_3 = 8.12$, $\rho_1 = -0.8$ and $\sigma = \pi$. Figure 8 depicts the 2-D and 3-D phase portraits for the perturbed system (62) for $A_1 = -1$, $A_2 = -1.056$, $A_3 = 8.12$, $\rho_1 = -0.8$ and $\sigma = \pi$. Figure 9 describes the 2-D and 3-D phase portraits for the perturbed system (62) for $A_1 = -0.93$, $A_2 = -1.056$, $A_3 = -2.5$, $\rho_1 = -0.8$ and $\sigma = \pi$.

(a) 2D-Phase portraits with perturbation term

(b) 3D-Phase portraits with perturbation term

Figure 8: For $A_1 = -1$, $A_2 = -1.056$, $A_3 = 8.12$, $\rho_1 = -0.8$ and $\sigma = \pi$, 2-D and 3-D phase portraits of the perturbed system (62).

(a) 2D-Phase portraits with perturbation term

(b) 3D-Phase portraits with perturbation term

Figure 9: For $A_1 = -0.93$, $A_2 = -1.056$, $A_3 = -2.5$, $\rho_1 = -0.8$ and $\sigma = \pi$, 2-D and 3-D phase portraits of the perturbed system (62).

6 Conclusion

In summary, we investigated the (2+1)-dimensional Konopelchenko-Dubrovsky (KD) model and constructed numerous exact closed-form solutions using GERF (generalized exponential rational function) approach with soft symbolic computations via Mathematica. The established soliton solutions exhibit that the KD equations (1) admits abundant exact closed-form solutions having arbitrary constant parameters. The findings depict rich dynamical formations of the generated closed-form solutions in the forms of multi-solitons, kink-wave profiles, the interaction of multi-soliton and kink-wave solutions, and multiple solitons in figures 1, 2, 3 and 4. The obtained solutions will be beneficial in the theory of solitons, nonlinear dynamics, fluid mechanics, applied physics, optic fiber, natural sciences, physics, and many other areas. The technique we have used is one of the powerful techniques for obtaining the exact analytical solutions of NLPDEs as this GERF method constructs an extensive

scale of established solitary wave solutions to the associated system. This method is very effective, trustworthy, and efficient. Moreover, the dynamics of KD equations (1) is also examined by using the bifurcation analysis in which we have found various equilibrium points, and different types of phase portraits have been discussed, see figures 5 and 6. Furthermore, we numerically studied the existence of quasi-periodic solutions for the perturbed model, which is obtained after inserting the periodic forces to the considered KD system (1). Also, $2D$ and $3D$ phase portraits shown by figures 7, 8 and 9 for the perturbed system were also plotted.

Availability of data and material

Data sharing is not applicable to this article as no data sets were created or analyzed in this study.

Conflict of interest

The authors declare that they have no known competing financial interests.

References

- [1] Kumar, H., Malik, A., Chand, F., Mishra, S.C.: Exact solutions of nonlinear diffusion reaction equation with quadratic, cubic and quartic nonlinearities. *Indian J. Phys.* **86**, 819–827 (2012). <https://doi.org/10.1007/s12648-012-0126-y>
- [2] Hirota, R.: *The Direct Method in Soliton Theory*, Cambridge University Press, Cambridge (2004)
- [3] Kumar, S., Mohan, B.: A study of multi-soliton solutions, breather, lumps, and their interactions for kadomtsev-petviashvili equation with variable time coefficient using hirota method. *Phys. Scr.* **96** (12), 125255 (2021). <https://doi.org/10.1088/1402-4896/ac3879>
- [4] Fu, Z.T., Liu, S.K., Liu, S.D., Zhao, Q.: New Jacobi elliptic function expansion and new periodic solutions of nonlinear wave equations. *Phys. Lett. A* **290**, 72–76 (2001). [https://doi.org/10.1016/S0375-9601\(01\)00644-2](https://doi.org/10.1016/S0375-9601(01)00644-2)
- [5] Kudryashov, N.A.: Simplest equation method to look for exact solutions of nonlinear differential equations. *Chaos Soliton Fractals* **24**(5), 1217–1231 (2005). <https://doi.org/10.1016/j.chaos.2004.09.109>
- [6] Malik, A., Chand, F., Kumar, H., Mishra, S.C.: Exact solutions of the Bogoyavlenskii equation using the multiple $(\frac{G'}{G})$ -expansion method. *Comput. Math. Appl.* **64**(9), 2850–2859 (2012). <https://doi.org/10.1016/j.camwa.2012.04.018>
- [7] Guan, X., Liu, W., Zhou, Q., Biswas, A.: Darboux transformation and analytic solutions for a generalized super-NLS-mKdV equation. *Nonlinear Dyn* **98**, 1491–1500 (2019). <https://doi.org/10.1007/s11071-019-05275-0>
- [8] Hu, X.B.: Nonlinear superposition formulae for the differential-difference analogue of the KdV equation and two-dimensional Toda equation. *J. Phys. A: Math. Gen.* **27**(1), 201 (1994). <https://doi.org/10.1088/0305-4470/27/1/014>
- [9] Wang, M.L.: Solitary wave solutions for variant Boussinesq equations. *Phys. Lett. A* **199**, 169–172 (1995). [https://doi.org/10.1016/0375-9601\(95\)00092-H](https://doi.org/10.1016/0375-9601(95)00092-H)

- [10] Kumar, S., Kumar, A.: Lie symmetry reductions and group invariant solutions of $(2 + 1)$ -dimensional modified Veronese web equation. *Nonlinear Dyn* **98**, 1891–1903 (2019). <https://doi.org/10.1007/s11071-019-05294-x>
- [11] Ghanbari, B., Inc, M.: A new generalized exponential rational function method to find exact special solutions for the resonance nonlinear Schrödinger equation. *Eur. Phys. J. Plus* **133**, 142 (2018). <https://doi.org/10.1140/epjp/i2018-11984-1>
- [12] Ghanbari, B., Osman, M.S., Baleanu, D.: Generalized exponential rational function method for extended Zakharov–Kuzetsov equation with conformable derivative. *Mod. Phys. Lett. B* **34**(20), 1950155 (2019). <https://doi.org/10.1142/S0217732319501554>
- [13] Kumar, S., Kumar, A., Wazwaz, A.M.: New exact solitary wave solutions of the strain wave equation in microstructured solids via the generalized exponential rational function method. *Eur. Phys. J. Plus* **135**, 870 (2020). <https://doi.org/10.1140/epjp/s13360-020-00883-x>
- [14] Kumar, S., Almusawa, H., Hamid, I., Abdou, M.A.: Abundant closed-form solutions and solitonic structures to an integrable fifth-order generalized nonlinear evolution equation in plasma physics. *Results in Physics* **26**, 104453 (2021). <https://doi.org/10.1016/j.rinp.2021.104453>
- [15] Konopelchenko, B. G., Dubrovsky, V. G.: Some new integrable nonlinear evolution equations in $2 + 1$ dimensions. *Phys. Lett. A* **102**(1-2), 15–17 (1984). [https://doi.org/10.1016/0375-9601\(84\)90442-0](https://doi.org/10.1016/0375-9601(84)90442-0)
- [16] Kumar, S., Hama, A., Biswas, A.: Solutions of Konopelchenko-Dubrovsky equation by traveling wave hypothesis and Lie symmetry approach. *Appl. Math. Inf. Sci.* **8**(4), 1533–1539 (2014). <https://doi.org/10.12785/amis/080406>
- [17] Alfalqi, S. H., Alzaidi, J. F., Lu, D., Khater, M.: On exact and approximate solutions of $(2 + 1)$ -dimensional Konopelchenko-Dubrovsky equation via modified simplest equation and cubic B-spline schemes. *Therm. Sci.* **23**1889–1899 (2019). <https://doi.org/10.2298/TSCI190131349A>
- [18] Khater, M.M.A., Lu, D., Attia, R.A.M.: Lump soliton wave solutions for the $(2 + 1)$ -dimensional Konopelchenko-Dubrovsky equation and KdV equation. *Modern Phys. Lett. B* **33**(18), 1950199 (2019). <https://doi.org/10.1142/S0217984919501999>
- [19] Ren, B., Cheng, X.P., Lin, J.: The $(2 + 1)$ -dimensional Konopelchenko–Dubrovsky equation: nonlocal symmetries and interaction solutions. *Nonlinear Dyn* **86**, 1855–1862 (2016). <https://doi.org/10.1007/s11071-016-2998-4>
- [20] Jhangeer, A., Hussain, A., Junaid-U-Rehman, M., Baleanu, D., Riaz, M.B.: Quasi-periodic, chaotic and travelling wave structures of modified Gardner equation. *Chaos Solitons Fractals* **143**, 110578 (2021). <https://doi.org/10.1016/j.chaos.2020.110578>
- [21] Hussain, A., Jhangeer, A., Tahir, S., Chu, Y.M., Khan, I., Nisar, K.S.: Dynamical behaviour of fractional Chen-Lee-Liu equation in optical fibers with beta derivatives. *Results in Physics* **18**, 103208 (2020). <https://doi.org/10.1016/j.rinp.2020.103208>
- [22] Chang, L., Liu, H., Xin, X.: Lie symmetry analysis, bifurcations and exact solutions for the $(2+1)$ -dimensional dissipative long wave system. *J. Appl. Math. Comput.* **64**, 807–823 (2020). <https://doi.org/10.1007/s12190-020-01381-0>

- [23] Jhangeer, A., Raza, N., Rezazadeh, H., Seadawy, A.: Nonlinear self-adjointness, conserved quantities, bifurcation analysis and travelling wave solutions of a family of long-wave unstable lubrication model. *Pramana - J Phys* **94**, 87 (2020). <https://doi.org/10.1007/s12043-020-01961-6>
- [24] Elbrolosy, M.E., Elmandouh, A.A.: Dynamical behaviour of nondissipative double dispersive microstrain wave in the microstructured solids. *Eur. Phys. J. Plus* **136**, 955 (2021). <https://doi.org/10.1140/epjp/s13360-021-01957-0>
- [25] Perko, L.: Differential equations and dynamical systems, third edition, Texts in Applied Mathematics, 7, Springer-Verlag, New York (2001)