Prevalence and associated factors of chlamydial infection among patients seeking clinic-based STI services in Shenzhen, China

Hong-Lin Wang

Shenzhen Center for Chronic Disease Control, Guangdong Province

Yu-Mao Cai (✉ 64165469@qq.com)

Shenzhen Center for Chronic Disease Control <https://orcid.org/0000-0002-8098-8244>

Xiang-Sheng Chen

Chinese Academy of Medical Sciences & Peking Union Medical College Hospital of Skin Diseases and Institute of Dermatology

Wei-Ye Yu

Baylor College of Medicine

Fu-Chang Hong

Shenzhen Center for Chronic Disease Control

Tie-Jian Feng

Shenzhen Center for Disease Control and Prevention

Jin-Hong Liu

Shenzhen Center for Chronic Disease Control

Chun-Lai Zhang

Shenzhen Center for Chronic Disease Control

Li-Zhang Wen

Shenzhen Center for Chronic Disease Control

Feng Wang

Shenzhen Center for Chronic Disease Control

Jian-Bin Ye

Shenzhen Center for Chronic Disease Control

Rong-Xing Weng

Shenzhen Center for Chronic Disease Control

Fen Tang

Shenzhen Center for Chronic Disease Control

Research article

Keywords: Chlamydia trachomatis, STI surveillance programming, intervention, resource allocation, China

Posted Date: July 16th, 2020

DOI: <https://doi.org/10.21203/rs.2.21284/v2>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

The authors have withdrawn this preprint due to erroneous posting.