Evaluation of The 10 Warning Signs In Immunodeficient Patients: Additional Suggestions

Fadime Ceyda Eldeniz

Necmettin Erbakan University

Yahya Gül (✉ yahya.palu@hotmail.com)

Necmettin Erbakan University <https://orcid.org/0000-0002-9973-4471>

Alaattin Yorulmaz

Selcuk University

Şükrü Nail Güner

Necmettin Erbakan University

Sevgi Keles

Necmettin Erbakan University

İsmail Reisli

Necmettin Erbakan University

Research Article

Keywords: Primary immunodeficiency, 10 warning signs, child

Posted Date: December 30th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-1189503/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Objective: Ten warning signs of primary immunodeficiency (PID) were suggested by the Jeffrey Modell Foundation (JMF), to increase physician awareness of PID. These warning signs have not yet been evaluated for patients with secondary immunodeficiency (SID). This study investigated whether the 10 warning signs used for the diagnosis of PID are sufficient for the diagnosis of SID, and explored the possibility of additional signs.

Methods: This prospective study was conducted between June and December 2020. The mothers of 162 patients with PID and SID, and mothers of 200 healthy children, were asked to complete a questionnaire about family and personal history in addition to the warning signs of PID developed by the JMF. A JMF score was created by giving one point for each “Yes” answer for the 10 warning signs of PID. Medical records of the patients were evaluated for possible additional warning signs for PID and SID.

Results: The JMF scores of the PID (3.36 ± 1.65) and SID (3.72 ± 1.12) groups were significantly higher than the scores of the control group (0.34 ± 0.61) ($p < 0.05$). A sign for immunological evaluation in two patients without warning signs in the PID group was found to be chronic diarrhea. In addition to the 10 JMF warning signs, we found that consanguinity and a family history of tuberculosis were statistically significant in our PID group, compared with the SID and control groups.

Conclusions: The JMF warning signs are important for early diagnosis of PID. Our study showed that these signs may also be used for the early diagnosis of SID in patients and, according to our results, in addition to the 10 JMF signs for PID, parental consanguinity, chronic diarrhea, and a family history of tuberculosis may also be considered warning signs for the early diagnosis of PID.

Introduction

Primary immunodeficiencies (PIDs) are a group of diseases comprised of more than 450 innate errors of immunity, and they are becoming more prevalent [1]. Although PIDs are rare diseases, they are more common than previously thought, following the use of modern diagnostic methods [2]. A recent evaluation indicated that at least 1–2% of the world’s population are affected by PIDs [3]. Studies from our country also reveal a high incidence of PIDs in children. In this regard, training family physicians, pediatricians and, in particular, infectious disease specialists about PIDs will allow early diagnosis of these patients; early and effective treatment may allow them to reach adulthood [4, 5].

Several warning signs have been developed to increase physician awareness about the early diagnosis of PIDs. Warning signs of PID were developed by an institution called the Jeffrey Modell Foundation (JMF), which tries to increase awareness of this issue (<http://www.info4pi.org/library/educational-materials/10-warning-signs>). A detailed history should be taken for children admitted with a history of frequent infections, and the 10 warning signs of PID described by the JMF should be evaluated in addition to a full physical examination. This approach will allow early diagnosis of PID patients and, hence, the possibility of early and effective treatment before the development of organ damage [6, 7]. These warning signs have not yet been evaluated for patients with secondary immunodeficiency (SID).

The objective of this study was to investigate whether the 10 warning signs used for the diagnosis of PID are sufficient for the diagnosis of SID, and to explore the possibility of additional signs.

Materials And Methods

This prospective study was conducted between June and December 2020. This research was conducted using data obtained for clinical purposes. The study was approved by Necmettin Erbakan University Meram Medical School Ethics Committee (Date: 06.26.2020/No: 2020/2599).

The mothers of 162 patients diagnosed with PID and SID were asked to complete a questionnaire about family and personal history, in addition to the 10 warning signs of PID developed by the JMF. The same survey was completed by the mothers of 200 children without any defined primary/secondary immunodeficiency (i.e., healthy), and they formed the control group.

The study group was divided into two groups based upon clinical and laboratory findings to form the PID and SID groups. In addition to the questionnaire, the medical records of patients in the study groups were evaluated; age at diagnosis, treatment of the immunodeficiency, duration of diagnostic delay, and patient characteristics during the follow-up period were recorded.

The questionnaire was provided to mothers of the study and control groups and the responses were recorded by the same research scientist. A JMF score was created by giving one point for each “Yes” answer for the 10 warning signs of PID.

Statistical Analyses

Analysis of the study data was performed by using the SPSS 25 program (IBM Corp. Released 2017. IBM SPSS Statistics for Windows, Version 25.0. Armonk, NY, USA). Frequency, ratio, mean, and standard deviation (SD) of different variables of the individuals were analyzed by descriptive statistics. The Independent Samples t-test was used for dual comparisons, whereas triple comparisons were performed using one-way analysis of variance. Mean $\pm$ SD values of the groups were reported for evaluation of distribution rates of different variables by the groups using chi-square analysis. Crosstabs were made, and numbers and ratios were reported in the crosstabs. For variables found to be significant as a result of chi-square analyses performed for triple groups, further dual comparisons were performed to find out the reason for this difference. The level of significance was determined to be $p < 0.05$.

Results

Of the patients included in the study, there were 98 (27.1%) in the PID group, and 64 (17.7%) in the SID group; the control group consisted of 200 (55.2%) healthy children. Overall, there were 200 (55.2%) males and 162 (44.8%) females. In the PID group, 48 (49%) were female and 50 (51%) were male, whereas in the SID group, 16 (25%) were female and 48 (75%) were male. In the control group, 98 (49%) were female and 102 (51%) were male. When sex distribution ratios were analyzed by group, the ratio of males in the SID group was significantly higher, compared with the PID and control groups ($p < 0.001$).

Ages of the participants varied between 1–216 months (mean $\pm$ SD: 74.93 $\pm$ 62.59). The mean age was 98.87 $\pm$ 66.37 months in the PID group, 67.89 $\pm$ 40.19 months in the SID group, and 65.46 $\pm$ 63.71 months in the control group. The demographic data of the patients are shown in Table 1. The mean age of the PID group was significantly higher compared with that of the control and SID groups ($p < 0.001$). There was no statistical difference between the control and SID groups.

Table 1
The demographic characteristics of the patients with PID and SID, and the control group

	Control		PID		SID		
	Mean±SD	Median (min-max)	Mean±SD	Median (min-max)	Mean±SD	Median (min-max)	p
Age (Month)	65.46±63.71	42 (1-204)	98.87±66.37	84 (7-216)	67.89±40.19	57 (3-204)	0.001
Age at first hospitalization (Month)	19.30±23.29	12.0 (1-120)	14.62±26.66	3.0 (1-162)	7.87±13.80	2.0 (1-60)	0.001
Age at diagnosis of immunodeficiency (Month)			48.58±53.74	21.00 (1-192)	36.64±38.87	24.00 (2-174)	0.127
Duration of diagnostic delay (Month)			21.04±32.71	10.50 (1-186)	21.67±25.79	12.0 (1-108)	0.897
Age at initiation of treatment (Month)			55.06±58.36	27.00 (1-210)	37.73±38.92	24.00 (2-174)	0.025
Duration of treatment (Month)			39.22±32.42	30.00 (1-162)	28.89±21.03	24.00 (1-120)	0.015
Age at discontinuation of treatment (Month)			97.50±60.57	84.00 (36-204)	39.00±4.24	39.00 (36-42)	0.228
Duration of follow-up (Month)			49.57±41.76	36.00 (5-192)	34.00±22.95	30.00 (1-120)	0.007
PID: primary immunodeficiency, SID: secondary immunodeficiency							

When the PID and SID groups were compared, age at initiation of therapy in the PID group was older ($p < 0.05$), and the duration of therapy and follow-up period were longer ($p < 0.05$) in the PID group compared with those of the SID group, with statistically significant differences.

JMF scores of the PID (3.36 ± 1.65) and SID (3.72 ± 1.12) groups were significantly higher than those of the control group (0.34 ± 0.61) ($p < 0.001$). The distribution of JMF scores of the PID, SID, and control groups is shown in Table 2. The JMF score was zero in two patients in the PID group; these two patients had been screened for PID due to chronic diarrhea. The IUIS classification (2020) [8] of patients with PID, and the distribution of JMF scores of the PID subgroups are shown in Table 3.

Table 2
Distribution of JMF scores of the PID, SID and Control Groups

JMF Score	PID Group (n/%)	SID Group (n/%)	Control Group (n/%)	Total (n/%)
0 point	2 (2.0%)	0	146 (73.0%)	148 (40.9%)
1 point	10 (10.2%)	1 (1.6%)	41 (20.5%)	52 (14.4%)
2 points	23 (23.5%)	8 (12.5%)	12 (6.0%)	43 (11.9%)
3 points	17 (17.3%)	18 (28.1%)	1 (0.5%)	36 (9.9%)
4 points	22 (22.4%)	21 (32.8%)	0	43 (11.9%)
5 points	14 (14.3%)	13 (20.3%)	0	27 (7.5%)
6 points	6 (6.1%)	3 (4.7%)	0	9 (2.5%)
7 points	4 (4.1%)	0	0	4 (1.1%)
8 points	0	0	0	0
9 points	0	0	0	0
10 points	0	0	0	0

JMF: Jeffrey Modell Foundation

Table 3
IUIS classification of the patients with PID (2020) and JMF scores of the PID subgroups

Diagnoses	n	%	JMF score Mean±SD	JMF score Median (Min-Max)
Immunodeficiencies affecting cellular and humoral immunity	18	18.4	3.28±1.60	3.00 (1-7)
Combined immunodeficiencies with associated syndromic features	21	21.4	3.90±1.44	4.00 (1-6)
Immunodeficiencies due to Antibody Deficiencies	39	39.8	2.95±1.58	2.00 (0-6)
Diseases of immune dysregulation	7	7.1	2.14±1.57	2.00 (0-5)
Congenital defects of phagocyte number or function or both	8	8.2	4.50±1.69	4.00 (2-7)
Defects in Intrinsic and Innate Immunity	1	1	7.00±0.00	7.00 (7-7)
Auto-inflammatory disorders	3	3,1	4.00±0.00	4.00 (4-4)
Complement deficiencies	1	1	3.00±0.00	3.00 (3-3)
Bone marrow failure	0	0.0	.	.
Phenocopies of PID	0	0.0	.	.
Total	98	100.0		

After evaluation of overall JMF scores, answers given to the questions were also compared separately. Regarding having ≥ 4 episodes of otitis in one year, ≥ 2 episodes of sinusitis in one year, and a family history of PID, the rates in the PID group were significantly higher than the control and SID groups ($p < 0.001$). No difference was found between the control and SID groups. Regarding oral antibiotic use for ≥ 2 months with little effect, recurrent deep tissue infections or organ abscesses, persistent thrush or cutaneous fungal infections, and ≥ 2 deep tissue infections including septicemia, the rates in the PID and SID groups were significantly higher compared with the SID group ($p < 0.001$). No difference was found between the PID and SID groups. Regarding failure to thrive, the need for IV antibiotics to clear infections, and having ≥ 2 lower respiratory tract infections in one year, the rates in the SID group were higher than the PID group. The rate in the PID group was significantly higher than the control group ($p < 0.001$). Distribution of the 10 warning signs by group is shown in Table 4.

Table 4
Distribution of the Warning Signs of the Groups According to Jeffrey Modell Foundation

Criteria			PID		SID		Control Group		<i>P</i>
			N	%	n	%	n	%	
Oral antibiotic	Two or more months on antibiotics with little effect.	No	53	54.1	33	51.6	192	96	<0.001
		Yes	45	45.9	31	48.4	8	4	
Abscess	Recurrent, deep skin or organ abscesses	No	89	90.8	60	93.8	199	99.5	<0.001
		Yes	9	9.2	4	6.3	1	0.5	
Growing	Failure of an infant to gain weight or grow normally	No	61	62.2	29	45.3	197	98.5	<0.001
		Yes	37	37.8	35	54.7	3	1.5	
Thrush	Persistent thrush in mouth or fungal infection on skin	No	55	56.1	34	53.1	188	94	<0.001
		Yes	43	43.9	30	46.9	12	6	
IV antibiotic	Need for intravenous antibiotics to clear infections	No	17	17.3	0	0	170	85	<0.001
		Yes	81	82.7	64	100	30	15	
Septicemia	Two or more deep-seated infections including septicemia	No	88	89.8	55	85.9	200	100	<0.001
		Yes	10	10.2	9	14.1	0	0	
PI in family	A family history of PI	No	73	74.5	60	93.8	196	98	<0.001
		yes	25	25.5	4	6.3	4	2	
Ear infection	Four or more new ear infections within 1 year	No	90	91.8	63	98.4	199	99.5	<0.001
		Yes	8	8.2	1	1.6	1	0,5	
Sinus infection	Two or more serious sinus infections within 1 year	No	89	90.8	63	98.4	198	99	<0.001
		Yes	9	9.2	1	1.6	2	1	
Pneumonia	Two or more pneumonias within 1 year	No	35	35.7	5	7.8	193	96.5	<0.001
		Yes	63	64.3	59	92.2	7	3.5	

Answers of the PID group to questions regarding the 10 warning signs of PID were estimated as observed and expected values. The criteria that exhibited a significant difference included the rates of recurrent deep cutaneous or organ abscesses, failure to thrive, need for IV antibiotics, ≥ 2 deep tissue infections including septicemia, a family history of PID, ≥ 4 episodes of otitis in one year, ≥ 2 episodes of sinusitis in one year, and ≥ 2 lower respiratory tract infections in one year. The criteria that exhibited the least differences between expected and observed values included the rates of being on oral antibiotics for ≥ 2 months with little effect, and a history of persistent thrush or cutaneous fungal infections. Among the warning signs of PID developed by the JMF, the criteria with the most statistically significant differences included the rates of ≥ 4 episodes of otitis in one year, recurrent deep cutaneous or organ abscesses, ≥ 2 episodes of sinusitis in one year, and ≥ 2 deep tissue infections including septicemia. These four warning signs seem to be less indicative of PID according to the statistics.

Odds ratios (ORs) for the 10 warning signs were estimated for both PID and SID groups versus the control group. Being on oral antibiotics was increased by 20.37-fold in the PID group compared with the control group, with a sensitivity of 45.91% and specificity of 96.00%. In the SID group, being on oral antibiotics was increased by 22.54-fold, with a sensitivity of 48.43% and specificity of 96.00%. OR, sensitivity, and specificity of the 10 warning signs are shown in Table 5.

Table 5
OR, sensitivity and specificity of the 10 warning signs developed by JMF

		OR (95% CI)	Sensitivity	Specificity	PPV	NPV	Accuracy
Oral antibiotic	PID	20.37 (9.05-45.86)	45.91	96.00	84.90	78.36	79.53
	SID	22.54 (9.53-53.30)	48.43	96.00	79.48	85.33	84.47
Abscess	PID	20.12 (2.51-61.24)	4.32	98.88	90.00	30.90	32.88
	SID	13.26 (1.45-120.96)	6.25	99.50	80.00	76.83	76.89
Growing	PID	39.83 (11.86-133.71)	37.75	98.50	92.50	76.35	78.52
	SID	79.25 (22.89-274.37)	54.68	98.50	92.10	87.16	87.87
Thrush	PID	12.24 (6.04-24.83)	43.87	94.00	78.18	77.36	77.51
	SID	13.82 (6.44-29.63)	46.87	94.00	71.42	84.68	82.57
IV antibiotic	PID	27.00 (14.07-51.78)	82.65	85.00	72.97	90.90	84.22
	SID	31.13 (2.33-42.21)	100.00	85.00	68.08	100.00	88.63
Septicemia	PID	47.57 (2.75-820.92)	10.10	100.00	100.00	69.20	70.23
	SID	68.63 (3.93-197.78)	14.06	100.00	100.00	78.43	79.16
PI in family	PID	16.78 (5.64-49.86)	25.51	98.00	86.20	72.86	74.16
	SID	3.26 (0.79-13.45)	6.25	98.00	50.00	76.56	75.75
Ear infection	PID	17.68 (2.18-143.54)	8.16	99.50	88.88	68.85	69.46
	SID	3.15 (0.19-51.23)	1.56	99.50	50.00	75.95	75.75
Sinus infection	PID	10.01 (2.12-47.28)	9.18	99.00	81.81	68.99	69.46
	SID	1.57 (0.14-17.62)	1.56	99.00	33.33	75.86	75.37
Pneumonia	PID	49.62 (21.00-117.26)	64.28	75.39	50.00	84.64	72.31
	SID	325.34 (99.56-1063.13)	92.18	96.50	89.39	97.47	95.45
OR: Odds ratio PPV: Positive predictive value NPV: Negative predictive value IV: Intravenous							

According to the receiver operating characteristic (ROC) results, the predictive area under the ROC curve (AUC) value for the JMF score in the PID and SID groups was determined to be 0.974 (95% confidence interval (CI), 0.959–0.989), which was statistically significant ($p < 0.001$). The ROC curve for the JMF score in the PID and SID

groups is shown in Figure 1. The cut-off for the JMF score in the PID and SID groups was determined to be 1.5. In accordance with the determined cut-off, sensitivity was found to be 0.920% and specificity was 0.935%.

When parental consanguinity was examined between the groups, it was revealed that 41 patients (41.8%) in the PID group, 16 patients (25%) in the SID group, and 18 patients (9%) in the control group had consanguineous parents. The distribution of family history features of the patients who participated in the study by group is shown in Table 6. A statistically significant difference was determined between the groups ($p < 0.001$). The rate of parental consanguinity of the PID group was higher than both SID and control groups. The rate of parental consanguinity in the PID group was 7.27-fold higher than that of the control group (OR: 7.27; 95% CI: 3.87–13.64). This rate was 3.37-fold higher in the SID group (95% CI: 1.60–7.09).

Table 6
Distribution of Family History Features of the Patients and the controls by the Groups

	Control Group		PID		SID		Total		P
	N	%	n	%	n	%	n	%	
Consanguinity	18	24	41	54.6	16	21.4	75	37.5	<0.001
First Degree	14	77.8	33	80.5	10	62.5	57	76	
Second Degree	1	5.6	3	7.3	1	6.3	5	6.7	
Third Degree	1	5.6	3	7.3	4	25	8	10.7	
Fourth Degree	2	11.1	2	4.9	1	6.3	5	6.7	
Family history of premature death (Yes/No)	11/189	5.5/94.5	66/32	67.3/32.7	38/26	59.4/40.6	115		<0.001
Family history of tuberculosis (Yes/No)	5/195	2.5/97.5	16/82	4.7/95.3	33/61	4.7/95.3	54		<0.001
Family history of rheumatic diseases (Yes/No)	23/177	11.5/88.5	28/70	28.6/71.4	22/42	34.4/65.6	73		<0.001
Family history of allergic diseases (Yes/No)	98/102	49/51	31/67	31.6/68.4	19/45	29.7/70.3	148		<0.001
Family history of Malignancy (Yes/No)	24/176	12/88	28/70	28.6/71.4	19/45	29.7/70.3	71		<0.001

In both the PID and SID groups, the rates of a family history of early death, rheumatic diseases, and malignancies were higher, compared with the control group ($p < 0.05$). The rate of a family history of tuberculosis was

significantly higher in the PID group, compared with the SID and control groups, whereas the rate of a family history of allergic diseases was significantly higher in the control group compared with the PID and SID groups.

The most common group of diseases in our PID patients was that of immunodeficiencies due to antibody deficiency, as shown in Table 3. Clinical characteristics of the patients with SID are shown in Figure 2

Discussion

Currently, the awareness level of physicians and healthcare professionals about PID, and their experience with a clinical approach to a patient with PID are still insufficient. For this reason, the history, features, and physical examination findings from patients, as well as expert opinions were combined, and the “10 Warning Signs of Primary Immunodeficiency Diseases”, which has significantly contributed to the diagnosis of PID, was defined by the JMF [9]. In our study, the JMF scores of PID and SID patients calculated via these warning signs were statistically significantly higher compared with the control group. In addition, we found that parental consanguinity and a family history of tuberculosis, chronic diarrhea may also be warning signs of PID.

Every patient with suspected PID should be asked in detail for information related to the “10 warning signs” checklist during history taking. It has been proposed that PID should be investigated when ≥ 2 warning signs are present [10]. In our study, the JMF scores of both the PID and SID groups were significantly higher than the control group. The cut-off for JMF score in terms of PID and SID was determined to be 1.5, with a sensitivity of 92% and specificity of 93.5%. According to our study results, we hypothesize that the JMF criteria are a guiding tool not only for PID patients but also for SID patients. In addition, in a study by Reda et al., at least one of the 10 warning signs was observed in all PID patients, whereas 28% of patients without PID had no warning sign [11]. In our study, all patients in the SID group had warning signs of immunodeficiency, whereas two of our patients in the PID group had no warning sign. The reason for these two patients with a JMF score of zero being investigated was a history of chronic diarrhea. Considering this, we propose that a history of chronic diarrhea should be included in the warning signs of PID. We attribute the presence of JMF warning signs in all patients in the SID group to the fact that they all had a more severe course that required IV immunoglobulin therapy.

Training programs aimed at increasing awareness of PID should target physicians who may discover a family history of PID, parental consanguinity, and a family history of early sibling deaths in societies where consanguineous marriages are common [11]. In a study from Egypt by Reda et al., 60% of PID patients had consanguineous parents [11]. In our country, the rate of parental consanguinity in PID patients has been found to be 14.3–37.5% [4]. In our study, the rate of parental consanguinity in PID cases was 41.8%. This rate was significantly higher compared with the control and SID groups, and we propose that parental consanguinity may be a warning sign of autosomal recessive-inherited PID for our region. According to a study conducted by Subbarayan et al. [6], one of the strongest identifiers of PID was a family history of immunodeficiency. In general, such a family history is 18 times more common in children with PID, compared to those without any identifiable PID. In our study, we determined a family history was 16.78 times more common in the PID patients compared with the control group. We propose that screening for PID would be important in the presence of a family history of PID, even when it exists alone.

In the Subbarayan et al. study, the most common of the 10 warning signs was the need for IV antibiotics. The second most common warning sign was a family history of PID (34%), followed by failure to thrive (31%) [6]. Similarly, in the study by Reda et al., the most common warning sign was the need for IV antibiotics (92%). In our

study, as in these two studies, the most common warning sign was the need for IV antibiotics (82.9%). The second most common warning sign was having ≥ 2 lower respiratory tract infections in one year (64.3%), and the third was being on oral antibiotics for longer than two months with little effect (45.9%). Our findings confirm that the 10 warning signs may be used for the diagnosis of PID, although in a different order of frequency, and that different frequencies may be reported in different studies.

Frequent infections, a more severe course than expected, long-lasting infections, the occurrence of unexpected or severe complications due to infections, incomplete recovery with antibiotic treatment, the need for prolonged use of antibiotics, chronic courses of infectious diseases, and the occurrence of infections with unusual agents may also be associated with PID diseases [12]. Infections usually recover rapidly and without complications in children with a healthy immune system and no other risk factors [13]. In our study, among the JMF warning signs, frequent recurrent upper respiratory tract infections were significant in the PID group, and frequent recurrent lower respiratory tract infections and failure to thrive were significant in the SID group. This may be due to the high rate of antibody deficiency, and the presence of accompanying conditions (tracheostomy, epilepsy) in our SID patients.

Comprehensive evaluation of family history and clinical features may be helpful for the early diagnosis of PID disease [14,15]. However, absence of a family history of immunodeficiency does not exclude the presence of PID. Since the majority of PID diseases are inherited, the presence of a similar disease, as well as the age and sex of affected individuals are important. In the study by Yorulmaz et al., 3.8% of PID patients were found to have a family member with PID [4]. This was higher in the PID and SID groups in our study, with rates of 25.5% and 6.3%, respectively. These higher rates in our study may be due to asking not only about siblings and parents but also about the siblings of the parents and their children. In our study, the rate in the PID group was significantly higher than both the control and SID groups. A patient with a history of frequent infections and a family history of PID should be evaluated for PID.

In a study conducted by Yorulmaz et al. in Konya, the rate of sibling death among patients with combined immunodeficiency (CID) was 7.5% [4]. However, in the Reda study from Egypt, 21.7% of the patients had sibling deaths. This 3-fold higher rate in Egypt may be related to a higher rate of consanguineous marriages and the level of community and economic development. The highest rate of sibling death was determined to be 50% in those with CID [11]. Rates of early death in the family history were also evaluated in our study. According to the results, 66 (67.3%) of PID cases, 38 (59.4%) of SID cases, and 11 (5.5%) in the control group had a family history of early death. In our study, the reason for the high rate of a family history of early death may be the inclusion of questions about siblings of the parents and their children. Therefore, we think that an extended family history of early sibling death may be an important warning sign for the diagnosis of immunodeficient patients.

Worldwide, the mean duration of diagnostic delay between the onset of symptoms and diagnosis in PID diseases is 4.08 years. The biggest factor in an 8–10-year delay in the diagnosis of PID diseases after the onset of symptoms was the low level of physician awareness of these diseases [9]. In our study, the mean duration of diagnostic delay for the PID and SID groups was 21 months, with no significant difference between the groups. Given that a delay in diagnosis can significantly increase morbidity and mortality, we may conclude that the index of suspicion for PID on the part of physicians in our region is similar to that in other centers.

In the literature, PID diseases have been reported to be more common in males than in females [6]. The predominance of males results not from PIDs inherited in an autosomal recessive manner, but from X-linked PIDs.

In our study of PID patients, 49% were female and 51% were male, with no statistically significant difference. We consider that this result was due to the high rate of consanguineous marriages in our study.

Antibody deficiencies are the most common subtype of PID [16]; they were also the most common PID group in our study. This is consistent with the European Society for Immunodeficiencies and the JMF databases, and our results are consistent with previous study results. However, given that the incidence of allergic, autoimmune, and hematological diseases, as well as the incidence of malignancies are high among PID patients, the medical history should be scrutinized in this respect [17]. Studies suggest the need for some additional warning signs to facilitate early diagnosis in such patients [18,19]. In our study, the rates of a family history of allergy, rheumatic diseases, and malignancy in the PID and SID groups were significantly higher than in the control group. Patients with immunodeficiencies may present with infectious diseases, and also with immune dysregulation diseases and malignancies; we consider that these diseases should also be considered as warning signs of immunodeficiency.

In conclusion, early diagnosis of PID will allow effective treatment of these diseases. We agree that the 10 warning signs of PID diseases defined by the JMF are important for the early diagnosis of PID. From our study results, a family history of parental consanguinity or tuberculosis may also be warning signs of PID, and a history of chronic diarrhea should be included. Studies from different immunology centers may clarify these additions. This approach will allow early diagnosis of PID and, thus, early and effective treatment, which will allow patients to reach adulthood before the development of organ injury.

Abbreviations

JMF: Jeffrey Modell Foundation

PID: Primary immunodeficiency

SID: Secondary immunodeficiency

CID: Combined immunodeficiency

IV: Intravenous

IUIS: International Union of Immunological Societies

Declarations

Author information

Affiliations

Necmettin Erbakan University, Meram Medical School, Division of Pediatrician, Konya, Turkey

Fadime Ceyda Eldeniz

Necmettin Erbakan University, Meram Medical School, Division of Pediatric Allergy and Immunology, 42080, Konya, Turkey

Yahya Gül, Şükrü Nail Güner, Sevgi Keleş, İsmail Reisli

Selcuk University, Medical Faculty Hospital, Pediatrician, Konya, Turkey

Alaattin Yorulmaz

Funding

No funding was received.

Conflicts of Interest

The authors declare that they have no conflicts of interest.

Data Availability

The data can be made available from the corresponding author on reasonable request

Code Availability

Not applicable

Contributions

Dr. Fadime Ceyda Eldeniz: analyzed and interpreted the study data, and wrote the paper.

Dr. İsmail Reisli : analyzed and interpreted the study data, and drafted the manuscript.

Dr. Sevgi Keleş, Dr. Şükrü Nail Güner: analyzed and interpreted the study data, and provided statistical support.

Dr. Yahya Gül, Dr. Alaattin Yorulmaz: acquisition and interpretation of the study data.

All authors read and approved the final manuscript.

Ethics Approval

This prospective research study was conducted using data obtained for clinical purposes. The study was approved by Necmettin Erbakan University Meram Medical School Ethics Committee (Date: 06.26.2020/No: 2020/2599).

Consent to participate

Not applicable

Consent to publication

Not applicable

References

1. Tangye SG, Al-Herz W, Bousfiha A, Chatila T, Cunningham-Rundles C, Etzioni A, Franco JL, Holland SM, Klein C, Morio T. Human inborn errors of immunity: 2019 update on the classification from the International Union

- of Immunological Societies Expert Committee. *J Clin Immunol*. 2020;40:24–64.
2. Bousfiha AA, Jeddane L, Ailal F, Benhsaien I, Mahlaoui N, Casanova JL, Abel L. Primary immunodeficiency diseases worldwide: more common than generally thought. *J Clin Immunol*. 2013;33:1–7.
 3. Modell V, Orange JS, Quinn J, Modell F. Global report on primary immunodeficiencies: update from the Jeffrey Modell Centers Network on disease classification, regional trends, treatment modalities, and physician reported outcomes. *Immunol Res*. 2018;66:367–80.
 4. Yorulmaz A, Artaç H, Kara R, Keleş S, Reisli İ. Retrospective evaluation of 1054 patients with primary immunodeficiency. *Asthma Allergy Immunol*. 2008;6:127–34.
 5. Kilic SS, Ozel M, Hafizoglu D, Karaca NE, Aksu G, Kutukculer N. The prevalences and patient characteristics of primary immunodeficiency diseases in Turkey-two centers study. *J Clin Immunol*. 2013;33:74–83.
 6. Subbarayan A, Colarusso G, Hughes SM, Gennery AR, Slatter M, Cant AJ, Arkwright PD. Clinical features that identify children with primary immunodeficiency diseases. *Pediatrics*. 2011;127:810–6.
 7. Tavakol M, Jamee M, Azizi G, Sadri H, Bagheri Y, Zaki-Dizaji M, et al. Diagnostic approach to the patients with suspected primary immunodeficiency. *Endocr Metab Immune Disord Drug Targets*. 2020;20:157–71.
 8. Bousfiha A, Jeddane L, Picard C, Al-Herz W, Ailal F, Chatila T, Cunningham-Rundles C, Etzioni A, Franco JL, Holland SM. Human inborn errors of immunity: 2019 update of the IUIS phenotypical classification. *J Clin Immunol*. 2020;40(1):66–81.
 9. Modell V, Quinn J, Ginsberg G, Gladue R, Orange J, Modell F. Modeling strategy to identify patients with primary immunodeficiency utilizing risk management and outcome measurement. *Immunol Res*. 2017;65:713–20.
 10. O'Sullivan MD, Cant AJ. The 10 warning signs: time for a change? *Current Opinion in Allergy and Clinical Immunology*. 2012; 12:588–594.
 11. Reda SM, El-Ghoneimy DH, Afifi HM. Clinical predictors of primary immunodeficiency diseases in children. *Allergy Asthma Immunol Res*. 2013;5:88–95.
 12. Woroniecka M, Ballow M. Office evaluation of children with recurrent infection. *Pediatr Clin North Am*. 2000;47:1211–24.
 13. Ballow M. Approach to the patient with recurrent infections. *Clin Rev Allergy Immunol*. 2008;34:129–40.
 14. Karakoc-Aydiner E, Ozen A, Baris S, Ercan H, Ozdemir C, Barlan I. Alteration in humoral immunity is common among family members of patients with common variable immunodeficiency. *J Investig Allergol Clin Immunol*. 2014;24:346–51.
 15. Aytekin G, Budh DP. X-Linked Immunodeficiency. In: StatPearls. StatPearls Publishing; 2021.
 16. Modell V, Gee B, Lewis DB, Orange JS, Roifman CM, Routes JM, et al. Global study of primary immunodeficiency diseases (PI)—diagnosis, treatment, and economic impact: an updated report from the Jeffrey Modell Foundation. *Immunol Res*. 2011;51:61–70.
 17. Aghamohammadi A, Moghaddam ZG, Abolhassani H, Hallaji Z, Mortazavi H, Pourhamdi S, Mohammadinejad P, Rezaei N. Investigation of underlying primary immunodeficiencies in patients with severe atopic dermatitis. *Allergol Immunopathol*. 2014;42:336–41.
 18. Arkwright PD, Gennery AR. Ten warning signs of primary immunodeficiency: a new paradigm is needed for the 21st century. *Ann N Y Acad Sci*. 2011; Nov 1238:7-14.

19. Bahrami A, Sayyahfar S, Soltani Z, Khodadost M, Moazzami B, Rezaei N. Evaluation of the frequency and diagnostic delay of primary immunodeficiency disorders among suspected patients based on the 10 warning sign criteria: A cross-sectional study in Iran. *Allergol Immunopathol.* 2020;48:711–9.

Figures

Figure 1

The ROC curve for JMF score in the PID and SID groups

Figure 2

The diagnosis and/or accompanying diseases* of the patients with SID