Psychopathology analysis of 23 cases with COVID-19 pneumonia accompanied by abnormal brain imagine

Fengli Sun (✉ sunfengli1980@163.com)

Zhejiang Province Mental Health Center

Xinsheng Chen

Tianyou hospital

Hongwei Ren

Tianyou Hospital

Jianning Liu

Xuhua CDC of Hangzhou

Weidong Jin

Zhejiang Chinese Medical University

Short report

Keywords: COVID-19, Brain CT, Psychopathology

Posted Date: December 4th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-119056/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background

COVID-19 broke out in Wuhan, China in late 2019. the persons involved maybe influenced in psychology. Abnormality of central nervous system (CNS) may promote appeared more psychological symptoms in patients with COVID-19 pneumonia.

Materials

All the cases were from the hospital where the author was located and diagnosed COVID-19 pneumonia with abnormality of cranial CT examination.

Result

All 23 patients performance mental symptoms of consciousness disturbance, cognitive impairment, anxiety and depression, sleep disorder. The more and serious mental symptoms correlated with dangerous and severe or severe type of COVID-19 pneumonia, more physical disease.

Conclusion

COVID-19 pneumonia patients with cranial CT abnormality always had mental symptoms.

Background

The novel coronavirus pneumonia (Coronavirus disease 2019, COVID-19) broke out in Wuhan, China in late 2019. Its propagation speed and severity have a significant impact on human beings.

COVID-19 is an emerging respiratory infectious disease caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which was first detected in early December 2019 in Wuhan, China. As of April 6, 2020, COVID-19 had quickly spread to the majority of countries worldwide, affected more than 1.1 million individuals, and caused nearly 63,000 deaths. Although studies have described the clinical characteristics of patients with COVID-19 [1, 2]. At same times, this kind of influence will also cause a kind of serious stress and cause a variety of psychological problems [3, 4]. In most cases, there are various manifestations, such as anxiety, depression, compulsion, somatization, etc., and more and more common is sleep disorder [3, 4, 5, 6]. The reason of these psychological problems during COVID-19 was both stress and some biological reasons. Because a certain number of new crown pneumonia patients have nervous system symptoms [7, 8], including three main aspects of nervous system symptoms. Obviously, these neurological symptoms promote the appearance of psychological symptoms to some extent. The pathological mechanism of neurological symptoms may be from the CNS invasion of SARS-CoV-2, similar to SARS and MERS viruses. As with other respiratory viruses, SARS-CoV-2 may enter the CNS through the hematogenous or retrograde neuronal route [7]. And the patients with

novel coronavirus pneumonia was more likely to develop mental symptoms when neurological symptoms appear. It is an important neurobiological basis when the nervous system, especially the central nervous system, has organic symptoms [8]. We report mental symptoms of 23 patients with novel coronavirus pneumonia and brain CT abnormality and analyze their, explore the relationship between the brain imaging characteristics and mental symptoms.

Materials

All the cases were from the hospital where the authors was located. They were Tianyou Hospital Affiliated to Wuhan University of science and technology and Tongde Hospital Affiliated to Zhejiang University of traditional Chinese medicine. All patients were diagnosed novel coronavirus pneumonia, and they were all asked for psychiatric consultation because of mental problems and they also had been completed the cranial CT examination. The male patients was 14 cases, female was 9 cases, The age was 51–87 years old.COVID-19 pneumonia is classified as dangerous and severe, severe, mild, and common type. The other physical diseases also had been diagnosed by physicians. All patients had positive results of CT. See Table 1 for all patient information.This clinical study was approved by ethic committee of Zhejiang Province Tongde Hospital

Results

1. Mental symptoms of patients: the most common are sleep disorders, anxiety and depression, and the most serious is the disturbance of consciousness, including coma and delirium. The most difficult to manage is psychomotor excitement, which brings great difficulties to treatment and management. There are other cognitive disorders. A few had compulsive symptoms and psychotic manifestations.

11 cases was company with disturbance of consciousness. 9 cases had cognitive disorders,2 cases of which had been diagnosed as Alzheimer's disease ,and 1 of which had been diagnosed as vascular dementia and 1 of which had been diagnosed as mixed dementia before hospitalization.

Most patients had sleep disorders, anxiety and depression.

2. Brain CT results: All patients had positive results. Three results are common. One is senile change, brain atrophy, including ventricular enlargement, widened sulcus, and leukoaraization. The second is cerebral ischemia, including infarct and softening focus, which occurs in the basal ganglia and around the ventricles. The most patients had both brain atrophy and cerebral ischemia.See Figure1-3.

3. COVID-19 pneumonia, psychiatric symptoms and brain imaging: (1) the influence of covid-19 pneumonia on mental symptoms: most of the patients with consciousness disturbance were critically ill,which type always was dangerous and heavy, almost all patients had sleep disorders, anxiety and depression, and some patients had delirium with psychomotor excitement. (2) The relationship between brain image and mental symptoms: the patients with severe mental symptoms usually occur in the old age and have more physical diseases,especially both brain atrophy and cerebral infarction.33The some patients had cognitive impairment,which mainly related the brain CT abnormality that may further aggravated in case of COVID-19 pneumonia.(4)The most patients had anxiety and depression related to both stress and biology impacts of COVID-19 pneumonia see table2.

Discussion

In a new study, of 214 patients included 88 (41.1%) had severe infection and 126 (58.9%) had nonsevere infection. Of these, 78 (36.4%) had various neurologic manifestations that involved CNS, PNS, and skeletal muscles[7]. And 40% patients with covid-19 pneumonia had various psychological manifestations that involved depression, anxiety, insomnia[9] ,suicide[10].These result showed that covid-19 pneumonia influenced both physical and psychological aspects.

Disturbance of consciousness is not only a symptom of central nervous system, but also a mental symptom, but more often, it is considered to be a more serious clinical manifestation. It strongly suggests that the central nervous system has been compromised. In our 23 case reports, nearly half of the cases (11 / 23) had disturbance of consciousness. Most of these patients had severe infection, were older and had a variety of physical diseases. These conclusions were similar to those of previous studies [7], but. There is something special about the cases we report, which is that the patients have already suffered from central nervous system injury that is brain atrophy, cerebral infarction or both, as seen by head CT scan. Therefore, this is also an important factor for these patients to have disturbance of consciousness.

Moreover, nervous system manifestations were significantly more common in severe infections compared with nonsevere infections [7]. They included acute cerebrovascular disease, patients with ischemic stroke or with cerebral hemorrhage or with ischemic, impaired consciousness and skeletal muscle injury. In the severe group, some patient had a seizure characterized by a sudden onset of limb twitching, foaming in the mouth, and loss of consciousness[7, 8],especially in acute stage of covid-19 [11].But we just report disturbance of consciousness,and also report the disorder of cognition,which also suggest more serious clinical manifestation.

The delirium was one kind of consciousness disturbance.The study showed that 28% patients with severe covid-19 presented confusion during the acute illness[11]. When data for patients with COVID-19 were examined (including preprint data), there was evidence for delirium (confusion in 26 [65%] of 40 intensive care unit patients and agitation in 40 [69%] of 58 intensive care unit patients in one study, and altered consciousness in 17 [21%] of 82 patients who subsequently died in another study)[11]. When patients presented delirium,they always had disturbance of behavior with agitation,which also was import reason of consultation liaison psychiatry.

As to disturbance of consciousness, the patents with impaired cognition and severe infections,were higher older,had more physical disease,more impaired CNS that be seen by brain CT.The study found 34.1% patients with severe infection of covid-19 had impaired memory during acute stage and 18.9% in the post-illness stage[11]. Generally,the impaired cognition was related to exited brain atrophy and cerebral ischemic.But we also do not know that neurologic manifestations are caused by the virus directly or by the pulmonary disease or other organ damage indirectly[8].

It was certain that impairment of CNS induce impaired cognition. In the patients reported,some patients had been diagnosed as Alzheimer's disease, mixed dementia which were disease characterized by disorder of cognition. And some patients ad been diagnosed as Parkinson's disease which were disease possibly induced disorder of cognition.But what cannot be ruled out is that covid-19 pneumonia can cause cognitive impairment in these elderly patients in a special way, and moreover these patients already have the damage that can be seen by head CT.

Symptoms that are less severe than consciousness and cognitive impairment are negative emotional symptoms.Prevalence of depression and anxiety symptoms were diagnosed in 35.9% and 38.5% of the patients, respectively[9].The most of patients we reported had the depression and anxiety.Coronavirus disease 2019 (COVID-19) has affected everyone around the world while often disrupting medical practices in non-COVID-19 cases and specifically in prenatal care. The systematic review revealed that during the acute illness, common symptoms among patients admitted to hospital for SARS or MERS included confusion (36 [27.9%; 95% CI 20.5–36.0] of 129 patients), depressed mood (42 [32.6%; 24.7–40.9] of 129), anxiety (46 [35.7%; 27.6–44.2] of 129), impaired memory (44 [34.1%; 26.2–42.5] of 129), and insomnia (54 [41.9%; 22.5–50.5] of 129)[11]. These results

showed that the patients with covid-19 experienced more negative affection, such as depression, anxiety. Our reports also showed most patients experienced depression and anxiety.

There was a findings showed a high rate of depression and anxiety in COVID-19 patients[9]. The prevalence of depression was 35.9%: 20.5% for mild depression, 12.8% for moderate depression, and 2.6% for severe depression. Almost four out of 10 (38.5%) patients were diagnosed with anxiety: 20.5% with mild anxiety, 12.8% with moderate anxiety, and 2.6% with severe anxiety. Nearly a quarter (24.3%) of the patients were diagnosed as having both depression and anxiety. Multivariate linear regression found having family members diagnosed with COVID-19 and family members who died from COVID-19 were independent predictors of both higher depression severity index and anxiety score. Being female was also independently associated with depression severity index in COVID-19 patients. It was very pity that we did not analyzed the risk with depression and anxiety. But ratio of our patients with depression and anxiety was more than 40%. It may also related to their special findings by head CT.

The patients with brain atrophy and cerebral ischemia always presented emotional regulation disturbance. There was a study revealed brain structural abnormalities in emotional regulation and sensory processing regions of anxious depression, which may suggested distinct neurobiological mechanisms of this MDD specifier and could help explain different clinical manifestations in anxious depression from pure depression[12]. But psychological distress of covid-19 may had importer role, as these findings of previous studies showing that public health emergencies can cause mental-health problems, such as the Wenchuan and Lushan earthquakes [13], the 2014 Ebola Outbreak [14, 15], and Severe Acute Respiratory Syndrome (SARS) [16].

Sleep disorder was very common psychological symptom without specificity. About 29.2% (95% CI, 28.8%-29.6%) general population had insomnia. The people with occupational exposure risks and residents in Hubei province had increased odds of symptoms of insomnia (adjusted ORs, 1.60 [95% CI, 1.45–1.77]. The confirmed patients with covid-19 had more increased odds of symptoms of insomnia (adjusted ORs, 3.06 [95% CI, 1.73–5.43][17]. The systematic review 42% patients had insomnia in acute illness, 100% patients had sleep disorder in the post-illness stage[11]. The patients we reported almost had sleep disorder. For patients, besides the impact of psychological factors[18], there may be more important biological factors[19], especially those with severe pneumonia, which induced insomnia, anxiety and depression.

COVID-19 may impact both CNS and psychology. Despite the occurrence of symptoms-such as shortness of breath(SOB)-associated with severe manifestations of COVID-19, only the severities of smell and taste loss were associated with depressed mood and anxiety. These results may raise the novel possibility of emotional disturbance as a CNS manifestation of COVID-19 given trans-olfactory tract penetration of the central nervous system (CNS) by coronaviruses[20].

Unfortunately, our report does not have such data. There is no evaluation and test of olfactory and taste sense, no correlation analysis with negative emotion, and no correlation analysis between this and CT results of head, which is insufficient.

Key Point

COVID-19 pneumonia patients with cranial CT abnormality were easier to had mental symptoms, which include severe symptoms of consciousness disturbance, cognitive impairment, and common symptom of anxiety and

depression,sleep disorder.The special attention should been to it. The consultation liaison psychiatry will play import role.

Declarations

Acknowledgment

We thank Mr Xing baoping give us study idea and Mr Zhu Jianfeng help us in literature retrieval. We thank Mr Ma Yongchun help us in the final revision of the article.

Ethic statement

1.This study was approved by Ethic Committee of Zhejiang Province Hospital.Ethic certificate (LL20200408-023)

2.Consent for publication

All participants agree to publish their data and no conflict in any interests.

Conflict of interest statement

There were not any financial and non-financial competing interests.

Funding

Not Available

Authou's contribution

Our authors have different contributions to this article. Sun Fengli participated in collection of data and the writing of the article, Zhongwei Guo ,Xinsheng Chen ,Hongwei Ren ,Jianning Liu participated in the collection of clinical data. Jin Weidong participated in the design , statistical processing and the final revision of the article.

References

[1]World Health Organization. Coronavirus disease 2019 (COVID-19) situation report-76. Accessed April 6,2020. https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200405-sitrep-76-covid-19.pdf?sfvrsn=6ecf0977_2

[2]An Pan,Li Liu, Chaolong Wang, et al.Association of Public Health Interventions With the Epidemiology of the COVID-19 Outbreak in Wuhan, China.JAMA. 2020;323(19):1915-1923. doi:10.1001/jama.2020.6130

[3]Tait Shanafelt, Jonathan Ripp, Mickey Trockel. Understanding and Addressing Sources of Anxiety Among Health Care Professionals During the COVID-19Pandemic.JAMA. 2020;323(21):2133-2134.doi:10.1001/jama.2020.5893

[4]Mark A. Reger, Ian H. Stanley,Thomas E. Joiner. Suicide Mortality and Coronavirus Disease 2019—A Perfect Storm?JAMA Psychiatry. Published online April 10, 2020. doi:10.1001/jamapsychiatry.2020.1060

- [5]Li W, Yang Y, Liu ZH. Progression of mental health services during the COVID-19 outbreak in China. *Int J Biol Sci.* 2020;16(10):1732-1738. doi:10.7150/ijbs.45120
- [6]Yutao Xiang, Yu Jin, Teris Cheung.Joint International Collaboration to Combat Mental Health Challenges During the Coronavirus Disease 2019 Pandemic. *JAMA Psychiatry.* 2020; doi: 10.1001/jamapsychiatry.2020.1057
- [7]Ling Mao, Huijuan Jin, Mengdie Wang ,et al.Neurologic Manifestations of Hospitalized Patients With Coronavirus Disease 2019 in Wuhan, China.*JAMA Neurol.* 2020;77(6):683-690. doi:10.1001/jamaneurol.2020.1127
- [8]Samuel J. Pleasure, Ari J. Green, S. Andrew Josephson.The Spectrum of Neurologic Disease in the Severe Acute Respiratory Syndrome Coronavirus 2 Pandemic Infection,Neurologists Move to the Frontlines.*JAMA Neurol.* 2020;77(6):679-680. doi:10.1001/jamaneurol.2020.1065
- [9]XueDan Nie , Qin Wang , MinNan Wang ,et al.Anxiety and depression and its correlates in patients with coronavirus disease 2019 in Wuhan.*Int J Psychiatry Clin Pract.* 2020 Jul 14;1-6.doi: 10.1080/13651501.2020.1791345.Online ahead of print.
- [10]Mark A. Reger,Ian H. Stanley, Thomas E. Joiner, Suicide Mortality and Coronavirus Disease 2019—A Perfect Storm?*JAMA Psychiatry.* Published online April 10, 2020. doi:10.1001/jamapsychiatry.2020.1060
- [11]Jonathan P Rogers , Edward Chesney , Dominic Oliver , et al.Psychiatric and neuropsychiatric presentations associated with severe coronavirus infections: a systematic review and meta-analysis with comparison to the COVID-19 pandemic. *Lancet Psychiatry.* 2020 ;7(7):611-627.doi:10.1016/S2215-0366(20)30203-0.
- [12]Wei Peng , Zhiyun Jia , Xiaoqi Huang , et al .Brain structural abnormalities in emotional regulation and sensory processing regions associated with anxious depression.*Prog Neuropsychopharmacol Biol Psychiatry.* 2019 ;30; 94:109676. doi: 10.1016/j.pnpbp.2019.109676.
- [13]Xie Z, Xu J, Wu Z. Mental health problems among survivors in hard-hit areas of the 5.12 Wenchuan and 4.20 Lushan earthquakes. *J Ment Health.* 2017;26:43–49. doi:10.1080/09638237.2016.1276525
- [14]Ji D, Ji YJ, Duan XZ, et al. Prevalence of psychological symptoms among Ebola survivors and healthcare workers during the 2014-2015 Ebola outbreak in Sierra Leone: a cross-sectional study. *Oncotarget.* 2017;8:12784–12791. doi:10.18632/oncotarget.14498
- [15]Shultz JM, Baingana F, Neria Y. The 2014 Ebola outbreak and mental health: current status and recommended response. *JAMA.* 2015, 313:567–568. doi:10.1001/jama.2014.17934
- [16] Mak IW, Chu CM, Pan PC, et al. Long-term psychiatric morbidities among SARS survivors. *Gen Hosp Psychiatry.* 2009.31:318–326. doi:10.1016/j.genhosppsych.2009.03.001
- [17]Le Shi , Zheng-An Lu , Jian-Yu Que , et al. Prevalence of and Risk Factors Associated With Mental Health Symptoms Among the General Population in China During the Coronavirus Disease 2019 Pandemic. *JAMA Netw*

Open. 2020 ,1;3(7):e2014053. doi:10.1001/jamanetworkopen. 2020.14053.

[18]Cuiyan Wang , Riyu Pan , Xiaoyang Wan , et al.Immediate Psychological Responses and Associated Factors during the Initial Stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the General Population in China.Int J Environ Res Public Health. 2020 , 6;17(5):1729.doi: 10.3390/ijerph17051729.

[19]Abebaw Mengistu Yohannes.COPD patients in a COVID-19 society: depression and anxiety.Expert Rev Respir Med. 2020 , 5;1-3.doi: 10.1080/17476348.2020.1787835.

[20]Marlene M Speth , Thirza Singer-Cornelius , Michael Oberle , et al .Mood, anxiety and olfactory dysfunction in COVID-19: evidence of central nervous system involvement?Laryngoscope.. 2020 , 2;10.1002/lary.28964.doi: 10.1002/lary.28964.

Tables

Table 1. Primary clinical data of 23 patients

No	sex	age	Type of covid-19	Brain CT	Other physical diagnosis	Mental symptoms or diagnosis
1	F	84	severe	1. Bilateral basal ganglia and right radial coronal lacunar infarction, part of the softening focus. 2. Leukoencephalopathy; brain atrophy. 3. Bilateral internal carotid atherosclerosis	1. Type 2 diabetes mellitus; 2. Anemia; 3. Electrolyte metabolism disorder; 4. Hyperuricemia; 5. Internal carotid atherosclerosis; 6. Lacunar cerebral infarction; 7. Aortic atherosclerosis; 8. Coronary atherosclerosis; 9. Pleural effusion; 10. Pleural thickening and adhesion	1.Dementia, 2.uncoordinated disorderly excitement
2	M	82	severe	1. Lacunar infarction in bilateral basal ganglia. 2. Brain atrophy, leukoencephalopathy.	1. Parkinson's syndrome; 2. Sequelae of cerebral infarction; 3. Mild anemia; 4. Abnormal liver function; 5. Electrolyte disorder (low sodium, low chloride, low calcium); 6. Hypoproteinemia; 7. Prostatic hyperplasia	1.Dementia, 2.disturbance of consciousness
3	M	87	Dangerous and severe	1. Lacunar infarctions and ischemic foci in pontine, bilateral basal ganglia, corona radiata and central semiovale were observed. 2. Brain atrophy, leukoencephalopathy. 3. Bilateral vertebral artery and internal carotid artery calcification.	1. Hypertension grade 2 (extremely high risk); 2. Diabetes mellitus; 3. Alzheimer's disease; 4. Left foot soft tissue infection	1.Alzheimer's disease, 2.disturbance of consciousness, 3.anxiety, 4.sleep disorder
4	M	76	severe	1. Cerebral infarction in bilateral frontal lobe, right occipital lobe, left pons and thalamus may occur. 2. Lacunar infarction in bilateral basal ganglia. 3. Brain atrophy.	1. Acute cerebral infarction; 2. Coronary atherosclerotic heart disease; 3. Pacemaker implantation status; 4. Pericardial effusion; 5. Pleural effusion; 6. Abnormal liver function	1.disturbance of consciousness, 2.anxiety, 3.sleep disorder
5	M	79	Dangerous and severe	1. Lacunar infarction in bilateral basal ganglia. 2. Brain atrophy. 3. Bilateral internal carotid atherosclerosis.	1. Type 2 diabetes; 2. Hypokalemia, 3.Alzheimer's disease;	1.Dementia, 2.disturbance of consciousness, 3.anxiety. 4.Psychomotor excitement
6	M	56	severe	1. Lacunar infarction in right basal ganglia. 2. Brain	1. Sudden deafness; 2. Abnormal liver	1.Anxiety, 2.sleep

				atrophy.	function; 3. Hypertension; 4. Type 2 diabetes mellitus; 5. Vertigo; 6. Cerebral atrophy; 7. Lacunar infarction in the right basal ganglia	disorders
7	M	82	severe	1. Lacunar infarction in bilateral basal ganglia. 2. Brain atrophy, 3. leukoencephalopathy	1. Parkinson's syndrome; 2. Sequelae of cerebral infarction; 3. Mild anemia; 4. Abnormal liver function; 5. Electrolyte disorder (low sodium, low chloride, low calcium); 6. Hypoproteinemia; 7. Prostatic hyperplasia. 8. Vascular demantia.	1. Dementia, 2. depression, 3. sleep disorders
8	M	69	mild		1. Hyperlipidemia; 2. Acute attack of chronic gastritis; 3. Hypertension	1. Anxiety, 2. depression, 3. sleep disorders
9	F	70	mild	1. Lacunar infarctions and ischemic foci in bilateral basal ganglia and around lateral ventricles. 2. Brain atrophy. 3. Calcification of the right vertebral artery and bilateral internal carotid artery. 4. Bilateral ethmoid sinus and right frontal sinus were slightly inflamed.	1. Type 2 diabetes mellitus; 2. Right hip arthroplasty; 3. Left femoral head necrosis	1. Anxiety, 2. depression, 3. sleep disorders
10	F	60	mild	1. Lacunar infarction in bilateral basal ganglia. 2. Leukoencephalopathy. 3. Mild brain atrophy. 4. Bilateral internal carotid atherosclerosis.	1. Coronary heart disease; 2. Hypertensive cerebral infarction; 3. Hypokalemia; 4. Hepatic cyst; 5. Renal cyst; 6. Aortic atherosclerosis; 7. Carotid atherosclerosis	1. Anxiety, 2. depression, 3. sleep disorders
11	F	84	severe	1. Lacunar infarction in bilateral basal ganglia. 2. Leukoencephalopathy. 3. Mild brain atrophy. 4. Bilateral internal carotid atherosclerosis.	1. Type 1 respiratory failure; 2. Hypertension grade 3 (extremely high risk); 3. Cardiac insufficiency.; 4. Mixed dementia.	1. Dementia, 2. disturbance of consciousness
12	M	51	mild	1. Cerebral ischemia in bilateral frontal and parietal lobes. 2. Calcification in the white matter area of the posterior horn of the right lateral ventricle. 3. Left maxillary sinusitis.		1. Anxiety, 2. depression, 3. sleep disorders

13	F	79	severe	1. Bilateral basal ganglia lacunar infarction. 2. Leukoencephalopathy 3. brain atrophy.	1. Type 2 diabetes mellitus; 2. Hypertension grade 3 (extremely high risk); 3. Hypokalemia; 4. Hypoproteinemia; 5. Lacunar cerebral infarction	disturbance of consciousness
14	F	62	mild	1. Lacunar infarction in bilateral basal ganglia. 2. Mild brain atrophy. 3. Bilateral internal carotid atherosclerosis.	1. Type 2 diabetes mellitus; 2. Hypertension	1. State of anxiety, 2. state of depression
15	F	69	Dangerous and severe	1. Lacunar infarction in bilateral basal ganglia. 2. Leukoencephalopathy; brain atrophy. 3. Bilateral internal carotid atherosclerosis.	1. Femoral head fracture; 2. Hypertension; 3. Type 2 diabetes mellitus	1. Disturbance of consciousness, 2. dementia, 3. sleep disorder
16	M	64	severe	1. Lacunar infarction in bilateral basal ganglia. 2. Brain atrophy. 3. Bilateral internal carotid atherosclerosis. 4. Bilateral ethmoid sinus, maxillary sinus, sphenoid sinus and left frontal sinusitis	1. Bacterial pneumonia; 2. Hypoxemia; 3. Hypoproteinemia; 4. Hypertension grade 2 (high risk); 5. Renal dysfunction; 6. Liver dysfunction; 7. Organic psychosis; 8. Bilateral carotid atherosclerosis	1. Dementia, 2. anxiety, 3. psychomotor excitement, 4. sleep disorders
17	F	66	severe	1. Cerebral ischemic changes should be considered, combined with MRI examination. 2. Brain atrophy. 3. Bilateral internal carotid atherosclerosis.	1. Cervical spondylosis; 2. Mixed hyperlipidemia; 3. Sleep disorder; 4. Moderate risk group of hypertension; 5. Bilateral internal carotid atherosclerosis	1. Dementia, 2. anxiety, 3. depression, 4. sleep disorders
18	M	61	mild	1. Cerebral ischemic changes	Left ethmoid sinusitis	sleep disorders
19	M	82	Dangerous and severe	1. Large area cerebral infarction in right basal ganglia and frontal parietal lobe, lacunar infarction in pontine and bilateral basal ganglia radiation coronal area, 2. White matter disease, brain atrophy. 3. The wall calcification of bilateral internal carotid artery.	Cerebral infarction	Coma
20	M	61	Dangerous and severe	1. Lacunar infarction in bilateral basal ganglia. 2. Small calcification under the right frontal lobe cranial plate.	1. Acute exacerbation of chronic obstructive pulmonary disease; 2. Aortic atherosclerosis; 3. Pleural hypertrophy; 4. Lacunar infarction	1. Disturbance of consciousness, 2. psychomotor excitement, 3. anxiety, 4. sleep disorder

21	M	69	severe	1. Lacunar infarction in bilateral basal ganglia. 2. The right lateral ventricle was slightly enlarged,	1. Acute coronary syndrome; 2. Benign prostatic hyperplasia; 3. Cervical spondylosis	1. Disturbance of consciousness, 2. psychomotor excitement, 3. anxiety, 4. sleep disorder
22	F	70	severe	Leukoencephalopathy Brain atrophy	1. Parkinson's syndrome; 2. Sequelae of cerebral infarction; 3. Mild anemia; 4. Vascular dementia. 5. Electrolyte disorder (low sodium, low chloride, low calcium); 6. Abnormal liver function	1. Disturbance of consciousness, 2. psychomotor excitement
23	M	57	mild	Lacunar infarction in bilateral basal ganglia.	1. Migraine; 2. Uremia	Anxiety, sleep disorders

Table 2 Information of patients with disturbance of consciousness and impairment of cognition

	N	male	female	age	Average Number of physical disease	Average Number of CNS disease ¹	Average Number of diagnosis by Brain CT	Type of COVID-19 pneumonia
disturbance of consciousness	11	7	4	76.2	4.2	4.2	3.0	S ² :6 DS ³ :5
impairment of cognition	9	5	4	77.1	5.7	4.1	3.1	S ² :6 DS ³ :3

1. Number of CNS disease = Number of diagnosis by Brain CT + clinical CNS disease

2. S = severe

3. DS = dangerous and severe

Figures

Figure 1

Brain atrophy

Figure 1

Brain atrophy

Figure 2

Cerebral ischemia

Figure 2

Cerebral ischemia

Figure 3

Cerebral ischemia and brain atrophy

Figure 3

Cerebral ischemia and brain atrophy