Arabic version of the Hayling sentence completion test: Validation and factors associated with executive functions in a sample of the Lebanese adults

Sarah Boutros

Universite Saint-Esprit de Kaslik Faculte de Medicine et des Sciences Medicales

Emilio Al Hachem

Universite Saint-Esprit de Kaslik Faculte de Medicine et des Sciences Medicales

Joseph Matar

Universite Saint-Esprit de Kaslik Faculte de Medicine et des Sciences Medicales

Souheil Hallit (✉ souheilhallit@hotmail.com)

Universite Saint-Esprit de Kaslik Faculte de Medicine et des Sciences Medicales <https://orcid.org/0000-0001-6918-5689>

Hanna Matar

Universite Saint-Esprit de Kaslik Faculte de Medicine et des Sciences Medicales

Research

Keywords: Hayling sentence completion test, Executive functions, inhibitory control, normative data, problematic mobile phone use, air pollution

Posted Date: January 21st, 2020

DOI: <https://doi.org/10.21203/rs.2.21399/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

To validate the Arabic version of the Hayling Sentence Completion Test (HSCT) in healthy community-dwelling Lebanese adults, and identify factors that might affect executive functions (EF), mainly inhibitory control, in these participants.

Methods

Between August-December 2019, 350 participants were randomly selected. The Arabic version of the HSCT, divided into automatic and inhibition conditions, was used; in each condition, participants' response-time and number of errors committed were recorded.

Results

None of the scale items was removed. For the automatic condition, response-time items converged over one factor (α Cronbach=0.905) and number of errors converged over seven factors (α Cronbach=0.334). For the inhibition condition, response-time converged over one factor (α Cronbach=0.943) and number of errors converged over four factors (α Cronbach=0.728). Using electricity as a heating method inside the house was significantly associated with a lower response-time, whereas higher problematic mobile phone use was associated with higher response-time. Using wood as a heating system inside the house and higher problematic mobile phone use were associated with higher number of errors, while using Arabian incense (bakhour) inside the house was associated with lower number of errors.

Conclusion

We were able to validate the HSCT Arabic version for use in the Lebanese population. Problematic mobile phone use was associated with lower executive functions in terms of response-time and number of errors.

Background

Executive functions (EFs) include the production, preservation, and customization of plans aimed to accomplish the objectives of a given assignment ¹. It includes three main features: (1) inhibitory control, including self-control and interference control, (2) cognitive flexibility, also called mental flexibility, linked to creativity, and (3) working memory ^{1,2}. EFs are crucial skills for mental and physical well-being, school and life success, and for cognitive, social, and psychological growth and evolution ².

Inhibition represents one of the key skills of EFs. It involves the capacity to control one's attention, behavior, thoughts, and emotions, to suppress a strong internal tendency or external temptation, in order to do what's more relevant or essential ². In the absence of inhibition, we would be controlled by our impulses, settled tendencies of thought or behavior, by the environment that pulls us this way or the other

². Consequently, inhibition makes each one of us somehow unique, and allows us to proceed and behave deliberately, instead of being unintentional and impulsive creatures of tradition and routine ².

Inhibition has become crucial in several research domains within psychology and neurology. It is broadly recognized that inhibitory control is a multidimensional procedure encompassing at least three features, as defined by Bayard S et al.: « (1) Prepotent response inhibition (the ability to deliberately suppress dominant, automatic, or prepotent responses), (2) Resistance to distractor interference (the ability to resist interference from distracting information in the external environment), (3) Resistance to proactive interference (the ability to resist the intrusions of irrelevant information, that were once considered relevant, into our memory). » ³

Taking into account inhibition as a multidetermined construct emphasizes the exigency to invent validated clinical tools for the evaluation of its features ³. The Hayling Sentence Completion Test (HSCT), originally developed by Burgess & Shallice in 1996, is one of the available tools that evaluate inhibition. It is a measure of prepotent response inhibition that detects frontal lobe dysfunction. In this test, participants hear sentences in which the last word is absent. In the automatic condition, subjects are asked to complete sentences correctly (i.e., by a word solidly related to the sentence), reflecting the initiation of a semantically supported automatic response. In the inhibition condition, participants are asked to restrain themselves from saying the cueing word, and to complete the sentence with a totally unrelated word. To perform this task appropriately, subjects have to inhibit the related word and its semantic associates ⁴.

As far as we know, only five validated languages have been published for the HSCT in adults and elderly people: (1) English ⁵, (2) Spanish ⁶, (3) French ³, (4) Italian ⁷, and (5) Swedish ⁸. There are no HSCT normative data for the Arabic speakers to date, so it can not be used in Arabic-speaking clinical practice and research groups. Therefore, this study intends to create an Arabic version of the HSCT that is standardized to the linguistic and cultural realities of the Lebanese population.

EFs are negatively influenced by many factors, among which cognitive aging, which usually weakens executive functioning ⁹, and low family socioeconomic status (SES), which predicts a worse performance on tasks that evaluate them ¹⁰. In contrast, high educational levels improve scores in the HSCT ³, and multilingualism is associated with greater amount of controlled attention and inhibitory control, and can play a crucial role in protecting against the deterioration of executive functions with aging by helping to achieve cognitive reserve ¹¹. Furthermore, there is a positive correlation between healthy diet (e.g. whole grains, fish, fruits and/or vegetables) and executive functions, unlike unhealthy diet (e.g. fast food, sweetened beverages and red meat) that is associated with a decrease in executive functioning ¹². Moreover, a meta-analysis revealed a wide EFs deficit in overweight participants compared to healthy weight controls, specifically a deficit in inhibition and working memory in overweight participants ¹³, while better physical fitness is shown to improve cognitive flexibility ¹⁴. On another note, a negative correlation was exhibited between EFs and constant exposure to polluted air and traffic noise ¹⁵.

Studies concerning the relationship between long term mobile phone use and EFs are equivocal. Previous findings showed no evidence of a harmful effect of smartphone use on cognitive functioning. Instead, it suggests there could possibly be a favorable effect of phone use on cognitive performance and executive functioning, but the results of longitudinal analyses were confusing ¹⁶. Another study indicated that a daily exposure (2 h/day for 4 weeks) to electromagnetic fields (EMF) emitted by mobile phones (MP) has no effect on executive function ¹⁷. One more research done in 2017 by Hayashi Y et al. reveals that participants who constantly text while driving have low levels of executive function and high levels of impulsivity ¹⁸.

Some studies show a relationship between the use of some medications and the increase or decline in executive functions; proton pump inhibitors (PPI's) have varying degrees of influence on different cognitive domains and have associations with Alzheimer's Disease (AD) ¹⁹. On another note, Angiotensin I receptor blockers (ARBs) improve memory and executive function in comparison to other antihypertensive drugs, and it attenuates the decline of cognition over time ²⁰. Moreover, aspirin use is also associated with a preventive effect against cognitive decline, particularly in people at risk for developing dementia ²¹.

This study was conducted for several reasons, primarily because of the lack of an Arabic version of the HSCT that could be beneficial in our clinical practice. Another reason is the need to find potential relationships between various factors with executive functions, especially problematic mobile phone use as suggested by many previous studies, since smartphones have become, nowadays, a daily companion of people from all generations. Thus, it is important to conduct this study in Lebanon to be adapted to the ideas, customs and social behaviour of the Lebanese citizens. Hence, the objectives of the current study are to use the Arabic version of the HSCT in healthy community-dwelling Arabic-speaking adults in Lebanon, to check its validity compared to other versions of the test, as well as to identify risk factors that might affect the executive functions in these adults.

Methods

Study design

This study was conducted between August and December 2019. Participants were randomly chosen from the general population across Lebanon. Individuals included in this study were aged above 18 years old, having no known history of neurological or psychological impairment. Excluded patients were those with abnormal performances on the HSCT, which have been described in a wide variety of neurological, psychiatric, and neurodevelopmental conditions, such as Alzheimer's disease and mild cognitive impairment ²²⁻²⁴, brain traumatic injury ²⁵, cerebrovascular accidents²⁶, Parkinson's disease ²⁷, amyotrophic later sclerosis ²⁸, frontotemporal dementia ²⁸, schizophrenia ²⁹, bipolar disorder ³⁰, and autism spectrum disorder ³¹. Patients aged above 55 years old ³² underwent a mini mental state examination (MMSE); those obtaining a score of 24 or more were included in the study ³³.

Minimal sample size calculation

According to Comrey and Lee³⁴, a minimal sample of 5-10 observations is needed per item of the scale in order to validate a scale. Therefore, a minimum of 300 participants was needed for adequate statistical power, since the assessment of the automatic and inhibition conditions included 15 items each.

Questionnaires and variables

Data was collected via a personal interview, and subjects were tested with only one examiner in a quiet office. The first section assesses the socio-demographic characteristics, including age, gender, region, the number of rooms in the household and the number of persons living in it, the level of education, the number of mastered languages, the monthly income, and the medical history and chronic treatments.

The second section evaluates the diet of each participant, taking into consideration vegetables, fruits, dairy products, breakfast, eating 5 meals per day, sweets consumption, red meat, beverages, and fast food. Each variable is scored between 1 and 5 to get a minimum score of 9 and a maximum score of 45; higher scores indicate a healthier diet. This scale was inspired by the constituents of the DASH diet (Dietary Approaches to Stop Hypertension)³⁵.

The third section is the Arabic translation of the short form of International Physical Activity Questionnaire (IPAQ)³⁶ that asks about three specific types of activities: walking, moderate-intensity activities, and vigorous-intensity activities. Then participants are classified to have low, moderate, or high physical activity.

To be classified as 'moderate', participants should match one of the following:

a) 3 days of vigorous-intensity activity for a minimum of 20 minutes per day.

OR

b) 5 days of moderate-intensity activity and/or walking for a minimum of 30 minutes per day.

OR

c) 5 days of any combination of walking, moderate-intensity or vigorous intensity activities, achieving at least a total physical activity of 600 minutes/week.

To be classified as 'high', participants should match one of the following:

a) vigorous-intensity activity 3 days, achieving at least 1500 minutes/week.

OR

b)7 days of any combination of walking, moderate-intensity or vigorous-intensity activities of at least 3000 minutes/week.

Participants who did not meet the above criteria were considered to have a 'low' physical activity level ³⁶.

The fourth section assesses the exposure to polluted air; participants are asked about the environment where they live, the heating system used (wood, gas, electricity), if they are living or working next to factories or power stations, and their exposure to cigarette smoking.

The fifth section represents the Arabic translation of the Short Version of the Problematic Mobile Phone Use Questionnaire (PMPUQ-SV), containing 15 items that are scored from 1 ('I strongly agree') to 4 ('I strongly disagree'), except for 8 items that are reversely scored. Overall scores range from 15 to 60, with higher scores indicating a higher risk for problematic mobile phone use ³⁷.

The Hayling Sentence Completion Test

The HSCT was translated to Arabic and adapted to the linguistic and cultural realities of the Lebanese population. The A-HSCT consists of two conditions (automatic and inhibition), for which two different groups of 15 sentences are given. In both conditions, the interviewer reads out loud the unfinished sentences, and the participant has to complete each sentence with one word ⁴.

Automatic condition

The participants are requested to give a word that is related to the beginning of the sentence, and should do so as quickly as possible. For example, "He mailed the letter without a . . . (participant says) stamp." Time latency in automatic condition measures the participant's rapidity in initiating an automatic response. According to the scoring system, three error points are scored when participants provide an incorrect word, one point when the answer is semantically related to the sentence, and no error point if the correct word is given. Higher error score corresponds to a lower performance ⁴.

Inhibition condition

The participants are requested to complete the sentence as fast as they can with a word that is completely unlinked to it, which makes no sense at all in the context of the sentence. For example, "The captain wanted to stay with the sinking . . . (participant says) apple." If at any time during this condition the subject completes the sentence correctly instead of using an unrelated word, s/he is told that the word is too linked to the sentence and is retold the task instructions. Time latency in inhibition condition gives information about the time needed to inhibit the correct response and find an incorrect one ⁴. According to the scoring system, three points are given when the sentence is completed with the answer that fits with it. One point is given when a subject gives an antonym, a semantically related word, or a

word that makes a vague reference to the sentence. Participants receive zero points when a totally unrelated word is provided. A higher error score indicates a lower performance ⁴.

The total time to complete both conditions of the A-HSCT was approximately five minutes. The automatic condition was tested prior to the inhibition section. Two practice sentences were initially presented prior to each condition. Time of response latency was measured and collected using a stopwatch in both conditions; the timing began soon after the tester finished the sentence, and was stopped soon after the participant began their answer. Response latencies were recorded in whole second units and were not rounded up. For instance, a time between 0 and 0.99 was scored as 0. An average response latency score of all the individual's latencies for each condition was then computed based on all responses, including errors. No time limit was given for responding. We also scored errors to evaluate the efficacy of the strategy elaborated by the participant to give an incorrect response ⁴. A simple guideline for the error scoring of the inhibition condition was created, containing a collection of several hundreds of errors for each sentence ⁵. All sections of the questionnaire were translated to Arabic by a translation specialist, then the translation was retranslated to its original language by another specialist. Upon fulfillment of this procedure, the translators compared the versions of every scale to determine whether the variables had the same meaning. No major incompatibilities were found between the two versions for all scales; they were resolved by consensus.

Statistical analysis

Statistical Package for Social Science (SPSS) version 23 was used for the statistical analyses. A principal component analysis, using a promax rotation since the questions of the scales were correlated, was conducted in order to validate the automatic and inhibition parts of the HSCT. Adequacy of the sample was confirmed through the Kaiser–Meyer–Olkin (KMO) index, Bartlett's Chi-square test of sphericity and scree plot. Factors with an Eigenvalue higher than one were retained. The Student t-test was used to compare two means, whereas the Pearson correlation was used to study the association between two continuous variables. Multivariable linear regression models were done to explore factors associated with the response-time and the number of errors taken as dependent variables and taking all variables that showed a $p < 0.05$ in the bivariate analysis as independent variables. A $p < 0.05$ was considered significant. Reliability was assessed using Cronbach's alpha.

Results

Out of 364 participants approached, 350 (96.15%) accepted to enroll in this study. The mean age of the participants was 45.72 years, with 177 (50.6%) of them being females. Other sociodemographic and characteristics of the participants are summarized in Table 1. In the automatic condition of the HSCT, 346 participants (98.9%) made ≤ 2 errors and 333 individuals (96%) responded in seconds. Accordingly, no additional calculations were conducted on these scores. The results of the inhibition condition of the

HSCT showed that 221 (63.5%) of the participants had normal response-time, whereas 205 (58.7%) had normal scores in terms of number of errors.

Table 1. Sociodemographic characteristics of the participants.	
Variable	Mean $\pm$ SD
Age (in years)	45.72 $\pm$ 21.15
Body Mass Index (kg/m ²)	24.64 $\pm$ 3.82
Years of study	16.49 $\pm$ 5.79
House crowding index	0.99 $\pm$ 0.45
Problematic mobile phone use	31.04 $\pm$ 10.84
Total food score	34.65 $\pm$ 4.53
	N (%)
Governorate	
Beirut	42 (12.0%)
Mount Lebanon	66 (18.9%)
North	106 (30.3%)
South	67 (19.1%)
Bekaa	69 (19.7%)
Classification executive function- time	
Pathologic	17 (4.9%)
Limit	45 (12.9%)
Normal	221 (63.5%)
Superior	55 (15.8%)
Very superior	10 (2.9%)
Classification executive function- number of errors	
Pathologic	16 (4.6%)
Limit	49 (14.0%)
Normal	205 (58.7%)
Superior	72 (20.6%)
Very superior	7 (2.0%)
Categories of the diet levels	

Table 1. Sociodemographic characteristics of the participants.	
Not healthy	2 (0.6%)
Moderately healthy	16 (4.6%)
Healthy	168 (48.0%)
Very healthy	164 (6.9%)

Validation of the HSCT scale

Exploratory factor analysis

The PCA for the automatic and inhibition response-time and number of errors questions was run over the whole sample ($n = 350$). None of the automatic condition response-time items was removed and converged over one factor (variance explained = 47.92%; KMO = 0.936; Bartlett sphericity $p < 0.001$; $\alpha_{\text{Cronbach}} = 0.905$). Two items (items 3 and 12) of the automatic condition number of errors items were removed since all participants had the same answer. The items converged over seven factors (variance explained = 71.41%; KMO = 0.464; Bartlett sphericity $p < 0.001$; $\alpha_{\text{Cronbach}} = 0.334$). None of the inhibition condition response-time items was removed and converged over one factor (variance explained = 56.59%; KMO = 0.962; Bartlett sphericity $p < 0.001$; $\alpha_{\text{Cronbach}} = 0.943$). One item (item 12) of the inhibitory condition number of errors items was removed since all participants had the same answer. The items converged over four factors (variance explained = 49.57%; KMO = 0.797; Bartlett sphericity $p < 0.001$; $\alpha_{\text{Cronbach}} = 0.728$) (Table 2).

Table 2. Principal component analysis results using the promax rotation.

Scale 1: Automatic condition response time.

Item number	Factor 1	Factor 2	Factor 3	Factor 4	Factor 5	Factor 6	Factor 7
12	0.818						
7	0.810						
1	0.732						
15	0.731						
11	0.728						
5	0.716						
10	0.716						
4	0.686						
8	0.682						
9	0.676						
14	0.635						
3	0.621						
13	0.614						
2	0.598						
6	0.569						

Scale 2: Automatic condition number of errors.

Item number	Factor 1	Factor 2	Factor 3	Factor 4	Factor 5	Factor 6	Factor 7
5	0.685						
7	0.634						
6	0.500						
4		0.572					
15		0.422					
14			0.579				
9			0.502				
13				0.728			

Table 2. Principal component analysis results using the promax rotation.							
2					0.709		
8					0.709		
1						0.729	
10						0.685	
11							0.816
Scale 3: Inhibitory condition response time.							
Item number	Factor 1	Factor 2	Factor 3	Factor 4	Factor 5	Factor 6	Factor 7
12	0.823						
6	0.811						
3	0.803						
4	0.779						
13	0.770						
11	0.769						
15	0.762						
5	0.758						
2	0.757						
7	0.750						
14	0.750						
10	0.738						
9	0.731						
8	0.666						
1	0.586						
Scale 4: Inhibitory condition number of errors.							
Item number	Factor 1	Factor 2	Factor 3	Factor 4	Factor 5	Factor 6	Factor 7
11	0.744						
9	0.713						

Table 2. Principal component analysis results using the promax rotation.		
10	0.669	
8	0.459	
6	0.791	
1	0.660	
5	0.442	
3	0.746	
2	0.718	
7	0.449	
4	0.425	
14	0.705	
13	0.665	
15	0.528	

Bivariate analysis

The results of the bivariate analysis of factors associated with the response time showed that diabetic patients (-0.42 vs 0.15), those with hypertension (-0.42 vs 0.20), those with cardiac diseases (-0.53 vs 0.08), those taking aspirin (-0.58 vs 0.11), and those who use electricity as a heating system in the house (-0.25 vs 0.21) had significantly lower response-time (better performance) compared to the opposite groups. In terms of number of errors, the results showed that those who do not use wood as a heating system in the house (-0.18 vs 0.29) and those who use bakhour (Arabian incense) (-0.25 vs 0.16) had significantly lower number of errors (better performance) compared to the opposite groups (Table 3).

Table 3. Bivariate analysis of factors associated with the response time and number of errors.		
Variable	Response time	Number of errors
Diabetes Mellitus		
No	0.15 ± 0.98	0.23 ± 1.02
Yes	-0.42 ± 0.93	-0.06 ± 0.95
P	< 0.001	0.301
Hypertension		
No	0.20 ± 0.97	0.07 ± 1.05
Yes	-0.42 ± 0.94	-0.13 ± 0.88
P	< 0.001	0.039
Cardiac diseases		
No	0.08 ± 0.98	0.02 ± 1.03
Yes	-0.53 ± 1.01	-0.15 ± 0.74
P	< 0.001	0.132
Proton pump inhibitors		
No	0.06 ± 1.01	0.04 ± 1.04
Yes	-0.27 ± 0.93	-0.20 ± 0.75
P	0.024	0.039
Aspirin		
No	0.11 ± 0.97	0.05 ± 1.03
Yes	-0.58 ± 0.94	-0.26 ± 0.79
P	< 0.001	0.015
Heating inside the house- wood		
No	-0.004 ± 1.01	-0.18 ± 1.04
Yes	0.007 ± 0.98	0.29 ± 0.86
P	0.796	< 0.001
Heating inside the house- Electricity		
No	0.21 ± 0.94	0.08 ± 0.94
Yes	-0.25 ± 1.01	-0.09 ± 1.06

Table 3. Bivariate analysis of factors associated with the response time and number of errors.		
P	< 0.001	0.111
Bakhour		
No	-0.04 ± 0.98	0.16 ± 0.95
Yes	0.06 ± 1.03	-0.25 ± 1.03
P	0.484	< 0.001
Living in a polluted area		
No	0.07 ± 0.88	-0.36 ± 1.09
Yes	-0.003 ± 1.03	0.10 ± 0.95
P	0.789	0.003
Exposed to sand and dust soil		
No	-0.75 ± 0.62	-0.15 ± 1.07
Yes	-0.16 ± 1.15	-0.05 ± 0.91
P	0.028	0.474
Factories		
No	0.003 ± 0.99	0.04 ± 0.99
Yes	-0.05 ± 1.17	-0.66 ± 0.78
P	0.692	0.002

Higher age was significantly associated with lower response-time (better performance), whereas higher problematic mobile phone use was significantly associated with higher response-time and higher number of errors (worse performance) (Table 4).

Table 4. Bivariate analysis of continuous variables associated with the response time and number of errors.		
Variable	Response time	Number of errors
Age	-0.345 ^a	-0.053
Body Mass Index	-0.153 ^b	-0.06
Number of years of study	0.145 ^b	-0.079
Total food score	-0.052	0.139 ^b
Number of smokers in the house	-0.135 ^b	-0.077
Problematic mobile phone use score	0.401 ^a	0.215 ^a
House crowding index	-0.115 ^c	0.012
^a p<0.001; ^b p<0.01; ^c p<0.05		

Multivariable analysis

The results of a first linear regression, taking the response-time as the dependent variable, showed that using electricity as a heating method inside the house (B=-0.3) was significantly associated with lower response-time (better performance), whereas a higher problematic mobile phone use (B = 0.03) was significantly associated with higher response-time (worse performance) (Table 5, Model 1).

The results of a second linear regression, taking the number of errors as the dependent variable, showed that using wood as a heating system inside the house (B = 0.41) and higher problematic mobile phone use (B = 0.02) were significantly associated with higher number of errors (worse performance), whereas using Arabian incense (bakhour) inside the house (B=-0.29) was significantly associated with a lower number of errors (better performance) (Table 5, Model 2).

Table 5. Multivariable analysis.					
Model 1: Linear regression taking the response time as the dependent variable.					
Variable	Unstandardized Beta	Standardized Beta	p	95% Confidence Interval	
Heating in the house- Electricity (yes vs no*)	-0.30	-0.15	0.003	-0.50	-0.10
Problematic mobile phone use	0.03	0.35	< 0.001	0.02	0.04
Model 2: Linear regression taking the number of errors as the dependent variable.					
Variable	Unstandardized Beta	Standardized Beta	p	95% Confidence Interval	
Heating in the house- Wood (yes vs no*)	0.41	0.20	< 0.001	0.20	0.63
Bakhour use in the house (yes vs no*)	-0.29	-0.14	0.007	-0.50	-0.08
Problematic mobile phone use	0.02	0.21	< 0.001	0.01	0.03

Discussion

To the best of the authors' knowledge, this is the first study to validate the Hayling test in Arabic, and to assess factors associated with executive functioning among a sample of the Lebanese population. These results constitute the exclusive source of norms for this test in the Lebanese population.

Validation of the A-HSCT

In this current study, we were able to validate the Arabic version of the Hayling Sentence Completion Test (A-HSCT) designed precisely for the Lebanese population. Results delivered primary evidence supporting the accuracy and validity of this test as a clinical instrument to measure prepotent response inhibition in Lebanese adults having a wide range of neurological or psychological disorders. In this study, we translated all the sentences to Arabic to be adapted to the Lebanese culture and habits. Consequently, the Lebanese protocol revealed that the new set of sentences of the HSCT preserved strong accuracy, based on Cronbach's alpha.

As compared to the Spanish version where the Cronbach's alpha was 0.864 for the response-time in the automatic condition, we got a slightly higher value of 0.905. In addition, Cronbach's alpha for the

response time in the inhibition condition in the Spanish version was 0.797, while in our version, its considerably higher with a value of 0.943.

The Cronbach's alpha for the number of errors in the inhibition condition of the Spanish version was 0.839. Our Arabic version got a slightly smaller value of 0.728. In this Arabic version, Cronbach's alpha for the number of errors in the automatic condition is 0.334 (where items were converged over eight factors). However, the one for the Spanish version in this same condition was not mentioned ⁶.

Factors associated with executive functions

Problematic mobile phone use

Many studies investigating the effect of Electromagnetic Field (EMF) exposure in mobile phone users report a slower electroencephalographic (EEG) activity in these individuals, along with a hypoactivation of a major participant in the regulation of executive functions, the Anterior Cingulate Cortex (ACC). Thus, worse performance on tasks requiring executive functions is associated with long-term mobile phone use ^{18,38}. In addition, a research studying the effect of EMF on the neurodevelopment of neonates and children highlights the importance of epigenetic mechanisms that can lead to altered attention, memory and cognition ³⁹. This goes hand in hand with the results of our study, which showed an increase in both response-time and number of errors in the Arabic version of the HSCT for individuals with problematic mobile phone use, indicating a poorer performance. On the other hand, some studies contradict these results, suggesting an enhancement in executive functions among mobile phone users. However, most of these studies reported limitations to their results ^{17,40}, emphasizing on the need to conduct further longitudinal studies to clarify the true effect of mobile phone use on executive functions.

Heating system and air pollution

The results of our study concerning the effects of outdoor air pollution on executive functions were inconclusive. However, indoor air pollution attributed to the use of wood burning as a heating system is linked to a lower performance of the interviewed participants, particularly in the number of errors committed in the inhibition condition of the A-HSCT. This negative correlation could be due to the exposure of individuals to air pollutants resulting from wood burning chemical reactions, as mentioned in various studies, particularly CO₂ inhalation that is anxiogenic and deleterious for executive functions ^{41,42}. In contrast, participants who use electricity as a source of heating in their households performed better in terms of response-time in the inhibition condition of our test, which goes in line with the result mentioned above.

On another note, Bakhour is a widespread spiritual practice in the Middle East region, composed of a vast range of chemical compounds and metals. Opposed to the upstated results concerning indoor air

pollution, the use of Arabian incense was associated to a decreased number of errors committed during the A-HSCT. We hypothesize that this could be attributed to the spirituality inferred by the use of Bakhour among the participants, which is the subject of many studies that elaborate the benefits of spirituality on executive functions, and its capacity in attenuating the severity of depression ⁴³.

Other factors

In the bivariate analysis, participants who are older, use aspirin, have cardiac diseases, diabetes, or hypertension exhibited better performance in terms of response-time in the inhibition condition of the A-HSCT. These results contradict the literature that underlines the negative correlation between executive functions and aging.⁹ However, they could be explained by the fact that many of the elderly participants reported taking metformin, aspirin, atorvastatin, and angiotensin converting enzyme inhibitors (ACEi) or ARBs, which, when taken as a combination, could potentiate the anti-oxidant effect of each drug, enhancing their neuroprotective effects on the hippocampus ⁴⁴.

Clinical implications

The A-HSCT is a clinical tool that measures inhibitory control as part of executive functions, reflecting mainly frontal cortex function. The obtained score could be used in patients with many neurological or psychological disorders to follow them up throughout the years, and to detect amelioration or deterioration of their scores. Moreover, by comparing the patients' scores, clinicians can detect the efficacy of their treatment and medications.

Limitations

Our study is based on the questionnaire that was applied on the participants during an interview. Thus, some of these participants might have felt pressured or shy while being interviewed, leading them to give wrong answers or to perform poorly in the given tasks. An information bias is also possible because of potential misunderstanding. The possibility of recall bias might be considered due to the need of the participants to remember activities of their daily lives. The effect of the recall bias could be differential and may lead to the overestimation or underestimation of effects for some factors, hence the need for prospective studies that overthrow the recall bias and can show more significant and precise association between problematic mobile phone use and executive functions. The extent of exposure to different air pollutants was subjectively evaluated by each participant. Unfortunately, there were no possible means to measure the quantity and time of exposure to each air pollutant. In future studies a bigger sample is needed to reinforce the correlation between problematic mobile phone use and the deterioration of executive functions.

Conclusion

The Arabic version of the Hayling Sentence Completion Test is now a valid tool that can be used by clinicians to measure response inhibition in adults and elderly patients in the Lebanese population. We were also able to suggest an association between many factors and executive functions. The most significant was problematic mobile phone use and the deterioration of executive functions, mainly inhibitory control, in terms of both response-time and number of errors in the A-HSCT. As far as we know, the present study is the first to create normative data, let alone validate the A-HSCT in the Lebanese population.

Furthermore, it should be noted that additional studies are needed to prove a possible harmful effect of mobile phone use on frontal cortex, since it became a primarily used feature in our daily life.

List Of Abbreviations

HSCT=Hayling Sentence Completion Test

EF=executive functions

SES= socioeconomic status

EMF=electromagnetic fields

MP=mobile phones

PPI=proton pump inhibitors

AD=Alzheimer's Disease

ARB=Angiotensin I receptor blockers

DASH=Dietary Approaches to Stop Hypertension

IPAQ= International Physical Activity Questionnaire

PMPUQ-SV=Problematic Mobile Phone Use Questionnaire

SPSS=Statistical Package for Social Science

KMO=Kaiser–Meyer–Olkin

EEG=electroencephalographic

ACC=Anterior Cingulate Cortex

ACEi=angiotensin converting enzyme inhibitors

Declarations

Ethics Approval and Consent to Participate

The study protocol was approved by the Holy Spirit University of Kaslik (USEK) ethics committee. A written informed consent was obtained from each participant.

Consent for publication

Not applicable.

Availability of data and materials

All data generated or analyzed during this study are not publicly available to maintain the privacy of the individuals' identities. The dataset supporting the conclusions is available upon request to the corresponding author.

Competing interests

The authors have nothing to disclose.

Funding

None.

Author contributions

SB was responsible for the data collection and designed the study; SB drafted the manuscript; SH carried out the analysis and interpreted the results, assisted in drafting and reviewing the manuscript; EAH and JM helped in the drafting the paper. SH and HM were the project supervisors. All authors reviewed the final manuscript and gave their consent.

Acknowledgments

We express our gratitude to the participants who donated their time in completing the questionnaires. We would also like to thank the research assistants and the investigators who contributed in the data collection, and the staff experts who helped building this study.

References

1. Barkley RA. *Executive functions: What they are, how they work, and why they evolved*. Guilford Press; 2012.
2. Sira C, Mateer C. Executive Function. 2014.

3. Bayard S, Gely-Nargeot MC, Raffard S, et al. French Version of the Hayling Sentence Completion Test, Part I: Normative Data and Guidelines for Error Scoring. *Arch Clin Neuropsychol*. 2017;32(5):585-591.
4. Burgess PW, Shallice T. Response suppression, initiation and strategy use following frontal lobe lesions. *Neuropsychologia*. 1996;34(4):263-272.
5. Bielak AA, Mansueti L, Strauss E, Dixon RA. Performance on the Hayling and Brixton tests in older adults: norms and correlates. *Arch Clin Neuropsychol*. 2006;21(2):141-149.
6. Pérez-Pérez A, Matias-Guiu JA, Cáceres-Guillén I, et al. The hayling test: development and normalization of the Spanish version. *Archives of Clinical Neuropsychology*. 2016;31(5):411-419.
7. Spitoni GF, Bevacqua S, Cerini C, et al. Normative Data for the Hayling and Brixton Tests in an Italian Population. *Arch Clin Neuropsychol*. 2018;33(4):466-476.
8. Vestberg S, Nordstrom EB, Waldo ML, Nilsson K, Santillo AF, Nilsson C. Swedish Version of the Hayling Test: Clinical Utility in Frontotemporal Dementia Syndromes. *J Int Neuropsychol Soc*. 2019;25(2):195-203.
9. Cervera-Crespo T, Gonzalez-Alvarez J. Age and Semantic Inhibition Measured by the Hayling Task: A Meta-Analysis. *Arch Clin Neuropsychol*. 2017;32(2):198-214.
10. Hackman DA, Gallop R, Evans GW, Farah MJ. Socioeconomic status and executive function: developmental trajectories and mediation. *Dev Sci*. 2015;18(5):686-702.
11. Quinteros Baumgart C, Billick SB. Positive Cognitive Effects of Bilingualism and Multilingualism on Cerebral Function: a Review. *Psychiatr Q*. 2018;89(2):273-283.
12. Cohen JF, Gorski MT, Gruber SA, Kurdziel LB, Rimm EB. The effect of healthy dietary consumption on executive cognitive functioning in children and adolescents: a systematic review. *Br J Nutr*. 2016;116(6):989-1000.
13. Yang Y, Shields GS, Guo C, Liu Y. Executive function performance in obesity and overweight individuals: A meta-analysis and review. *Neurosci Biobehav Rev*. 2018;84:225-244.
14. Berryman N, Bherer L, Nadeau S, et al. Executive functions, physical fitness and mobility in well-functioning older adults. *Exp Gerontol*. 2013;48(12):1402-1409.
15. Tzivian L, Dlugaj M, Winkler A, et al. Long-term air pollution and traffic noise exposures and cognitive function: A cross-sectional analysis of the Heinz Nixdorf Recall study. *J Toxicol Environ Health A*. 2016;79(22-23):1057-1069.
16. Ng TP, Lim ML, Niti M, Collinson S. Long-term digital mobile phone use and cognitive decline in the elderly. *Bioelectromagnetics*. 2012;33(2):176-185.
17. Besset A, Espa F, Dauvilliers Y, Billiard M, de Seze R. No effect on cognitive function from daily mobile phone use. *Bioelectromagnetics*. 2005;26(2):102-108.
18. Hayashi Y, Rivera EA, Modico JG, Foreman AM, Wirth O. Texting while driving, executive function, and impulsivity in college students. *Accid Anal Prev*. 2017;102:72-80.
19. Akter S, Hassan MR, Shahriar M, Akter N, Abbas MG, Bhuiyan MA. Cognitive impact after short-term exposure to different proton pump inhibitors: assessment using CANTAB software. *Alzheimers Res*

Ther. 2015;7:79.

20. Ho JK, Nation DA. Memory is preserved in older adults taking AT1 receptor blockers. *Alzheimer's research & therapy.* 2017;9(1):33.
21. Group VTC. Effect of vitamins and aspirin on markers of platelet activation, oxidative stress and homocysteine in people at high risk of dementia. *Journal of internal medicine.* 2003;254(1):67-75.
22. Martyr A, Boycheva E, Kudlicka A. Assessing inhibitory control in early-stage Alzheimer's and Parkinson's disease using the Hayling Sentence Completion Test. *Journal of neuropsychology.* 2019;13(1):67-81.
23. Cervera Crespo T, González Álvarez J, Rosell Clari V. Semantic inhibition and dementia severity in Alzheimer's disease. 2019.
24. Ramanan S, Bertoux M, Flanagan E, et al. Longitudinal Executive Function and Episodic Memory Profiles in Behavioral-Variant Frontotemporal Dementia and Alzheimer's Disease. *J Int Neuropsychol Soc.* 2017;23(1):34-43.
25. Rabinowitz AR, Levin HS. Cognitive sequelae of traumatic brain injury. *Psychiatr Clin North Am.* 2014;37(1):1-11.
26. Povroznik JM, Ozga JE, Haar CV, Engler-Chiurazzi EB. Executive (dys) function after stroke: special considerations for behavioral pharmacology. *Behavioural pharmacology.* 2018;29(7):638-653.
27. Obeso I, Wilkinson L, Casabona E, et al. Deficits in inhibitory control and conflict resolution on cognitive and motor tasks in Parkinson's disease. *Exp Brain Res.* 2011;212(3):371-384.
28. Matías-Guiu JA, Cabrera-Martín MN, Valles-Salgado M, et al. Inhibition impairment in frontotemporal dementia, amyotrophic lateral sclerosis, and Alzheimer's disease: clinical assessment and metabolic correlates. *Brain imaging and behavior.* 2019;13(3):651-659.
29. Giraldo-Chica M, Rogers BP, Damon SM, Landman BA, Woodward ND. Prefrontal-Thalamic Anatomical Connectivity and Executive Cognitive Function in Schizophrenia. *Biol Psychiatry.* 2018;83(6):509-517.
30. Lima IMM, Peckham AD, Johnson SL. Cognitive deficits in bipolar disorders: Implications for emotion. *Clin Psychol Rev.* 2018;59:126-136.
31. Demetriou EA, Lampit A, Quintana DS, et al. Autism spectrum disorders: a meta-analysis of executive function. *Mol Psychiatry.* 2018;23(5):1198-1204.
32. Ronnlund M, Nyberg L, Backman L, Nilsson LG. Stability, growth, and decline in adult life span development of declarative memory: cross-sectional and longitudinal data from a population-based study. *Psychol Aging.* 2005;20(1):3-18.
33. Bernier PJ, Gourdeau C, Carmichael PH, et al. Validation and diagnostic accuracy of predictive curves for age-associated longitudinal cognitive decline in older adults. *CMAJ.* 2017;189(48):E1472-E1480.
34. Comrey AL, Lee HB. *A first course in factor analysis.* Psychology press; 2013.
35. Lung NH, Institute B. Your guide to lowering your blood pressure with DASH. *Bethesda (MD).* 2006.

36. Hagströmer M, Oja P, Sjöström M. The International Physical Activity Questionnaire (IPAQ): a study of concurrent and construct validity. *Public health nutrition*. 2006;9(6):755-762.
37. Lopez-Fernandez O, Kuss DJ, Pontes HM, et al. Measurement Invariance of the Short Version of the Problematic Mobile Phone Use Questionnaire (PMPUQ-SV) across Eight Languages. *Int J Environ Res Public Health*. 2018;15(6).
38. Ng TP, Lim ML, Niti M, Collinson S. Long-term digital mobile phone use and cognitive decline in the elderly. *Bioelectromagnetics*. 2012;33(2):176-185.
39. Sage C, Burgio E. Electromagnetic Fields, Pulsed Radiofrequency Radiation, and Epigenetics: How Wireless Technologies May Affect Childhood Development. *Child Dev*. 2018;89(1):129-136.
40. Arns M, Van Luijckelaar G, Sumich A, Hamilton R, Gordon E. Electroencephalographic, personality, and executive function measures associated with frequent mobile phone use. *Int J Neurosci*. 2007;117(9):1341-1360.
41. Savulich G, Hezemans FH, van Ghesel Grothe S, et al. Acute anxiety and autonomic arousal induced by CO2 inhalation impairs prefrontal executive functions in healthy humans. *Transl Psychiatry*. 2019;9(1):296.
42. Siponen T, Yli-Tuomi T, Tiittanen P, et al. Wood stove use and other determinants of personal and indoor exposures to particulate air pollution and ozone among elderly persons in a Northern Suburb. *Indoor air*. 2019;29(3):413-422.
43. Herren OM, Burris SE, Levy SA, et al. Influence of Spirituality on Depression-Induced Inflammation and Executive Functioning in a Community Sample of African Americans. *Ethn Dis*. 2019;29(2):267-276.
44. Paseban M, Mohebbati R, Niazmand S, Sathyapalan T, Sahebkar A. Comparison of the Neuroprotective Effects of Aspirin, Atorvastatin, Captopril and Metformin in Diabetes Mellitus. *Biomolecules*. 2019;9(4).