Ancient CO₂ Levels Favor Nitrogen Fixing Plants Over a Broader Range of Soil N Compared to Present

Haoran Chen (✉ Chenh318@myumanitoba.ca)

Department of Biological Sciences, University of Manitoba, Winnipeg, Canada

John Markham

University of Manitoba

Research Article

Keywords: ancient CO₂, N deposition, N fixation, Alnus, Betula

Posted Date: December 8th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-120004/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on February 4th, 2021.
See the published version at <https://doi.org/10.1038/s41598-021-82701-7>.

Abstract

Small increases in CO₂ stimulate nitrogen fixation and plant growth. Increasing soil N can inhibit nitrogen fixation. However, no studies to date have tested how nitrogen fixing plants perform under ancient CO₂ levels (100 MYA) when nitrogen fixers evolved, with different levels of N additions. The aim of this study was to assess if ancient CO₂, compared to present, favors nitrogen fixers over a range of soil nitrogen concentrations. Nitrogen fixers (*Alnus incana* ssp. *rugosa*, *Alnus viridis* ssp. *crispa*, and *Alnus rubra*) and their close non-nitrogen fixing relatives (*Betula pumila*, *Betula papyrifera*, *Betula glandulosa*) were grown at ancient CO₂ (1600 ppm) or present CO₂ (400 ppm) over a range of soil N levels, equivalent to 0, 10, 50, and 200 kg N ha⁻¹ year⁻¹. The growth of non-N fixing plants increased more than N fixing plants in response to the increasing N levels. When grown at an ancient CO₂ level, the N level at which non-nitrogen fixing plant biomass exceeded nitrogen fixing plant biomass was twice as high (61 kg N ha⁻¹ y⁻¹) as the N level when plants were grown at an ambient CO₂ level. Specific nodule activity was also reduced with an increasing level of soil N. Our results showed there was a greater advantage in being a nitrogen fixer under ancient levels of CO₂ compared with the present CO₂ level.

Introduction

Fossil evidence shows that nitrogen fixing plants first evolved in the late Cretaceous, when atmospheric CO₂ was *ca.* four times higher than at present^{1,2}. Although there is some question as to whether the evolution of symbiotic nitrogen fixation arose once, or multiple times, in a small number of related clades³, it is clear that symbiotic nitrogen fixation is a trait that has been lost repeatedly. This has resulted in most taxa within the nitrogen fixing clade lacking the ability to fix nitrogen⁴. It has been suggested that the decrease in CO₂ levels over geological time provided the global selection pressure to allow for this loss of the nitrogen fixing trait across the entire nitrogen fixing clade³.

There is also, at present, a low abundance of nitrogen fixing plants in many parts of the globe⁵. Symbiotic nitrogen fixing plant species abundance, and to a lesser degree species richness, decreases with latitude, with legume species having a higher abundance at lower latitudes (< 35°), and actinorhizal species having a higher abundance at higher latitudes^{6,7}. The actinorhizal plants are mainly pioneer species, colonizing N poor soils, where they have access to high light intensity in the early stages of succession⁸. One reason for the present lack of success of nitrogen fixing plants may relate to the high energy cost of nitrogen fixation, especially at higher latitudes⁹. In all but N poor soils, nitrogen fixing plant plants spend more energy in acquiring nitrogen from fixation than non-N fixers taking up soil inorganic N. In addition, N fixers indirectly provide fixed nitrogen to their non-N fixing competitors, which may then shade them⁸. Menge and Crews⁵ have also argued that actinorhizal plants are obligate N fixers (unlike legumes), and thus cannot down-regulate symbiotic nitrogen fixation. There are however a lack of studies comparing the performance of N-fixers to non-N fixers, especially to closely related non-N fixers with the same growth form, under environmental conditions that are predicted to alter their performance.

The loss of the nitrogen fixing trait with decreasing CO₂ levels over geological time may relate to the energetic cost of fixation, with decreased CO₂ limiting energy capture, and to plants ability to balance carbon and nitrogen acquisition. The recent industrial era increases in atmospheric CO₂ show that increasing CO₂ increases plant N demand in two ways. First, increasing CO₂ concentration increases C sequestration resulting in a simultaneous increase in N uptake from the soil¹⁰. Secondly, increased CO₂ levels increase the C:N ratio of leaves, and consequently the C:N ratio of the soil, which decreases soil N availability^{10,11}. Elevated CO₂ therefore increases both symbiotic and free living nitrogen fixation compared with ambient CO₂ levels^{11,12}. Although many studies have focused on the effect of increasing atmospheric CO₂ on biological nitrogen fixation, they have been restricted to less than 800 ppm CO₂ levels in order to understand future scenarios^{11,12}. There is no study comparing nitrogen fixing plant growth at ambient and ancient CO₂ levels.

Another factor that may negatively affect nitrogen fixing plants, especially obligate N fixers, is increasing N deposition. Low levels of available N can critically limit the establishment and growth of non-N fixers. In such N-poor areas, actinorhizal plants are favored¹³. At present human activities have resulted in increased N deposition at a global scale. N deposition increases have arisen from the unintentional production of NO_x in the atmosphere and the intentional production of reduced N (NH_x) for synthetic fertilizer¹⁴. The global rate of anthropogenic N inputs has increased from *ca.* 100 Tg N year⁻¹ to *ca.* 200 Tg N year⁻¹ during the past 20 years¹⁵⁻¹⁷. Currently, 11% of natural vegetation in North America, Europe, south and east Asia receives anthropogenic atmospheric nitrogen deposition of over 10 kg N ha⁻¹ yr⁻¹, a level that can have significant effects on ecosystem processes¹⁴. The annual total N deposition in some areas is projected to further increase in the future, ranging from 20 to 50 kg N ha⁻¹ yr⁻¹ by 2100^{14,18}. Since the net production of many terrestrial ecosystems is limited by N availability, N deposition should enhance ecosystem productivity. On the other hand, N deposition can threaten soil quality through soil acidification, altering the ground flora, leading to a decline of productivity and biodiversity loss in some ecosystems^{19,20}. Nitrogen deposition has been repeatedly reported to reduce the biodiversity in some ecosystems^{21,22}. Global rates of biological nitrogen fixation have also fallen from 100–290 Tg year⁻¹ to 44 Tg year⁻¹ during the past 20 years^{15,23}, which is largely due to anthropogenic N inputs^{16,23}. It has also been reported that plants confined to growing in low N soils and those depending on N fixation are more negatively affected by increased N deposition²¹. It was found that N fixing plants are more susceptible to declines in performance than non-nitrogen fixing plants, with increased N deposition due to a reduced competitive ability²⁴. Some nitrogen fixers are able to adjust soil N additions by reducing nitrogen fixation. For example, a two-year field study found that 180 kg N ha⁻¹ application reduced biological nitrogen fixation of soybean (*Glycine max* (L.) Merr) compared with non-fertilizer treatments²⁵. While it has been argued that actinorhizal plants cannot down-regulate symbiotic nitrogen fixation⁵, lab studies have shown that actinorhizal plants can shift from using fixed-N to using N-derived fertilizer with an increasing level of N fertilization²⁶. This may help them to cope with increased N deposition.

We know little about the effects of ancient levels of CO₂ and soil nitrogen levels on the relative performance of nitrogen fixing and non-nitrogen fixing plants. We know elevated CO₂ leads to the increase of plant C:N ratio across all plants^{27–29}. Therefore, increases in N deposition could potentially buffer the effect increases in atmosphere CO₂ on plant tissue C:N ratio³⁰.

The clade that includes actinorhizal plants also contains non-nitrogen fixing species. This makes them ideal for comparing species with and without the ability to form a nitrogen fixing symbiosis. The purpose of this study was to look at the effect of soil nitrogen levels on nitrogen fixing plants under present and ancient levels atmospheric CO₂ levels by comparing them to their close non-nitrogen fixing relatives. We hypothesized nitrogen fixing plants will maintain greater growth with increasing N levels under ancient CO₂ conditions compared with present, helping to explain the evolution of symbiotic nitrogen fixation in the Late Cretaceous. We also hypothesized that actinorhizal N-fixing plants are partially facultative N-fixers and so can downregulate N fixation when N deposition is high at all level of CO₂.

Results

Plant mass

The effect of nitrogen on plant mass varied with CO₂ level ($P = 0.006$, $F = 8.03$ for the nitrogen by CO₂ level interaction in the least squares model), and differed between the nitrogen fixing and non-nitrogen fixing plants ($P < 0.0001$, $F = 28.22$ for the plant group by nitrogen level interaction) but the model showed no effect of species within plant groups ($P = 0.20$, $F = 1.55$, **Supplementary Figure S1**). We therefore performed a least square fits of the responses of nitrogen fixing and non-nitrogen fixing plants to soil nitrogen additions at each CO₂ level separately (Fig. 1). At both ambient and ancient CO₂ levels, non-nitrogen fixing plant mass increased faster than nitrogen fixing plants mass in response to N addition. However, because there was a greater difference in the rate of plant mass increase between the plant types at 400 ppm than at 1600 ppm CO₂, the N level at which the two plant groups performed equally well (i.e., the mass of plants was equal) differed between the two CO₂ levels. At 400 ppm CO₂, plant biomass of the nitrogen fixing and non-nitrogen fixing plants was equal at a soil N deposition rate of 27 kg N ha⁻¹ year⁻¹, while at 1600 ppm CO₂, the N level at which plant biomass was equal was at 61 kg N ha⁻¹ year⁻¹, more than double the N when plants were grown at 400 ppm CO₂.

Nitrogen Fixiation

Increasing soil N reduced all parameters associated with nitrogen fixation, regardless of CO₂ level, with significant differences between plant species for some parameters (Fig. 2). Biomass allocation to nodules, including plants that failed to form nodules, decreased from $0.98 \pm 0.21\%$ ($\pm$ standard error) of total biomass at no N addition to $0.17 \pm 0.05\%$ at an addition rate of 200 kg N ha⁻¹ y⁻¹ ($P < 0.0001$, $F = 20.99$ for the N addition effect on square root transformed data), with a lower allocation for *A. viridis* spp.

crispa than for *A. incana* spp. *rugosa* ($P = 0.04$, $F = 3.35$ for the species effect). Also, at the $200 \text{ kg N ha}^{-1} \text{ y}^{-1}$ addition rate, 33% of the plants failed to form nodules, whereas only 16% of the plants failed to form nodule at all other nitrogen addition levels combined. An ANOVA model using only plants grown at this highest N addition rate indicated that there was no difference in plant mass between plants that formed nodules and plants that did not ($P = 0.61$, $F = 0.27$). For the plants with nodules, specific nodule activity (SNA) decreased from $18.48 \pm 4.10 \mu\text{mol C}_2\text{H}_4 \text{ g}^{-1} \text{ h}^{-1}$ at no N addition to $3.85 \pm 0.77 \mu\text{mol C}_2\text{H}_4 \text{ g}^{-1} \text{ h}^{-1}$ at a N addition rate of $200 \text{ kg N ha}^{-1} \text{ y}^{-1}$ ($P = 0.02$, $F = 5.61$ for the N addition effect on log transformed data) with no differences between the three species. Because both allocation to nodules and their physiological activity declined with increases N addition, there was an overall reduction in nitrogenase activity per plant mass from $193.57 \pm 60.67 \text{ nmol C}_2\text{H}_4 \text{ g}^{-1} \text{ h}^{-1}$ at no N addition to $6.0.4 \pm 1.68 \text{ nmol C}_2\text{H}_4 \text{ g}^{-1} \text{ h}^{-1}$ at a N addition rate of $200 \text{ kg ha}^{-1} \text{ y}^{-1}$ ($P < 0.0001$, $F = 20.98$ for the N addition effect on log transformed data) with *A. viridis* spp. *crispa* having significantly lower rates than *A. incana* spp. *rugosa* ($P = 0.03$, $F = 3.64$).

Discussion

Nitrogen is the primary limiting nutrient for plant growth. Our study found that increasing N deposition did not bring as great benefits to N-fixers (*Alnus* spp.) compared with closely related non-N fixers (*Betula* spp.) in two ways. First, we found that biomass production of non-N fixing plants responded more strongly to increasing levels of soil N at both ambient and ancient CO_2 , compared with N fixing plants. N fixing plants grew more than non-N fixing plants at low levels of soil N, while high levels of soil N resulted in greater increases in growth of non-N fixing plants. Our results support the previous studies suggesting that N fertilization has limited effect on N fixing plant growth but increases the growth and abundance of unrelated non-N fixers^{31–34}. For example, Skogen²⁴ found the ratio of aboveground mass of a non-fixer (*Solidago canadensis*) to an N fixer (*Desmodium cuspidatum*) increased with the increasing levels of N deposition. This suggested that N fixing plants are not competitive under high level of N when they grow with non-N fixers. Our study showed that under present atmospheric CO_2 conditions N-fixing plants will only have an advantage over non-N fixing plants at low levels of soil N. Second, our study found that increasing levels of N decrease nodule allocation, specific nodule activity and nitrogen fixation rate per plant mass. This shows that actinorhizal plants can be facultative N fixers, downregulating N fixation at high levels of soil N, as has been found in other lab studies²⁶. This would seem to contradict Menge and Crews⁵ hypothesis that actinorhizal plants are unable to maintain their dominance in high latitude regions because they are unable to switch from nitrogen fixation to inorganic soil N uptake, and so they become less competitive as soil N levels increase. We agree that under most conditions where actinorhizal plants are found, there is little regulation of nitrogen fixation since soil N levels are not high enough to affect a response in the plants. More importantly, even though actinorhizal plants can reduce nodule formation and activity as soil N increases, they are not able to increase growth to the same degree that closely related non-nitrogen fixing plants can, making them less competitive.

Atmospheric CO₂ levels have been decreasing since nitrogen fixing plants first evolved. Our study suggests there is a greater advantage in being a nitrogen fixer under ancient levels of CO₂, in that it takes a higher level of soil N for non-nitrogen fixing plants to have greater growth than nitrogen fixing plants. This implies that there was greater selection pressure in the past for the evolution of symbiotic nitrogen fixation. It was proposed that higher CO₂ levels in the late Cretaceous might favor the nitrogen fixing symbiosis. Our results support the hypothesis that the decreasing CO₂ constrains favoring the evolution of nitrogen fixing nodules³. At present the clade of plants that includes the actinorhizal plants has many non-nitrogen fixing species as well ^{35,36}. It is likely that the ability to fix nitrogen has been subsequently lost numerous times since it first evolved^{2,4,37}. As atmospheric CO₂ levels dropped since the Late Cretaceous, there is likely to have been increasing selective pressure to lose the ability to fix nitrogen but utilize nitrogen from the soil³. Under current rising CO₂ levels, there is some evidence that nitrogen fixation is becoming a more favorable trait, at least in the absence of other nutrient limitations (e.g., P, Mo)^{12,38}. At the same time, anthropogenic inputs of N tend to counteract this advantage of being a nitrogen fixer^{39,40}. It is unclear from our data as to why nitrogen fixing plants do not have the same rate of growth increase in response to nitrogen additions as non-nitrogen fixing plants do. Skogen²⁴ has suggested that nitrogen fixing plants cannot uptake inorganic N as efficiently as non-N fixing plants, but the mechanism for this has not been described.

Our study did not show a stimulation of nitrogen fixation and plant growth at an ancient CO₂ level. Previous study has shown that stimulation of N fixation under elevated CO₂ is limited by the availability of other soil nutrients, especially P³⁸ since symbiotic N fixation has a high demand of soil P⁴⁰. Lower P (0.03 mM) reduced cell division in the cortex and prenodules and emerging nodules number as compared with medium P concentration (0.1 mM)⁴¹. The soil we used was low in P and therefore likely prevented a response in nitrogen fixation as elevated CO₂ levels. Future studies are therefore needed to test the interactive effect of elevated CO₂ and N deposition on N-fixing and non-N fixing plants in the absence of other nutrient limitations (e.g., P).

Overall, our study found that increasing N level could disadvantage N fixers relative to non-N fixers. Although actinorhizal plants can reduce nitrogen fixation in response to high soil N, this does not allow them to match the increasing growth of non-N fixers. This may help to explain the lack of N fixing plant dominance at present.

Materials And Methods

Three nitrogen fixing and three non-fixing plant species were grown at two levels of CO₂ and four levels of N deposition. Three-month old seedlings of nitrogen fixers (*Alnus incana* ssp. *rugosa*, *Alnus viridis* ssp. *crispa*, and *Alnus rubra*) and the non-nitrogen fixers (*Betula pumila*, *Betula papyrifera*, *Betula glandulosa*) were grown in six growth chambers, three of which were maintained at 400 ppm, and three of which were maintained at 1600 ppm CO₂. The concentration of CO₂ was monitored continuous and adjusted by

injecting CO₂ into the chambers as needed using a microcontroller-based system⁴². The species are all in the Betulaceae family. *Betula* spp. are the closest relative to *Alnus* spp., and possibly still retain the precursor state of nitrogen fixation⁴³. The plants were grown in *ca.* 0.5 liters pots in a sandy soil collected from a forest with *A. viridis* ssp. *crispa* growing in the understory of mature *Pinus banksiana* trees. Previous study has shown that the soil from the site has an inorganic N level of 10.2 ± 0.6 mg/kg, and an extractable phosphate level of 0.98 ± 0.33 mg/kg⁴⁴. All of the plants, including the non-nitrogen fixers, were inoculated with 10 mL of *Frankia* strain Cpl1. The strain Cpl1 was cultured for four months in the sterilized liquid nitrogen-free *Frankia* Defined Minimal Medium (FDM) containing sodium succinate as carbon source⁴⁵. The cultures were dispersed with a tissue grinder and then diluted 10 times in sterilized distilled water, using for inoculation⁴⁶. Six weeks after inoculation, plants were fertilized with one of four levels of N fertilization (0, 0.42, 2.28, 8.4 mM ammonium nitrate) once a week for a total of nine weeks. These four N fertilization treatments simulate a range of N deposition levels (0, 10, 50, and 200 kg N ha⁻¹ year⁻¹), as per Skogen²⁴. A modified Rorison nutrient solution²⁶ with the N concentration varied was used in the first fertilization. For the remaining eight weeks plants received different levels of ammonium nitrate. After 9 weeks fertilization, plants were inoculated again with crushed nodules (3 mg/plant) of *Alnus*⁴⁶. At week 24, Nitrogen fixation rate was estimated using the acetylene reduction assay²⁶. Fresh nodules were placed into 50 mL bottles and sealed with a rubber septum with a 10% acetylene atmosphere. The bottles were incubated for 1 h at 22 °C. A 5 mL gas sample from each bottle was collected and analyzed for ethylene with gas chromatograph (Varian 450) with a Haysep T column, FID and gas-sampling valve. The specific nodule activity (SNA) was expressed as the ethylene production rate per nodule dry mass. Nitrogen fixation rate per total plant mass was also calculated. Nodule allocation was calculated as the percentage of total plant dry mass.

Statistics Analysis

The data analysis used a three-factor least square model with CO₂ level, N deposition level, and plant group (N fixers or non-N fixers), with species nested within plant group. The N deposition treatment was considered as a continuous variable, and the CO₂ level and plant group were considered nominal variables. When there was a significant interaction between N deposition and the CO₂ level or plant group, the effect of N deposition on the response variables was modelled using linear regressions, with the data transformed as needed in order to maintain homogeneity of variances. The data was analyzed using JMP Pro 14 (SAS Institute).

Declarations

Acknowledgements This research was financed by University of Manitoba Entrance Scholarship, University of Manitoba and Natural Sciences, Engineering Research Council of Canada (NSERC), and Graduate Education Teaching stipend. Special thanks go to my supervisor Dr. John Markham and my lab mate Paige Anderson.

Author contributions statement H.C. conceived and conducted the experiment with the guidance of J.M. H.C. and J.M. analysed the results. All authors reviewed the manuscript.

Additional information

Ethics declarations The authors declare no competing interests.

References

1. Tajika, E. Climate change during the last 150 million years: reconstructing from a carbon cycle model. *Earth Planet Sci Lett.* **160**, 659–707 (1998).
2. Li, H.-L. *et al.* Large-scale phylogenetic analyses reveal multiple gains of actinorhizal nitrogen-fixing symbioses in angiosperms associated with climate change. *Sci Rep.* **5**, 14023 (2015).
3. van Velzen, R., Doyle, J. J. & Geurts, R. A resurrected scenario: single gain and massive loss of nitrogen-fixing nodulation. *Trends Plant Sci.* **24**, 49–57 (2019).
4. Griesmann, M. *et al.* Phylogenomics reveals multiple losses of nitrogen-fixing root nodule symbiosis. *Science.* **361**, eaat1743 (2018).
5. Menge, D. N. L. & Crews, T. E. Can evolutionary constraints explain the rarity of nitrogen-fixing trees in high-latitude forests? *New Phytol.* **211**, 1195–1201 (2016).
6. Menge, D. N. L. *et al.* Nitrogen-fixing tree abundance in higher-latitude North America is not constrained by diversity. *Ecol Lett.* **20**, 842–851 (2017).
7. Tedersoo, L. *et al.* Global database of plants with root-symbiotic nitrogen fixation: NodDB. *J Veg Sci.* **29**, 560–568 (2018).
8. Swensen, S. M. & Benson, D. R. Evolution of actinorhizal host plants and *Frankia* endosymbionts. In *Nitrogen-fixing Actinorhizal Symbioses*. Springer Netherlands. 73–104 (2008).
9. Houlton, B. Z., Wang, Y.-P., Vitousek, P. M. & Field, C. B. A unifying framework for dinitrogen fixation in the terrestrial biosphere. *Nature* **454**, 327–330 (2008).
10. Luo, Y. *et al.* Progressive nitrogen limitation of ecosystem responses to rising atmospheric carbon dioxide. *BioScience.* **54**, 731 (2004).
11. Cheng, W., Inubushi, K., Yagi, K., Sakai, H. & Kobayashi, K. Effects of elevated carbon dioxide concentration on biological nitrogen fixation, nitrogen mineralization and carbon decomposition in submerged rice soil. *Biol Fertil Soils.* **34**, 7–13 (2001).
12. Ainsworth, E. A. & Long, S. P. What have we learned from 15 years of free-air CO₂ enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂. *New Phytol.* **165**, 351–372 (2004).
13. Pawlowski, K. & Newton, W. E. *Nitrogen-fixing Actinorhizal Symbioses*. (Springer Netherlands, 2008). doi:10.1007/978-1-4020-3547-0.
14. Dentener, F. *et al.* Nitrogen and sulfur deposition on regional and global scales: a multimodel evaluation. *Global Biogeochemical Cycles.* **20**, GB4003-n/a (2006).

15. Vitousek, P. M. *et al.* Human alteration of the global nitrogen cycle: sources and consequences. *Ecol Appl.* **7**, 737–750 (1997).
16. Benjamin, W. S. *et al.* Spatially robust estimates of biological nitrogen (N) fixation imply substantial human alteration of the tropical N cycle. *PNAS.* **111**, 8101–8106 (2014).
17. Li, X. *et al.* Seasonal and spatial variations of bulk nitrogen deposition and the impacts on the carbon cycle in the arid/semiarid grassland of inner Mongolia, China. *PloS one.* **10**, e0144689 (2015).
18. Lamarque, J. F. *et al.* Assessing future nitrogen deposition and carbon cycle feedback using a multimodel approach: analysis of nitrogen deposition. *J Geophys Res Atmos.* **110**, D19303-n/a (2005).
19. Tian, D. & Niu, S. A global analysis of soil acidification caused by nitrogen addition. *Environ Res Lett.* **10**, 24019 (2015).
20. Binkley, D. & Högberg, P. Tamm review: revisiting the influence of nitrogen deposition on Swedish forests. *Forest Ecol Manag.* **368**, 222–239 (2016).
21. Lee, J. A. Unintentional experiments with terrestrial ecosystems: ecological effects of sulphur and nitrogen pollutants. *J Ecol.* **86**, 1–12 (1998).
22. Southon, G. E., Field, C., Caporn, S. J. M., Britton, A. J. & Power, S. A. Nitrogen deposition reduces plant diversity and alters ecosystem functioning: field-scale evidence from a nationwide survey of UK heathlands. *PloS one.* **8**, e59031 (2013).
23. Vitousek, P. M., Menge, D. N. L., Reed, S. C. & Cleveland, C. C. Biological nitrogen fixation: rates, patterns and ecological controls in terrestrial ecosystems. *Phil Trans R Soc B.* **368**, 20130119 (2013).
24. Skogen, K. A., Holsinger, K. E. & Cardon, Z. G. Nitrogen deposition, competition and the decline of a regionally threatened legume, *Desmodium cuspidatum*. *Oecologia.* **165**, 261–269 (2011).
25. Salvagiotti, F. *et al.* Growth and nitrogen fixation in high-yielding soybean: impact of nitrogen fertilization. *Agron J.* **101**, 958–970 (2009).
26. Markham, J. H. & Zekveld, C. Nitrogen fixation makes biomass allocation to roots independent of soil nitrogen supply. *Can J Bot.* **85**, 787–793 (2007).
27. Coley, P. D. Possible effects of climate change on plant/herbivore interactions in moist tropical forests. *Clim Change.* **39**, 455–472 (1998).
28. Coley, P. D., Massa, M., Lovelock, C. E. & Winter, K. Effects of elevated CO₂ on foliar chemistry of saplings of nine species of tropical tree. *Oecologia.* **133**, 62–69 (2002).
29. Roger, M. G., Damian, J. B. & Jason, L. L. The effects of elevated [CO₂] on the C:N and C:P mass ratios of plant tissues. *Plant soil* **224**, 1–14 (2000).
30. McLauchlan, K. K., Williams, J. J., Craine, J. M. & Jeffers, E. S. Changes in global nitrogen cycling during the Holocene epoch. *Nature.* **495**, 352–355 (2013).
31. Hossain, M. A., Ishimine, Y., Akamine, H. & Kuramochi, H. Effect of nitrogen fertilizer application on growth, biomass production and N-uptake of torpedograss (*Panicum repens* L.). *Weed Biol Manag.* **4**,

- 86–94 (2004).
32. Thomas, R. B., Bashkin, M. A. & Richter, D. D. Nitrogen inhibition of nodulation and N₂ fixation of a tropical N₂-fixing tree (*Gliricidia sepium*) grown in elevated atmospheric CO₂. *New Phytol.* **145**, 233–243 (2000).
 33. Dordas, C. A. & Sioulas, C. Safflower yield, chlorophyll content, photosynthesis, and water use efficiency response to nitrogen fertilization under rainfed conditions. *Ind Crops Prod.* **27**, 75–85 (2008).
 34. Xu, D. *et al.* Interactive effects of nitrogen and silicon addition on growth of five common plant species and structure of plant community in alpine meadow. *Catena.* **169**, 80–89 (2018).
 35. Roy, a & Bousquet, J. The evolution of the actinorhizal symbiosis through phylogenetic analysis of host plants. *Acta Bot Gall* **143**, 635–650 (1996).
 36. Swensen, S. M. The evolution of actinorhizal symbioses: Evidence for multiple origins of the symbiotic association. *Am J Bot.* **83**, 1503–1512 (1996).
 37. van Velzen, R. *et al.* Comparative genomics of the nonlegume *Parasponia* reveals insights into evolution of nitrogen-fixing rhizobium symbioses. *Proc Natl Acad Sci.* **115**, E4700–E4709 (2018).
 38. Rogers, A., Ainsworth, E. A. & Leakey, A. D. B. Will elevated carbon dioxide concentration amplify the benefits of nitrogen fixation in legumes? *Plant Physiol.* **151**, 1009–1016 (2009).
 39. DeLuca, T. H., Zackrisson, O., Gundale, M. J. & Nilsson, M. C. Ecosystem feedbacks and nitrogen fixation in boreal forests. *Science.* **320**, 1181 (2008).
 40. Zheng, M., Zhou, Z., Luo, Y., Zhao, P. & Mo, J. Global pattern and controls of biological nitrogen fixation under nutrient enrichment: a meta-analysis. *Glob Chang Biol.* **25**, 3018–3030 (2019).
 41. Gentili, F., Wall, L. G. & Huss-Danell, K. Effects of phosphorus and nitrogen on nodulation are seen already at the stage of early cortical cell divisions in *Alnus incana*. *Ann. Bot.* **98**, 309–315 (2006).
 42. Chen, H. & Markham, J. Using microcontrollers and sensors to build an inexpensive CO₂ control system for growth chambers. *Appl. Plant Sci.* **8**, e11393 (2020).
 43. Werner, G. D. A., Cornwell, W. K., Sprent, J. I., Kattge, J. & Kiers, E. T. A single evolutionary innovation drives the deep evolution of symbiotic N₂-fixation in angiosperms. *Nat Commun.* **5**, 4087 (2014).
 44. Chen, H., Renault, S. & Markham, J. The effect of *Frankia* and multiple ectomycorrhizal fungal species on *Alnus* growing in low fertility soil. *Symbiosis.* **80**, 207–215 (2020).
 45. Noridge, N. A. & Benson, D. R. Isolation and nitrogen-fixing activity of *Frankia* sp. strain Cpl1 vesicles. *J. Bacteriol.* **166**, 301–305 (1986).
 46. Markham, J. H. Does *Dryas integrifolia* fix nitrogen? *Botany.* **87**, 1106–1109 (2009).

Figures

Figure 1

The response of nitrogen fixing and non-N fixing plant groups to the increasing soil N level interactive with two level of CO₂ (400 ppm and 1600 ppm CO₂). $P = 0.006$, $F = 8.03$ came from the interactive effect between N level and CO₂ in the least square model. $P < 0.0001$, $F = 28.22$ came from the interactive effect between the plant group and nitrogen level in the least squares model. Green line represented N fixers, and orange line represented non-N fixers. The solid round symbol represented *A. rugosa*. The solid triangle symbol represented *A. rubra*. The solid square symbol represented *A. crispa*. The open round symbol represented *B. glandulosa*. The open triangle symbol represented *B. papyrifera*. The open square symbol represented *A. pumila*.

Figure 2

The effect of increasing soil N level on the proportion biomass allocated to nodules, acetylene reduced per nodule mass (SNA) and per plant mass.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryFigureS1.pdf](#)