

Musculoskeletal pain in other body sites is associated with new-onset low back pain: A longitudinal study among survivors of the Great East Japan Earthquake

Yutaka Yabe

Tohoku University

Yoshihiro Hagiwara (✉ hagi@med.tohoku.ac.jp)

Takuya Sekiguchi

Tohoku University

Yumi Sugawara

Tohoku University

Masahiro Tsuchiya

Tohoku University

Shinichirou Yoshida

Tohoku University

Yasuhito Sogi

Tohoku University

Toshihisa Yano

Tohoku University

Takahiro Onoki

Tohoku University

Tadahisa Takahashi

Tohoku University

Jun Iwatsu

Tohoku University

Ichiro Tsuji

Tohoku University

Eiji Itoi

Tohoku University

Research article

Keywords: Great East Japan Earthquake, Low back pain, Musculoskeletal pain, natural disaster

Posted Date: March 23rd, 2020

DOI: <https://doi.org/10.21203/rs.2.21418/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Musculoskeletal Disorders on April 13th, 2020. See the published version at <https://doi.org/10.1186/s12891-020-03234-0>.

Abstract

Background: Low back pain (LBP) is a common health problem experienced after natural disasters. LBP is often concurrent with other musculoskeletal pain; however, the effects of preexisting musculoskeletal pain on the development of LBP are not clear. The purpose of this study was to elucidate the association of musculoskeletal pain in other body sites with new-onset LBP among survivors of the Great East Japan Earthquake (GEJE). **Methods:** A longitudinal study was conducted with survivors of the GEJE. The survivors who did not have LBP at the three year time period after the GEJE were followed up one year later (n = 1,782). Musculoskeletal pain, such as low back, hand and/or foot, knee, shoulder, and neck pain, were assessed with self-reported questionnaires. The outcome of interest was new-onset LBP, which was defined as LBP absent at three years but present at four years after the disaster. The main predictor was musculoskeletal pain in other body sites three years after the GEJE, which was categorized according to the number of pain sites (0, 1, ≥ 2). Multiple regression analyses were performed to calculate the odds ratio (OR) and 95% confidence interval (CI) for new-onset LBP due to musculoskeletal pain in other body sites. **Results:** The incidence of new-onset LBP was 14.1% (251/1,782). Musculoskeletal pain in other body sites was significantly associated with new-onset LBP. Including people without other musculoskeletal pain as a reference, the adjusted OR and 95% CI for new-onset LBP were 1.73 (1.16-2.57) for people with one musculoskeletal pain site and 3.20 (2.01-5.09) for people with ≥ 2 sites (p < 0.001). **Conclusions:** Preexisting musculoskeletal pain in other body sites was associated with new-onset LBP among survivors in the recovery period after the GEJE .

Background

Low back pain (LBP) is one of the most frequent health problem worldwide, and the ranges of prevalence for point, 1-year, and lifetime are reported to be 12-33%, 22-65%, and 51-84%, respectively [1, 2]. Moreover, LBP increases disability-adjusted life years [3, 4]; therefore, gaining an understanding of the factors related to LBP are important. Risk factors for LBP include age, sex, obesity, smoking, psychological distress, and sleep disturbance in the adult population [3, 5-8]. Further, musculoskeletal pain often occurs at multiple sites, and single-site pain has been shown to increase the risk of pain at other sites [9]. Indeed, some reports have found that LBP occurs concurrently with other musculoskeletal pain [10-12]. Most of these studies were cross-sectional; therefore, the association of preexisting musculoskeletal pain with new onset of LBP is not clear.

Musculoskeletal pain, including LBP, is reported to increase after natural disasters [13]. Disasters create severe stress in survivors due to fear of death, deprivation of sleep, the loss of housing and social relationships, all of which are considered to lower their pain threshold [13]. The Great East Japan Earthquake (GEJE), accompanied by a devastating tsunami, attacked the north-eastern coastal areas of Japan on March 11, 2011 [14]. This terrible disaster resulted in serious damage to these areas, and required a long period of reconstruction. A high prevalence of LBP has also been reported after the GEJE [15, 16], and previous longitudinal studies have revealed associated factors, such as subjective economic hardship and sleep disturbance [7, 16]. The prevalence of musculoskeletal pain apart from LBP (hereafter

referred to as “other musculoskeletal pain”) was also increased in the recovery phase after the GEJE, and almost half of the survivors had musculoskeletal pain at multiple sites [17]. Since musculoskeletal pain could co-exist at multiple sites, we speculated that increased other musculoskeletal pain could be associated with new onset of LBP, and could lead to a high prevalence of LBP after natural disasters. Clarifying this association can be helpful to prevent and treat LBP after a disaster. The aim of this study was to examine the association of other musculoskeletal pain with new-onset LBP in the recovery period after the GEJE longitudinally.

Methods

Participants

We hypothesized that other musculoskeletal pain could be associated with new onset of LBP after natural disasters. A panel study was conducted with the GEJE survivors living in the severely damaged coastal areas, including Ogatsu and Oshika areas in Ishinomaki City, and Wakabayashi Ward in Sendai City, Miyagi prefecture, Japan [18]. The main purpose of the study was to comprehend the physical, mental, and social problems experienced by the survivors to provide effective support. The surveys began three months after the GEJE and were administered every six months. The first study population included residents registered in the Residential Registry of the Ogatsu and Oshika areas and survivors living in prefabricated housing in the Wakabayashi Ward [19]. Because the number of responders increased up to three years after the GEJE and remained constant after that period, we decided to examine the data at the three and four year time periods after the GEJE. From November 2013 to February 2014, three years after the GEJE, the residents (aged 18 years or over) who were registered in the Residential Registry of Ogatsu and Oshika areas, and the survivors who had participated in the previous survey in Wakabayashi Ward, were recruited (n = 6,396). Self-reported questionnaires and informed consent forms were mailed to these residents and a 44.6% (2,853/6,396) response rate was obtained. Among those, the participants who already had LBP were excluded (n = 663). The remaining participants were followed from November 2014 to February 2015, four years after the GEJE, and an 81.4% (1,782/2,189) follow-up rate was obtained for this period. Finally, a total of 1,782 participants were included in this study (Fig. 1). This study was approved by the institutional review board of our university (approval number: 201192) and was performed in accordance with the ethical standards as laid down in the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards.

Musculoskeletal pain

Musculoskeletal pain was assessed using self-reported questionnaires based on the Comprehensive Survey of Living Conditions. The questions were: “Have you had symptoms in the last few days? If yes, please place a mark next to all your symptoms.” The examples of choices were palpitation, dizziness, diarrhoea, and musculoskeletal symptoms such as low back, hand and/or foot, knee, shoulder, and neck pain [17]. The outcome of interest was new-onset LBP, which was defined as LBP absent at three years (first period), and present at four years (second period) after the GEJE. The main predictor was other

musculoskeletal pain at the first period which included hand and/or foot, knee, shoulder, and neck pain. Other musculoskeletal pain was categorized into three groups according to the number of painful sites (0, 1, ≥ 2).

Covariates

The following variables were included in the analysis because they were considered potential confounding factors in previous reports [16, 20]: sex, age, body mass index (BMI), living area, smoking habits, drinking habits, comorbid conditions (hypertension, diabetes mellitus, myocardial infarction, and cerebral stroke), working status, walking time per day, living status, subjective economic conditions, psychological distress (Kessler Psychological Distress Scale) [21], sleep disturbance (Athens Insomnia Scale) [22], and social isolation (Lubben Social Network Scale) [20].

Statistical analysis

Univariate and multivariate logistic regression models were used to calculate odds ratios (OR) and 95% confidence intervals (95% CI) for new-onset LBP according to the number of other musculoskeletal pain sites in the first period. Variables included in the analysis were sex (male or female), age (continuous variable), BMI (continuous variable), living area (Ogatsu, Oshika, or Wakabayashi), smoking habits (non-smoker, smoker, or unknown), drinking habits (non-drinker, < 45.6 grams of alcohol per day, ≥ 45.6 grams of alcohol per day, or unknown), comorbid conditions (absence or presence of each comorbid condition), working status (unemployed, employed, or unknown), walking time per day (< 30 min, 30min to < 1 h, ≥ 1 h, or unknown), living status (living in the same house as before the GEJE, prefabricated housing, new house, others, or unknown), subjective economic conditions (normal, a little bit hard, hard, very hard, or unknown), Kessler Psychological Distress Scale (continuous variable), Athens Insomnia Scale (continuous variable), and Lubben Social Network Scale (continuous variable). We further divided the participants into subgroups by sex (male or female), and ORs and 95% CIs for new-onset LBP were calculated in the same manner. For the stratified analysis, multiplicative interaction between other musculoskeletal pain and sex were tested using the Wald test. In addition, the ORs and 95% CIs for new-onset LBP according to each musculoskeletal pain except LBP in the first period were evaluated. We included the same variables (Model 1) and added each musculoskeletal pain such as hand and/or foot, knee, shoulder, and neck pain as covariates (Model 2). All statistical analyses were performed using SPSS 24.0 (SPSS Japan Inc, Tokyo, Japan). A p value of < 0.05 was accepted as statistically significant.

Results

Baseline characteristics of the participants are presented in Table 1. Among the 1,782 participants, 1,343 (75.4%) had 0, 283 (15.9%) had one, 156 (8.8%) had two or more other musculoskeletal pain regions in the first period, respectively. The participants who reported having other musculoskeletal pain were more likely to be female and older. They were also more likely to have high BMI, comorbid conditions such as hypertension and myocardial infarction, short walking time, subjective economic hardship, higher scores on the Kessler Psychological Distress and Athens Insomnia Scales, and a lower score on the Lubben

Social Network Scale (Table 1). The rate of new-onset LBP was 14.1% (251/1,782). The crude and adjusted ORs and 95% CIs for new-onset LBP according to the number of other musculoskeletal pain regions are shown in Table 2. Other musculoskeletal pain was significantly associated with new-onset LBP in the crude and adjusted analyses. Including people without other musculoskeletal pain as a reference, adjusted ORs and 95% CIs for new-onset LBP were 1.73 (1.16–2.57) for people with one musculoskeletal pain site, and 3.20 (2.01–5.09) for people with ≥ 2 (p for trend < 0.001) (Table 2). The results for the stratified analysis are shown in Table 3. Other musculoskeletal pain was significantly associated with new-onset LBP in each group. The association was stronger in males compared with females (adjusted OR 3.16 (1.68–5.95) for people with one musculoskeletal pain site and 4.60 (2.06–10.30) for people with ≥ 2 , p for trend: < 0.001 in males and 1.24 (0.71–2.14), 2.54 (1.38–4.68), 0.011 in females). There was no statistically significant multiplicative interaction between other musculoskeletal pain regions and sex (Table 3). For each musculoskeletal pain site, hand and/or foot, knee, shoulder, and neck pain were all associated with new-onset LBP in Model 1, and the association was also significant for knee and neck pain in Model 2. The adjusted ORs and 95% CIs (p value) for new-onset LBP were 2.04 (1.30–3.21, 0.002) for Model 1 and 1.33 (0.80–2.19, 0.27) for Model 2 where hand and/or foot pain were exposure variables; 2.56 (1.66–3.96, < 0.001) for Model 1 and 1.87 (1.16–3.01, 0.01) for Model 2 where knee pain was an exposure variable; 2.41 (1.32–4.39, 0.004) for Model 1 and 1.56 (0.82–2.95, 0.18) for Model 2 where shoulder pain was an exposure variable; and 2.61 (1.77–3.85, < 0.001) for Model 1 and 2.10 (1.40–3.17, < 0.001) for Model 2 where neck pain was an exposure variable, respectively (Table 4).

Discussion

The present study revealed that preexisting other musculoskeletal pain was associated with new-onset LBP among the survivors in the recovery period after the GEJE. Further, the effect was stronger with musculoskeletal pain that occurred at multiple sites.

Some cross-sectional studies have shown that musculoskeletal pain often occurs at multiple sites, such as shoulder, elbow, knee, and low back [23, 24]. Further, other authors reported a significant association between LBP and neck or knee pain [10-12]. A small number of longitudinal studies have investigated the effect of musculoskeletal symptoms on the onset of LBP. Smith et al. reported that preexisting pain resulting from arthritis or injury was associated with new-onset LBP [25]. Papageorgiou et al. showed that musculoskeletal pain history was a predictor of subsequent LBP [26]. The results of the present study reveal that the existence of musculoskeletal pain is associated with subsequent onset of LBP, which corresponds with these previously published reports. There has been speculation in the literature about the association between concurrent pain at different sites. Pain at one site can negatively affect motion or posture and place additional burden on the other parts of the body [27]. The factors associated with one pain can also be related to the other pain [28]. In addition, pain at one site can cause central sensitization which can result in the development of pain at other sites [11]. These conditions may explain the association between preexisting musculoskeletal pain and new-onset LBP.

To our knowledge, this is the first study to report that the effect of musculoskeletal pain on onset of LBP becomes stronger with multisite musculoskeletal pain, which may be considered a dose-response relationship. Nordstoga et al. reported that LBP with an increasing number of musculoskeletal pain sites tends to have a worse recovery rate, which also supports our results [29]. The association of musculoskeletal pain with LBP is stronger due to an increased number of pain sites. A high prevalence of musculoskeletal pain was reported after the GEJE, and many survivors experienced pain at multiple sites [17]. This is presumed to be one explanation for our finding of increased LBP after the GEJE. Attention should be paid to other musculoskeletal pain sites to treat and prevent LBP after natural disasters.

The stratified analysis according to sex categories revealed that the association of other musculoskeletal pain with new-onset LBP was also significant among categories in each group, which likely highlights the robustness of the association in this study. The rate of musculoskeletal pain was higher in females compared with males; however, the association of musculoskeletal pain with LBP was stronger in males. However, musculoskeletal pain, especially multisite pain, is more common among females [23, 24], and various factors, such as menopause and loss of oestrogen, may affect such pain [30], which is assumed to lower the association of musculoskeletal pain with LBP in females.

Some authors reported the association between LBP and hand or foot [31], knee [12, 28], shoulder [31], and neck pain [11] in cross-sectional studies. There have also been a small number of longitudinal studies regarding the association between LBP and each musculoskeletal pain, and preexisting LBP was reported to be associated with onset of knee [27] and neck pain [10]. To our knowledge, the present study was the first to report that preexisting knee and neck pain were individually associated with onset of LBP, even if the effect of the other musculoskeletal pain was considered. There is a closed kinetic relationship between the knee and lower back [12], and dysfunction of the knee joint due to pain may result in compensation and LBP. The spine undergoes a similar aging process, including genetic influences and risk factors to pain in the neck and lower back [10], which can cause LBP following neck pain. Some twin studies showed the important roles of genetic factors on pain, however, the effects depended on age [32-34]. Further study is needed to clarify the mechanism of the relationships between LBP and neck pain. On the other hand, the association of hand and/or foot, and shoulder pain with LBP was not significant when considering the effect of the other musculoskeletal pain. Other musculoskeletal pain may also be associated with other pain, and that association may affect the results. Further, survivors who experienced LBP in the first period were excluded from this study, because the purpose of this study was to assess the effects of other musculoskeletal pain on LBP onset. The survivors who already had both LBP and other musculoskeletal pain were excluded, which could reduce the association.

The strength of this study was its longitudinal design that allowed us to investigate the research question. Further, this study included a large sample size of 1,782 participants along with a high follow-up rate of 81.4% at the second time-point. In addition, stratified analyses were performed to highlight the robustness of the results. On the other hand, this study had several limitations. First, the questionnaires and informed consent forms were mailed to the participants, and the response rate for the first time-point was low at 44.6%. Although there is no information regarding non-responders, it might be that responders

are healthier than non-responders, or that more severely affected persons may pay more attention to their situation, and may therefore be more likely to respond. These issues could affect the reported rate of musculoskeletal pain, and strengthen or weaken the association between other musculoskeletal pain and new-onset LBP. Further, this study examined the data at the three and four year time periods after the GEJE. The other time period had a different response rate, which could also affect the results. Second, musculoskeletal pain was assessed using a self-report questionnaire based on the Comprehensive Survey of Living Conditions. Although this survey is widely accepted in Japan to assess the symptoms of participants, the reliability and validity of this method were not evaluated in this study. Further, the questionnaire included five pain sites but did not include other pain sites such as hip or elbow. Pain at these sites could also affect the onset of LBP and were not assessed in this study. In addition, pain severity and frequency were not assessed. The difference in severity and frequency of LBP might depend on the number of other musculoskeletal pain sites, which could not be assessed. Finally, this study was conducted with disaster survivors and generalizability of the results might be unclear. Future studies are required to determine if the findings of this study can be extrapolated to the general population.

Conclusions

Preexisting musculoskeletal pain at other sites (especially knee and neck) was associated with new-onset LBP among survivors in the recovery period after the GEJE. Further research is needed to clarify whether this association is truly causal and to identify the mechanisms that could offer an explanation for why people with other musculoskeletal pain are at a higher risk of developing LBP, which could offer useful recommendations for clinicians and public health policies for LBP.

Abbreviations

BMI: Body mass index, GEJE: Great East Japan Earthquake, LBP: Low back pain, OR: Odds ratio, 95%CI: 95% Confidence interval

Declarations

Ethics approval and consent to participate:

The study protocol was reviewed and approved by the Ethics Committee of Tohoku University Graduate school of Medicine (approval number: 201192). Written informed consent was obtained from all the participants.

Consent to publish

Not applicable

Availability of data and materials

All relevant data are contained in this article.

Competing interests

The authors declare no conflict of interest.

Funding:

This work was supported by Health Sciences Research Grants (H24, 25-Kenki-Shintei-002 [Fukkou]) from the Ministry of Health, Labor, and Welfare of Japan.

Authors' contributions

YY, YH, and IT contributed to the design of the study. YS, MT, SY, YS, TY, TT and TO were responsible for data collection and supervised the study. TS and YY performed statistical analysis. YY, YH, and EI wrote the manuscript. YS, MT, IT, and JI helped to analyse the data and draft the manuscript. All authors have read and approved the manuscript.

Acknowledgements

Not applicable

Authors' details

¹Department of Orthopaedic Surgery, Tohoku University School of Medicine, 2-1 Seiryomachi, Aoba-ku, Sendai, Miyagi 980-8574, Japan. ²Division of Epidemiology, Department of Health Informatics and Public Health, Tohoku University, Graduate School of Public Health, 2-1 Seiryomachi, Aoba-ku, Sendai, Miyagi 980-8575, Japan. ³Department of Nursing, Faculty of Health Science, Tohoku Fukushi University, 1-8-1, Kunimi, Aoba-ku, Sendai, Miyagi 981-8522, Japan.

References

1. McBeth J, Jones K. Epidemiology of chronic musculoskeletal pain. *Best Pract Res Clin Rheumatol*. 2007;21:403-425. doi: 10.1016/j.berh.2007.03.003.
2. Hoy D, Bain C, Williams G, March L, Brooks P, Blyth F, Woolf A, Vos T, Buchbinder R. A systematic review of the global prevalence of low back pain. *Arthritis Rheum*. 2012;64:2028-2037. doi: 10.1002/art.34347
3. Fujii T, Matsudaira K. Prevalence of low back pain and factors associated with chronic disabling back pain in Japan. *Eur Spine J*. 2013;22:432-438. doi: 10.1007/s00586-012-2439-0.
4. Vos T, Abajobir AA, Abate KH, Abbafati C, Abbas KM, Abd-Allah F, Abdulkader RS, Abdulle AM, Abebo TA, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990-2016: a systematic analysis for the Global Burden of

- Disease Study 2016. *Lancet* (London, England). 2017;390:1211-1259. doi: 10.1016/S0140-6736(17)32154-2.
5. Deyo RA, Weinstein JN. Low back pain. *N Engl J Med*. 2001;344:363-370. doi: 10.1056/NEJM200102013440508.
 6. Pincus T, Burton AK, Vogel S, Field AP. A systematic review of psychological factors as predictors of chronicity/disability in prospective cohorts of low back pain. *Spine (Phila Pa 1976)*. 2002;27:E109-120.
 7. Yabe Y, Hagiwara Y, Sekiguchi T, Sugawara Y, Tsuchiya M, Koide M, Itaya N, Yoshida S, Sogi Y, Yano T, Tsuji I, Itoi E. Sleep disturbance is associated with new onset and continuation of lower back pain: a longitudinal study among survivors of the Great East Japan Earthquake. *Tohoku J Exp Med*. 2018;246:9-14. doi: 10.1620/tjem.246.9.
 8. Angst F, Angst J, Ajdacic-Gross V, Aeschlimann A, Rössler W. Epidemiology of back pain in young and middle-aged adults: a longitudinal population cohort survey from age 27–50 years. *Psychosomatics*. 2017;58:604-613.
 9. Butera KA, Roff SR, Buford TW, Cruz-Almeida Y. The impact of multisite pain on functional outcomes in older adults: biopsychosocial considerations. *J Pain Res*. 2019;12:1115-1125. doi: 10.2147/jpr.S192755.
 10. Kaaria S, Solovieva S, Leino-Arjas P. Associations of low back pain with neck pain: a study of industrial employees with 5-, 10-, and 28-year follow-ups. *Eur J Pain (London, England)*. 2009;13:406-411. doi: 10.1016/j.ejpain.2008.05.006.
 11. Fernandez-de-las-Penas C, Hernandez-Barrera V, Alonso-Blanco C, Palacios-Cena D, Carrasco-Garrido P, Jimenez-Sanchez S, Jimenez-Garcia R. Prevalence of neck and low back pain in community-dwelling adults in Spain: a population-based national study. *Spine (Phila Pa 1976)*. 2011;36:E213-219. doi: 10.1097/BRS.0b013e3181d952c2.
 12. Rahbar M, Shimia M, Toopchizadeh V, Abed M. Association between knee pain and low back pain. *J Pak Med Assoc*. 2015;65:626-631.
 13. Angeletti C, Guetti C, Papola R, Petrucci E, Ursini ML, Ciccozzi A, Masi F, Russo MR, Squarcione S, Paladini A, Pergolizzi J, Taylor R, Jr., Varrassi G, Marinangeli F. Pain after earthquake. *Scand J Trauma Resusc Emerg Med*. 2012;20:43. doi: 10.1186/1757-7241-20-43.
 14. Ishigaki A, Higashi H, Sakamoto T, Shibahara S. The Great East-Japan Earthquake and devastating tsunami: an update and lessons from the past Great Earthquakes in Japan since 1923. *Tohoku J Exp Med*. 2013;229:287-299.
 15. Yabuki S, Ouchi K, Kikuchi S, Konno S. Pain, quality of life and activity in aged evacuees living in temporary housing after the Great East Japan earthquake of 11 March 2011: a cross-sectional study in Minamisoma City, Fukushima prefecture. *BMC Musculoskelet Disord*. 2015;16:246. doi: 10.1186/s12891-015-0711-2.
 16. Yabe Y, Hagiwara Y, Sekiguchi T, Sugawara Y, Sato M, Kanazawa K, Koide M, Itaya N, Tsuchiya M, Tsuji I, Itoi E. Influence of living environment and subjective economic hardship on new-onset of low

- back pain for survivors of the Great East Japan Earthquake. *J Orthop Sci.* 2017;22:43-49. doi: 10.1016/j.jos.2016.11.003.
17. Yabe Y, Hagiwara Y, Sekiguchi T, Sugawara Y, Tsuchiya M, Koide M, Itaya N, Yoshida S, Sogi Y, Yano T, Tsuji I, Itoi E. Musculoskeletal pain is associated with new-onset psychological distress in survivors of the Great East Japan Earthquake. *Disaster Med Public Health Prep.* 2019;13:295-300. doi: 10.1017/dmp.2018.57.
 18. Hagiwara Y, Yabe Y, Sugawara Y, Sato M, Watanabe T, Kanazawa K, Sonofuchi K, Koide M, Sekiguchi T, Tsuchiya M, Tsuji I, Itoi E. Influence of living environments and working status on low back pain for survivors of the Great East Japan Earthquake. *J Orthop Sci.* 2016;21:138-142. doi: 10.1016/j.jos.2015.12.015.
 19. Yabe Y, Hagiwara Y, Sekiguchi T, Sugawara Y, Tsuchiya M, Itaya N, Yoshida S, Sogi Y, Yano T, Onoki T, Tsuji I, Itoi E. Musculoskeletal pain and new-onset poor physical function in elderly survivors of a natural disaster: a longitudinal study after the great East Japan earthquake. *BMC Geriatr.* 2019;19:274. doi: 10.1186/s12877-019-1283-z.
 20. Sone T, Nakaya N, Sugawara Y, Tomata Y, Watanabe T, Tsuji I. Longitudinal association between time-varying social isolation and psychological distress after the Great East Japan Earthquake. *Soc Sci Med.* 2016;152:96-101. doi: 10.1016/j.socscimed.2016.01.037.
 21. Suzuki Y, Fukasawa M, Obara A, Kim Y. Mental health distress and related factors among prefectural public servants seven months after the great East Japan Earthquake. *J Epidemiol.* 2014;24:287-294.
 22. Soldatos CR, Dikeos DG, Paparrigopoulos TJ. Athens Insomnia Scale: validation of an instrument based on ICD-10 criteria. *J Psychosom Res.* 2000;48:555-560.
 23. Coggon D, Ntani G, Palmer KT, Felli VE, Harari R, Barrero LH, Felknor SA, Gimeno D, Cattrell A, Vargas-Prada S, Bonzini M, Solidaki E, Merisalu E, Habib RR, Sadeghian F, Masood Kadir M, Warnakulasuriya SS, Matsudaira K, Nyantumbu B, Sim MR, Harcombe H, Cox K, Marziale MH, Sarquis LM, Harari F, Freire R, Harari N, Monroy MV, Quintana LA, Rojas M, Salazar Vega EJ, Harris EC, Serra C, Martinez JM, Delclos G, Benavides FG, Carugno M, Ferrario MM, Pesatori AC, Chatzi L, Bitsios P, Kogevinas M, Oha K, Sirk T, Sadeghian A, Peiris-John RJ, Sathiakumar N, Wickremasinghe AR, Yoshimura N, Kelsall HL, Hoe VC, Urquhart DM, Derrett S, McBride D, Herbison P, Gray A. Patterns of multisite pain and associations with risk factors. *Pain.* 2013;154:1769-1777. doi: 10.1016/j.pain.2013.05.039.
 24. Kamaleri Y, Natvig B, Ihlebaek CM, Benth JS, Bruusgaard D. Change in the number of musculoskeletal pain sites: a 14-year prospective study. *Pain.* 2009;141:25-30. doi: 10.1016/j.pain.2008.09.013.
 25. Smith BH, Elliott AM, Hannaford PC, Chambers WA, Smith WC. Factors related to the onset and persistence of chronic back pain in the community: results from a general population follow-up study. *Spine (Phila Pa 1976).* 2004;29:1032-1040. doi: 10.1097/00007632-200405010-00016.
 26. Papageorgiou AC, Croft PR, Thomas E, Ferry S, Jayson MI, Silman AJ. Influence of previous pain experience on the episode incidence of low back pain: results from the South Manchester Back Pain Study. *Pain.* 1996;66:181-185. doi: 10.1016/0304-3959(96)03022-9.

27. Ito H, Tominari S, Tabara Y, Nakayama T, Furu M, Kawata T, Azukizawa M, Setoh K, Kawaguchi T, Matsuda F, Matsuda S. Low back pain precedes the development of new knee pain in the elderly population; a novel predictive score from a longitudinal cohort study. *Arthritis Res Ther.* 2019;21:98. doi: 10.1186/s13075-019-1884-0.
28. Croft P, Dunn KM, Von Korff M. Chronic pain syndromes: you can't have one without another. *Pain.* 2007;131:237-238. doi: 10.1016/j.pain.2007.07.013.
29. Nordstoga AL, Nilsen TIL, Vasseljen O, Unsgaard-Tondel M, Mork PJ. The influence of multisite pain and psychological comorbidity on prognosis of chronic low back pain: longitudinal data from the Norwegian HUNT Study. *BMJ open.* 2017;7:e015312. doi: 10.1136/bmjopen-2016-015312.
30. Pieretti S, Di Giannuario A, Di Giovannandrea R, Marzoli F, Piccaro G, Minosi P, Aloisi AM. Gender differences in pain and its relief. *Ann 1st Super Sanita.* 2016;52:184-189. doi: 10.4415/ANN_16_02_09.
31. Suri P, Morgenroth DC, Kwoh CK, Bean JF, Kalichman L, Hunter DJ. Low back pain and other musculoskeletal pain comorbidities in individuals with symptomatic osteoarthritis of the knee: data from the osteoarthritis initiative. *Arthritis Care Res (Hoboken).* 2010;62:1715-1723. doi: 10.1002/acr.20324.
32. Ståhl MK, El-Metwally AA, Mikkelsen MK, Salminen JJ, Pulkkinen LR, Rose RJ, Kaprio JA. Genetic and environmental influences on non-specific neck pain in early adolescence: a classical twin study. *Eur J Pain (London, England).* 2013;17:791-798. doi: 10.1002/j.1532-2149.2012.00247.x.
33. Nielsen CS, Knudsen GP, Steingrímssdóttir ÓA. Twin studies of pain. *Clin Genet.* 2012;82:331-340. doi: 10.1111/j.1399-0004.2012.01938.x.
34. Hartvigsen J, Petersen HC, Frederiksen H, Christensen K. Small effect of genetic factors on neck pain in old age: a study of 2,108 Danish twins 70 years of age and older. *Spine.* 2005;30:206-208. doi: 10.1097/01.brs.0000150481.84048.08.

Tables

Table 1. Baseline characteristics of the participants.

		Number of musculoskeletal pain sites except low back pain			P value
		0	1	≥ 2	
		1,343	283	156	
Sex, n (%)	Male	678 (50.5)	97 (34.3)	51 (32.7)	< 0.001
	Female	665 (49.5)	186 (65.7)	105 (67.3)	
Age (18-97, years)**, mean (SD)	continuous variable	59.8 (18.3)	63.3 (16.1)	65.9 (13.8)	< 0.001
Body mass index (9.2-45.0)**, mean (SD)	continuous variable	23.9 (3.6)	23.9 (3.8)	24.8 (3.6)	< 0.001
Living area, n (%)	Ogatsu	573 (42.7)	117 (41.3)	80 (51.3)	0.101
	Oshika	535 (39.8)	112 (39.6)	45 (28.8)	
	Wakabayashi	235 (17.5)	54 (19.1)	31 (19.9)	
Smoking habits*, n (%)	Non-smoker	1,011 (75.3)	204 (72.1)	121 (77.6)	0.154
	Smoker	263 (19.6)	55 (19.4)	24 (15.4)	
Drinking habits*, n (%)	Non-drinker	794 (59.1)	178 (62.9)	101 (64.7)	0.13
	< 45.6g of alcohol/day	273 (20.3)	54 (19.1)	24 (15.4)	
	≥ 45.6g of alcohol/day	148 (11.0)	20 (7.1)	11 (7.1)	
Comorbid conditions, n (%)	Hypertension	499 (37.2)	123 (43.5)	86 (55.1)	< 0.001
	Diabetes mellitus	122 (9.1)	28 (9.9)	19 (12.2)	0.444
	Myocardial infarction	70 (5.2)	18 (6.4)	20 (12.8)	0.001
	Cerebral stroke	21 (1.6)	8 (2.8)	4 (2.6)	0.282
Working status*, n (%)	Unemployed	696 (51.8)	160 (56.5)	95 (60.9)	0.123
	Employed	619 (46.1)	115 (40.6)	59 (37.8)	
Walking time/day*, n (%)	≥ 1h	423 (31.5)	69 (24.4)	27 (17.3)	< 0.001
	30min to < 1h	503 (37.5)	92 (32.5)	64 (41.0)	
	< 30min	394 (29.3)	117 (41.3)	62 (39.7)	
Living status*, n (%)	Same house as before the GEJE	393 (29.3)	90 (31.8)	47 (30.1)	0.494
	Prefabricated house	530 (39.5)	103 (36.4)	52 (33.3)	
	New house	163 (12.1)	34 (12.0)	27 (17.3)	
	Others	246 (18.3)	55 (19.4)	30 (19.2)	
Economic condition*, n (%)	Normal	653 (48.6)	114 (40.3)	60 (38.5)	< 0.001

	A little hard	347 (25.8)	65 (23.0)	43 (27.6)	
	Hard	207 (15.4)	77 (27.2)	41 (26.3)	
	Very hard	98 (7.3)	26 (9.2)	11 (7.1)	
Kessler Psychological Distress Scale (0-24)**, mean (SD)	continuous variable	3.2 (4.0)	5.5 (5.0)	6.7 (5.0)	< 0.001
Athens Insomnia Scale (0-23)**, mean (SD)	continuous variable	3.3 (3.1)	5.3 (3.9)	6.2 (4.0)	< 0.001
Lubben Social Network Scale (0-30)**, mean (SD)	continuous variable	15.3 (6.0)	14.2 (5.6)	14.3 (6.2)	0.004

*Because each item has a limited number of respondents, the actual number is not necessarily in accordance with the total.

**Indicates the range of the participants. GEJE; Great East Japan

Earthquake, SD; standard deviation

Table 2. Association of musculoskeletal pain except LBP with new-onset LBP.

	Number of musculoskeletal pain sites except LBP				P for trend
	total	0	1	≥ 2	
Participants	1,782	1,343	283	156	
New-onset LBP, n (%)	251 (14.1)	155 (11.5)	53 (18.7)	43 (27.6)	
Crude OR (95% CI)		1	1.77 (1.25-2.49)	2.92 (1.98-4.30)	< 0.001
Adjusted OR (95% CI)		1	1.73 (1.16-2.57)	3.20 (2.01-5.09)	< 0.001

Adjusted for sex, age, body mass index, living area, smoking habits, drinking habits, complications, working status, walking time, living status, subjective economic condition, Kessler Psychological Distress Scale, Athens Insomnia Scale, and Lubben Social Network Scale. LBP: low back pain, OR: Odds ratio, CI: Confidence interval

Table 3. Stratified analysis for each age and sex group.

Number of musculoskeletal pain sites except LBP						
	total	0	1	≥ 2	P for trend	P-interaction
	1,782	1,343	283	156		
Sex						
Male (n = 826)						
New-onset LBP/participants	117/826 (14.2%)	75/678 (11.1%)	26/97 (26.8%)	16/51 (31.4%)		
Adjusted OR (95% CI)		1	3.16 (1.68-5.95)	4.60 (2.06-10.30)	< 0.001	
Female (n = 956)						
New-onset LBP/participants	134/956 (14.0%)	80/665 (12.0%)	27/186 (14.5%)	27/105 (25.7%)		
Adjusted OR (95% CI)		1	1.24 (0.71-2.14)	2.54 (1.38-4.68)	0.011	0.12

Adjusted for age, body mass index, living area, smoking habits, drinking habits, complications, working status, walking time, living status, subjective economic condition, Kessler Psychological Distress Scale, Athens Insomnia Scale, and Lubben Social Network Scale. LBP: low back pain, OR: Odds ratio, CI: Confidence interval

Table 4. Association of each musculoskeletal pain with new-onset LBP.

		Absence	Presence	P value
Hand or foot pain	Participants	1,616	166	
	New-onset LBP, n (%)	212 (13.1)	39 (23.5)	
	Model 1 OR (95% CI)	1	2.04 (1.30-3.21)	0.002
	Model 2 OR (95% CI)	1	1.33 (0.80-2.19)	0.27
Knee pain	Participants	1,595	187	
	New-onset LBP	206 (12.9)	45 (24.1)	
	Model 1 OR (95% CI)	1	2.56 (1.66-3.96)	< 0.001
	Model 2 OR (95% CI)	1	1.87 (1.16-3.01)	0.01
Shoulder pain	Participants	1,708	74	
	New-onset LBP	229 (13.4)	22 (29.7)	
	Model 1 OR (95% CI)	1	2.41 (1.32-4.39)	0.004
	Model 2 OR (95% CI)	1	1.56 (0.82-2.95)	0.18
Neck pain	Participants	1,558	224	
	New-onset LBP	190 (12.2)	61 (27.2)	
	Model 1 OR (95% CI)	1	2.61 (1.77-3.85)	< 0.001
	Model 2 OR (95% CI)	1	2.10 (1.40-3.17)	< 0.001

Adjusted for sex, age, body mass index, living area, smoking habits, drinking habits, complications, working status, walking time, living status, subjective economic condition, Kessler Psychological Distress Scale, Athens Insomnia Scale, and Lubben Social Network Scale (Model 1). Additionally, adjusted for hand or foot pain, knee pain, shoulder pain, and neck pain (Model 2). LBP: low back pain, OR: Odds Ratio, CI: Confidence interval.

Figures




Figure 1

Flowchart of this study