

Children's Adherence to Antiretroviral Therapy and Associated Factors: Multicenter Cross-Sectional Study

Getahun B. Gemechu (✉ getahunbeyene1@gmail.com)

Jimma University

J Habtamu

Jimma University

K Zarihun

Jimma University

Research Article

Keywords: adherence, ART, children, Ethiopia

Posted Date: April 11th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1202199/v4>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background- Suboptimal adherence to antiretroviral therapy will lead to drug resistance, treatment failure, clinical deterioration, death and failure to thrive in children. Studies conducted among children below 15 years old were limited in Ethiopia in general and in the study area. Therefore, this study aimed to assess the status of children's adherence to ART and associated factors in the study area.

Methods- We conducted a facility-based cross-sectional study by including 282 children <15 years who received anti-retroviral therapy for at least one month. All children/caregivers who were attending the ART clinic during the data collection period were consecutively recruited for the study. Both bivariate and multivariate logistic regression were performed.

Result- Out of 282 caregivers included with their children, 226 (80.2%) were females (mean age= 38.6 and SD = 12.35). In similar, among total number of children included in the study, half (50%) were female and 246 (87.2%) were between the ages 5–14 years (mean age= 8.5 and SD = 2.64). Two hundred and forty six (87.2%) children had an adherence status of $\geq 95\%$ in the month prior to the interview.

Children whose caregivers were residing in urban areas were 3.3 (95% CI: 1.17, 9.63) times more adherent to ART when compared to caregivers in rural areas. Children whose caregivers were biological parents, were 2.37(95% CI: 1.59, 3.3) times more adherent when compared with caregivers with non-biological parents. Additionally, children of caregivers who were knowledgeable about ART treatment, were 4.5(95% CI: 1.79, 9.8) times more adherent to ART than their counterparts.

Conclusion and recommendation- The adherence status of children in our study area was comparable. Biological caregivers, residing in urban areas and knowledgeable about ART treatment facilitate adherence to ART. Ongoing education about treatment and further study with multiple adherence assessment methods were recommended.

Background

According to a global HIV/AIDS report, the status of adherence to ART among HIV infected children varied from 65.0% to 90.0% (1), 75.0% in developing countries(2) and 88.8% in Ethiopia(3).

An important factor in achieving treatment success is maintaining optimal adherence to ART (4–6). Adherence to antiretroviral treatment is defined as taking 95% or more of the prescribed doses on time and in the correct way, either with or without food (5,7).

Poor adherence to ART will lead to an increased risk of drug resistance, treatment failure and clinical deterioration with increased premature death in children(9,10,12).

The causes of poor adherence to ART among pediatric patients are complex and interrelated factors. Total dependency of the child on caregiver, pill burden and dosing frequency, treatment side effects, poor

health literacy, limited access to ART(10–12), socioeconomic status of the caregiver, substance use and access to service delivery) (13) (14) are among some factors.

The emergence of drug resistant HIV virus among children adds an enormous period. It may create an introduction of resistant HIV virus to the general population as the child with HIV grows to adolescence and young adulthood(15)(16).

Therefore, designing strategies for maintaining the optimal status of adherence among children, is essential to ensure treatment success. However, the task requires careful assessment of the status of adherence and age specific factors (17). Studies conducted among children below 15 years of age are limited in Ethiopia, in general and in the study area. Additionally, adherence is not static and can vary with time on treatment, making assessing adherence status crucial(14). Moreover, viral load was added as one variable in this study, which has not been researched in previous studies. Therefore, this study aimed to examine the status of adherence and associated factors among children under the age of 15 years in the study area.

Methods And Materials

A facility based cross sectional study was conducted from April 1 to May 10, 2020 in the Sebeta, Teji, Tullu Bollo and Waliso towns which are located 20 km, 50 km, 65 km and 114 km away from Addis Ababa, respectively. These towns were, in the southwestern direction along the main road from Addis Ababa to Jimma. In total, there are 8 public ART sites included in all 4 towns. Three of the public ART sites are within Sebeta town (Sebeta HC provide ART services for 41 under 15 children, while 21 and 09 are served in Alemgena and Tefki HC respectively. Two public ART sites are found in Tullu Bolo town. Tullu Bolo General Hospital serves 25 HIV positive children and Tullu Bolo HC provides services for 17 HIV positive children. Teji town

also, has Teji HC who provide services for 18 under 15 children. The rest are found in Waliso town, (St. Lukes Referral Hospital serves 130 and Waliso HC serves 29 under 15 children). In total, 290 children are on ART in those health facilities. These health facilities have case managers and adherence supporters.

Sample size was calculated by considering the ratio of unexposed to exposed 0.6, outcome in the exposed group 49.0%, and outcome in the unexposed group 29.5%(18) for both objectives. Then after, largest sample was selected, with a 10.0% nonresponse rate, giving a sample size of 254. There were 290 children on ART in these health facilities, and all were include.

Pretested and structured questionnaires, using face- to - face interviews, were held for data collection. The questionnaires were adapted from previously conducted related studies. Some changes were made after reviewing the relevant literature. The questionnaire was translated to Afan Oromo and Amharic for better understanding by the respondents.

Pretest of the questionnaire was performed on 5.0% (14) of the caregivers in Awash Health Center. Medical records were reviewed to collect clinical data such as WHO clinical stages, regimen type, viral loads and the CD4 counts of children.

Data were collected by 8 clinical nurses. The data collection process was supervised by one health officer and one BSc nurse. Adherence and exposure variables were measured by the caregivers' or children's report of a one-month recall of missed doses prior to the date of the interview. The questionnaires were coded and entered into the EPI data statistical software and thereafter, exported to SPSS Windows version 23 for further analysis. Data were summarized and presented using descriptive and analytic statistics. Bivariate and multiple logistic regressions were computed to identify the presence and strength of associations. Variables with a p value <0.25, in binary logistic regression, were subjected to multiple logistic regression for analysis. However, before multiple logistic regression, collinearity diagnostic test was done and no multicollinearity (maximum Variance Inflation Factor (VIF) =3.883). On the other hand, Breslow-Day test showed, there significant synergetic interaction between two variables (i.e. child relationship with caregiver vs caregiver ART history). As a result, the stratum specific odds ratio was considered to measure the strength of the association. At the end, multiple logistic regression (odds ratios with 95% CIs were computed) and variables with p-values less than 0.05 were considered significantly associated with the dependent variable. Finally, the model was fitted using the backward elimination method, and the Hosmer-Lemeshow goodness of fit test indicated a p- value of 0.47).

Operational Definition

Adherent to ART – when the child has missed ≤ 1 dose or takes $\geq 95\%$ of ART in the month prior to the interview.

Non adherent to ART- missing at least two doses or takes <95% of ART in the month prior to interview.

Care givers- Parent/guardian or person in charge of routinely administering antiretroviral drugs to children on ART.

Baseline CD4 count- CD4 count done when a child started ART(7).

Current CD count- CD4 count done within 6 months of data collection(7).

Knowledge about ART- Those respondents who scored greater than or equal to the mean for the knowledge questions were considered knowledgeable; otherwise, they were not.

Care giver/child substance use- if caregiver/child use either of alcohol, khat, cigarettes or other substance in the month prior to data collection.

Ethical considerations

Ethical clearance letters were obtained from the Ethical Review Committee, Jimma University, Institute of Health. Each health facility was contacted after the official letter was submitted. The interviews were held

after verbal informed consent and assurance of confidentiality.

Results

Sociodemographic characteristics of the study participants

Of the 290 study participants, 282 children, along with their caregivers, were included in the analysis, yielding a response rate of 97.2%. Four caregivers were excluded because they did not appear in the health facility during the data collection period. Three questionnaires were discarded as they were not complete (caregivers did not know the dosing history for the past month) and one caregiver was not willing to respond to the questionnaire. Among the respondents, 226 (80.2%) of the caregivers were females, and the mean age of the respondents was 38.6 years (SD =12.35). Out of the total respondents, 176 (62.5%) were married and 31 (11.0%) were single. One hundred and fifty-eight (56.0%) of the caregivers attended primary education. Two hundred and nine (74.1%) of the caregivers were urban dwellers. Among the children included in the study, half (50.0%) were female, 246 (87.2%) were 5–14 years old with the mean age of 8.5 years (SD = 2.64). One hundred and eighty three (64.9%) children are currently attending 1-4 grades. (**Table 1**).

Table 1- Socio demographic characteristics of study participants, central Ethiopia, April 1- May 10, 2020. (n=282)

Variables		Frequency	Percentage
Caregivers Sex	female	226	80.2
	male	56	19.8
Caregivers age	<30yrs old	60	21.3
	>=30yrs old	222	78.7
Caregivers marital status	married	176	62.5
	widowed	53	18.8
	single	31	11.0
	Divorced	22	7.7
Caregivers educational status	Not educated	67	23.7
	Primary	158	56.0
	Secondary	48	17.4
	Tertiary	8	2.8
caregivers religion	Orthodox	216	76.6
	Protestant	40	14.2
	Muslim	26	9.2
Caregivers ethnicity	Oromo	231	81.9
	Amhara	38	13.4
	Gurage	13	4.7
caregivers occupation	Self Employed	147	52.1
	Farmer	76	27.0
	Merchant	45	15.9
	Gov't employee	11	3.8
	Unemployed	3	1.1
caregivers residence area	Urban	209	74.1
	Rural	73	25.9
Caregivers monthly income	< 500 ETB	35	12.4
	≥ 500 ETB	247	87.6
Sex of child	Male	141	50.0
	Female	141	50.0
Age of child	0-4 yrs. old	36	12.8
	5-14 yrs. old	246	87.2
Educational status of child	Not eligible	33	11.7
	Not started	45	15.9
	KG	6	2.0
	1-4 grades	183	64.9
	5-8 grades	15	5.4

Clinical characteristics of study participants

Among the children who participated in this study, 102 (36.2%) started ART at WHO clinical stage I and 273 (96.8%) were currently on WHO stage I. At initiation of ART, 142 (50.4%) children had a CD4 count of >500 cells/mm³. During data collection, viral load was done for 206 (73%) children and out of these, high viral load (≥1000 copies/ml) was detected in 55 (19.5%) children. (**Table 2**)

Variables		N ₀	%
Baseline CD4 count of the child	≤200	26	9.2
	201-499	54	19.2
	≥500	142	50.4
	Not done	60	21.2
Current CD4 count of the child	≤200	3	1.0
	201-499	13	4.6
	≥500	109	38.7
	Not done	157	55.7
WHO stage at Initiation of ART	WHO stage I	102	36.2
	WHO stage II	87	30.9
	WHO stage III	82	29.0
	WHO stage IV	11	3.9
Current WHO stage	WHO stage I	273	96.8
	WHO stage II	4	1.4
	WHO stage III	4	1.4
	WHO stage IV	1	0.4
Viral RNA Copies per ml	<1000	151	53.5
	≥1000	55	19.5
	Not done	76	26.9

Table 2- Clinical characteristics of study participants, in central Ethiopia, April 1- May 10, 2020. (n=282)

ART and other drug history of the children

Among the children included in the study, 279 (95.8%) were on 1st line ART medication and 195 (69.2%) of the children were taking 2 pills per day (morning and night), while 38 (13.4%) of the children were taking 3 doses daily. The current study also showed that among children who took part in the study, 247 (87.6%) had been on ART for more than one year. Concerning CPT, 200 (70.9%) of the children were not on CPT at the time of data collection. **(Table 3)**.

Variables		No	%
Duration of child on ART	<1yrs	35	12.3
	>=1yrs	247	87.6
Line of treatment	1 st line	270	95.8
	2 nd line	12	4.2
Number of pills child is taking/day	1 pills	38	13.4
	2 pills	195	69.2
	≥ 3pills	49	17.3
Is child taking CPT	Yes	82	29.0
	No	200	70.9

Table 3- ART and other drug history of study participants in central Ethiopia, April 1-May 10, 2020. (n=282).

Among children on ART in study the area, 176 (62.4%) are on AZT-3TC-NVP and 47 (15.6 %) are on AZT-3TC-EFV regimen. Only 6 (2.0%) of the children who are currently on ART have been taking a combination of AZT based kaletra (boosted drug with Protease inhibitor).

Adherence status of the children to ART

Based on the caregivers' report, a total of 246 (87.2%, (95% CI: 83.3%- 91.1%) children had an adherence status of ≥95% in the month prior to interview. The status of adherence to ART among children in the past three and seven days before the interview date was 98.9% and 95.8%, respectively.

Reasons for non-adherence

According to the current study, 36 (12.8%) of the children missed at least 2 doses out of their prescribed dose preceding for the month before data collection. The commonly mentioned reasons were: fear of stigma (21.3%), caregiver's forgetfulness (14.2%), child's refusal to take the drugs (13.8%) and run out of medication (13.8%). (Fig 1)

Distance from health facility and mode of transportation

Out of the total caregivers/children who participated in the study, the majority, 219 (77.7%), came from less than 10 km and 113 (40.1%) of the participants came to the health facility on foot to refill medication. (Table 4)

Variables		No	%
Distance	< 10 km	219	77.7
	≥ 10 km	63	22.3
Mode of presentation	On foot	113	40.1
	By public bus	104	36.9
	By cart	22	7.8
	By Bajaj	43	15.3

Table 4- Distance from health facility and mode of transportation of study participant to refill medication, central Ethiopia, May 10, 2020.

Factors associated with adherence to ART

In multiple logistic regression, the residence area of caregivers, blood relationship between children and caregivers and knowledge of caregivers about ART treatment, were significantly and independently associated with adherence. Children whose caregivers resided in urban areas were 3.3 times more likely adherent to ART than their counterparts [AOR = 3.3(95% CI: 1.17, 9.63)]. Children whose caregivers were biological parents were 2.37 times more likely to be adherent than those whose caregivers were non-biological parents [AOR = 2.37(95% CI: 1.59, 3.2)]. The present study also showed that if the caregiver was knowledgeable about ART treatment, the child was 4.5 times more likely to be adhere to ART than their counterpart [AOR = 4.5(95% CI: 1.79, 9.8)]. (**Table 5**)

Table 4- Crude and adjusted odds ratios and 95% Confidence Intervals (CIs) of explanatory variables of adherence to HAART in the study area, central Ethiopia, April 1 May 10, 2020. (n=282).

Variables		Adherence status		Bivariate Logistic regression		Multiple logistic regression
		Adherent: No (%)	Non-adherent: No (%)	COR(95% CI)	P-value	AOR(95% CI)
Sex of care givers	male	46(82.1)	10(17.9)	0.6(0.27-1.32)	0.206	1.56 (0.51-4.73)
	female	200(88.5)	26(11.5)	1		1
Educational status of care givers	Not educated	48(71.6)	19(28.4)	1		1
	Primary	144(91.1)	14(8.9)	0.25(0.114-0.53)	0.000	0.3(0.45-1.97)
	≥ 9 Grade	54(94.7)	3(5.3)	0.14(0.04-0.504)	0.003	0.78(0.14-4.35)
Residence of care giver	Urban	187(89.5)	22(10.5)	2.02(0.97-4.19)	0.060	3.2(1.17-9.63)**
	rural	59(80.8)	14(19.2)	1		1
Monthly income of caregiver	<500 ETB	22(62.9)	13(37.1)	0.17(0.077-0.39)	0.000	0.45(0.15-1.34)
	≥500 ETB	224(90.7)	23(9.3)	1		1
Educational status of child	Not eligible	26(78.8)	7(21.2)	1		1
	Not started	41(91.1)	4(8.9)	0.36(0.097-1.36)	0.133	0.35(0.005-24.0)
	KG-4 grades	165(87.3)	24(12.7)	0.54(0.211-1.38)	0.198	0.12(0.02-7.24)
	5-8 grades	14(93.3)	1(6.7)	0.27(0.03-2.38)	0.276	
Biological parent	Yes	213(95.1)	11(4.9)	14.67(6.6-32.59)	0.000	2.37(1.59-3.2)**
	No	33(56.9)	25(43.1)	1		1
Is care givers taking ART	Yes	203(94.9)	11(5.1)	10.73(4.9-23.45)	0.000	0.61(0.07-5.51)
	No	43(63.2)	25(36.8)	1		1
Knowledgeable	Yes	206(94.1)	13(5.9)	9.11(4.26-19.48)	0.000	4.9(1.7-9.8)**
	No	40(63.5)	23(36.5)	1		1
Viral RNA Copies per ml	<1000	136(90.1)	15(9.9)	1		1
	≥1000	42(76.4)	13(23.6)	2.8(1.24-6.37)	0.014	0.51(0.14-1.88)
	Not done	68(89.5)	8(10.5)	1.07(0.43-2.64)	0.889	

**----- significant variable

Discussion

According to this study, 87.2% of children had an adherence status of ≥95% in the month prior to the interview. This is higher than the finding of global pooled estimation, which is 65.0% -75.0% (9). This difference could be due to the context of the study area (i.e., specific study area among children living in

almost similar socio-demographic characteristics **vs** global context). In addition, the global pooled estimate used secondary data. However, our finding was comparable with an studies in East Africa 90.0%(19); in Southeastern Nigeria, 91.0%(20); Ethiopian national pooled prevalence of optimal HAART adherence 88.8% (3); Eastern Ethiopia, 90.7% (21), in Mekelle Hospital, Tigray, 90.7% (22); in Gondar University Hospital and Gondar Poly Clinic, 90.4%(23); and in Addis Ababa, 86.9%(24). However, the finding of the current study was higher than the finding of study conducted in India, which revealed an overall adherence of 82.1% (25). This difference might be due to the study design (cross-sectional vs cohort). It was also higher than that in a study in Uganda which reported an overall adherence status of 79.1%(26), South Wallo zone, provided an adherence status of 78.6%(18), and in Fiche Hospital, showed an adherence rate of 74.0%(27). This difference could be due to the small sample size in Fiche Hospital ((120 vs 282). Again, higher adherence rate was observed in our study when compared with finding of study conducted in the ART clinic of Tikur Anbessa Hospital, (i.e. an adherence rate of children to ART was 34.8%) (28). This difference was due to adherence assessment methods (caregiver's report vs unannounced home based pill count). In general, the current study finding of the adherence status of children to ART is below the recommendation of the WHO, which recommended that $\geq 90\%$ of people on ART should be adherent to ART to fully benefit from ART(5).

Regarding associated risk factors for non-adherence, the current study showed that children whose caregivers were biological parents were 2.37 times more adherent than those whose caregivers were non-biological parents. [AOR= 2.37, 95 CI= (1.59-3.2)]. This finding is similar to the cross-sectional study finding conducted in Hiwot Fana and Dil-Chora Referral Hospital in eastern Ethiopia, which revealed that children whose caregivers were biological parents were more adherent.

The other independently associated risk factor for adherence to ART among children identified by this study was caregivers' knowledge of ART. Accordingly, we found that children with knowledgeable caregivers were 4.5 times more adherent than those children whose caregivers were not knowledgeable [AOR 4.5 95% CI (1.7-9.8)]. Our finding is similar to the study finding conducted in Hiwot Fana and Dil-Chora Hospital, which revealed that caregivers of children with good knowledge about the disease were 7.3 times [AOR = 7.3] adherent than their counterparts(21).

Again, our finding is also similar to the finding of a study conducted in India that came up with, children whose caregivers had good knowledge about ART treatment were 14.7 times more adherent (25). A similarly study conducted in south Wollo zone justified our finding [AOR=2.72] (34). Our finding is again in line with the study finding conducted at University of Gondar Hospital and Gondar Poly Clinic, which showed that children of knowledgeable caregivers had 4.7 times more adherent to ART than their counterparts [AOR=4.7](23). This implies that making caregivers knowledgeable about the importance of ART treatment through ongoing and extensive counseling is mandatory to make children more likely to benefit from good adherence to ART.

The current study also showed that children residing in urban areas were 3.2 times more adherent to ART than those residing in rural. [AOR=3.2, (95% CI 1.17-9.63)]. This finding is supported by a study conducted in Tanzania that reported, suboptimal to ART was observed among children in the rural context of the Mwanza region(29). However, our study is contradicts a comparative study conducted between rural and urban children, in South Africa that revealed, rural children were more adherent

to ART (93.1% vs 88.0%)(30). This difference might be due to the study design (cross-sectional vs cohort). In general, this implies that urban dwellers have more access to information and education when compared with rural dwellers.

Our study is not free of limitations. There might be an overestimation of adherence status in our study because of recall and social desirability bias. Thus, to minimize this bias we used non ART providers as data collectors. Additionally, we did not assess adherence related to the correct timing of ART drug administration. Again, the cross-sectional nature of the study also hinders the causal relationship between adherence status and associated factors.

Conclusion And Recommendation

The adherence status of the children to ART in our study area was comparable. Children residing in urban areas, with biological caregivers and children of knowledgeable caregivers were more adherent to ART. Ongoing counseling of non-biological parent and for those who come from rural areas is mandatory. Other studies should be conducted using multiple and more objective adherence assessment tools.

Declarations

Acknowledgment

My heartfelt thanks to Jimma University Institute of Public Health Department of Epidemiology for giving us this opportunity. My deepest gratitude again goes to my advisors Habtemu Jarso (assistant professor) and Mr.Zarihun Kura for their corrective comments

Competing of Enterers Disclaimer

Authors have declared that no competing of interest exist. The product used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of between authors and producers of the product because we do not intent to use these product as an avenue for litigation but for the advancement of knowledge. Also, the research was not funded by producing company rather it was funded by the personal efforts of the authors.

References

1. Iacob SA, Iacob DG, Jugulete G. Improving the Adherence to Antiretroviral Therapy , a Difficult but Essential Task for a Successful HIV Treatment – Clinical Points of View and Practical Considerations. *Frontiers Pharmacol.* 2017;8(doi:10.3389/fphar. 2017.00831 Edited November):1–12.
2. Rachel C. Vreeman, MD, MS, Sarah E. Wiehe, MD, MPH, Emily C. Pearce, MD, and Winstone M. Nyandiko, MbChB Mm. A Systematic Review of Pediatric Adherence to Antiretroviral Therapy in Low- and Middle-Income Countries. *Pediatr Infect J.* 2008;27(8):686–91.
3. Endalamaw A, Tezera N, Eshetie S, Ambachew S, Dejenie T. Adherence to Highly Active Antiretroviral Therapy Among Children in Ethiopia : A Systematic Review and Meta - analysis. *AIDS Behav [Internet].* 2018;22(8):2513–23. Available from: <https://doi.org/10.1007/s10461-018-2152-z>
4. Federal Democratic Republic of Ethiopia F. NATIONAL GUIDELINES FOR HIV PREVENTION TREATMENT AND CARE NATIONAL AIDS AND STI ' s CONTROL. 2016.
5. WHO. The use of antiretroviral drugs for treating and preventing HIV infection. 2016.
6. PEPFAR. Differentiated Care in Ethiopia. The way forward. 2017;(March).
7. Federal Democratic republic of Ethiopia F. Nationa Guide Lines for Comprehensive HIV Prevention, Care and Treatment. 2014.
8. Shargie, Mulatu Biru; Jerene, Degu; Lundqvist, Pia; Molla, Mitikie; Abebe WK, Hallström K. Caregiver-reported antiretroviral therapy non-adherence during the first week and after a month of treatment initiation among children diagnosed with HIV in Ethiopia. *AIDS Care.* 2018;(DOI: 10.1080/09540121.2016.1257098):0–14.
9. Butler LM, Brickley DB, Kennedy G, Rutherford G, Chan J, Asia E. Rates and determinants of adherence to antiretroviral therapy (ART) in infants , children and adolescents : A systematic review. 2010.
10. Teju D. First-line antiretroviral treatment failure and associated factors in HIV patients at the University of Gondar Teaching Hospital , Gondar , Northwest Ethiopia. *HIV/AIDS - Res Palliat Care.* 2016;8(<http://dx.doi.org/10.2147/HIV.S112048>):141–6.
11. Malede Mequanent, Tadesse Awoke, Yalemzewod Assefa AT and et al. Incidence and risk factors of first-line antiretroviral treatment failure among human immunodeficiency virus-infected children in Amhara regional state, Ethiopia a retrospective follow-up study. *BMJ Open.* 2016;10(<http://dx.doi.org/10.1136/bmjopen-2017-019181>):1–10.
12. Zvanaka Sithole, Elizabeth Mbizvo, Prosper Chonzi, More Mungati, Tsitsi Patience Juru, Gerald Shambira, Notion Tafara Gombe and MT. Virological failure among adolescents on ART, Harare City,

2017- a case-control study. BMC Infect Dis. 2018; 18:(doi: [10.1186/s12879-018-3372-6]).

13. Nichols J, Steinmetz A, Paintsil E. Impact of HIV-Status Disclosure on Adherence to Antiretroviral Therapy Among HIV-Infected Children in Resource-Limited Settings: A Systematic Review. AIDS Behav. 2016;(DOI 10.1007/s10461-016-1481-z):1–11.
14. Schaecher KL. The Importance of Treatment Adherence in HIV. AJMC. 2019;19(2):2019.
15. USAID/AIDSTAR-One. IMPLEMENTATION OF WORLD HEALTH ORGANIZATION ' S 2008 PEDIATRIC HIV TREATMENT GUIDELINES TECHNICAL BRIEF. 2010.
16. WHO. The consequences of poor adherence to long-term therapies are poor health outcomes and increased health care costs. 2018.
17. Kassi T, Eboua F, Nouaman M, Bleu Y, Kouadio B, Niamien E, et al. Factors Associated with Non-Adherence to Antiretroviral Therapy among HIV-Infected Adolescents at Pediatric Department of Yopougon University Hospital , Côte d ' Ivoire. Open J Pediatr 8, 238-248. 2018;8(<http://www.scirp.org/journal/ojped>):238–48.
18. Arage G, Tessema GA, Kassa H. Adherence to antiretroviral therapy and its associated factors among children at South Wollo Zone Hospitals , Northeast Ethiopia : a cross-sectional study. BMC Public Health. 2014;14 (<http://www.biomedcentral.com/14712458/14/365>):1–7.
19. Rachel C. Vreeman and etal. Adherence to antiretroviral therapy in a clinical cohort of HIV-infected children in East Africa. PLoS One. 2018;13(2)(e0191848.).
20. Akahara C, Nwolisa E, Odinaka K, Okolo S. Assessment of Antiretroviral Treatment Adherence among Children Attending Care at a Tertiary Hospital in Southeastern Nigeria. J Trop Med. 2017;10(<https://doi.org/10.1155/2017/3605850> Research):1–5.
21. Zegeye S, Gemechu E, Sendo. Adherence to Antiretroviral Therapy among Hiv-Infected Children Attending Hiwot Fana and Dil-Chora Art Clinic at Referral Hospitals in Eastern Ethiopia. J HIV Clin Sci Res. 2015;2(Res 2(1): 008-014):8–14.
22. Gultie T, Amlak TG, Sebsibie G. Factors Affecting Adherence to Pediatrics Antiretroviral Therapy in Mekelle Factors Affecting Adherence to Pediatrics Antiretroviral Therapy in Mekelle Hospital , Tigray Ethiopia. Int J Public Heal Sci. 2015;Vol.4, No.(April):1–6.
23. Dachew BA, Tesfahunegn TB, Birhanu AM. Adherence to highly active antiretroviral therapy and associated factors among children at the University of Gondar Hospital and Gondar Poly Clinic , Northwest Ethiopia : a cross-sectional institutional based study. BMC PublicHealth. 2014;14:875(<http://www.biomedcentral.com/1471-2458/14/875>):1–6.

24. Naod Firdu, Fikre Enquselassie and Degu Jerene¹. HIV-infected adolescents have low adherence to antiretroviral therapy a cross-sectional study in Addis Ababa, Ethiopia. *Pan Afr Med J.* 27,2017.
25. Kendre GM, Gabhale YR, Shah ND, Jadhav VM, Nath K, Manglani M V. Adherence to antiretroviral therapy and factors affecting adherence among paediatric HIV patients. *Int J Contemp Pediatr* Kendre. 2017;4(DOI: <http://dx.doi.org/10.18203/2349-3291.ijcp20174154>):1962–8.
26. Wadunde I, Tuhebwe D, Ediau M, Okure G, Mpimbaza A, Wanyenze RK. Factors associated with adherence to antiretroviral therapy among HIV infected children in Kabale district , Uganda: a cross sectional study. *BMC Res Notes [Internet]*. 2018;(https://doi.org/10.1186/s13104-018-3575-3):1–6. Available from: <https://doi.org/10.1186/s13104-018-3575-3>
27. Feyissa A. Magnitude and Associated Factors of Non-Adherence to Highly Active Antiretroviral Therapy among Children in Fiche Hospital, North Shewa, Ethiopia, 2016. *J Pharm Care Heal Syst.* 2016(4:170. doi: 10.4172/2376-0419.1000170).
28. Biressaw S, Abegaz WE, Abebe M, Taye WA, Belay M. Adherence to Antiretroviral Therapy and associated factors among HIV infected children in Ethiopia : unannounced home-based pill count versus caregivers ' report. *BMCPediatr.*2013;13(<http://www.biomedcentral.com/14712431/13/132>):1471–2431.
29. Id GM, Antonucci R, Mukurasi A, Zepherine H. Adherence to antiretroviral treatment among children and adolescents in Tanzania : Comparison between pill count and viral load outcomes in a rural context of Mwanza region. *PLoS One.* (| <https://doi.org/10.1371/journal.pone.0214014> March):1–15.
30. Musiime V, Kayiwa J, Kiconco M, Tamale W, Alima H, Mugerwa H, et al. Response to Antiretroviral Therapy of HIV Type 1-Infected. *AIDSRESEARCH ANDHUMANRETROVIRUSES.* 2017;38(April):1–13.

Figures




Figure 1

Reason for non-adherence to ART drugs among children in study area, central Ethiopia, April 1- May 10, 2020