

# Children's Adherence to Antiretroviral Therapy and Associated Factors: Multi-center Cross-sectional Study

Getahun B. Gemechu (✉ [getahunbeyene1@gmail.com](mailto:getahunbeyene1@gmail.com))

Jimma University

J Habtamu

Jimma University

K Zarihun

Jimma University

---

## Research Article

**Keywords:** adherence, ART, children, Ethiopia

**Posted Date:** February 2nd, 2022

**DOI:** <https://doi.org/10.21203/rs.3.rs-1202199/v2>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# CHILDREN'S ADHERENCE TO ANTIRETROVIRAL THERAPY AND ASSOCIATED FACTORS: MULTI-CENTER CROSS-SECTIONAL STUDY

*Getahun B. Gemechu<sup>1</sup>, Habtamu J<sup>2</sup>, Zarihun K<sup>3</sup>*

*<sup>1</sup>Field epidemiologist at Ilu Woreda Health Office, <sup>2</sup>Assistant professor at Epidemiology Department, Jimma University, <sup>3</sup> lecturer at Epidemiology Department, Jimma University.*

*Email-getahunbeyene1@gmail.com.*

*Mobile-0912855854/0978143352*

**Background-** Sub-optimal adherence to antiretroviral therapy will lead drug resistance, treatment failure, clinical deterioration, death and failure to thrive in children. Studies conducted among children below 15 years old were limited in Ethiopia in general and in study area in particular. Therefore, this study was aimed to assess status of children's adherence to ART and associated factors in study area.

**Methods-** We conduct a facility-based cross-sectional study by including total of 282 children <15 years, who received Anti retro viral therapy for at least one month. All children/caregivers who were attending ART clinic during data collection period were consecutively recruited to the study. Both bivariate and multivariate logistic regression were performed.

**Result-** Out of 282 caregivers included with their children, 226(80.2%) were females (mean age= 38.6 and SD = 12.35) and out of the total children, half (50%) were female and 246(87.2%) were between the ages 5–14 years (mean age= 8.5 and SD = 2.64). Two hundred forty six (87.2%) children had adherence status of  $\geq 95\%$  in the month prior to interview. Children whose caregivers were residing in urban were 3.3 (95% CI: 1.17, 9.63) times more adherent to ART than those whose caregivers were residing in rural. Children whose caregivers were biological parent were 2.37(95% CI: 1.59, 3.3) times more adherent than those whose caregivers were non biological parent. Also children of caregivers who were knowledgeable about ART treatment, were 4.5(95% CI: 1.79, 9.8) times more adherent to ART than their counter parts

**Conclusion and recommendation-** Adherence status of children in our study area was comparable. Being biological caregivers, residing in urban and knowledgeable about ART treatment were facilitate adherence to ART. Ongoing education about treatment and further study with multiple adherence assessment method were recommended.

**Key words-** adherence, ART, children, Ethiopia

## **Background**

According to global HIV/AIDS report, the status of adherence to ART among HIV infected children varied from 65% to 90% (1), 75% in developing countries(2) and 88.8% in Ethiopia(3).

An important factor in achieving treatment success is maintaining optimal adherence to ART (4–6). Adherence to antiretroviral treatment is defined as taking 95% or more of the prescribed doses on time and in the correct way, either with or without food (5,7).

Poor adherence to ART will lead to increased risk of drug resistance, treatment failure and clinical deterioration with increased pre-mature death in children(9,10,12).

The causes of poor adherence to ART among pediatric are complex and interrelated factors (i.e. total dependency of the child on caregiver for procurement and drug administration, pill burden and dosing frequency, treatment side effects, poor health literacy, poor patient- physician relationship limited access to ART as a result of formulary restrictions(10–12), challenge related age, socioeconomic status of the caregiver, substance use and access to service delivery) (13) (14).

Emergence of drug resistance HIV virus among children adds an enormous burden since it may create an introduction of resistant HIV virus to the general population as the child with HIV growing to adolescent and young adult(15)(16).

As a result, designing strategies for maintaining optimal status of adherence among children is an essential to ensure treatment success. However, this task requires careful assessment of the status of adherence and age specific factors (17). But studies conducted among children below 15 years is limited in Ethiopia in general and in study area in particular. Additionally adherence is not static and can vary with time on treatment making assessing adherence status is crucial(14). Moreover, viral load was added as one variable in this study which is not researched in previous studies. Therefore, this study was aimed to look into the status of adherence and associated factors among children under the age of 15 years in the study area.

## **Methods and Materials**

Facility based cross sectional study was conducted from April 1 to May 10, 2020 in Sebeta, Teji, Tullu Bollo and Waliso towns that are found at 20km,50km, 65km and 114km away from Addis Ababa respectively, in south western direction along the main road from Addis Ababa to Jimma. In these four towns, there are 8 public ART sites and all were included. From those, three are within Sebeta town (Sebeta HC that serves 1135 HIV positive patients, of which 41 were children less than 15 years, Alemgena HC that serves 395 patients, out of which 21 were children under the age of 15 and Tefki HC that give services for 112 patients in which 09 were

children under the age of 15). Two public ART sites are found in Tullu Bolo town, Tullu Bolo General Hospital that serves 245 HIV positive patients of which 25 were children, Tullu Bolo HC that give services for 17 HIV positive children. Teji town has also, Teji HC that provide services for 230 HIV positive patients of which 18 were children under the age of 15. The rest are found in Waliso town, St. Lukes Referral Hospital which serves 1562 HIV infected patients of which 130 are children under the age of 15 and Waliso HC which serves 801 HIV infected patients of which 29 are children below 15 years. Totally 290 children are on ART in those health facilities. All those health facilities have case managers and adherence supporters. After we calculated sample size for both objectives (assessing adherence status and identifying factors), we selected largest sample calculated by considering ratio of unexposed to exposed 0.6, outcome in exposed group 49%, outcome in unexposed group 29.5%(18), with 10% non -response rate giving sample size of 254. However, there were 290 children were on ART in these health facilities and all were included.

Pre-tested and structured questionnaires using face to face interview was held for data collection. The questionnaires was adapted from previously conducted related studies and some change was made after reviewing relevant literature according to local context and translated to Afan Oromo and Amharic for better understanding by the respondents. Pre-test of questionnaire was done

on 5% (14) caregivers in Awash Health Center. Medical records was reviewed to collect clinical data such as WHO clinical stage, regimen type child is on, viral load and the CD4 count of children. Data was collected by 8 clinical nurse. The data collection process was supervised by one health officer and one BSc nurse. Adherence and exposure variables were measured by the caregivers' or children's report of a one-month recall of missed doses prior to the date of the interview. The questionnaires was coded and entered into EPI data statistical software and then exported to SPSS windows version 23 for further analysis. Data was summarized and presented using descriptive and analytic statistics. Bivariate and multiple logistic regressions were computed to identify the presence and strength of associations. Variables with p value <0.25 in binary logistic regression were taken to multiple logistic regression for further analysis. But, before taking variables to multiple logistic regression, multicollinearity was checked for all candidate variables using collinearity diagnostic test and no multicollinearity (maximum Variance Inflation Factor (VIF) =3.883). Interaction was also checked by Breslow-Day test and significant synergetic interaction was detected between two variables (i.e. blood relationship of the child with caregiver and caregiver taking ART). As a result, stratum specific odds ratio was considered to measure strength of association. At the end multiple logistic regression, odds ratio with 95% CI was computed and variables having p-values less than 0.05 in the multiple logistic regression

models were considered significantly associated with the dependent variable. Finally model was fitted using backward elimination method and Hosmer-Lemeshow goodness of fit test indicated that model is fitted well (p- value 0.47).

### **Operational Definition**

**Adherent to ART** – when the child has missed  $\leq 1$  dose or takes  $\geq 95\%$  of ART in the month prior to the interview.

**Non adherent to ART**- missing at least two doses or takes  $<95\%$  of ART in the month prior to interview.

**Care givers**- Parent/guardian or person in charge of routinely administering antiretroviral drugs to children on ART.

**Baseline CD4 count**- CD4 count done when a child started ART(7).

**Current CD count**- CD4 count done within 6 months of data collection(7).

**Knowledge about ART**- Those respondents who scored greater than or equal to the mean for the knowledge questions were considered as knowledgeable, otherwise not.

**Care giver/child substance use**- if caregiver/child use either of alcohol, khat, cigarettes and other substance in the month prior to data collection.

### **Ethical considerations**

Ethical clearance letter is obtained from the Ethical review committee, Jimma University,

institute of health. Each health facilities was contacted after official letter submitted. Interview had been held after informed verbal consent and assurance of confidentiality.

## **Results**

### **Socio-demographic characteristics of the study participants**

Of the 290 study participants, 282 children along with their caregivers were included in the analysis, yielding a response rate of 97.2%. Four caregivers were excluded as they did not appear in the health facility during data collection period. Three questionnaires were discarded as they were not complete (caregivers did not know the dosing history for the past month) and one caregiver was not willing to respond to the questionnaires. Among respondents, 226(80.2%) of the caregivers were females and the mean age of the respondents was 38.6 and SD =12.35. Out of the total respondent, 176 (62.5%) were married and only 31(11%) were single. One hundred fifty eight (56%) of the caregivers have attended primary education. Two hundred nine (74.1%) of the caregivers were urban dwellers. Among the children included in the study, half (50%) were female and 246(87.2%) were in the age of 5–14 years with the mean age of 8.5 and SD = 2.64). One hundred eighty three (64.9%) of the children are currently attending grade 1-4. (**Table 1**)

Table 1- Socio demographic characteristics of study participants, central Ethiopia, April 1- May 10, 2020. (n=282)

Variables		Frequency	Percentage
Sex of care giver	female	226	80.2
	male	56	19.8
Care giver age	<30yrs old	60	21.3
	>=30yrs old	222	78.7
Marital status of care giver	married	176	62.5
	widowed	53	18.8
	single	31	11.0
	Divorced	22	7.7
Educational status of care givers	Not educated	67	23.7
	Primary	158	56.0
	Secondary	48	17.4
	Tertiary	8	2.8
Religions of care givers	Orthodox	216	76.6
	Protestant	40	14.2
	Muslim	26	9.2
Care givers ethnicity	Oromo	231	81.9
	Amhara	38	13.4
	Gurage	13	4.7
Occupation of care giver	Self Employed	147	52.1
	Farmer	76	27.0
	Merchant	45	15.9
	Gov't employee	11	3.8
	Unemployed	3	1.1
Residence of care giver	Urban	209	74.1
	rural	73	25.9
Monthly income of care giver	<500 ETB	35	12.4
	>=500 ETB	247	87.6
Sex of child	Male	141	50.0
	Female	141	50.0
Age of child	0-4 yrs. old	36	12.8
	5-14 yrs. old	246	87.2
Educational status of child	Not eligible	33	11.7
	Not started	45	15.9
	KG	6	2.0
	1-4 grade	183	64.9
	5-8 grade	15	5.4

#### Clinical characteristics of study participants

Among the children participated in this study, 102(36.2%) started ART at WHO clinical stage I and 273(96.8%) were on WHO stage I currently. At initiation of ART, 142(50.4%) children had a

CD4 count of >500 cells/mm<sup>3</sup>. From 206 (73%) children who had viral load count, RNA of HIV virus were <1000 copies /ml in 151(73.3%) children. (Table 2)

*Table 2- Clinical characteristics of study participants at, central Ethiopia, April 1- May 10, 2020. (n=282)*

Variables	No	%	
Base line CD4 count of the child	≤200	26	9.2
	201-499	54	19.2
	≥500	142	50.4
	Not done	60	21.2
Current CD4 count of the child	≤200	3	1.0
	201-499	13	4.6
	≥500	109	38.7
	Not done	157	55.7
WHO stage at Initiation of ART	WHO stage I	102	36.2
	WHO stage II	87	30.9
	WHO stage III	82	29.0
	WHO stage IV	11	3.9
Current WHO stage	WHO stage I	273	96.8
	WHO stage II	4	1.4
	WHO stage III	4	1.4
	WHO stage IV	1	0.4
Viral RNA Copies per ml	<1000	151	53.5
	≥1000	55	19.5
	Not done	76	26.9

#### **ART and other drug history of the children**

Among children included in the study, 279(95.8%) of the children are on 1<sup>st</sup> line

ART medication and 195(69.2%) of the children are taking 2 pills per day (morning and night) while, 38(13.4%) of the children are taking 3 doses daily. The current study also showed that, among children who took part in the study, 247(87.6%) have been on

ART for more than one year. Concerning CPT, 200(70.9%) of the children were not on CPT at the time of data collection. (**Table 3**).

*Table 3- ART and other drug history of study participants at central Ethiopia, April 1-May 10, 2020. (n=282).*

Variables		No	%
Duration of child on ART	<1yrs	35	12.3
	≥1yrs	247	87.6
Line of treatment	1 <sup>st</sup> line	270	95.8
	2 <sup>nd</sup> line	12	4.2
Number of pills child is taking/day	1 pills	38	13.4
	2 pills	195	69.2
	≥ 3pills	49	17.3
Is child taking CPT	Yes	82	29.0
	No	200	70.9

Out of the total children who are on ART in study area, 176(62.4%) of the children are prescribed with AZT-3TC-NVP followed by, AZT-3TC-EFV based regimen which account 47(15.6 %). Only 6(2%) of the children who are currently on ART have been taking a combination of AZT based kaletra (boosted drug with Protease Inhibitor).


#### **Adherence status of the children to ART**

Based on the caregivers' report, a total of 246 (87.2%, (95% CI: 83.3%- 91.1%) children had adherence status of ≥95% in the month prior to interview. The status of adherence to ART among children in the past three and seven days before the interview date was 98.9% and 95.8%, respectively.

#### **Reasons for non-adherence**

According to the current study, 36(12.8%) of children missed at least 2 doses out of their

prescribed dose preceding for the month before data collection. The commonly mentioned reasons were: fear of stigma (21.3%), caregiver's forgetfulness (14.2%), and child's refusal to take the drugs (13.8%) and run out of medication (13.8%). (**Fig 1**)



*Figure 1- Reason for non- adherence to ART drugs among children at study area, central Ethiopia, April 1- May 10, 2020*

#### **Distance from health facility and mode of transportation**

Out of total care givers/children participated in the study, majority, 219(77.7%) were came from less than 10km and 113(40.1%) of participants were came to health facility on foot to refill medication. (**Table 4**)

*Table 4-distance from health facility and mode of presentation to refill medication central Ethiopia, May 10, 2020.*

Variables		No	%
Distance	< 10km	219	77.7
	≥ 10km	63	22.3
Mode of presentation	On foot	113	40.1
	By public bus	104	36.9
	By cart	22	7.8
	By Bajaj	43	15.3

#### **Factors associated with adherence to ART**

In multiple logistic regression, residence area of caregivers, blood relationship between children and caregivers and knowledge of caregivers about ART treatment, were significantly and independently associated with adherence. Children whose caregivers were residing in urban area were 3.3 times more likely to be adherent to ART than their counterparts [AOR = 3.3(95% CI: 1.17, 9.63)]. Children whose caregivers were biological parent were 2.37 times more likely to be adherent than those whose caregivers were non biological parent [AOR = 2.37(95% CI: 1.59, 3.2)]. The present study also showed that, if the caregiver was knowledgeable about ART treatment, the child was 4.5 times more likely to be adherent to ART than their counter part [AOR = 4.5(95% CI: 1.79, 9.8)]. (**Table 5**)

Table 4- Crude and adjusted odds ratio and 95% confidence interval (CI) of explanatory variables of adherence to HAART at study area, central Ethiopia, April 1 May 10, 2020. (n=282).

Variables		Adherence status		Bivariate Logistic regression		Multiple logistic regression
		Adherent: No (%)	Non-adherent: No (%)	COR(95% CI)	P-value	AOR(95% CI)
Sex of care givers	male	46(82.1)	10(17.9)	0.6(0.27-1.32)	0.206	1.56 (0.51-4.73)
	female	200(88.5)	26(11.5)	1		1
Educational status of care givers	Not educated	48(71.6)	19(28.4)	1		1
	Primary	144(91.1)	14(8.9)	0.25(0.114- 0.53)	0.000	0.3(0.45-1.97)
	≥ 9 Grade	54(94.7)	3(5.3)	0.14(0.04-0.504)	0.003	0.78(0.14-4.35)
Residence of care giver	Urban	187(89.5)	22(10.5)	2.02(0.97-4.19)	0.060	3.2(1.17-9.63)**
	rural	59(80.8)	14(19.2)	1		1
Monthly income of caregiver	<500 ETB	22(62.9)	13(37.1)	0.17(0.077-0.39)	0.000	0.45(0.15-1.34)
	≥500 ETB	224(90.7)	23(9.3)	1		1
Educational status of child	Not eligible	26(78.8)	7(21.2)	1		1
	Not started	41(91.1)	4(8.9)	0.36(0.097-1.36)	0.133	0.35(0.005-24.0)
	KG-4 grade	165(87.3)	24(12.7)	0.54(0.211-1.38)	0.198	0.12(0.02-7.24)
	5-8 grade	14(93.3)	1(6.7)	0.27(0.03-2.38)	0.276	
Biological parent	Yes	213(95.1)	11(4.9)	14.67(6.6-32.59)	0.000	2.37(1.59-3.2)**
	No	33(56.9)	25(43.1)	1		1
Is care givers taking ART	Yes	203(94.9)	11(5.1)	10.73(4.9-23.45)	0.000	0.61(0.07-5.51)
	No	43(63.2)	25(36.8)	1		1
Knowledgeable	Yes	206(94.1)	13(5.9)	9.11(4.26-19.48)	0.000	4.9(1.7-9.8)**
	No	40(63.5)	23(36.5)	1		1
Viral RNA Copies per ml	<1000	136(90.1)	15(9.9)	1		1
	≥1000	42(76.4)	13(23.6)	2.8(1.24-6.37)	0.014	0.51(0.14-1.88)
	Not done	68(89.5)	8(10.5)	1.07(0.43-2.64)	0.889	

\*\*----- significant variable

## Discussion

According to this study, 87.2% of children have an adherence status of  $\geq 95\%$  in the month prior to interview. This is higher than the finding of global pooled estimation (9). This difference could be due to the context of study area (i.e. specific study area among children living almost similar socio demographic characteristics vs global context). Besides, global pooled estimate used secondary data. But our finding was comparable with study conducted in East Africa showed 90%(19), Southeastern Nigeria 91%(20), Ethiopian national pooled prevalence of optimal HAART adherence was 88.8% (3), Eastern Ethiopia that provided adherence status of 90.7% (21), in Mekelle hospital, Tigray 90.7% (22), in Gondar University Hospital and Gondar Poly Clinic, 90.4%(23), Addis Ababa 86.9%(24). But the finding of current study was higher than the finding of the study conducted in India, that revealed an overall adherence of 82.1% (25). This difference might be due to study design (cross sectional vs cohort). It was also higher than study in Uganda (overall adherence status of 79.1%)(26), South Wallo zone which provided an adherence status of 78.6%(18), in Fiche Hospital that showed adherence rate of 74%(27). This difference could be due to the fact small sample size (120) in Fiche Hospital as compared to ours (282). Again the current study finding was also greater than the finding of study conducted in ART clinic of Tikur Anbessa Hospital that showed adherence rate of children to ART was 34.8%(28). This difference due to adherence

assessment methods (caregiver's report vs unannounced home based pill count). In general the current study finding of adherence status of children to ART is below the recommendation of WHO which recommended that  $\geq 90\%$  of people on ART should be adherent to ART in order to fully benefit from ART(5).

Regarding associated risk factors for non-adherence, current study provided that children whose caregivers were biological parent were 2.37 times more adherent than those whose caregivers were non biological parent. [AOR= 2.37, 95 CI= (1.59-3.2)]. This finding is similar to the cross sectional study finding conducted in Hiwot Fana and Dil-Chora referral hospital eastern Ethiopia that revealed children whose caregivers were biological parent were more adherent.

The other independently associated risk factor for adherence to ART among children identified by this study was care givers knowledge of ART. Accordingly, we found that children with knowledgeable caregivers were 4.5 times more adherent than those children whose caregivers were not knowledgeable [AOR 4.5 95% CI (1.7-9.8)]. Our finding is similar to the study finding conducted in Hiwot Fana and Dil-Chora hospital that revealed caregivers of children with good knowledge about the disease were 7.3 times [AOR = 7.3] adherent than their counterparts(21). Again our finding is also similar to the finding of study conducted in India that came up

with, children whose caregivers have good knowledge about ART treatment were 14.7 times more adherent (25). Similarly study conducted in south Wollo zone was justify our finding [AOR=2.72] (34). Our finding is again in line with the study finding conducted at University of Gondar Hospital and Gondar Poly Clinic, showed that, children of knowledgeable caregivers had 4.7 times more adherent to ART than their counterparts [AOR=4.7](23). This implies that making caregivers to be knowledgeable about the importance of ART treatment through ongoing and extensive counselling is mandatory to make a children more beneficiary from a good adherence to ART.

Current study also provided that children residing in urban area were 3.2 times more adherent to ART than those who residing in rural. [AOR=3.2, (95% CI 1.17-9.63)]. This finding is supported by the study conducted in Tanzania that reported us suboptimal to ART was observed among children in rural context of Mwanza region(29). But our study is contradicting with comparative study conducted between rural and urban children, in rural area, children whose caregivers were not biological parent and children whose caregivers not knowledgeable. Adherence by using multiple and more objective adherence assessment tool good adherence to ART especially for non-biological parent and for those who come from rural area at each follow up. Another researchers should also conduct another study by using multiple and more objective adherence assessment tool.

south Africa toward ART adherence revealed that rural children were more adherent to ART (93/1% vs 88%)(30). This difference might be due to study design (cross sectional vs cohort). In general, this implies that urban dwellers have more access to information and education when compared to rural dwellers.

Our study is not free of limitation. There might be overestimation of adherence status in our study because of recall and social desirability bias. But we tried to minimize social desirability bias by using non ART providers as data collectors. Also we did not to assess adherence related to correct timing of ART drug administration. Again the cross sectional nature of the study also hinders causal relationship between adherence status and associated factors.

### **Conclusion and recommendations**

Adherence status of the children to ART in our study area was comparable. Children residing in urban area, whose caregivers were biological parent and children of knowledgeable caregivers were more adherent to ART compared to those residing

**Acknowledgment-** My heartfelt thanks to Jimma University Institute of Public Health Department of Epidemiology for giving us this opportunity. My deepest gratitude again goes to my advisors Habtemu Jarso (assistant professor) and Mr.Zarihun Kura for their corrective comments.

### **Competing of enterers**

There is no competing of interest.

## References

1. Iacob SA, Iacob DG, Jugulete G. Improving the Adherence to Antiretroviral Therapy , a Difficult but Essential Task for a Successful HIV Treatment — Clinical Points of View and Practical Considerations. *frontiers Pharmacol.* 2017;8(doi:10.3389/fphar.2017.00831 Edited November):1–12.
2. Rachel C. Vreeman, MD, MS, Sarah E. Wiehe, MD, MPH, Emily C. Pearce, MD, and Winstone M. Nyandiko, MbChB Mm. A Systematic Review of Pediatric Adherence to Antiretroviral Therapy in Low- and Middle-Income Countries. *Pediatr Infect J.* 2008;27(8):686–91.
3. Endalamaw A, Tezera N, Eshetie S, Ambachew S, Dejenie T. Adherence to Highly Active Antiretroviral Therapy Among Children in Ethiopia : A Systematic Review and Meta - analysis. *AIDS Behav* [Internet]. 2018;22(8):2513–23. Available from: <https://doi.org/10.1007/s10461-018-2152-z>
4. Federal Democratic Republic of Ethiopia F. NATIONAL GUIDELINES FOR HIV PREVENTION TREATMENT AND CARE NATIONAL AIDS AND STI ' s CONTROL. 2016.
5. WHO. The use of antiretroviral drugs for treating and preventing HIV infection. 2016.
6. PEPFAR. Differentiated Care in Ethiopia. The way forward. 2017;(March).
7. Federal Democratic republic of Ethiopia F. National Guide Lines for Comprehensive HIV Prevention, Care and Treatment. 2014.
8. Shargie, Mulatu Biru; Jerene, Degu; Lundqvist, Pia; Molla, Mitikie; Abebe WK, Hallström K. Caregiver-reported antiretroviral therapy non-adherence during the first week and after a month of treatment initiation among children diagnosed with HIV in Ethiopia. *AIDS Care.* 2018;(DOI: 10.1080/09540121.2016.1257098):0–14.
9. Butler LM, Brickley DB, Kennedy G, Rutherford G, Chan J, Asia E. Rates and determinants of adherence to antiretroviral therapy ( ART ) in infants , children and adolescents : A systematic review. 2010.
10. Teju D. First-line antiretroviral treatment failure and associated factors in HIV patients at the University of Gondar Teaching Hospital , Gondar , Northwest Ethiopia. *HIV/AIDS - Res Palliat Care.* 2016;8(<http://dx.doi.org/10.2147/HIV.S112048>):141–6.
11. Malede Mequanent, Tadesse Awoke, Yalemzewod Assefa AT and et al. Incidence and risk factors of first-line antiretroviral treatment failure among human immunodeficiency virus-infected children in Amhara regional state, Ethiopia a retrospective follow-up study. *BMJ Open.* 2016;10(<http://dx.doi.org/10.1136/bmjopen>

- 2017-019181):1–10.
12. Zvanaka Sithole, Elizabeth Mbizvo, Prosper Chonzi, More Mungati, Tsitsi Patience Juru, Gerald Shambira, Notion Tafara Gombe and MT. Virological failure among adolescents on ART, Harare City, 2017- a case-control study. *BMC Infect Dis.* 2018; 18:(doi: [10.1186/s12879-018-3372-6]).
  13. Nichols J, Steinmetz A, Paintsil E. Impact of HIV-Status Disclosure on Adherence to Antiretroviral Therapy Among HIV-Infected Children in Resource-Limited Settings : A Systematic Review. *AIDS Behav.* 2016;(DOI 10.1007/s10461-016-1481-z):1--11.
  14. Schaecher KL. The Importance of Treatment Adherence in HIV. *AJMC.* 2019;19(2):2019.
  15. USAID/AIDSTAR-One. IMPLEMENTATION OF WORLD HEALTH ORGANIZATION ' S 2008 PEDIATRIC HIV TREATMENT GUIDELINES TECHNICAL BRIEF. 2010.
  16. WHO. The consequences of poor adherence to long-term therapies are poor health outcomes and increased health care costs. 2018.
  17. Kassi T, Eboua F, Nouaman M, Bleu Y, Kouadio B, Niamien E, et al. Factors Associated with Non-Adherence to Antiretroviral Therapy among HIV-Infected Adolescents at Pediatric Department of Yopougon University Hospital , Côte d ' Ivoire. *Open J Pediatr* 8, 238-248. 2018;8(<http://www.scirp.org/journal/ojped>):238–48.
  18. Arage G, Tessema GA, Kassa H. Adherence to antiretroviral therapy and its associated factors among children at South Wollo Zone Hospitals , Northeast Ethiopia : a cross-sectional study. *BMC Public Health.* 2014;14 (<http://www.biomedcentral.com/14712458/14/365>):1–7.
  19. Rachel C. Vreeman and etal. Adherence to antiretroviral therapy in a clinical cohort of HIV-infected children in East Africa. *PLoS One.* 2018;13(2)(e0191848.).
  20. Akahara C, Nwolisa E, Odinaka K, Okolo S. Assessment of Antiretroviral Treatment Adherence among Children Attending Care at a Tertiary Hospital in Southeastern Nigeria. *J Trop Med.* 2017;10(<https://doi.org/10.1155/2017/3605850> Research):1–5.
  21. Zegeye S, Gemechu E, Sendo. Adherence to Antiretroviral Therapy among Hiv-Infected Children Attending Hiwot Fana and Dil-Chora Art Clinic at Referral Hospitals in Eastern Ethiopia. *J HIV Clin Sci Res.* 2015;2(Res 2(1): 008-014):8–14.
  22. Gultie T, Amlak TG, Sebsibie G. Factors Affecting Adherence to Pediatrics Antiretroviral Therapy in Mekelle Factors Affecting Adherence to Pediatrics Antiretroviral Therapy in Mekelle Hospital , Tigray Ethiopia. *Int J Public Heal Sci.*

- 2015;Vol.4, No.(April):1–6.
23. Dachew BA, Tesfahunegn TB, Birhanu AM. Adherence to highly active antiretroviral therapy and associated factors among children at the University of Gondar Hospital and Gondar Poly Clinic , Northwest Ethiopia : a cross-sectional institutional based study. *BMC PublicHealth*. 2014;14:875(<http://www.biomedcentral.com/1471-2458/14/875>):1–6.
24. Naod Firdu, Fikre Enquesselassie and Degu Jerene1. HIV-infected adolescents have low adherence to antiretroviral therapy a cross-sectional study in Addis Ababa, Ethiopia. *Pan Afr Med J*. 27,2017.
25. Kendre GM, Gabhale YR, Shah ND, Jadhav VM, Nath K, Manglani M V. Adherence to antiretroviral therapy and factors affecting adherence among paediatric HIV patients. *Int J Contemp Pediatr* Kendre. 2017;4(DOI: <http://dx.doi.org/10.18203/2349-3291.ijcp20174154>):1962–8.
26. Wadunde I, Tuhebwe D, Edia M, Okure G, Mpimbaza A, Wanyenze RK. Factors associated with adherence to antiretroviral therapy among HIV infected children in Kabale district , Uganda : a cross sectional study. *BMC Res Notes* [Internet]. 2018;(https://doi.org/10.1186/s13104-018-3575-3):1–6. Available from: <https://doi.org/10.1186/s13104-018-3575-3>
27. Feyissa A. Magnitude and Associated Factors of Non-Adherence to Highly Active Antiretroviral Therapy among Children in Fiche Hospital, North Shewa, Ethiopia, 2016. *J Pharm Care Heal Syst*. 2016(4:170. doi: 10.4172/2376-0419.1000170).
28. Biressaw S, Abegaz WE, Abebe M, Taye WA, Belay M. Adherence to Antiretroviral Therapy and associated factors among HIV infected children in Ethiopia : unannounced home-based pill count versus caregivers ' report. *BMC Pediatr*. 2013;13(<http://www.biomedcentral.com/14712431/13/132>):1471–2431.
29. Id GM, Antonucci R, Mukurasi A, Zepherine H. Adherence to antiretroviral treatment among children and adolescents in Tanzania : Comparison between pill count and viral load outcomes in a rural context of Mwanza region. *PLoS One*. (| <https://doi.org/10.1371/journal.pone.0214014> March):1–15.
30. Musiime V, Kayiwa J, Kiconco M, Tamale W, Alima H, Mugerwa H, et al. Response to Antiretroviral Therapy of HIV Type 1- Infected. *AIDS RESEARCH AND HUMAN RETROVIRUSES*. 2017;38(April):1–13.