

# Protective Effect of Phytogenic based Additives in Enterotoxigenic *Escherichia Coli* Challenged Piglets

**Valentina Caprarulo**

Università degli Studi di Milano: Università degli Studi di Milano

**Lauretta Turin**

Università degli Studi di Milano: Università degli Studi di Milano

**Monika Hejna**

Università degli Studi di Milano: Università degli Studi di Milano

**Serena Reggi**

Università degli Studi di Milano: Università degli Studi di Milano

**Matteo Dell'Anno** (✉ [matteo.dellanno@unimi.it](mailto:matteo.dellanno@unimi.it))

Università degli Studi di Milano: Università degli Studi di Milano <https://orcid.org/0000-0002-2171-864X>

**Pietro Riccaboni**

Università degli Studi di Milano: Università degli Studi di Milano

**Paolo Trevisi**

Alma Mater Studiorum Università di Bologna: Università di Bologna

**Diana Luise**

Alma Mater Studiorum Università di Bologna: Università di Bologna

**Antonella Baldi**

Università degli Studi di Milano: Università degli Studi di Milano

**Luciana Rossi**

Università degli Studi di Milano: Università degli Studi di Milano <https://orcid.org/0000-0003-1178-4683>

---

## Research Article

**Keywords:** phytogenic, feed additives, alternatives to antibiotics, short chain fatty acids, medium chain fatty acids, O138 E. coli, diarrhea, intestinal health, gut histopathology, intestinal gene expression.

**Posted Date:** February 16th, 2022

**DOI:** <https://doi.org/10.21203/rs.3.rs-1207181/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)



# Abstract

Post Weaning Diarrhea (PWD) is the most important multifactorial gastroenteric disease of the weaning in pig livestock. Phytogetic (PHY) natural extracts are largely studied as alternatives to antibiotic treatments in combating the global concern of the antimicrobial resistance. The aim of this study was to evaluate the protective effect of innovative phytogetic premix with or without short and medium chain fatty acids (SCFA and MCFA) in O138 *E. coli* challenged piglets. Twenty-seven weaned piglets were allotted into four groups receiving different diets according to the following experimental doses: PHY1 (n=7) fed basal diet supplemented with 0.2% of phytogetic premix, PHY2 (n=7) fed basal diet supplemented with 0.2% of phytogetic premix added with 2000 ppm of SCFA and MCFA. After 6 days of feeding experimental diets, animals were challenged (day 0) with  $2 \times 10^9$  CFU/dose of *E. coli* and CTRL group fed basal diet was divided at day 0 into positive (CTRL+; n=6) and negative control group (CTRL-; n=7). Body weights were recorded at -14, -6, 0, 4 and 7 days and the feed intake was recorded daily. *E. coli* shedding was monitored for 4 days post-challenge by plate counting. Fecal consistency was registered daily by a four-point scale (0-3; diarrhea >1) during the post-challenge period. Tissue samples were obtained for gene expression and histological evaluations at day 7 from four animals per group. Lower average feed intake was observed in CTRL+ compared to PHY2 and CTRL during the post-challenge period. Infected groups showed higher *E. coli* shedding compared to CTRL- during the 4 days post-challenge ( $p < 0.01$ ). PHY2 showed lower frequency of diarrhea compared to PHY1 and CTRL+ from 5 to 7 days post-challenge. No significant alterations among groups were observed in histopathological evaluation. Duodenum expression of occludin tended to be lower in challenged groups compared to CTRL- at 7 days post-challenge ( $p = 0.066$ ). In conclusion, dietary supplementation of PHY plus SCFA and MCFA revealed encouraging results for diarrhea prevention and growth performance in weaned piglets.

## 1 Introduction

Post weaning diarrhea (PWD) is a gastrointestinal multifactorial disease that generally occurs during the first two weeks after the weaning phase. It is one of the most economically-relevant diseases in swine husbandry due to the costs of treatments, reduced growth, and increased of mortality (Bonetti et al. 2021). Although many factors are involved in the development of this disease, PWD is often exacerbated by many enterotoxigenic *Escherichia coli* pathotypes characterized by the presence of virulence factors such as toxins and adhesive fimbriae (Sun and Kim 2017). Bacterial resistance to a wide range of commonly used antibiotics is a global concern and a recent increase in prevalence and severity of PWD required alternative measures for their control (Renzhammer et al. 2020; Dell'Anno et al. 2021a; Dell'Anno et al. 2021b). Reducing and possibly replacing antimicrobials is a crucial aim of European policies, even though it is still unclear how the use of antibiotics in farming can decrease the prevalence of resistant bacteria related to human medicine (Rossi et al. 2014a; Cormican et al. 2017; Tang et al. 2017).

The aim of nutrition is no longer simply to satisfy the nutritional requirements, but also play a key role in the health and welfare of humans and animals (Domínguez Díaz et al. 2020; Grossi et al. 2021). Functional feed additives, which sustain the health status and reduce the risk of pathologies, have thus

become fundamental in replacing or reducing antimicrobials in food-producing animals. The dietary inclusion of phytochemicals (PHYs), represented by plant secondary metabolites, are largely studied as alternative growth promoters because of their biological properties which include antimicrobial, antioxidant, and nutrigenomic effects on the development of immunity (Durmic and Blache 2012; Yang et al. 2015; Lillehoj et al. 2018; Reyes-Camacho et al. 2020). In particular Yan and Kim (Yan and Kim 2012) observed a significant reduction in fecal *E. coli* count after 1 g/kg of eugenol supplementation in pigs. A blend of oregano, anise, and citrus peel (40 mg/kg diet) supplementation to piglets' diet has been demonstrated to evolve anti-inflammatory effect by reducing the gene expression of NF- $\kappa$ B and TNF $\alpha$  (Upadhaya et al. 2016). The dietary supplementation of thymol, cinnamaldehyde and menthol have been reported to positively affect the feed digestibility in swine (Maenner et al. 2011; Li et al. 2012). The *in vivo* effects, resulting from the various biological activities of the PHYs, depend on their structure, dosage, and pharmaco-kinetics, as well as the animal species, productive phase and administration period. For this reason, several combinations of natural extracts are currently studied in order to promote their possible synergistic or complementary effect on animal health. Although PHYs show antimicrobial activity in the gastrointestinal tract against specific pathogens such as *Escherichia coli*, *Clostridium perfringens* and *Salmonella* spp. (Thacker 2013; Mohammadi Gheisar and Kim 2018), their effectiveness can vary due to the presence and the location of functional hydroxyl and phenolic terpenoids (Dubreuil 2013). Rational combinations of PHYs have been studied in order to increase the spectrum of beneficial activities. In addition, the synergistic or complementary effect of PHYs with other compounds leads to various beneficial activities of several compounds, especially organic acid (OA). Amongst feed additives with antimicrobial activities, organic acids, in particular short-chain fatty acids (SCFAs) and medium-chain fatty acids (MCFAs), have a strong antimicrobial activity and are key to modulating intestinal health and improving animal performance (Ferronato and Prandini 2020; Jackman et al. 2020). SCFAs and MCFAs regulate the growth and virulence of enteric pathogens, such as enterohemorrhagic *E. coli*, *Klebsiella* and *Salmonella* (Zhang et al. 2020). They damage the bacterial structure and in some cases separate the inner and outer membranes (Hanczakowska 2017) and thus increase the concentration of IgG and IgM in piglets challenged with enterotoxigenic *Escherichia coli* (ETEC) strains (Han et al. 2020). A synergistic antimicrobial effect has been observed in the combination of PHYs and organic acids *in vitro* (Costa et al. 2013). However, the effect of their dietary supplementation on pigs' growth and the optimization of the inclusion level for diarrhea prevention against major pathogens of weaned piglets has not been fully investigated.

The aim of this study was thus to evaluate the protective effect against O138 *E. coli* F18+ infection of an innovative phytochemical premix composed by caraway oil, lemon oil, clove, cinnamon, nutmeg, onion, pimento, orange peel, peppermint and chamomile powder with and without short and medium chain fatty acids.

## 2 Materials And Methods

### 2.1 Animal Selection Criteria

The trial was performed at the Experimental Animal Research and Application Centre of University of Milan and was authorized by the Italian Health Ministry (authorization n° 711/-PR) in accordance with EU regulations (Directive 2010/63/EU).

Animals enrolled in the experimental trial were selected from a conventional herd free from contagious diseases (Ex A-list International Office of Epizootic, porcine reproductive and respiratory syndrome, atrophic rhinitis, Aujeszky's disease, transmissible gastroenteritis, salmonellosis) and without a history of PWD or oedema disease. Sows were assessed for genetic susceptibility to *Escherichia coli* carrying F18 adhesive fimbriae (F18 *E. coli*) by screening the fucosyltransferase 1 (FUT1) genotypes using polymerase chain reaction (PCR) reaction according to Luise et al. (2019a, b). Briefly, genomic DNA was extracted from hair samples of sows and genotyped to identify polymorphic variants. Sows carrying the GG genotypes at FUT1 gene were considered for piglet enrolment. A further selection criterion was the absence of hemolytic *E. coli* in piglets feces. Microbiological analyses of selective mediums (Agar MacConkey) (Hayer et al. 2020; Li et al. 2020; Remfry et al. 2020) were thus carried out before transport and upon arrival on fecal samples collected from enrolled piglets.

## 2.2 Animals and Experimental Design

Twenty-seven weaned piglets ( $28 \pm 2$  days) balanced per weight ( $9.79 \pm 1.25$  kg) and sex, after a 7-day adaptation period, were randomly allotted in four experimental groups in randomized complete block design and fed *ad libitum* for the entire experimental period: phytogenic additive 1 (PHY1, n=7), phytogenic additive 2 (PHY2, n=7), positive control (CTRL+, n=6), and negative control (CTRL -, n=7). After the adaptation period, the basal diet was added by 200g/100kg phytogenic additive (PHY1) and by phytogenic plus 2000 ppm of short and medium chain fatty acids premix (PHY2). Control groups received basal diet supplemented with a premix carrier (95% wheat meal and 5% of coconut oil) to achieve the same nutrient concentrations. The iso-energetic and iso-proteic diets (Table S1) were formulated (Plurimix; Fabermatica, CR, Italy) according to animal requirements for the post weaning phase defined by the US National Research Council (National Research Council 2012). The phytogenic feed additive (FRESTA®F, Delacon Biotechnik GmbH), approved by EU regulation (Reg. CE 1831/2003), as zootechnical additive, was composed of essential oil from caraway oil (d-carvone 3.5-6.0 mg/g) and lemon (limonene: 2.3 - 9.0 mg/g), dried herbs and spices (1.5% clove powder, 10% cinnamon powder, 1.5% nutmeg powder, 5% onion powder, 2% pimento powder, 5% orange peel powder, 12.5% peppermint powder and 12.5% chamomile powder). The SCFA and MCFA premix was composed by butyric (C4), caprylic (C8), capric (C10) and lauric acid (C12). The phytogenic products (with or without SCFA and MCFA) or premix carrier were mixed with the basal diet for 30 minutes.

Piglets were housed in two environmentally controlled rooms, in individual pens, with a plastic slatted floor and constant temperature ( $27^\circ$  C) and humidity (60%) for the entire experimental period. The trial was divided into a pre- and post-challenge, considering the challenge as day 0 (Figure 1).

PHY1: treatment group fed basal diet supplemented with 200g/100kg of phytogenic additive; PHY2: treatment group fed basal diet supplemented with 200g/100kg of phytogenic additive supplemented with

2000 ppm of short and medium chain fatty acids premix; phytogetic additive composed by caraway oil, lemon oil, clove, cinnamon, nutmeg, onion, pimento, orange peel, peppermint and chamomile powder; short chain and medium chain fatty acids blend included butyric (C4), caprylic (C8), capric (C10) and lauric acid (C12) at dose levels of 2000 ppm. CTRL: group fed basal diet supplemented with premix carrier divided into negative control (CTRL-) and positive control (CTRL+) challenged at day 0. Diets were analyzed for proximate analysis, including moisture, crude protein (CP), crude fibre (CF), ether extract (EE), and ash. The moisture determination was performed by oven-drying at 65°C for 24 h (Regulament EC 152/2009). Crude protein content was measured according to the Kjeldahl method (Association of Official Analytical Chemists method 2001.11). Crude fiber was determined by the filter bags technique (American Oil Chemistry Society 2009). Ether extract content was determined in a Soxhlet system after hydrolysis (Association of Official Analytical Chemists method 2003.05). Ash was measured using a muffle furnace at 550°C (Association of Official Analytical Chemists method 942.05).

## 2.3 Experimental Challenge

*E. coli* challenger strain was genetically characterized by polymerase chain reaction (PCR) (Applied Biosystem 7500) in order to detect the presence of the two important virulence profile: subunit B of verocytotoxin type 2 and F18 adhesive fimbriae (Table 1).

Twenty piglets, except for piglets in CTRL- group (n = 7), on day 0 (challenge day) were orally infected with O138 *Escherichia coli* F18+ strain obtained from a permanent collection of the University of Milan and previously characterized (Rossi et al. 2014b; Dell'Anno et al. 2020; Rossi et al. 2021).

Sixty minutes before the challenge, the piglets were sedated with azaperone (Stresnil TM, Janssen Cilag Spa, 2 mL/pig), thereafter 30 mL of a 10% bicarbonate solution was orally administered to neutralize gastric acid and to increase the survival rate of the challenger strain in the stomach. After 10-15 min, the inoculum was given orally in a single dose of 5 mL of bacterial medium with  $2 \times 10^9$  colony-forming unit (CFU) of challenger strain, using a 16G catheter (Rossi et al. 2021). Animals were fasted 3h before and 3h after the challenge. At the same time, piglets in CTRL- were orally inoculated with 5 mL of Luria Bertani (LB) medium to balance the level of stress associated with the oral challenge.

Table 1

PCR conditions and oligonucleotide sequences of F18 adhesive fimbriae and VTe2 (B-subunit) encoding genes.

Gene	Accession number (GenBank)	Size (pb)	Primer sequence (5' to 3')	PCR conditions
F18 adhesive fimbriae	AJ308332.1	519	5'GATCCATGAAAAGACTAGTGTTTATTTCTTTTG 3'CGAATGCGCCAATGAATGTTTCATTCTCGAG	Den.95°C 1' ann.56°C 1'20" ext.72°C 1'30"  35 cycles
VTe2 (B-subunit)	GU459254.1	270	5'GGATCCATGAAGAAGATGTTTATAGCGG 3'AACGGGTCCACTTCAAATGATTCTCGAG	Den.95°C 1' ann.50°C 1'20" ext.72°C 1'30"  35 cycles

## 2.4 Zootechnical Performance, Clinical and Fecal Score

Average daily feed intake (ADFI) was recorded daily from day -6 to day 7 by measuring the refusals. Body weight (BW) was recorded on day -6 (first day of experimental diets), day 0 (challenge day), day 4 and day 7 (sacrifice day). Average daily gain (ADG) and feed efficiency were also calculated.

Piglets were individually evaluated throughout the trial by clinical examination, including observation of behavioral disturbances. In particular, oedema, epiphora, respiratory and hair scores were evaluated through three-point scales (oedema score: 0=normal, 1=mild, 2=severe; epiphora score: 0=normal, 1=mild, 2=severe; respiratory score: 0=normal, 1=slightly quick, 2=quick; hair/bristles score: 0=smooth, 1=lightly brushy, 2=highly brushy) (Rossi et al. 2021). In addition, cyanosis, a blue or red discoloration of the skin, which may or may not be localized to small areas, was considered not as a specific skin condition but as a symptom of disease. From day -6 to day 7, all piglets were evaluated for the fecal score. Clinical signs of the disease were identified according to the point scale score described by Rossi et al. (Rossi et al. 2014b). A four-point scale was adopted to score fecal consistency: 0=normal, 1=soft consistency, 2=mild diarrhea, 3=severe diarrhea; considering >1 as an indicative of diarrhea. Fecal color was evaluated using a three-point scale: 1 = yellow, 2 = green; 3 = brown.

## 2.5 Microbiological Evaluation of Fecal Samples

Individual fecal samples were collected from rectal ampulla from each piglet, on days -1, 1, 2, 3 and 4 to perform microbiological analysis and evaluate the challenger strain shedding. For each sample, 1 g of feces was homogenized with 1 ml of saline solution and incubated overnight at 37°C on sheep blood agar plates 5% (Blood Agar Base No. 2-Oxoid) in order to examine the presence of hemolytic colonies.

The total hemolytic bacteria count was performed by counting the number of colonies cultured from serial dilutions of each fecal sample in order to evaluate the presence of hemolytic *E. coli* in relation to the total bacteria population.

## 2.6 Necropsy, Intestinal Samples, and Histopathology

At day 7 post-challenge, sixteen animals (n=4/treatment) were sacrificed, tissue samples were collected for histopathological and molecular analyses of intestinal tissues. The intestine of each animal was weighed, and intestinal samples of ileum were harvested. For the histological evaluation, intestinal samples were fixed in 10% neutral formalin buffer and stored at 4°C. Tissues were rinsed with sterile saline solution and transferred into 2 mL cryotubes, snap-frozen in liquid nitrogen and stored at -80°C until further analysis.

Histological examinations of collected intestinal samples for each piglet were carried out. The fixed samples fixed were embedded in paraffin, and 5µm thick histological sections were performed with a microtome. Cross sections were stained with hematoxylin and eosin and were blind evaluated by light microscopy. A four-point scale was adopted for inflammatory infiltrates, epithelial regeneration, fusion of villi, oedema, hyperemia, necrosis of mucosa, T atrophy, stroma, and follicular hyperplasia; considering: 0=no evidence; 1=slight presence; 2=moderate; 3=severe. Samples of duodenum were collected and frozen in liquid nitrogen for gene expression analysis.

## 2.7 Duodenum Gene Expression

Total RNA was extracted from the duodenum using FastGene Scriptase Basic (Nippon genetics) according to the manufacturer's instructions. The integrity of total RNA was assessed by gel electrophoresis to detect the 18S and 28S rRNA bands. A combination of oligo-dT and random primers was used to reverse transcribe 100 ng of total duodenal RNA to cDNA (cDNA synthesis kit, FastGene Scriptase Basic, Nippon Genetics). Primer pairs were first tested for their specificity in qualitative PCR, using the pooled cDNA as a template. The cycling profile for the assay consisted of initial denaturation of RNA (65°C x 5'), then the annealing of random primers (25°C x10'), followed by the annealing of oligo-dT and transcription (42°C x 60'). At the end of the cycle, the enzyme deactivation (90°C x 5') was performed. The abundance of cytochrome c oxidase subunit I (COX1), cytochrome c oxidase subunit II (COX2), interleukin 10 (IL-10), interleukin 6 (IL-6), lysyl oxidase (LOX), glutathione peroxidase 2 (GPX2), NAD (P) H quinone dehydrogenase 1 (NQO1) claudin domain containing 1 (CLDND1) and occludin (OCLN) (Table 2) mRNA was determined using SYBR Green-based real-time quantitative PCR assays (7500 Fast Dx, Applied Biosystems). Only reaction efficiencies that were near to 100% were considered for further analysis. The mean values for the transcripts were normalized to the arithmetic mean of mRNA abundance of  $\beta$ actin as the reference gene within each sample. The comparative CT method was used to determine fold changes in gene expression, calculated as  $2^{-\Delta\Delta CT}$ . The final results were presented as the fold changes of target gene expression in a target sample relative to a reference sample, normalized to  $\beta$ actin rRNA (Livak and Schmittgen, 2001). The  $\beta$ actin rRNA was used to calculate the threshold cycles, since it previously showed constant values under all the conditions adopted.

Table 2  
Primer sequences and relative amplicon dimensions.

Gene <sup>1</sup>	Accession number (GenBank)	Size (pb)	Primer sequence (5' to 3')
βactin F	DQ845171	76 bp	CTACGTCGCCCTGGACTTC
βactin R	DQ845172		GCAGCTCGTAGCTCTTCTCC
IL-6 F	JQ839263	112 bp	TGGGTTCAATCAGGAGACCT
IL-6 R	JQ839264		CAGCCTCGACATTTCCCTTA
IL-10 F	L2001	105 bp	TGAAGAGTGCCTTTAGCAAGCTC
IL-10 R	L2002		CTCATCTTCATCGTCATGTAGGC
COX1 F	EF568726	102 bp	GGAGCGGGTACTGGATGAAC
COX1 R	EF568726		CACCTGCAAGGGTGTAGGGAGL
COX2 F	AF304201	141 bp	AAGACGCCACTTCACCCATC
COX2 R	AF304201		TCCATTGTGCTAGTGTGTGTCA
GPx2 F	DQ898282	103 bp	GGAGATCCTGAACAGCCTCA
GPx2 R	DQ898282		GCGAAGACAGGATGCTCATT
LOX F	NM_001164001	112 bp	GTGGAGCACGAAAGCAAGACCC
LOX R	NM_001164001		AAGGTGGGGTATGCATCGACAC
NQ01 F	NM_001159613	118 bp	ATCACAGGTAAGTGAAGGACCC
NQ01 R	NM_001159613		GCGGCTTCCACCTTCTTTTG
CLAUDIN1 F	NM_001244539	90 bp	TCTTTCTTATTTTCAAGTCTGGCT
CLAUDIN1 R	NM_001244539		ACTGGGGTCATGGGGTCATA
OCCLUDIN F	NM_001163647	106 bp	GTCCACCTCCTTATAGGCCTGATG
OCCLUDIN R	NM_001163647		CGCTGGCTGAGAAAGCATTGG

<sup>1</sup>CTB: actin beta; IL-6: interleukin-6; IL-10: interleukin 10; COX1: cytochrome c oxidase subunit I; COX2; cytochrome c oxidase subunit II; LOX: lysyl oxidase; GPX2: glutathione peroxidase 2; NQ01: NAD (P) H quinone dehydrogenase 1; CLDND1: claudin domain containing 1; OCLN: occludin.

## 2.8 Blood Samples, Serum Metabolite Profile and Serum Acute Phase Proteins

Blood was collected from the jugular vein of each animal on day -1, day 3 and day 7 through vacutainer tubes containing ethylenediaminetetraacetic acid (EDTA) and heparin as anticoagulants. Plasma was collected after centrifugation (3000 rpm, 10 min, 4°C), aliquoted and stored at -20°C for further analysis.

Hematocrit was evaluated on whole blood using the microhematocrit method. The concentration of total protein (g/L), albumin (g/L), globulin (g/L), albumin/globulin (A/G ratio), alanine aminotransferase (ALT-GPT; IU/L), aspartate aminotransferase (AST-GOT; IU/L), phosphatase alkaline (ALP; IU/L), glucose (mmol/L), urea (mmol/L), creatinine ( $\mu\text{mol/L}$ ), total bilirubin ( $\mu\text{mol/l}$ ), total cholesterol (mmol/L), triglycerides (mmol/L), high-density lipoprotein (HDL; mmol/L), low-density lipoproteins (LDL; mmol/L), calcium (mmol/L), phosphorus (mmol/L) and magnesium (mmol/L) were analyzed via standard enzymatic colorimetric analysis through a multiparametric autoanalyzer for clinical chemistry (ILab 650; Instrumentation Laboratory Company, Lexington, MA, USA) at 37°C by the Lombardy and Emilia Romagna Experimental Zootechnic Institute (IZSLER). Porcine C-reactive protein (CRP) concentration was determined with a commercial sandwich immunoassay Kit (Mybiosource, San Diego, CA, USA) following the manufacturer's instructions. The results were read at 450 nm using a microplate reader (Model 680, Bio-Rad Laboratories, CA, USA). Haptoglobin (HP) concentrations were measured through a colorimetric kit (Phase<sup>TM</sup> Range porcine Haptoglobin Assay; Tridelta Development Ltd) according to the manufacturer's instructions. The results were read at 630 nm on a microplate reader (Model 680, Bio-Rad Laboratories, CA, USA).

## 2.9 Statistical Analysis

Zootechnical performance and fecal microbiological analysis were analyzed using a linear model after testing the normality of data through Shapiro-Wilk test using JMP Pro 15® (SAS Inst. Inc., Cary, NC, USA). The model included the fixed effect of treatments (Trt), the effect of time (Time), and the interaction between treatment and time (Trt x Time).

Serum metabolites were evaluated performing analysis of covariance (ANCOVA) to adjust the initial variability of the pre-challenge period after testing the normality of data through Shapiro-Wilk test using JMP Pro 15® (SAS Inst. Inc., Cary, NC, USA).

Clinical score data were converted into a dichotomous variable (normal/pathological), and observed frequencies were assessed using the Chi-squared Test. Histological scores, intestinal weight and relative gene expression were analyzed using Kruskal-Wallis test (PROC NPAR1WAY of SAS 9.4 software) for non-parametric data due to the small sample size of euthanized animals at day 7. Multiple comparisons for parametric statistics were evaluated with the Tukey's Honestly Significant Difference test (Tukey's HSD) or Tukey-Kramer test and Steel-Dwass test was used for non-parametric multiple comparisons. The results were presented as least square means (LSMEANS)  $\pm$  standard error (SE) for parametric data and as medians and range (minimum-maximum) for non-parametric results. Means or medians were considered statistically different when  $p \leq 0.050$  and statistical tendency was considered when  $p < 0.100$ .

## 3 Results

### 3.1 Chemical Composition of the Experimental Diets

Proximate analysis of the experimental diets showed comparable contents of the principal nutrients. The inclusion of phytogenic based additives with or without MCFA and SCFA did not affect the nutrient balance of feed (Table S1).

## **3.2 Zootechnical Performance**

During the pre-challenge period, no statistically significant differences among experimental groups were observed. Considering the entire post-challenge period, ADFI of CTRL+ was lower than PHY2 and CTRL- ( $p < 0.005$ ; Table 3).

Table 3  
Zootechnical performance of experimental groups during the post-challenge period.

	PHY1	PHY2	CTRL+	CTRL-	<i>p-value</i>		
	(n=7)	(n=7)	(n=6)	(n=7)	Trt	Time	Trt×Time
<b>BW, kg</b>							
d 0	10.39±0.68	10.46±0.68	10.23±0.73	10.67±0.68	0.342	<0.001	0.963
d 4	11.29±0.68	11.73±0.68	11.25±0.73	11.89±0.68			
d 7	12.33±0.68	12.96±0.68	11.55±0.73	13.42±0.68			
<b>ADG, kg/d</b>							
d 1-4	0.22±0.09	0.32±0.09	0.26±0.10	0.35±0.09	0.083	0.323	0.262
d 5-7	0.35±0.09	0.41±0.09	0.10±0.10	0.51±0.09			
d 1-7	0.29±0.06	0.36±0.06	0.18±0.07	0.41±0.06			
<b>ADFI, kg/d</b>							
d 1-4	0.42±0.04	0.46±0.04	0.35±0.05	0.44±0.04	<0.005	0.040	0.294
d 5-7	0.42±0.04	0.56±0.04	0.37±0.05	0.58±0.04			
d 1-7	0.42±0.03 <sup>ab</sup>	0.51±0.03 <sup>b</sup>	0.36±0.03 <sup>a</sup>	0.51±0.03 <sup>b</sup>			
<b>FCR, kg/kg</b>							
d 1-4	1.84±0.41	1.59±0.38	1.60±0.41	1.41±0.41	0.054	0.248	0.069
d 5-7	1.57±0.04	1.40±0.38	3.60±0.58	1.32±0.38			
d 1-7	1.70±0.31	1.49±0.27	2.60±0.36	1.36±0.28			
Data are presented as least squared means (LSMEANS) and standard errors (SE).							
<sup>a-b</sup> Different lowercase letters indicate statistically significant differences between treatment groups.							
BW: body weight, ADFI: average daily feed intake, ADG: average daily gain, FCR: feed conversion ratio, PHY1: basal diet plus 200g/100kg of phytogenic additive; PHY2: basal diet plus 200g/100kg of phytogenic additive plus 2000 ppm of short and medium chain fatty acids premix; phytogenic additive composed by caraway oil, lemon oil, clove, cinnamon, nutmeg, onion, pimento, orange peel, peppermint and chamomile powder; CTRL: basal diet plus premix carrier divided per negative control (CTRL-) and positive control (CTRL+) challenged at day 0.							

### 3.3 Influence of Phytogenic Treatments on Clinical Score, Fecal Consistency and Color

During the pre-challenge period and at day 0, the piglets did not show significant differences among clinical scores, indicating a general good health status. After experimental infection, the frequencies of clinical score (considered as altered clinical conditions with a score of  $\geq 1$ ) revealed that the experimental procedures influenced the clinical status of the piglets (Table 4). However, from 5 to 7 days post challenge, a non-normal hair score frequency tended to increase in CTRL+ compared to the other experimental groups (9.52% for PHY1, 14.81% for PHY2, 33.33% for CTRL+ and 14.29% for CTRL-;  $p=0.071$ ).

Although statistical differences among treatments were not identified, several altered scores were registered from 1 to 4 days post challenge. Fecal score and color were the most informative indicators during the post-challenge period (Table 5). Significant higher frequencies of altered fecal color were recorded in challenged groups compared to CTRL- from 1 to 4 days post-challenge (57.14% PHY1, 37.96% PHY2, 39.29% CTRL+, 17.86% CTRL-;  $p<0.050$ ). Significant differences in the manifestations of diarrhea (fecal consistency  $\geq 2$ ) were observed from 5 to 7 days after the challenge. In particular, PHY1 had higher number of diarrhea cases compared to PHY2, CTRL+ and CTRL-, and PHY2 had a lower incidence compared to CTRL+ and PHY1 (80.95% PHY1, 28.57% PHY2, 61.11% CTRL+, 38.10% CTRL-;  $p<0.010$ ).

Table 4

Frequencies (expressed as percentages) of clinical score  $\geq 1$  from 1 to 7 days post-challenge.

<b>Treatments</b>					
<b>Days 1-4</b>	<b>PHY1 (n=7)</b>	<b>PHY2 (n=7)</b>	<b>CTRL+ (n=6)</b>	<b>CTRL- (n=7)</b>	<b>p-value</b>
Hair	28.57	10.71	16.67	14.29	0.325
Respiratory	3.57	0.00	4.17	0.00	0.528
Oedema	3.57	3.70	8.33	0.00	0.471
Epiphora	3.57	14.29	8.33	7.14	0.536
<b>Days 5-7</b>	<b>PHY1 (n=7)</b>	<b>PHY2 (n=7)</b>	<b>CTRL+ (n=6)</b>	<b>CTRL- (n=7)</b>	
Hair	9.52	14.81	33.33	14.29	0.071
Respiratory	0.00	0.00	0.00	0.00	-
Oedema	0.00	0.00	0.00	0.00	-
Epiphora	0.00	0.00	0.00	0.00	-
Data are presented as a percentage of clinical score $\geq 1$ registered from day 1 to day 7 post-challenge.					
PHY1: basal diet plus 200g/100kg of phytogenic additive; PHY2: basal diet plus 200g/100kg of phytogenic additive plus 2000 ppm of short and medium chain fatty acids premix; phytogenic additive composed by caraway oil, lemon oil, clove, cinnamon, nutmeg, onion, pimento, orange peel, peppermint and chamomile powder; CTRL: basal diet plus premix carrier divided per negative control (CTRL-) and positive control (CTRL+) challenged at day 0.					

Table 5

Frequencies (expressed as percentages) of fecal consistency  $\geq 2$  and fecal color =1 registered 1 to 7 days post-challenge.

Treatments					
Days 1-4	PHY1 (n=7)	PHY2 (n=7)	CTRL+ (n=6)	CTRL- (n=7)	p-value
Fecal consistency	42.86	28.57	25.00	21.43	0.319
Fecal color	57.14 <sup>a</sup>	39.29 <sup>a</sup>	37.50 <sup>a</sup>	17.86 <sup>b</sup>	0.027
Days 5-7	PHY1 (n=7)	PHY2 (n=7)	CTRL+ (n=6)	CTRL- (n=7)	
Fecal consistency	80.95 <sup>a</sup>	28.57 <sup>b</sup>	61.11 <sup>c</sup>	38.10 <sup>b</sup>	0.003
Fecal color	90.48	71.43	94.44	71.43	0.114
Data are presented as a percentage of fecal consistency $\geq 2$ and fecal color =1 registered from day 1 to day 7 post-challenge.					
<sup>a-b-c</sup> Different lowercase letters indicate statistically significant differences among treatment groups.					
PHY1: basal diet plus 200g/100kg of phytogenic additive; PHY2: basal diet plus 200g/100kg of phytogenic additive plus 2000 ppm of short and medium chain fatty acids premix; phytogenic additive composed by caraway oil, lemon oil, clove, cinnamon, nutmeg, onion, pimento, orange peel, peppermint and chamomile powder; CTRL: basal diet plus premix carrier divided per negative control (CTRL-) and positive control (CTRL+) challenged at day 0.					

### 3.4 Microbiological Evaluation of Feces and Challenger Strain Shedding

Weaned piglets did not show the presence of challenger *E. coli* in feces during the adaptation period and on day 0. Total bacterial count did not show statistically significant differences among groups from 1 to 4 days after the challenge ( $8.37 \pm 0.47 \log_{10}$  CFU/g for PHY1,  $8.04 \pm 0.47 \log_{10}$  CFU/g for PHY2,  $7.73 \pm 0.51 \log_{10}$  CFU/g for CTRL+ and  $7.71 \pm 0.47 \log_{10}$  CFU/g for CTRL-). Also after the challenge, all the experimental groups (except for negative control, CTRL-) registered fecal shedding of challenger *E. coli* strain (Figure 2). Statistically significant increased fecal shedding of hemolytic *E. coli* was observed in challenged groups compared to CTRL- from day 1 to day 4 post-challenge ( $4.09 \pm 0.01 \log_{10}$  CFU/g for PHY1,  $5.25 \pm 1.10 \log_{10}$  CFU/g for PHY2,  $5.95 \pm 1.09 \log_{10}$  CFU/g for CTRL+ and  $0.00 \pm 1.01 \log_{10}$  CFU/g for CTRL-;  $p < 0.001$ ).

Data are presented as least squared means (LSMEANS) and standard errors (SE).

PHY1: basal diet plus 200g/100kg of phytogenic additive; PHY2: basal diet plus 200g/100kg of phytogenic additive plus 2000 ppm of short and medium chain fatty acids premix; phytogenic additive

composed by caraway oil, lemon oil, clove, cinnamon, nutmeg, onion, pimento, orange peel, peppermint and chamomile powder; CTRL: basal diet plus premix carrier divided per negative control (CTRL-) and positive control (CTRL+) challenged at day 0.

### **3.5 Histological Evaluation and Gene Expression**

Samples were examined for the presence of inflammation both in villi and in lamina propria, epithelial regeneration, fusion of villi, oedema in deep lamina propria, T atrophy, stroma (fibroconnective and histiocytes), and follicular hyperplasia.

Intestinal weight results did not reveal significant differences between treatment groups after 7 days post-challenge (Table S4). Phytogetic dietary treatments did not significantly affect ileum inflammatory infiltrates, epithelial regeneration, oedema and hyperemia after 7 days (Table 6).

Table 6

Histological examination of ileum and lymphoid of weaned piglets fed experimental diets on day 7.

Treatments									
Score	PHY1 (n=4)		PHY2 (n=4)		CTRL+ (n=3)		CTRL- (n=4)		p-value
	median	min-max	median	min-max	median	min-max	median	min-max	
Ileum									
Inflammatory infiltrates	1	0-2	2	2-3	2	1-2	1	1-2	0.068
Epithelial regeneration	0	0-0	0	0-0	0	0-0	0	0-0	1.000
Fusion of villi	1	0-3	3	2-3	2	1-3	1	0-2	0.223
Oedema	0	0-1	0	0-1	0	0-1	0	0-1	1.000
Hyperemia	1	0-1	1	1-2	0	0-2	1	0-1	0.382
Necrosis of mucosa	0	0-0	0	0-1	0	0-0	0	0-0	1.000
Lymphoid									
T atrophy	0	0-1	1	0-1	0	0-1	0	0-0	0.253
Stroma	0	0-1	1	1-2	2	0-3	0	0-1	0.073
Follicular hyperplasia	0	0-2	2	1-3	1	0-3	0	0-1	0.186
Data are presented as medians and minimum and maximum value (min-max).									
PHY1: basal diet plus 200g/100kg of phytogenic additive; PHY2: basal diet plus 200g/100kg of phytogenic additive plus 2000 ppm of short and medium chain fatty acids premix; phytogenic additive composed by caraway oil, lemon oil, clove, cinnamon, nutmeg, onion, pimento, orange peel, peppermint and chamomile powder; CTRL: basal diet plus premix carrier divided per negative control (CTRL-) and positive control (CTRL+) challenged at day 0.									

Relative expressions of IL-10, IL-6, LOX, GPX2, NQO1 and CLDND1 were not affected by phytogenic dietary treatments (Table 7). The relative expression of occludin was downregulated at day 7 post-challenge ( $p < 0.012$ ). Pairwise comparisons revealed only a tendency to increase in challenged groups compared to CTRL- ( $p = 0.066$ ).

Table 7

Duodenum expression of the main genes related to the intestinal integrity, inflammation and health of weaned piglets fed experimental diets on day 7 post-challenge.

Relative expression <sup>1</sup>	Treatment								p-value
	PHY1 (n=4)		PHY2 (n=4)		CTRL+ (n=4)		CTRL- (n=4)		
	median	min-max	median	min-max	median	min-max	median	min-max	
IL-6	0.32	0.22-0.43	0.48	0.08-0.87	0.34	0.23-1.10	1.00	1.00-1.00	0.139
IL-10	0.26	0.14-0.61	0.94	0.11-1.71	0.24	0.09-0.83	1.00	1.00-1.00	0.110
COX1	0.74	0.58-8.15	4.24	0.51-15.31	3.36	0.56-11.36	1.00	1.00-1.00	0.671
COX2	0.46	0.24-1.77	1.57	0.10-4.94	1.34	0.14-2.36	1.00	1.00-1.00	0.734
LOX	0.61	0.25-1.37	2.33	0.21-3.96	1.19	0.40-1.39	1.00	1.00-1.00	0.331
GPX2	1.04	0.38-1.93	2.98	0.31-3.53	2.11	0.25-6.33	1.00	1.00-1.00	0.426
NQ01	0.68	0.27-5.73	6.21	0.28-8.62	5.61	0.46-8.21	1.00	1.00-1.00	0.315
CLDND1	1.37	0.31-6.33	14.40	0.39-22.20	6.05	0.69-22.67	1.00	1.00-1.00	0.619
OCLN	0.42	0.02-0.96	0.34	0.09-0.72	0.55	0.14-0.96	1.00	1.00-1.00	0.012

<sup>1</sup> Relative expressions of selected genes are presented as  $2^{-\Delta\Delta CT}$ .

Data are presented as medians and minimum and maximum value (min-max).

IL-6: interleukin-6; IL-10: interleukin 10; COX1: cytochrome c oxidase subunit I; COX2; cytochrome c oxidase subunit II; LOX: lysyl oxidase; GPX2: glutathione peroxidase 2; NQ01: NAD (P) H quinone dehydrogenase 1; CLDND1: claudin domain containing 1; OCLN: occludin.

PHY1: basal diet plus 200g/100kg of phytogetic additive; PHY2: basal diet plus 200g/100kg of phytogetic additive plus 2000 ppm of short and medium chain fatty acids premix; phytogetic additive composed by caraway oil, lemon oil, clove, cinnamon, nutmeg, onion, pimento, orange peel, peppermint and chamomile powder; CTRL: basal diet plus premix carrier divided per negative control (CTRL-) and positive control (CTRL+) challenged at day 0.

## 3.6 Influence of Phytogetic Treatments on Hematological and Serum Metabolites

The serum metabolic profile did not show statistically significant differences between the experimental groups at day 3 after the challenge (Table S2). After 7 days post-challenge, a significantly higher level of total protein content was observed in CTRL+ compared to CTRL- ( $62.32 \pm 2.89$  and  $50.83 \pm 2.67$  mg/mL, respectively;  $p=0.050$ ) (Table 8). Globulin content tended to be higher in CTRL+ than CTRL- at 7 days post-challenge ( $35.31 \pm 2.86$  and  $24.43 \pm 2.61$  g/L respectively,  $p=0.055$ ). PHY2 had a higher level of AST-GOT at day 7 compared to the other experimental groups ( $39.11 \pm 6.79$  IU/L for PHY1,  $72.02 \pm 7.42$  IU/L for PHY2,  $37.82 \pm 7.05$  IU/L for CTRL+,  $43.59 \pm 6.74$  IU/L for CTRL-;  $p<0.050$ ). Acute phase proteins were not affected by dietary treatments and experimental challenge and showed no statistically significant differences after 3- and 7-days post-challenge (Table S3).

Table 8  
Serum metabolites of weaned piglets fed experimental diets on day 7 post-challenge.

Blood	Treatments				p-value
	PHY1 (n=7)	PHY2 (n=7)	CTRL+ (n=6)	CRTL- (n=7)	
Total protein, mg/ml	56.95±2.67 <sup>ab</sup>	54.04±2.67 <sup>ab</sup>	62.32±2.89 <sup>a</sup>	50.83±2.67 <sup>b</sup>	0.050
Hematocrit, %	26.03±1.05	26.09±1.05	25.00±1.16	25.43±1.05	0.880
Albumin, g/L	30.76±2.85	32.05±2.72	28.70±3.06	26.11±2.71	0.452
Globulin, g/L	28.31±2.63	26.20±2.61	35.31±2.86	24.43±2.61	0.055
A/G ratio	1.06±0.09	1.09±0.09	0.89±0.09	1.14±0.09	0.280
Urea, mmol/L	2.51±0.30	2.09±0.30	2.90±0.33	1.75±0.30	0.093
ALT-GPT, IU/L	26.51±2.51	31.09±2.54	24.52±2.81	26.66±2.54	0.375
AST-GOT, IU/L	39.11±6.79 <sup>a</sup>	72.02±7.42 <sup>b</sup>	37.82±7.05 <sup>a</sup>	43.59±6.74 <sup>ab</sup>	0.014
ALP, UI/L	170.30±17.78	197.15±17.71	149.83±19.35	195.84±18.32	0.262
Total bilirubin, µmol/l	2.25±0.16	1.86±0.16	1.86±0.17	1.70±0.16	0.123
Glucose, mmol/L	4.91±0.28	5.52±0.28	5.12±0.31	4.91±0.28	0.421
Total cholesterol, mmol/L	2.17±0.11	2.21±0.11	2.20±0.12	2.19±0.11	0.992
Calcium, mmol/L	2.55±0.15	2.89±0.15	2.49±0.17	2.54±0.15	0.243
Phosphorus, mmol/L	2.78±0.11	3.00±0.11	2.87±0.12	3.04±0.13	0.426
Magnesium, mmol/L	0.85±0.04	0.91±0.04	0.96±0.04	0.87±0.04	0.174
Data are presented as least squared means (LSMEANS) and standard errors (SE).					
A/G: albumin/globulin; ALT-GPT: alanine aminotransferase; AST-GOT: aspartate aminotransferase; ALP: alkaline phosphatase; HDL: high-density lipoprotein; LDL: low density lipoprotein; PHY1: basal diet plus 200g/100kg of phytogenic additive; PHY2: basal diet plus 200g/100kg of phytogenic additive plus 2000 ppm of short and medium chain fatty acids premix; phytogenic additive composed by caraway oil, lemon oil, clove, cinnamon, nutmeg, onion, pimento, orange peel, peppermint and chamomile powder; CTRL: basal diet plus premix carrier divided per negative control (CTRL-) and positive control (CTRL+) challenged at day 0.					

## 4 Discussion

Weaning is a critical period where piglets need to adapt to a new diet, environment and to develop their own immunity (Tretola et al. 2019). During this phase, PWD is one of the major causes of gastrointestinal disorders leading to high morbidity, antibiotic use and economic losses. Several natural extracts have

been investigated for their functional proprieties to decrease diarrhea occurrence in piglets, with discordant results. The general aim of this study was to evaluate the protective effect of innovative phytogetic premix with or without MCFA and SCFA against O138 *E. coli* in weaned piglets. Genetic characterization of the sows led to the enrollment of piglets that were potentially susceptible to F18 fimbriae. In fact, the presence of F18 receptor (F18R) on porcine intestinal epithelium is crucial for the development of *E. coli* infections.

During the pre-challenge period (day -6 to day 0), the piglets showed comparable growth performance, demonstrating that the supplementation of additives did not influence their growth and feed consumption or feed palatability. Even if the effect on zootechnical performance was limited by the short experimental period (European Food Safety Authority 2018), ADFI was affected by the treatment.

However, ADFI of CTRL+ was significantly lower compared to PHY2 and CTRL- groups. The observed increase in feed intake of PHY2 compared to CTRL+ suggests that phytogetic premix plus SCFA and MCFA positively affected feed palatability. In addition, higher dietary intake is often related to a better health status (Czech et al. 2021), indicating that the treatment with PHY and organic acids could have supported animals' health resulting in increased feed intake during the entire post-challenge period. PHY2 group showed a similar performance to CTRL- (uninfected), suggesting that dietary supplementation with the phytogetic premix, MCFA and SCFA was very effective in dealing with O138 *E. coli* infection, thus supporting intestinal health of animals. The addition of MCFA and SCFA may enhance animal growth by several mechanisms as previous studies described (e.g. inhibitory activity, mucosal epithelium integrity support) (Royce et al. 2013; Ferrara et al. 2017; Diao et al. 2019). In addition, phytogetic feed additives derived from spices and herbs are commonly used in animal nutrition as an alternative to in-feed antibiotics due to their antibacterial, antiviral and antioxidant properties. These effects are generally due to the presence of different bioactive compounds such as alkaloids, flavonoids, glycosides, mucilage, saponins, tannins, phenolics, polyphenols, terpenoids, and polypeptides (Upadhaya et al. 2016; Nowak et al. 2017; Caprarulo et al. 2020a; Dell'Anno et al. 2020; Reggi et al. 2020). Our results are in line with other studies demonstrating the antibacterial activity of PHYs, MCFA and SCFA on a wide range of pathogens (Dibner and Buttin 2002; Salsali et al. 2008).

In terms of clinical examination, from day 1, clinical scores were affected by experimental infection, confirming that disease development impaired the clinical status of challenged animals compared to the pre-challenge period. Moreover, significant differences in pathological hair, respiratory, oedema and epiphora scores were not detected in infected groups. This was probably due to the individual variability and the small sample size that could prevented to observe differences among groups. The O138 *Escherichia coli* challenger strain can impair gut health due to its capacity to adhere to the intestinal epithelium by specific fimbriae which could be followed by verocytotoxin production (Rossi et al. 2012; Rossi et al. 2013) and in consequence may show systemic symptoms.

A slightly different situation was found during the evaluation of the fecal score and incidence of diarrhea. Experimental challenge affected transitory the fecal color and consistency during the 7 days post-

challenge. Firstly, from day 1 to day 4 post-challenge, feces of yellowish color were registered more frequently in challenged group compared to CTRL- typically related to gastrointestinal disorders (Rossi et al. 2012). From day 5 to day 7 the highest diarrhea frequencies were registered in all infected animals, suggesting a late effect of challenge on fecal consistency compared to fecal color. These data are confirmed by a previous study by Rossi et al. (Rossi et al. 2021) showing that O138 *E. coli* experimental infection increased the sum of fecal score from 3 to 9 days post-challenge. Particularly, the highest diarrhea occurrence was observed in PHY1 compared to other groups, while PHY2 showed a fecal consistency comparable to CTRL- suggesting the counteracting activity of the phytogenic additives, SCFA and MCFA against experimental infection. Even if antibacterial activity of phytogenic additives was reported (Namkung et al. 2004), the observed effect on diarrhea incidence was probably related to their combined effect with SCFA and MCFA. It has been demonstrated, that SCFA and MCFA can exert an inhibitory activity (Lei et al. 2017; Swanson et al. 2018; Zhang et al. 2020) or enhance the functional properties of phytogenic additives (McKnight et al. 2019).

In addition, dietary supplementation of organic acids can modulate the intestinal environment, creating undesirable environmental conditions for pathogenic bacteria, thus also influencing the intestinal microbiota (Verstegen and Williams 2002). Even if is difficult to establish the exact mechanisms for the enhancing antimicrobial effect by the combination of PHYs with organic acids (SCFA and MCFA) in pigs, we can suppose that PHYs can act as a permeabilizing complex and modify pores of the bacterial wall, thus facilitating the entrance of organic acids with antimicrobial activity (Tugnoli et al. 2020). In addition, the reduction in undigested feed protein by organic acids reduces the negative fermentative processes, increases growth performance and repairment of damaged intestinal tissues (Jia et al. 2020). Our results suggest that the addition of MCFA and SCFA to the phytogenic premix significantly inhibited enterotoxigenic *E. coli* diarrhea, thus supporting intestinal health of animals.

Considering the challenger strain shedding, the proliferation started gradually from the day of challenge in line with clinical observations. Compared with the uninfected control group infected animals showed hemolytic *E. coli* shedding from day 1 post-challenge, thus confirming the success of the experimental infection.

Histopathological examination of the ileum, jejunum and large intestine is thus used to highlight clinical signs of *E. coli* infection (Luppi 2017). In our study, histological evaluation of the ileum and lymphoid of intestinal tissues did not reveal significant lesions. The animals in the experimental trial thus did not show severe signs of intestinal lesions. Although more frequent lesions were registered in the PHY2 group, these did not impair animal performance and there was a comparable growth curve to CTRL-. This was probably due to the supplementation of phytogenic with SCFA and MCFA which could have supported intestinal health.

Gene expressions showed a high individual variability in terms of inflammatory parameters and tight junctions (TJs), probably due to the limited number of animals. We thus analyzed the expression of the TJ transmembrane protein (occludin) and the observed data were in line with morphological analyses. Our

findings suggested that tight junction integrity tended to be disrupted seven days after infection in challenged groups compared to the CTRL-. Intestinal permeability is regulated by the tight junctions which are a primary determinant of epithelial paracellular permeability (Zhang et al. 2021). Disruption of occludin regulation is related to many diseases. During the inflammation process, specific domains of occludin are in fact thought to mediate the transepithelial migration of neutrophils across the TJ (Feldman et al. 2005). Inflammation produces effects on epithelial barriers, increasing the leakiness of occludin, and decreasing the barrier function of this protein. Occludin responds earlier to oxidative stress than claudin, which responds later to reactive oxygen species (ROS) (Blasig et al. 2011). Intestinal bacterial infection is associated with intestinal epithelial and crypt architectural irregularity and with barrier dysfunction, leading to an increase in intestinal mucosal permeability. The observed slight downregulation of occludin after seven days in challenged groups could be due to the harmful activity of the challenger strain. Further investigations are required to better understand the effect of PHYs, SCFA and MCFA on the modulation of genes involved in inflammation and intestinal integrity.

Considering the biochemical parameters of the experimental groups (PHY1, PHY2, CTRL+ and CTRL-), the values were within the reference range of weaned pigs (Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia Romagna 2017; Klem et al. 2010), thus confirming that phytogenic additive supplementation had no detrimental effect on serum metabolism. The metabolite profile showed an increased level of total protein and a higher globulin content in CTRL+ compared to CTRL-. However, globulin together with albumin are the two major constituents of serum proteins, which play a crucial role in the inflammatory process (Balan et al. 2020; Wang et al. 2020). The increase in globulin could be associated with an inflammatory process probably due to the experimental *E. coli* infection leading to an increased concentration of total serum proteins. The serum aspartate aminotransferase (AST-GOT) level is a specific marker for liver tissue and represents a valuable indicator for acute hepatocyte injury or cell membranes damage (Kim 2020; Amirabagya et al. 2021). Although our results are in line with the proper range of pig physiology parameters (Klem et al. 2010; Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia Romagna 2017; Caprarulo et al. 2020b), AST-GOT was probably higher in the PHY2 group due to the presence of SCFA and MCFA which are immediately available for hepatic metabolism. In fact, short-chain fatty acids can activate lipid and glucose metabolism independently of the pig gut microbiota (Zhou et al. 2021).

## 5 Conclusions

Our study showed that phytogenic additive dietary supplementation limited the detrimental effect of experimental challenge. Phytogenic premix plus SCFA and MCFA revealed a complementary effect improving ADFI and fecal consistency during the post-challenge period compared to infected control group, suggesting that the combination of PHYs and organic acids can be considered as effective against pathogenic *E. coli* strains of weaned piglets. Future studies will be necessary to extensively investigate how phytogenic additives and organic acids affect gene expression over time.

# Declarations

**Acknowledgments:** We would like to thank the technical and veterinary staff and professors for participating in this experimental trial, and Dr. Giancarlo Selmini for his help in formulating the diets.

**Author contributions:** Conceptualization: L. Rossi, A. Baldi; Methodology: V. Caprarulo, L. Turin, M. Hejna, S. Reggi, M. Dell'Anno, P. Riccaboni, P. Trevisi, D. Luise, A. Baldi, L. Rossi; Writing-original draft preparation: M. Dell'Anno, V. Caprarulo, M. Hejna, L. Rossi; Review and editing: M. Dell'Anno, V. Caprarulo; M. Hejna, L. Rossi, A. Baldi; Formulation and preparation of all tested diets: L. Rossi; Statistical analysis: M. Dell'Anno, V. Caprarulo. All authors drafted and approved the final version of the manuscript.

**Funding:** This study received no funding.

**Data Availability:** All data generated or analyzed in this study are available from the corresponding author upon reasonable request. Data are stored on Google Drive at the following link:  
<https://drive.google.com/drive/folders/1fpkkuaELR6a5ZleTh3gszVoMrQJcGcMF?usp=sharing>.

**Ethics approval:** This study was performed in line with the principles of the Declaration of Helsinki.

The procedures and protocols used in this study were designed in accordance with the guidelines for animal welfare and the use of animals regulated under Directive 2010/63/EU on the protection of animals used for scientific purposes. The protocol was approved by the Animal Welfare Organization of the University of Milan and by the Italian Ministry for project (authorization number: 711/2017-PR, 28/09/2017).

**Consent to participate:** All authors agreed to participate in this work. They also approved the content of the research and the submission of the manuscript to Veterinary Research Communication journal.

**Consent to publish:** All authors agree to the content of the paper for publication.

**Conflicts of Interest:** The authors declare no conflict of interest.

# References

1. American Oil Chemistry Society (2009) Crude fiber analysis in feeds by filter bag technique. Official Methods and Recommended practices, Champaign, IL
2. Amirabagya F, Hapsari RAF, Wulandari E (2021) The Effect of *Jatropha curcas* L Seed Extract on AST/ALT Activity and The Central Vein Thickness in Liver. *Pharmacogn J* 13:66–72.  
<https://doi.org/10.5530/pj.2021.13.10>
3. Association of Official Analytical Chemists (2019) Official Methods of Analysis. Washington, DC, USA
4. Balan P, Staincliffe M, Moughan PJ (2020) Effects of spray-dried animal plasma on the growth performance of weaned piglets—A review. *J of Anim Physiol and Anim Nutr* 105:699–714.

<https://doi.org/10.1111/jpn.13435>

5. Blasig IE, Bellmann C, Cording J, Del Vecchio G, Zwanziger D, Huber O, Haseloff RF (2011) Occludin protein family: oxidative stress and reducing conditions. *Antiox Redox Signaling* 15:1195–1219. <https://doi.org/10.1089/ars.2010.3542>
6. Bonetti A, Tugnoli B, Piva A, Grilli E (2021) Towards Zero Zinc Oxide: Feeding Strategies to Manage Post-Weaning Diarrhea in Piglets. *Animals* 11:1–24. <https://doi.org/10.3390/ani11030642>
7. Caprarulo V, Giromini C, Rossi L (2020a) Chestnut and quebracho tannins in pig nutrition: the effects on performance and intestinal health. *Animal* 15:1–10. <https://doi.org/10.1016/j.animal.2020.100064>
8. Caprarulo V, Hejna M, Giromini C, Liu Y, Dell'Anno M, Sotira S, Reggi S, Sgoifo-Rossi CA, Callegari ML, Rossi L (2020b) Evaluation of Dietary Administration of Chestnut and Quebracho Tannins on Growth, Serum Metabolites and Fecal Parameters of Weaned Piglets. *Animals* 10:1–15. <https://doi.org/10.3390/ani10111945>
9. Cormican M, Hopkins S, Jarlier V, Reilly J, Simonsen G, Strauss R, Vandenberg O, Zabicka D, Zarb P, Catchpole M, Heuer O, Iosifidis E, Monnet D, Plachouras D, Weist K, Ricci A, Allende A, Bolton D, Chemaly M, Davies R, Escamez P, Girones R, Herman L, Koutsoumanis K, Lindqvist R, Norrung B, Robertson L, Ru G, Sanaa M, Simmons M, Skandamis P, Snary E, Speybroeck N, Ter Kuile B, Threlfall J, Wahlstrom H, Tenhagen B, Teale C, Schuepbach G, Beloeil P, Liebana E, Stella P, Murphy D, Hauser B, Urbain B, Kozhuharov E, Bozic F, Michaelidou-Patsia A, Bures J, Vestergaard E, Baptiste K, Tiirats T, Nevalainen M, Rouby J, Hahn G, Schlumbohm W, Malemis I, Kulcsar G, Lenhardsson J, Beechinor J, Breathnach R, Pasquali P, Auce Z, Maciulskis P, Schmit M, Spiteri S, Schefferlie G, Hekman P, Bergendahl H, Swiezicka A, Da Silva J, Taban L, Hederova J, Straus K, Madero C, Persson E, Jukes H, Weeks J, Kivilahti-Mantyla K, Moulin G, Wallmann J, Grave K, Greko C, Munoz C, Bouchard D, Catry B, Moreno M, Pomba C, Rantala M, Ruzauskas M, Sanders P, Schwarz C, van Duijkeren E, Wester A, Ignate K, Kunsagi Z, Torren-Edo J (2017) ECDC, EFSA and EMA Joint Scientific Opinion on a list of outcome indicators as regards surveillance of antimicrobial resistance and antimicrobial consumption in humans and food-producing animals. *EFSA J* 15. <https://doi.org/10.2903/j.efsa.2017.5017>
10. Costa LB, Luciano FB, Miyada VS, Gois FD (2013) Herbal extracts and organic acids as natural feed additives in pig diets. *S Afr J Anim Sci* 43:181–193. <https://doi.org/10.4314/sajas.v43i2.9>
11. Czech A, Grela ER, Kiesz M (2021) Dietary fermented rapeseed or/and soybean meal additives on performance and intestinal health of piglets. *Sci Rep* 11:1–10. <https://doi.org/10.1038/s41598-021-96117-w>
12. Dell'Anno M, Sotira S, Rebucci R, Reggi S, Castiglioni B, Rossi L (2020) *In vitro* evaluation of antimicrobial and antioxidant activities of algal extracts. *Ital J Anim Sci* 19:103–113. <https://doi.org/10.1080/1828051X.2019.1703563>
13. Dell'Anno M, Callegari ML, Reggi S, Caprarulo V, Giromini C, Spalletta A, Coranelli S, Sgoifo Rossi CA, Rossi L (2021) *Lactobacillus plantarum* and *Lactobacillus reuteri* as Functional Feed Additives to

- Prevent Diarrhoea in Weaned Piglets. *Animals* 11:1–19. <https://doi.org/10.3390/ani11061766>
14. Dell'Anno M, Hejna M, Sotira S, Caprarulo V, Reggi S, Pilu R, Miragoli F, Callegari ML, Panseri S, Rossi L (2020) Evaluation of leonardite as a feed additive on lipid metabolism and growth of weaned piglets. *Anim Feed Sci Technol* 266:1–12. <https://doi.org/10.1016/j.anifeedsci.2020.114519>
  15. Dell'Anno M, Reggi S, Caprarulo V, Hejna M, Sgoifo Rossi CA, Callegari ML, Baldi A, Rossi L (2021b) Evaluation of Tannin Extracts, Leonardite and Tributyrin Supplementation on Diarrhoea Incidence and Gut Microbiota of Weaned Piglets. *Animals* 11:1–18. <https://doi.org/10.3390/ani11061693>
  16. Diao H, Jiao A, Yu B, Mao X, Chen D (2019) Gastric infusion of short-chain fatty acids can improve intestinal barrier function in weaned piglets. *Genes Nutr* 14:1–16. <https://doi.org/10.1186/s12263-019-0626-x>
  17. Dibner J, Buttin P (2002) Use of organic acids as a model to study the impact of gut microflora on nutrition and metabolism. *J Appl Poult Res* 11:453–463. <https://doi.org/10.1093/japr/11.4.453>
  18. Directive (2010) 63/EU of the European Parliament and of the Council of 22 September 2010 on the protection of animals used for scientific purposes Text with EEA relevance
  19. Domínguez Díaz L, Fernández-Ruiz V, Cámara M (2020) The frontier between nutrition and pharma: The international regulatory framework of functional foods, food supplements and nutraceuticals. *Crit Rev Food Sci Nutr* 60:1738–1746. <https://doi.org/10.1080/10408398.2019.1592107>
  20. Dubreuil JD (2013) Antibacterial and antidiarrheal activities of plant products against enterotoxinogenic *Escherichia coli*. *Toxins* 5:2009–2041. <https://doi.org/10.3390/toxins5112009>
  21. Durmic Z, Blache D (2012) Bioactive plants and plant products: Effects on animal function, health and welfare. *Anim Feed Sci Technol* 176:150–162. <https://doi.org/10.1016/j.anifeedsci.2012.07.018>
  22. European Food Safety Authority, Rycken G, Aquilina G, Azimonti G, Bampidis V, Bastos MdL., Bories G, Chesson A, Cocconcelli PS, Flachowsky G (2018) EFSA Panel on Additives and Products or Substances used in Animal Feed: Guidance on the assessment of the efficacy of feed additives. *EFSA J* 16:e05274. <https://doi.org/10.2903/j.efsa.2018.5274>
  23. Feldman GJ, Mullin JM, Ryan MP (2005) Occludin: structure, function and regulation. *Adv Drug Delivery Rev* 57:883–917. <https://doi.org/10.1016/j.addr.2005.01.009>
  24. Ferrara F, Tedin L, Pieper R, Meyer W, Zentek J (2017) Influence of medium-chain fatty acids and short-chain organic acids on jejunal morphology and intra-epithelial immune cells in weaned piglets. *J Anim Physiol Anim Nutr* 101:531–540. <https://doi.org/10.1111/jpn.12490>
  25. Ferronato G, Prandini A (2020) Dietary supplementation of inorganic, organic, and fatty acids in pig: A review. *Animals* 10:1–27. <https://doi.org/10.3390/ani10101740>
  26. Grossi S, Rossi L, De Marco M, Sgoifo Rossi CA (2021) The Effect of Different Sources of Selenium Supplementation on the Meat Quality Traits of Young Charolaise Bulls during the Finishing Phase. *Antioxidants* 10:1–14. <https://doi.org/10.3390/antiox10040596>
  27. Han Y, Zhan T, Zhao Q, Tang C, Zhang K, Han Y, Zhang J (2020) Effects of mixed organic acids and medium chain fatty acids as antibiotic alternatives on the performance, serum immunity, and


- intestinal health of weaned piglets orally challenged with *Escherichia coli* K88. Anim Feed Sci Technol 269:1–13. <https://doi.org/10.3390/ani11051292>
28. Hanczakowska E (2017) The use of medium-chain fatty acids in piglet feeding-a review. Ann Anim Sci 17:967–977. <https://doi.org/10.1515/aoas-2016-0099>
29. Hayer SS, Rovira A, Olsen K, Johnson TJ, Vannucci F, Rendahl A, Perez A, Alvarez J (2020) Prevalence and trend analysis of antimicrobial resistance in clinical *Escherichia coli* isolates collected from diseased pigs in the USA between 2006 and 2016. Transbound Emerg Dis 67:1930–1941. <https://doi.org/10.1111/tbed.13528>
30. Istituto Zooprofilattico Sperimentale della Lombardia e dell'Emilia Romagna (2017) Parametri di Chimica Clinica: Valori Osservati in Suini di Diversa Età. Available online: [https://www.izsler.it/pls/izs\\_bs/v3\\_s2ew\\_consultazione.mostra\\_pagina?id\\_pagina=1494](https://www.izsler.it/pls/izs_bs/v3_s2ew_consultazione.mostra_pagina?id_pagina=1494)
31. Jackman JA, Boyd RD, Elrod CC (2020) Medium-chain fatty acids and monoglycerides as feed additives for pig production: towards gut health improvement and feed pathogen mitigation. J Anim Sci Biotechnol 11:1–15. <https://doi.org/10.1186/s40104-020-00446-1>
32. Jia M, Zhang Y, Gao Y, Ma X (2020) Effects of Medium Chain Fatty Acids on Intestinal Health of Monogastric Animals. Curr Protein Pept Sci 21:777–784. <https://doi.org/10.2174/1389203721666191231145901>
33. Kim HK (2020) The Effects of Anti-Inflammatory and Liver Function using Heat-Treated Cabbage. Int J Internet Broadcast Commun 12:131–138. <https://doi.org/10.7236/IJIBC.2020.12.3.131>
34. Klem TB, Bleken E, Morberg H, Thoresen SI, Framstad T (2010) Hematologic and biochemical reference intervals for Norwegian crossbreed grower pigs. Vet Clin Pathol 39:221–226. <https://doi.org/10.1111/j.1939-165X.2009.00199.x>
35. Lei XJ, Park JW, Baek DH, Kim JK, Kim IH (2017) Feeding the blend of organic acids and medium chain fatty acids reduces the diarrhea in piglets orally challenged with enterotoxigenic *Escherichia coli* K88. Anim Feed Sci Technol 224:46–51. <https://doi.org/10.1016/j.anifeedsci.2016.11.016>
36. Li P, Piao X, Ru Y, Han X, Xue L, Zhang H (2012) Effects of adding essential oil to the diet of weaned pigs on performance, nutrient utilization, immune response and intestinal health. Asian-Australas J Anim Sci 25:1617–1626. <https://doi.org/10.5713/ajas.2012.12292>
37. Li S, Wang L, Zhou Y, Miao Z (2020) Prevalence and characterization of virulence genes in *Escherichia coli* isolated from piglets suffering post-weaning diarrhoea in Shandong Province, China. Vet Med Sci 6:69–75. <https://doi.org/10.1002/vms3.207>
38. Lillehoj H, Liu Y, Calsamiglia S, Fernandez-Miyakawa M, Chi F, Cravens R, Oh S, Gay C (2018) Phytochemicals as antibiotic alternatives to promote growth and enhance host health. Vet Res 49:1–18. <https://doi.org/10.1186/s13567-018-0562-6>
39. Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta CT}$  method. Methods 25:402–408. <https://doi.org/10.1006/meth.2001.1262>

40. Luise D, Lauridsen C, Bosi P, Trevisi P (2019a) Methodology and application of Escherichia coli F4 and F18 encoding infection models in post-weaning pigs. *J Anim Sci Biotechnol* 10:1–20. <https://doi.org/10.1186/s40104-019-0352-7>
41. Luise D, Motta V, Bertocchi M, Salvarani C, Clavenzani P, Fanelli F, Pagotto U, Bosi P, Trevisi P (2019b) Effect of Mucine 4 and Fucosyltransferase 1 genetic variants on gut homeostasis of growing healthy pigs. *J Anim Physiol Anim Nutr* 103:801–812. <https://doi.org/10.1111/jpn.13063>
42. Luppi A (2017) Swine enteric colibacillosis: diagnosis, therapy and antimicrobial resistance. *Porcine Health Manag* 3:1–18. <https://doi.org/10.1186/s40813-017-0063-4>
43. Maenner K, Vahjen W, Simon O (2011) Studies on the effects of essential-oil-based feed additives on performance, ileal nutrient digestibility, and selected bacterial groups in the gastrointestinal tract of piglets. *J Anim Sci* 89:2106–2112. <https://doi.org/10.2527/jas.2010-2950>
44. McKnight LL, Peppler W, Wright DC, Page G, Han Y (2019) A blend of fatty acids, organic acids, and phytochemicals induced changes in intestinal morphology and inflammatory gene expression in coccidiosis-vaccinated broiler chickens. *Poult Sci* 98:4901–4908. <https://doi.org/10.3382/ps/pez241>
45. Mohammadi Gheisar M, Kim IH (2018) Phytobiotics in poultry and swine nutrition—a review. *Ital J Anim Sci* 17:92–99. <https://doi.org/10.1080/1828051X.2017.1350120>
46. Namkung H, Li M, Gong J, Yu H, Cottrill M, De Lange C (2004) Impact of feeding blends of organic acids and herbal extracts on growth performance, gut microbiota and digestive function in newly weaned pigs. *Can J Anim Sci* 84:697–704. <http://dx.doi.org/10.4141/A04-005>
47. Nowak P, Kasprowicz-Potocka M, Zaworska A, Nowak W, Stefańska B, Sip A, Grajek W, Juzwa W, Taciak M, Barszcz M (2017) The effect of eubiotic feed additives on the performance of growing pigs and the activity of intestinal microflora. *Arch Anim Nutr* 71:455–469. <https://doi.org/10.1080/1745039x.2017.1390181>
48. National Research Council (2012) Nutrient Requirements of Swine. The National Academies Press, Washington, DC. <https://doi.org/10.17226/13298>
49. Reg, COMMISSION REGULATION (EC) EC 152/2009, No 152/2009 of 27 January 2009 laying down the methods of sampling and analysis for the official control of feed (Text with EEA relevance)
50. Reggi S, Giromini C, Dell’Anno M, Baldi A, Rebucci R, Rossi L (2020) *In Vitro* Digestion of Chestnut and Quebracho Tannin Extracts: Antimicrobial Effect, Antioxidant Capacity and Cytomodulatory Activity in Swine Intestinal IPEC-J2 Cells. *Animals* 10:1–14. <https://dx.doi.org/10.3390%2Fani10020195>
51. Remfry SE, Amachawadi RG, Shi X, Bai J, Woodworth JC, Tokach MD, Dritz SS, Goodband RD, DeRouchey JM, Nagaraja TG (2020) Polymerase Chain Reaction-Based Prevalence of Serogroups of Escherichia coli Known to Carry Shiga Toxin Genes in Feces of Finisher Pigs. *Foodborne Pathog Dis* 17:782–791. <https://doi.org/10.1089/fpd.2020.2814>
52. Renzhammer R, Loncaric I, Roch F, Pinior B, Kasbohrer A, Spargser J, Ladinig A, Unterweger C (2020) Prevalence of Virulence Genes and Antimicrobial Resistances in *E. coli* Associated with Neonatal

- Diarrhea, Postweaning Diarrhea, and Edema Disease in Pigs from Austria. *Antibiotics* 9:1–13. <https://doi.org/10.3390/antibiotics9040208>
53. Reyes-Camacho D, Vinyeta E, Pérez JF, Aumiller T, Criado L, Palade LM, Taranu I, Folch JM, Calvo MA, Van der Klis JD (2020) Phytogenic actives supplemented in hyperprolific sows: effects on maternal transfer of phytogenic compounds, colostrum and milk features, performance and antioxidant status of sows and their offspring, and piglet intestinal gene expression. *J Anim Sci* 98:1–13. <https://dx.doi.org/10.1093%2Fjas%2Fskz390>
54. Rossi L, Di Giancamillo A, Reggi S, Domeneghini C, Baldi A, Sala V, Dell'Orto V, Coddens A, Cox E, Fogher C (2013) Expression of verocytotoxic *Escherichia coli* antigens in tobacco seeds and evaluation of gut immunity after oral administration in mouse model. *J Vet Sci* 14:263–270. <https://dx.doi.org/10.4142%2Fjvs.2013.14.3.263>
55. Rossi L, Dell'Orto V, Vagni S, Sala V, Reggi S, Baldi A (2014b) Protective effect of oral administration of transgenic tobacco seeds against verocytotoxic *Escherichia coli* strain in piglets. *Vet Res Commun* 38:39–49. <https://doi.org/10.1007/s11259-013-9583-9>
56. Rossi L, Pinotti L, Agazzi A, Dell'Orto V, Baldi A (2014a) Plant bioreactors for the antigenic hook-associated flgK protein expression. *Ital J Anim Sci* 13:23–29. <https://doi.org/10.4081/ijas.2014.2939>
57. Rossi L, Turin L, Alborali GL, Demartini E, Filipe JFS, Riva F, Riccaboni P, Scanziani E, Trevisi P, Dall'Ara P (2021) Translational Approach to Induce and Evaluate Verocytotoxic *E. coli* O138 Based Disease in Piglets. *Animals* 11:1–17. <http://dx.doi.org/10.3390%2Fani11082415>
58. Rossi L, Vagni S, Polidori C, Alborali GL, Baldi A, Dell'Orto V (2012) Experimental Induction of *Escherichia coli* Diarrhoea in Weaned Piglets. *Open J Vet Med* 2:1–8. <http://dx.doi.org/10.4236/ojvm.2012.21001>
59. Royce L, Liu P, Stebbins M, Hanson B, Jarboe L (2013) The damaging effects of short chain fatty acids on *Escherichia coli* membranes. *Appl Microbiol Biotechnol* 97:8317–8327. <https://dx.doi.org/10.1007%2Fs00253-013-5113-5>
60. Salsali H, Parker WJ, Sattar SA (2008) The effect of volatile fatty acids on the inactivation of *Clostridium perfringens* in anaerobic digestion. *World J Microbiol Biotechnol* 24:659–665. <http://dx.doi.org/10.1007%2Fs11274-007-9514-4>
61. Sun Y, Kim S (2017) Intestinal challenge with enterotoxigenic *Escherichia coli* in pigs, and nutritional intervention to prevent postweaning diarrhea. *Anim Nutr* 3:322–330. <https://doi.org/10.1016/j.aninu.2017.10.001>
62. Swanson A, Cochrane R, Amachawadi R, Remfry S, Lerner A, Nagaraja T, Pluske J, Niederwerder M, Stark C, Paulk C (2018) Determination of the Minimum Inhibitory Concentration of Various Medium Chain Fatty Acid-Based Products in *E. coli*, Enterotoxigenic *E. coli*, and *Campylobacter coli*. *J Anim Sci* 96:258–258. <https://doi.org/10.1093/jas/sky073.479>
63. Tang KL, Caffrey NP, Nóbrega DB, Cork SC, Ronksley PE, Barkema HW, Polachek AJ, Ganshorn H, Sharma N, Kellner JD (2017) Restricting the use of antibiotics in food-producing animals and its associations with antibiotic resistance in food-producing animals and human beings: a systematic

- review and meta-analysis. *Lancet Planet Health* 1:e316–e327. [https://doi.org/10.1016/s2542-5196\(17\)30141-9](https://doi.org/10.1016/s2542-5196(17)30141-9)
64. Thacker PA (2013) Alternatives to antibiotics as growth promoters for use in swine production: a review. *J Anim Sci Biotechnol* 4:1–12. <https://doi.org/10.1186/2049-1891-4-35>
65. Tretola M, Ottoboni M, Luciano A, Rossi L, Baldi A, Pinotti L (2019) Former food products have no detrimental effects on diet digestibility, growth performance and selected plasma variables in post-weaning piglets. *Ital J Anim Sci* 18:987–996. <https://doi.org/10.1080/1828051X.2019.1607784>
66. Tugnoli B, Giovagnoni G, Piva A, Grilli E (2020) From acidifiers to intestinal health enhancers: How organic acids can improve growth efficiency of pigs. *Animals* 10:1–18. <https://dx.doi.org/10.3390%2Fani10010134>
67. Upadhaya SD, Kim SJ, Kim IH (2016) Effects of gel-based phytogenic feed supplement on growth performance, nutrient digestibility, blood characteristics and intestinal morphology in weanling pigs. *J Appl Anim Res* 44:384–389. <https://doi.org/10.1080/09712119.2015.1091334>
68. Verstegen MW, Williams BA (2002) Alternatives to the use of antibiotics as growth promoters for monogastric animals. *Anim Biotechnol* 13:113–127. <https://doi.org/10.1081/abio-120005774>
69. Wang D, Lindemann MD, Estienne MJ (2020) Effect of Folic Acid Supplementation and Dietary Protein Level on Growth Performance, Serum Chemistry and Immune Response in Weanling Piglets Fed Differing Concentrations of Aflatoxin. *Toxins* 12:1–13. <https://dx.doi.org/10.3390%2Ftoxins12100651>
70. Yan L, Kim I (2012) Effect of eugenol and cinnamaldehyde on the growth performance, nutrient digestibility, blood characteristics, fecal microbial shedding and fecal noxious gas content in growing pigs. *Asian-Australas J Anim Sci* 25:1178–1183. <https://dx.doi.org/10.5713%2Fajas.2012.12111>
71. Yang C, Chowdhury M, Huo Y, Gong J (2015) Phytogenic compounds as alternatives to in-feed antibiotics: potentials and challenges in application. *Pathogens* 4:137–156. <https://doi.org/10.3390/pathogens4010137>
72. Zhang S, Dogan B, Guo C, Herlekar D, Stewart K, Scherl EJ, Simpson KW (2020) Short Chain Fatty Acids Modulate the Growth and Virulence of Pathosymbiont *Escherichia coli* and Host Response. *Antibiotics* 9:1–20. <https://doi.org/10.3390/antibiotics9080462>
73. Zhang Y, Li X, Qiao S, Yang D, Li Z, Xu J, Li W, Su L, Liu W (2021) Occludin degradation makes brain microvascular endothelial cells more vulnerable to reperfusion injury *in vitro*. *J Neurochem* 156:352–366. <https://doi.org/10.1111/jnc.15102>
74. Zhou H, Yu B, Sun J, Liu Z, Chen H, Ge L, Chen D (2021) Short-chain fatty acids can improve lipid and glucose metabolism independently of the pig gut microbiota. *J Anim Sci Biotechnol* 12:1–14. <https://doi.org/10.1186/s40104-021-00581-3>


## Figures



**Figure 1**

Experimental trial design from arrival (-14) to 7 days post-challenge.

PHY1: treatment group fed basal diet supplemented with 200g/100kg of phytogenic additive; PHY2: treatment group fed basal diet supplemented with 200g/100kg of phytogenic additive supplemented with 2000 ppm of short and medium chain fatty acids premix; phytogenic additive composed by caraway oil, lemon oil, clove, cinnamon, nutmeg, onion, pimento, orange peel, peppermint and chamomile powder; short chain and medium chain fatty acids blend included butyric (C4), caprylic (C8), capric (C10) and lauric acid (C12) at dose levels of 2000 ppm. CTRL: group fed basal diet supplemented with premix carrier divided into negative control (CTRL-) and positive control (CTRL+) challenged at day 0. Diets were analyzed for proximate analysis, including moisture, crude protein (CP), crude fibre (CF), ether extract (EE), and ash. The moisture determination was performed by oven-drying at 65°C for 24 h (Regulament EC 152/2009). Crude protein content was measured according to the Kjeldahl method (Association of Official Analytical Chemists method 2001.11). Crude fiber was determined by the filter bags technique (American Oil Chemistry Society 2009). Ether extract content was determined in a Soxhlet system after hydrolysis (Association of Official Analytical Chemists method 2003.05). Ash was measured using a muffle furnace at 550°C (Association of Official Analytical Chemists method 942.05).



**Figure 2**

*Escherichia coli* fecal shedding during the four days post-challenge where A) presents daily hemolytic *E. coli* fecal shedding from day 1 to day 4 post-challenge; B) presents average fecal hemolytic *E. coli* fecal shedding from 1 to 4 days post-challenge.

Data are presented as least squared means (LSMEANS) and standard errors (SE).

PHY1: basal diet plus 200g/100kg of phytogetic additive; PHY2: basal diet plus 200g/100kg of phytogetic additive plus 2000 ppm of short and medium chain fatty acids premix; phytogetic additive composed by caraway oil, lemon oil, clove, cinnamon, nutmeg, onion, pimento, orange peel, peppermint and chamomile powder; CTRL: basal diet plus premix carrier divided per negative control (CTRL-) and positive control (CTRL+) challenged at day 0.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterials10.01.22.docx](#)