Using Bayesian state-space models to understand the population dynamics of the dominant malaria vector, *Anopheles funestus* in rural Tanzania

Halfan Ngowo (✉ hngowo@ihi.or.tz)

Ifakara Health Institute <https://orcid.org/0000-0003-2789-0261>

Fredros Oketch Okumu

Ifakara Health Institute

Emmanuel Elirehema Hape

Ifakara Health Institute

Issa H Mshani

Ifakara Health Institute

Heather M Ferguson

University of Glasgow

Jason Matthiopoulos

University of Glasgow

Research Article

Keywords: *Anopheles funestus*, State space model, population dynamic, seasonality, abundance, density dependence

Posted Date: January 10th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1209027/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

1 Using Bayesian state-space models to understand the population dynamics of the
2 dominant malaria vector, *Anopheles funestus* in rural Tanzania

3
4 Halfan S. Ngowo^{1,2*}, Fredros O. Okumu^{1,2,3,4}, Emmanuel E. Hape^{1,2}, Issa H. Mshani^{1,2},
5 Heather M. Ferguson^{1,2}, Jason Matthiopoulos²

6 * Corresponding author

7
8 **Affiliations**

- 9 1. Department of Environmental Health & Ecological Sciences, Ifakara Health
10 Institute, Tanzania
11 2. Institute of Biodiversity, Animal Health and Comparative Medicine, University
12 of Glasgow, United Kingdom
13 3. School of Public Health, University of the Witwatersrand, Republic of South
14 Africa
15 4. School of Life Science and Bioengineering, Nelson Mandela African Institution of
16 Science & Technology, Tanzania

17
18 **Email address:**

- 19 1. Halfan Ngowo: hngowo@ihi.or.tz
20 2. Fredros Okumu: fredros@ihi.or.tz
21 3. Emmanuel Hape: ehape@ihi.or.tz
22 4. Issa Mshani: imshani@ihi.or.tz
23 5. Heather Ferguson: Heather.Ferguson@glasgow.ac.uk
24 6. Jason Matthiopoulos: Jason.Matthiopoulos@glasgow.ac.uk

Abstract

Background: It is often assumed that the population dynamics of the malaria vector *Anopheles funestus*, its role in malaria transmission and the way it responds to interventions are similar to the more elaborately characterized *An. gambiae*. However, *An. funestus* has several unique ecological features that could generate distinct transmission dynamics and responsiveness to interventions. The objectives of this work were to develop a model which will; 1) reconstruct the population dynamics, survival, and fecundity of wild *An. funestus* populations in southern Tanzania, 2) quantify impacts of density dependence on the dynamics, and 3) assess seasonal fluctuations in *An. funestus* demography. Through quantifying the population dynamics of *An. funestus*, this model will enable analysis of how their stability and response to interventions may differ from that of *An. gambiae* s.l.

Methods: A Bayesian State Space Model (SSM) based on mosquito life history was fit to time series data on the abundance of female *An. funestus* s.s. collected over 2 years in southern Tanzania. Prior values of fitness and demography were incorporated from empirical data on larval development, adult survival and fecundity from laboratory-reared first generation progeny of wild caught *An. funestus*. The model was structured to allow larval and adult fitness traits to vary seasonally in response to environmental covariates (i.e. temperature and rainfall), and for density dependency in larvae. We measured the effects of density dependence and seasonality through counterfactual examination of model fit with or without these covariates.

Results: The model accurately reconstructed the seasonal population dynamics of *An. funestus* and generated biologically-plausible values of their survival larval, development and fecundity in the wild. This model suggests that *An-funestus* survival and fecundity annual pattern was highly variable across the year, but did not show consistent seasonal trends either rainfall or temperature. While the model fit was somewhat improved by inclusion of density dependence, this was a relatively minor effect and suggests that this process is not as important for *An. funestus* as it is for *An. gambiae* populations.

Conclusion: The model's ability to accurately reconstruct the dynamics and demography of *An. funestus* could potentially be useful in simulating the response of these populations to vector control techniques deployed separately or in combination. The observed and simulated dynamics also suggests that *An. funestus* could be playing a role in year-round malaria transmission, with any apparent seasonality attributed to other vector species.

Key words: *Anopheles funestus*, State space model, population dynamic, seasonality, abundance, density dependence.

Introduction

Anopheles funestus is one of the major malaria vectors in Africa and is widely distributed across the continent [1,2]. With the exception of *Anopheles gambiae* s.s, the species appears to have higher vectorial capacity than many other members of the *An. gambiae* s.l. complex [3-8]. *Anopheles funestus* makes a higher contribution to transmission than *An. gambiae* s.l. in numerous parts in sub-Saharan Africa [6,9-12]; particularly in settings where *An. gambiae* s.l. abundance has plummeted due to either effective indoor-based vector control interventions [13,14] or environmental change. It is hypothesized that *An. funestus* persistence despite the recent scale-up of insecticide-treated nets may have been facilitated by their earlier development of strong physiological resistance [15].

Anopheles funestus is typically grouped with *An. gambiae* s.l. when modelling transmission and formulating policies for malaria vector control [16-18]. The lack of explicit consideration of *An. funestus* ecology and transmission potential may be partially due to this species having been relatively neglected compared to *An. gambiae* s.l. Comparatively the ecology of *An. funestus* s.l is less well understood, and it is much more difficult to maintain under insectary or semi-field conditions [19]. However, this species has several unique ecological features, such as its different larval habitat and dry season persistence [20], that could give rise to distinct population dynamics and differentiate its response to core and supplementary

interventions. For example, *Anopheles funestus* prefers larger aquatic habitats that are semi-permanent or permanent throughout the year, and contain clear water with some emergent vegetation [20]. This differs from *An. gambiae* s.s which generally prefer small temporary habitats such as puddles, ditches and animal hoof prints etc. [20,21], or *An. arabiensis*, which can breed extensively in rice fields and other sunlit open pools [1]. The use of more permanent larval habitats means that *An. funestus* has greater persistence through the driest periods of the year compared to *An. gambiae* sl. [22] whose habitats evaporate quickly in the absence of rainfall [21,23]. This ecological feature means that the seasonal phenology of *An. funestus* and its response to aquatic microclimate differs from *An. gambiae* sl. [21,22,24];and could thus generate differential response to seasonally-targeted interventions such as Indoor Residual Spraying (IRS) and larviciding.

Differential use of aquatic habitats may also impact the relative importance of key intrinsic drivers of mosquito population dynamics such as density dependence. Density dependence in malaria vectors occurs during larval development as a product of competition for space and nutritional resources [25,26]. In space-limited habitats, high larval densities can influence larval development rates and survival but also subsequent adult fitness traits such as body size, survival, fecundity and mating success [27-30]. While there is evidence that density dependence is an important driver of *An. gambiae* s.l population dynamics [25], the relative importance of this process for *An. funestus* is unknown. Given that larval crowding and competition are less likely within the larger habitats preferred by *An. funestus*, we hypothesize that density dependence may be less pronounced for this vector species. Quantifying the strength of density dependence is important to inform the ease with which vector populations can be suppressed and how quickly they can recover [26,27,31].

Models of vector population dynamics and their response to interventions must be parameterised by reliable estimates of their demography and fitness. For vectors in the *An. gambiae* s.l. group, such estimates are often acquired from insectary and semi-field studies [32-37] as well as field studies. Similar data has been difficult to

obtain for *An. funestus* because of its poorly understood ecology and the difficulties of creating laboratory colonies; which so far has been achieved on only two occasions [11,19,38]. State-space models (SSM) provide an alternative approach to indirectly estimate these parameters by fitting a population dynamics model to observed time series data [39,40]. These models are widely used in other fields of ecology and conservation biology to investigate the population dynamics of other animals (e.g. [39,41]) and guide management decisions [41]. However, these models have so far had limited uptake in medical entomology. Given data on population fluctuations are available, these models can infer and estimate plausible demographic rates that could generate the observed dynamics [42].

SSMs are time-series models that distinguish between two stochastic components, namely, process (i.e. biological), which captures sequential dependencies between population components (e.g. eggs, larvae, pupae etc.) and an observation component, which captures and corrects for biases and imprecisions in the data-collection process. Prior knowledge of the model parameters is used to bolster the information content of the time series data with existing expert or lab data and uncertainty in estimates. Population projections are then quantified on the basis of posterior probability distributions for parameters and population states. SSM models have recently been used to elucidate the dynamics and impacts of interventions on malaria vectors in laboratory and semi field populations [32,33], but have not yet been applied to estimate *An. funestus* vector demographics in the wild. Here, we present an innovative SSM application to describe the population dynamics of wild *An. funestus* populations in Tanzania, and use it to assess extrinsic (environmental) and intrinsic (density dependence) drivers of their fitness and demography for the first time. We incorporate empirical data from laboratory experiments on *An. funestus* colonization [19] together with the wild population data to develop an SSM. Time series field data collected in 2015 [6] and 2018 south-eastern Tanzania, and corresponding environmental information were used to validate the model. Specific aims were: 1) to accurately reconstruct the population dynamics, survival and fecundity of wild *An. funestus* populations in southern Tanzania, 2) quantify the

effects of density dependence on the dynamics, and 3) to identify and quantify seasonal variations in *An. funestus* demography.

Methods

Time series data on wild An. funestus populations

Indoor densities of female *An. funestus* s.l adults were recorded over 12 months of entomological surveys conducted in three villages (Tulizamoyo, Ikwambi and Sululu) in Kilombero (8.1539°S, 36.6870°E) and Ulanga (8.3124°S, 36.6879°E) districts, south-eastern Tanzania from June 2018 to May 2019. The villages were selected because of the high abundance of *An. funestus* s.l; within which *An. funestus* s.s. is the dominant sibling species (93%) [1]. Annual rainfall was 1200mm-1800mm, and temperature, 20 °C-32 °C. CDC Light traps [43] were used to sample host-seeking mosquitoes from 6pm to 6am for 5 days per week, 4 weeks a month for 12 months in 10-15 houses per village. The houses were randomly selected and consent obtained from household heads. The mosquitoes were sorted by taxa and sex, and females further classified as unfed, blood-fed or gravid. Daily climatic data (rainfall and temperature) were obtained from a weather station, approximately ~20km from the farthest village.

To complement this data, we extracted *An. funestus* data from a 2015 dataset from three other villages in Ulanga district (Mavimba, Minepa and Kivukoni) [6]. These data were collected five days per week for a period of 12 months. This data allowed us to fit the model simultaneously to multiple time series so that it could learn hierarchically from *An. funestus* trajectories enfolding in different years and locations. This additional data has previously been described elsewhere and used to demonstrate the epidemiological dominance of *An. funestus*, which now contributes >85% of all malaria infections in the region [6].

Prior information on life-history and gonotrophic cycle stages

Female *An. funestus* adults collected from the same three villages in 2018 were maintained in insectary conditions for one generation to estimate baseline fitness traits as already described in Ngowo et al [19] and in Figure 1. Data collected from this 1st-generation laboratory progeny included: a) proportions of eggs that hatched

into larvae, larvae that transitioned to pupae, and of pupae that emerged into adults, b) the length of the transition periods (days) between life stages ((i) Eggs to 1st instar larvae, (ii) 1st instar larvae to pupae, and (iii) pupae to unfed adult female (one day post emergence)), c) transition period of adult females between three different stages of their gonotrophic cycle , i.e. unfed, blood-fed and gravid.

The gonotrophic cycle starts with ‘unfed’ females who transition to ‘blood-fed’ after obtaining a blood meal. In the wild, the first gonotrophic cycle usually starts after unfed females have mated [44]; which is assumed to happen soon after emergence. In insectary experiments, females had access to males immediately on emergence. As the blood meal is digested, blood-fed females transition into the ‘gravid’ state during which eggs develop. Gravid females then oviposit their eggs into aquatic habitats and return to the ‘unfed’ stage with the cycle begins again (Figure 1). In the wild, the rate of transition between these gonotrophic stages is governed by both intrinsic and extrinsic environmental conditions including the availability of blood-meals and oviposition sites [45]. In the insectary, the first blood-meal was offered 5 days post emergence to ensure individuals had sufficient time for mating.

Per capita fecundity was defined as the number of eggs laid per fully bloodfed adult female. The proportions surviving between life-history stages or gonotrophic stages were calculated as the inverse of the number of days required to transit from one stage to the next:

$$s_m = (j_m)^{1/r} \quad (1)$$

Here, m is the life cycle stage, j is proportion survived as the percentage of the total from preceding life stage, r is the average number of days it took to transit from one life stage to another, and s is the daily survival within the stage (Table 1).

Biological process components of the Bayesian SSMs

Daily survival: The daily survival of larval stages was assumed to be the same for all instar stages. Adult survival was assumed to be the same for unfed, blood-fed and

gravid females. Survival probabilities (s_l, s_a, s_f, s_v) were linked to their covariates through a *logit* transformation of linear predictors (here, subscripts l, a, f, v refer to larvae, unfed, blood-fed and gravid respectively). Pupal (S_p) and egg (S_e) survival probabilities were considered to be independent of any climatic and density-dependent covariates, and were treated as a binomial distribution, with baseline rates assigned priors as described in eq. 16-17.

A range of covariates hypothesized to be associated with the demography of *An. funestus* were incorporated to allow baseline larval and adult survival to vary with environmental conditions. Rainfall (current and 1 week-lagged) and temperature were incorporated into the larval survival model. Rainfall regulates the availability and permanence of aquatic habitats, thus influencing both survival and carrying capacity of larval habitats [46]. We also incorporated density dependence into the model of larval survival [25] to assess whether this could improve the fit of the adult population dynamics model. Additionally, the speed of larval development was modelled as a function of temperature based on its known importance [47,48]. The daily survival of larvae was thus defined as a function of daily rainfall (current and lagged), daily temperature and density dependence. We estimated the daily survival rates (lowercase $s_l(t)$) of larvae through a logit transformation of linear predictors (uppercase $S_l(t)$).

$$s_l(t) = \frac{\exp(S_l(t))}{1 + \exp(S_l(t))} \quad (2)$$

Specifically, $S_l(t)$ is written as a function of both intrinsic and extrinsic drivers:

$$S_l(t) = \beta_0 - \beta_1 R_{(t-1)} - \beta_2 D_{(t-1)} \left(1 - \frac{\beta_3 Q_{(t-1)}}{\max(Q)}\right) + \beta_4 T_{(t-1)} - \beta_5 T_{(t-1)}^2 + \varepsilon_{l,t} \quad (3)$$

$$\varepsilon_l \sim \text{Normal}(0, \sigma_{l,t}) \quad (4)$$

Here, β_0 is the baseline daily larval survival on the linear scale. We assigned a prior to survival probability under zero rainfall and average temperature (i.e. 27°C) and then calculated the intercept of β_0 to reflect this prior information. When there is no effect of any environmental covariates (prior takes values between 0.80-1, Table 1). The coefficient β_1 quantifies the effect of current rainfall (R); with the envisioned scenario being that higher R (i.e. flooding) tends to wash away larvae hence reducing the baseline survival [49]. This β_1 was defined by an informative gamma prior with shape = 5.382 and rate of 46.4 (Table 2) which permits anything from no rain effects to 100% mortality. The coefficient β_2 quantifies the effect of larval density at time t on larval survival. We assume a monotonic negative relationship based on the biologically-plausible hypothesis that larval survival is reduced at high larval density because of resource competition and intraspecific cannibalism [29,50]. This coefficient β_2 was defined by an uninformative gamma prior with shape of 0.5 and rate of 1 (Table 2), which allows the impact of density to range from no effect to complete annihilation.

The term inside brackets in Eq. (3) represents the fact that density dependence needs to be modulated by the availability of larval habitat. The availability of suitable aquatic habitats for oviposition will increase with rainfall; thus potentially reducing the crowding of larvae into the remaining habitats that persist during the dry season. This hypothesis has been supported for *An. gambiae* sl., where their seasonal population dynamics can be explained by models incorporating a rainfall-dependent carrying capacity [25]. Here, the coefficient β_3 was a proportion that captures the potential interaction between larval habitat availability (defined as the cumulative rainfall (Q) over the past week) and larval density (D). When rain in the recent week has been the maximum observed (i.e. $Q = \max(Q)$), then $(1 - \frac{\beta_3 Q_{(t-1)}}{\max(Q)})$ would be the smallest amount of density dependency experienced by *An. funestus*. The prior distribution for β_3 was defined by an upward-biased beta prior with mean 0.9 and variance of 0.01 allowing β_3 to have positive impact on larvae survival.

Additional covariates were incorporated to assess the role of temperature on larval survival (via the coefficients β_4, β_5). The parameter β_4 captures the potentially positive effects of temperature on daily larval survival, which were defined by an uninformative gamma prior with mean of 1 and variance of 0.1 considering 27°C as the optimal temperature (ρ) for maximum survival [51]. This prior allows temperature to vary from having no impact, to high positive impact on larval survival. Alternatively, the relationship between larval survival and temperature may be characterized by survival being reduced at low or very high temperature, and peaking in the middle [52]. The coefficient β_5 was incorporated to capture this potential curvilinear relationship but was dependent on β_4 , to ensure that the optimum temperature was fixed at 27°C, ($\beta_5 = \frac{\beta_4}{2\rho}$). The prior of β_4 allowed extreme temperatures away from the optimum to range from having no effect to generating 100% mortality. The parameter ε_l is capturing the unexplained stochasticity associated with larval survival. This error term was defined by a normal prior with mean of 0 and a precision σ_l from a gamma distribution with both shape and rate of 10.

The linear predictors for survival of unfed, blood-fed, gravid (uppercase $S_a(t), S_f(t), S_v(t)$) and daily probabilities of survival (lowercase $s_a(t), s_f(t), s_v(t)$) were structured similarly to eq 3. The daily survival probabilities of adult stages were thus defined as the functions of daily temperature; such that an increase in temperature would result in an increase in the survival of all three adults stages and reduction in survival when temperature become lethal [22,53]. The biological relationship between adult survival and temperature was assumed to be curvilinear [22,53,54].

$$S_a(t) = \varphi_0 + \varphi_1 T_{(t-1)} - \varphi_2 T_{(t-1)}^2 + \varepsilon_{a,t} \quad (5)$$

$$S_f(t) = \theta_0 + \theta_1 T_{(t-1)} - \theta_2 T_{(t-1)}^2 + \varepsilon_{f,t} \quad (6)$$

$$S_v(t) = \alpha_0 + \alpha_1 T_{(t-1)} - \alpha_2 T_{(t-1)}^2 + \varepsilon_{v,t} \quad (7)$$

$$\varepsilon_t^* \sim \text{Normal}(0, \sigma_t^*) \quad (8)$$

Here, φ_0 , θ_0 , and α_0 refer to the baseline survival of unfed, blood-fed and gravid females respectively on a linear scale, under fixed temperature conditions of $27 \pm 2^\circ \text{C}$ (insectary standard under which *An. funestus* have maximum survival [52,54]), and assumes no blood meal limitation. The positive impact of temperature on all three life stages was represented by the coefficients φ_1 , θ_1 and α_1 with an uninformative gamma prior with mean 12.5 and variance of 6.25. The coefficients φ_2 , θ_2 and α_2 correspond to the curvilinear effect of temperature on the survival of all three adult stages with their priors derived from the ratio between the linear coefficient and twice optimum temperature. This formulation ensured that the optimum temperature is fixed at (27°C). The parameters ε_a , ε_f , ε_v capture unexplained variation associated with survival during the distinct gonotrophic stages. These error terms (ε^*) were defined by normal priors with mean of 0 and a precision σ^* from a gamma distribution with both shape and rate of 10 for unfed, blood-fed and gravid females.

Development between stages: The daily development probability from one life stage to the next was defined as the reciprocal of the development time (days) between the stages (assuming that all development times take longer than a day). An increase in temperature was assumed to reduce the development period of larvae [47,53,55,56].

$$l(t) = \frac{\exp(L(t))}{1 + \exp(L(t))} \quad (9)$$

Specifically, $L(t)$ is written as the function of temperature covariates:

$$L(t) = C_0 + C_1 T_{(t-1)} \quad (10)$$

Here C_0 corresponds to the baseline daily development period on a linear scale defined by an informative beta prior with range defined in eq 16-17 (Table 1). The coefficient C_1 explains the positive effect of temperature on larval development period, with its prior values derived from an uninformative gamma prior with mean 0.001 and standard deviation 0.001.

The development time for other life history stages (eggs and pupae) and the time between gonotrophic stages were assumed to be independent of temperature and other environmental covariates. The numbers of individuals ($\mathcal{K}_m$) graduating from one stage to the next each day were modelled as a binomial process Eq 11.

$$\mathcal{K}_m(t) \sim \text{Binomial}(r_m, \mathcal{W}_{m-1}(t)) \quad (11)$$

Here the rate r_m is a development probability as defined in eq 11 for m stage, with assigned informative prior values as described through a generic prior in eq 16-17 (Table 1). Parameter $\mathcal{W}_{m-1}(t)$ refers to the number surviving the preceding life stage.

Fecundity

The number of eggs laid at each time step was drawn from a Poisson distribution whose rate was the product of per-capita fecundity (number of eggs laid by blood-fed *An. funestus* under insectary conditions (b_0), a penalized rate for the egg survival (s_e), the number of gravid mosquitoes ($V_{(t-1)}$) and ratio of females-males (assumed to be 0.5) as assessed at the pupae stage [19].

$$b_t = \exp(b_0 + \varepsilon_{b,t}) \quad (12)$$

$$B(t) \sim \text{Poisson}(0.5b_t s_e V_{(t-1)}) \quad (13)$$

The error term ε_b was defined by normal prior with mean of 0 and a precision from a gamma distribution with both shape and rate of 10

Observation-derived *components of the Bayesian SSMs*

Observations of the abundance of adults (unfed, bloodfed, gravid) A at time t were modelled as a normal distribution with varying daily means $\bar{a}$ determined by the biological model and a precision τ representing observation error. We assumed a fixed coefficient of variation (ξ) for the daily observation process which was assigned an uninformative prior with values between 0.1-0.9 (Table 1). The CDC light trap typically samples mosquitoes from populations of unknown size, for which the daily catch rates are difficult to quantify independently. A parameter ϑ was therefore incorporated both into the precision τ and daily varying means to account for an observed weekly periodicity in adult abundance, which was otherwise hard to interpret. This parameter was allowed to vary both by day of the week j and between the two populations k (2015 and 2018-19 datasets). The ϑ values were derived from a logit function $\exp(\rho)/(1 + \exp(\rho))$, with ρ defined from the uninformative normal prior with mean and standard deviation of 0 and 10 respectively. Therefore precision τ can be written as $\frac{1}{(\xi_t \bar{a}_t \vartheta_{jk})^2}$ for all the adult stages. This mean the observation abundance was estimated as follows:

$$A(t) \sim Normal\left(\bar{a}_t \vartheta_{jk}, \frac{1}{(\xi_t \bar{a}_t \vartheta_{jk})^2}\right) \quad (14)$$

Trap bias: The trapping method (CDC Light traps) primarily targets unfed host-seeking mosquitoes [57]. Blood fed and gravid mosquitoes are assumed to no longer host-see, and represent a small proportion 0.5-3% of total collections of females caught [6,22]). To account for these biases in sampling, we added a new parameter of “trap-biasness” ω in the observation model for both precision τ and varying daily means $\bar{a}_\eta$. The prior values for ω were estimated from independent studies from the

same locations [6,58], and ranged from 0.05-0.15 (Table 1), with variations between the two life stages η . Therefore, the observation model for blood-fed and gravid (A_η) was rewritten by modifying eq 14 as follows

$$A_\eta(t) \sim Normal\left(\bar{a}_{\eta t} \vartheta_{jk} \omega_\eta, \frac{1}{(\xi_t \bar{a}_{\eta t} \vartheta_{jk} \omega_\eta)^2}\right) \quad (15)$$

Prior distributions: Since this model contains a large number of parameters, use of un-scaled informative priors restricted model convergence and mixing. We therefore opted for rescaled beta distributions of the informative priors [59], calibrated as follows:

$$Y = Beta(5, 5) \quad (16)$$

$$X = X_{min} + Y(X_{max} - X_{min}) \quad (17)$$

Where Y is a dummy variable that takes values in the interval $[0, 1]$ with mean of 0.5 and standard deviation of 0.15, selected to provide low likelihood at the values 0 and 1. The values of X_{min} and X_{max} define the range of the parameter of interest as dictated by the prior information. Since the information on priors was provided in form of mean (μ) and standard deviation (σ), the values were defined as $X_{min}, X_{max} = \mu \pm 2\sigma$. Survival, development period, trap-biasness, variability in daily catches and fecundity parameters were all assigned priors according to Eq. 17.

Model selection, model fitting and outputs

Model fitting was done using the R statistical software version 4.0.5 [60]. Population models were fitted using a Markov Chain and Monte Carlo sampling (MCMC) algorithm via the JAGS software [61] interfaced to R via the *runjags* package [62] (code provided in the supplementary material S1). To achieve convergence, the model with 6 chains was run in parallel for 10^5 samples with a burn-in of 10^5 , keeping every 10^{th} iteration for memory-saving reasons. Convergence was assessed by visual investigation of the trace plots, prior-posterior distribution using the *coda* package [63], effective

sample sizes and the Gelman Rubin diagnostic [64]. Model comparisons were done using the deviance information criterion (DIC) [65], and the ones with the lowest DIC selected as the most preferred. The predicted and observed densities of *An. funestus* adult females were plotted to evaluate consistent prediction biases visually (Figure S2). Posterior means and 95% credible intervals for the key survival parameters, development period, density dependence, environmental covariates (temperature and rainfall) and fecundity were also reported to reveal different dynamical aspects of the system.

Results

Population trajectories and seasonal trends: We used a Bayesian state-space model to describe the population dynamics of wild populations of *An. funestus*. The full results, including summaries of posterior means for all the fitness and demographic parameters are reported in Table 1. The most parsimonious model (model-7, Table 3) included density dependence, and temperature and rainfall (current and lagged) impacts on larval survival, and the effect of temperature on larval development period. The only covariate that was not retained in the “model-7” was temperature impacts on adult survival. This model satisfactorily reconstructed the population dynamics of *An. funestus* in the study villages, with all environmental covariate relaxation applied based on DIC selection. Population trajectories were estimated for all six *An. funestus* life history and gonotrophic stages after accounting for potential impacts of environmental covariates and density dependence (Figure 2).

These trajectories reflect the annual trend in abundance spanning in periods from low or no rainfall to high rainfall. All trajectories show a relatively high abundance of *An. funestus* right after the rainy season, followed by reduced but sustained abundance during the dry period for all the life stages. After accounting for observation biases during sampling, the observed abundance of unfed, gravid and blood-fed groups largely falls within the credible intervals of the predicted values (Figure 3).

Survival and fecundity: Estimated *An. funestus* larval survival trajectories demonstrate substantial mean variability during the two seasons, with no clear pattern of seasonality (Figure 4a&b), Table 1). Similarly, the survival trajectories of the adult stages (all gonotrophic states) were variable throughout the year, with daily survival rate ranging from 0.2 to 1.0 and not consistently differing between wet and dry seasons (Figure 4c-h, Table 1). Per capita fecundity was estimated to be between 75 and 81 eggs per female *An. funestus* (Table 1). While the abundance of this species fluctuated seasonally, per capita fecundity remained consistent throughout the year (Figure 4k&l).

Temperature was an important predictor of larval survival with a curvilinear relationship ($\Delta\text{DIC} = 138$, Table 3, Figure 5b), and that temperature has a positive monotonic relationship with larval development period ($\Delta\text{DIC} = 336$, Table 3, Figure 5a); with the larval development period estimated to last about 16 days on average. Additionally, daily rainfall was an important driver for the dynamics of *An. funestus* by reducing larval survival ($\Delta\text{DIC} = 5605$, Table 3, Figure 5c) with a negative monotonic relationship.

Effects of density dependence: Density dependence was the only intrinsic feature incorporated in this dynamic model of *An. funestus*. The model was able to converge efficiently without crashing when density dependency was removed, suggesting this process plays a detectable but relatively minor role population regulation when compared with extrinsic factors ($\Delta\text{DIC} = 222$, Table 3, Figure 5e). To verify this, we run a simulation and discovered that the estimated density dependence was actually quite low, even when simulating with a single dataset. The model fitting process also suggested the interaction parameter (β_3) between larvae density (D) and one week cumulative rainfall (Q) contributes to *An. funestus* dynamics by positively increasing larval survival ($\Delta\text{DIC} = 38$, Table 3, Figure 5d).

Discussion

We have adapted advanced methods used in mainstream ecological studies and fitted a state-space model (SSM) to field and laboratory data to accurately reconstruct population dynamics of wild population of *An. funestus*. The SSM inferred the trajectories of multiple life-cycle and gonotrophic stages of wild *An. funestus* females. This allowed us to reconstruct the observed trajectories of larvae and adult females for the wild *An. funestus* in Tanzania for the first time. This analysis indicated that the dynamics of *An. funestus* were best explained in a model that included density dependency, temperature (curvilinear relationship) and rainfall (negative monotonic relationship) on larval survival, and temperature on the larval development period (positive monotonic relationship). In contrast, model fit was not improved by incorporating temperature dependency into adult survival (all gonotrophic stages). *Anopheles funestus* abundance vary seasonally between wet and dry but demographics rates (i.e. survivals, fecundity and development period) did not vary after accounting for the impact of environmental covariates and density dependence. These results are very useful for generating hypotheses about the nature and relative magnitude of drivers of *An. funestus* population dynamics in the wild. This model can be extended to include a component on malaria dynamics in humans; or to compare the efficacy and effectiveness of different interventions in combination or singly. This would allow more sophisticated evaluation of suitability of *An. funestus*-specific interventions; including prediction of the potential combined effect of strategies that acting at different life-cycle stages and/or target different demographic processes (e.g. survival versus fecundity).

Extrinsic covariates such as rainfall and temperature were all hypothesised to be the main drivers for the dynamics of this vector species. This study supports the hypothesis that rainfall is a significant driver of the population dynamics of wild *An. funestus*. Overall, the abundance of all life stages were relatively higher in rainy compared to dry periods of the year as previously documented [22,66-68]. Rainfall covariates were directly included in the larval survival model since it is the only stage on which rainfall was hypothesized to have a significant impact. Daily larval survival

as estimated by the SSM model showed high variability both within seasons and across the year. There was support for a monotonic association between rainfall and larval survival; characterized as reduction in larval survival during periods of heavy rainfall [69]. Time lags have been used to assess rainfall impacts on the dynamics of these vectors (e.g.[22]). *Anopheles funestus* abundance have been shown to be positively associated with the cumulative lag rainfall [22]. Here we modelled a one week cumulative rainfall to account for its effect on survival. Similar to other vectors of malaria transmission such as *An. gambiae* s.l, rainfall have always been considered as the main factor regulating the dynamics, despite ecological differences between the two vector species [22,46,70,71].

The SSM also provided support for the hypothesis that temperature is an important driver of *An. funestus* dynamics; although the nature of temperature effects was complex and variable between life history stages. For example, temperature was associated with both larval survival and development, but not adult survival or fitness. Furthermore the estimated impacts of temperature on larval ecology were complex; with the SSM suggesting a curvilinear relationship with survival but a positive monotonic impact with the larval development period. Our findings validate prior studies that demonstrated that temperature had a curvilinear influence on *Anopheles* larval survival, with a rise in temperature above/below the optimum lowering survival [47,72,73]. The larval period of *An. gambiae* s.l. is temperature dependent [55,73]; thus our model incorporated a positive monotonic effect such that development is fastest when temperature is high and just below maximum threshold for larval development [73,74]. In our final model the effect of temperature on gonotrophic stages was not found to be an important driver for the dynamic of this species thus left out during model selection process.

Little is known about the effect of density dependence on *An. funestus* due to its ecology and reliance on the large semi-permanent and permanent breeding habitats [20,21]. However, density dependence is already well-known to be an important driver for dynamic of other malaria vectors like *An. gambiae* [25,27-29,31,67,75] and other non-malaria vectors like *Aedes aegypti* [76,77]. Variations in densities during

the aquatic stages of *An. gambiae* s.l. have been found to affect adult fitness [28,78]. The SSM model fit better when density dependence of larval survival was included, though the relative magnitude of this process was quite small and likely to have minor impact on the overall dynamics of *An. funestus* populations (Figure 5e). These findings suggest that *An. funestus* populations are likely to be regulated more by extrinsic than intrinsic processes. These findings corroborate our original hypotheses about density dependence having a weaker regulatory role in this species on the basis of the types of larval habitats (i.e. larger and more permanent habitat [20,79]), which can likely sustain higher resources and thus reduced competition than in *An. gambiae* s.l. To our knowledge, this is the first report documenting the role of density dependence on the dynamics of the wild populations of *An. funestus*. Now that colonies are becoming more feasible, more thorough investigation on the role of density dependence in the dynamic of *An. funestus* is prerequisite.

In addition to highlighting potential drivers of *An. funestus* populations, the SSM here generated plausible estimates of key demographic and life-history process in the wild. Our model estimated that *An. funestus* larvae takes an average of 15.6-16.1 days to grow from first instar larvae to pupae; which is relatively long compared to the other major vectors in the *An. gambiae* s.l complex (9-11 days [55,72]). This apparently longer development period of *An. funestus* may be a product of their adaptation to more permanent, year-round breeding habitats that are unlikely to dry up; thus reducing selection for rapid development. The SSM estimated that the daily survival of wild *An. funestus* larvae could be as high as 0.95, compared to the 0.83 [0.80, 0.86] mean daily survival rate of the known vector of malaria transmission *An. gambiae* s.l. [72,80]. This matches observations from insectary experiments in which *An. funestus* larvae have higher survival than *An. gambiae* s.l. [19,38,81]. Given the apparently higher rates of survival in *An. funestus* than in *An. gambiae* s.l., these findings suggest that more lethal intervention may be required to control *An. funestus* both at larvae and adults stages.

The impact of any vector control largely depends on the ecology of the specific vector species. Differences in ecology between *An. gambiae* s.l. and *An. funestus* are likely

to affect the relative impact of interventions. For instance, *An. gambiae* s.l. prefers breeding in small and temporary habitats which dry up quickly when there is no rainfall which is opposite to *An. funestus* habitats. Despite the fact that *An. funestus* habitats are "few, fixed, and findable" and might be easily targeted for larviciding [20] during the dry season, treating habitats such as rivers or bigger ponds could pose logistical challenges. The persistence of *An. funestus* throughout the year even during the driest periods suggest this vector is less seasonal compared to *An. gambiae* s.l. which experience much more dramatic "boom and bust" dynamics in relation to seasonal rains [6,22,68]. If interpreted together with the observation that survival estimates were not seasonal, the model suggests that this species is likely responsible for year-round malaria transmission throughout the year, while other species, which mostly occupy temporary habitats may be responsible for any apparent seasonality in transmission.

Models of vector population dynamics can provide a useful guide for the selection of optimal vector control strategies; particular through enabling more focal investigation of the benefits of seasonal or spatial targeting and use of combined versus single interventions. Despite its complexity, this population dynamics model provides a useful framework for investigation of the stability of *An. funestus* populations. With additional data, this model can be further refined to include additional modifications related to vector ecology and behaviour that may impact intervention impact (e.g. host choice and its impacts on fitness, predation during larval or adult phase). Such further elucidation may increase the predictive accuracy of this SSM in specific contexts, but even the more general framework developed here have flexibility to introduce stage-specific mortality effects expected from different types of vector control [16,32,33]. For example, this framework could be used to model the impact of combined interventions including those that target adult females (insecticide-treated nets (ITNs), IRS) and larviciding; and assessment of how mortality varies with different coverage [16,32]. An important limitation of this study is we have no idea what percentage of *An. funestus* mosquito population is sampled by the trap, which is important for understanding the relative magnitude of demographic stochasticity in

modelled dynamics. This highlights the need to explicitly incorporate this source of uncertainty into vector and transmission dynamics; including the need for further calibration and standardization of the efficiency and biases associated with particular mosquito trapping methods.

Conclusions

This study used Bayesian State Space Models (SSM) parameterized with empirical data to quantify key demographic and fitness processes underpinning the population dynamics of *An. funestus* in Tanzania. We report the first use of SSM to understand the population dynamic of the wild vector of residual malaria transmission, *An. funestus* in Tanzania. The model structure allowed investigation of the relative importance of seasonally-varying environmental covariates (i.e. rainfall and temperature) and density dependence; providing some support for both processes although the magnitude of the former was much greater than the latter. The ability of this model to accurately reconstruct the seasonal dynamics and demography of *An. funestus* indicate its value for simulating the response of these populations to vectors control measures applied either individually or in combination. Additionally, the relatively limited evidence of seasonality in key fitness and demographic rates further corroborate evidence that this vector species can facilitate efficient year-round transmission of malaria. Finally, this model also highlights the clear importance of accounting for regional and daily observation biases when modelling mosquito population dynamics.

Table 1: Priors as used in the state-space population models of *Anopheles funestus* and the estimated posteriors mean and 95% credible intervals.

Parameter		Prior distribution				Posterior distribution	
Notation	Description	Type	Source	Mean	95-percentiles	Mean	95-percentiles
s_e	Eggs daily survival rate	Beta	This study	0.794	[0.619, 1]	0.789	[0.776, 0.804]
λ	Eggs development period	Beta	This study	0.5	[0.4, 0.6]	0.499	[0.485, 0.514]
β_0	Baseline larval daily survival	Beta	This study	0.923	[0.801, 1]	0.950	[0.943, 0.956]
C_0	Baseline larval development period	Beta	This study	0.063	[0.055, 0.071]	0.063	[0.062, 0.064]
s_p	Pupae daily survival rate	Beta	This study	0.941	[0.874, 1]	0.944	[0.930, 0.950]
p_r	Pupae development period	Beta	This study	0.522	[0.253, 0.792]	0.525	[0.506, 0.546]
φ_0	Baseline unfed daily survival	Beta	This study	0.935	[0.877, 0.992]	0.937	[0.933, 0.941]
a_r	Unfed development period	Beta	This study	0.20	[0.19, 0.21]	0.200	[0.198, 0.201]
θ_0	Baseline blood-fed daily survival	Beta	This study	0.807	[0.654, 0.961]	0.810	[0.799, 0.820]
f_r	Blood-fed daily	Beta	This study	0.25	[0.05, 0.45]	0.269	[0.256, 0.280]

	transition rate						
α_0	Baseline gravid daily survival	Beta	This study	0.904	[0.848, 0.961]	0.903	[0.899, 0.907]
v_r	Gravid daily transition rate	Beta	This study	0.333	[0.133, 0.533]	0.311	[0.297, 0.324]
b_0	No. eggs/female (Per capita fecundity)	Beta	This study	80	[60, 100]	78	[75, 81]
ξ	Coefficient of variability	Beta	Uninformative prior	0.5	[0.1, 0.9]	0.79	[0.810, 0.825]
$\omega_{\eta=f}$	Coefficient of “Trap biasness” for the blood-fed	Beta	Msugupakyula et al 2020 [58] and Kaindoa et al 2017 [6]	0.1	[0.05, 0.15]	0.122	[0.117, 0.127]
$\omega_{\eta=v}$	Coefficient of “Trap biasness” for the gravid			$0.505^* \omega_f$	[0.025, 0.076]	0.062	[0.059, 0.064]

Table 2: Priors for the intrinsic and extrinsic drivers of the population dynamic as used in the state-space model of *Anopheles funestus* and the estimated posteriors mean and 95% credible intervals.

Parameter		Prior distribution				Posterior distribution	
Notation	Description	Type	Source	Mean	sd	Mean	95-percentiles
β_1	Linear coefficient for rainfall on larvae survival	Gamma	Uninformative prior	0.1	0.05	0.01681	[0.00604, 0.0308]
β_2	Density dependent coefficient for larvae on larvae survival	Gamma	Uninformative prior	0.5	0.7	1.0283e-4	[1.0e-4, 1.1205e-4]
β_3	Coefficient of interaction between larvae and rainfall on larvae survival	Gamma	Uninformative prior	0.9	0.1	0.9601	[0.80810, 0.99999]
β_4	Linear coefficient for temperature on larvae survival	Gamma	Uninformative prior	1	0.316	0.487	[0.203, 0.806]
β_5	Quadratic coefficient for temperature on larvae survival		a function of β_4	$\beta_5 = \frac{\beta_4}{2 * \rho}$		-0.00902	[-0.01492, -0.00376]
c_1	Linear coefficient for temperature on larvae development period	Gamma	Uninformative prior	0.001	0.001	5.362e-4	[2.51e-8, 2.311e-3]

φ_1	Linear coefficient for temperature on unfed, survival	Gamma	Uninformative prior	1	0.316	0.074	[0.068, 0.081]
φ_2	Quadratic coefficient for temperature on unfed survival	Gamma	a function of φ_1	$\varphi_2 = \frac{\varphi_1}{2 * \rho}$		-1.378e-3	[-1.50e-3, -1.26e-3]
θ_1	Linear coefficient for temperature on bloodfed survival	Gamma	Uninformative prior	1	0.316	0.074	[0.068, 0.081]
θ_2	Quadratic coefficient for temperature on bloodfed survival	Gamma	a function of θ_1	$\theta_2 = \frac{\theta_1}{2 * \rho}$		-1.378e-3	[-1.50e-3, -1.26e-3]
α_1	Linear coefficient for temperature on gravid survival	Gamma	Uninformative prior	1	0.316	0.074	[0.068, 0.081]
α_2	Quadratic coefficient for temperature on gravid survival	Gamma	a function of α_1	$\alpha_2 = \frac{\alpha_1}{2 * \rho}$		-1.378e-3	[-1.50e-3, -1.26e-3]

Table 3: Model selection: Description of all models fitted with and without environmental covariates and their corresponding delta-Deviance Information Criterion Δ DIC

Model	Removed covariate(s)	Fitness measure	Penalized Deviance (pD)/DIC	Δ pD/DIC
Model 1-Full	None		35500	25339
Model 2	Temperature	Larval survival	10555	394
Model 3	Rainfall	Larval survival	10277	116
Model 4	One week cummulative rainfall*density dependency	Larval survival	11886	1725
Model 5	Density dependency	Larval survival	11011	850
Model 6	Temperature	Larval development period	10873	712
Model 7^	Temperature	Adult survival	10161	0
Model 8	Model 7 - Temperature	Larval survival	10299	138
Model 9	Model 7 - Rainfall	Larval survival	15766	5605
Model 10	Model 7 - 1 week rainfall:density dependency	Larval survival	10199	38
Model 11	Model 7 - Density dependency	Larval survival	10383	222
Model 12	Model 7 - Temperature	Larval development	10497	336

^ The best model (lowest DIC/Penalized Deviance) value-model-7 followed by model-10. **Model 8-12** consists of model-7 minus one more environmental covariate. **Model 4** involved the removal of the interaction term.

Figures:

Figure 1: Schematic representation of the state-space population model showing different life stages compartment (circles) and flows (arrows) of *Anopheles funestus*. Abundance data were only available for unfed, blood-fed and gravid stages. The model assumes that once a gravid mosquito has laid eggs, they return to the unfed stage. The annotations are described in Table 1. The model incorporates six life stages (eggs, larvae, pupae, unfed, bloodfed and gravid) of *An. funestus*.

Figure 2: Reconstruction of the abundance trajectories for all the six life-stages. The red line indicates the mean posterior values and the respective 95% confidence intervals are shown in “sky-blue”. Left column (a,c,e,g,i,k) is data collected from June 2018 to May 2019 and right column (b,d,f,h,j,l) is data collected from Jan-Dec 2015. The grey area indicates the period with rainfall.

Figure 3: Observed vs. model estimated values for the three adult stages with data collected using CDC light trap both in May 2018 -June 2019 (left column- a,c,d) and Jan-Dec 2015 (right column- b,d,e). Red lines are the model estimated trajectories with “sky-blue” showing their 95% credible intervals. The blue circles are the observed values from the Light trap catches. Grey areas are the periods with rainfalls episodes.

Figure 4: Reconstruction of the survival trajectories for all the four stages (larvae, unfed, bloodfed and gravid) which were affected by the environmental covariates with the two bottom row showing the larval development period and fecundity trends. Left column (a,c,e,g,i,k) is trajectories from June 2018 to May 2019 and right column (b,d,f,h,j,l) is from Jan-Dec 2015. Grey area is the period with rainfall. Y-axis shows the survival rates of different life stages and the bottom row (k&l) shows per-capita fecundity

Figure 5: Relationship between environmental covariates and fitness parameters as estimated from the SSMs of population dynamic of *Anopheles funestus*.

Supplementary Figures

Figure S1: SSMs model development R codes

Figure S2: Goodness-of-fit: Observed versus predicted unfed, bloodfed and gravid densities across all populations. Adjusted R-squared, intercept and slope values are from a linear model of the predicted against observed values. Dotted lines correspond to 1:1 line. Left column (a,c,d) is data collected from June 2018 to May 2019 and right column (b,d,e) is data from Jan-Dec 2015. Grey area is the period with rainfall.

Figure S3.1: Prior (orange histogram) and posterior (blue histogram) distribution of the main baseline and observational parameters in the state-space model.

Figure S3.2: Prior (orange histogram) and posterior (blue histogram) distribution of the main environmental covariates parameters in the state-space model.

Declarations:

Ethics approval and consent to participate

This study was approved by Ifakara Health Institute Review Board (Ref: IHI/IRB/No: 007-2018) and the Medical Research Coordinating Committee (MRCC) at the National Institute for Medical Research-NIMR (Ref: NIMR/HQ/R.8a/Vol.IX/2895). Individual verbal and written consent was also obtained from household owner where CDC light traps were places for collecting mosquitoes.

Consent for publication

Permission to publish was obtained from Medical Research Coordinating Committee (MRCC) at the National Institute for Medical Research (Ref. No: NIMR/HQ/P.12 VOL XXXIII/116).

Availability of data and materials

The R-codes used for model development are freely available in the supplementary file.

Competing interests

The authors declare no conflicts of interest

Funding

This work was supported by Howard Hughes Medical Institute (HHMI)-Gates Foundation (Grants: OPP1099295) and Gates Foundation (Grants: INV-002138) both awarded to FOO and HMF which support HSN.

Author's contributions

HSN, FOO, HMF and JM designed the study. HSN, IHM and EEH conducted the lab experiment and collected data used in model development. HSN and JM conducted the formal model development, fitting and validation. HSN drafted the manuscript. FOO, HMF and JM reviewed the manuscript and provide supervision. All authors accept and approve the final manuscript for publication.

Acknowledgments:

We are grateful to member of IHI-VectorSphere insectary's Joseph Mgando, Dickson, Mwasheshi, Godfrey Matanila, Neema Nombo, Rukiyah Mohammed, Johnson Kyeba and Khamis Bwanally for their contribution in different occasion during mosquito rearing, colonization and collection of wild population data. The earlier version of the model development benefited from the contributions of Luca Nelli, Fergus Chadwick, Heather McDevitt, Crinan Jarrett, Yacob Haddou and Prashanth Selvaraj.

Reference:

1. Gillies M, Meillon D. The Anophelinae of Africa south of the Sahara (Ethiopian zoogeographical region). Sahara Ethiop Zoogeographical. 1968;343 pp.
2. Sinka ME, Bangs MJ, Manguin S, Rubio-Palis Y, Chareonviriyaphap T, Coetzee M, et al. A global map of dominant malaria vectors. Parasit Vectors. 2012;5:69.
3. Russell TL, Govella NJ, Azizi S, Drakeley CJ, Kachur SP, Killeen GF, et al. Increased proportions of outdoor feeding among residual malaria vector populations following increased use of insecticide-treated nets in rural Tanzania. Malar J. BioMed Central; 2011;10:80.
4. Hunt RH, Brooke BD, Pillay C, Koekemoer LL, Coetzee M. Laboratory selection for and characteristics of pyrethroid resistance in the malaria vector *Anopheles funestus*. Med Vet Entomol. 2005;19:271-5.
5. Braack LE, Coetzee M, Hunt RH, Biggs H, Cornel A, Gericke A. Biting pattern and host-seeking behavior of *Anopheles arabiensis* (Diptera: Culicidae) in northeastern South Africa. J Med Entomol. 1994;31:333-9.
6. Kaindoa EW, Matowo NS, Ngowo HS, Mkandawile G, Mmbando A, Finda M, et al. Interventions that effectively target *Anopheles funestus* mosquitoes could significantly improve control of persistent malaria transmission in south-eastern Tanzania. PLoS One. 2017;12.
7. Lwetoijera D, Harris C, Kiware SS, Dongus S, Devine GJ, McCall PJ, et al. Increasing role of *Anopheles funestus* and *Anopheles arabiensis* in malaria transmission in the Kilombero Valley, Tanzania. Malar J. 2014;13:331.
8. Garrett-Jones C, Shidrawi GR. Malaria vectorial capacity of a population of *Anopheles gambiae*: an exercise in epidemiological entomology. Bull World Health Organ. 1969;40:531-45.
9. Seyoum A, Sikaala CH, Chanda J, Chinula D, Ntamatungiro AJ, Hawela M, et al. Human exposure to anopheline mosquitoes occurs primarily indoors, even for users of insecticide-treated nets in Luangwa Valley, South-east Zambia. Parasit Vectors. 2012;5:101.
10. Riveron JM, Chiumia M, Menze BD, Barnes KG, Irving H, Ibrahim SS, et al. Rise of multiple insecticide resistance in *Anopheles funestus* in Malawi: A major concern for malaria vector control. Malar J. 2015;
11. Hunt RH, Edwardes M, Coetzee M. Pyrethroid resistance in southern African *Anopheles funestus* extends to Likoma Island in Lake Malawi. Parasites and Vectors. 2010;3.
12. McCann RS, Ochomo E, Bayoh MN, Vulule JM, Hamel MJ, Gimnig JE, et al. Reemergence of *Anopheles funestus* as a vector of *Plasmodium falciparum* in Western

Kenya after long-term implementation of insecticide-treated bed nets. *Am J Trop Med Hyg.* 2014;

13. Bayoh MN, Mathias DK, Odiere MR, Mutuku FM, Kamau L, Gimnig JE, et al. *Anopheles gambiae*: historical population decline associated with regional distribution of insecticide-treated bed nets in western Nyanza Province, Kenya. *Malar J.* 2010;9:62.

14. Killeen GF, Seyoum A, Sikaala C, Zomboko AS, Gimnig JE, Govella NJ, et al. Eliminating malaria vectors. *Parasit Vectors.* 2013;6:172.

15. Okumu F, Finda M. Key Characteristics of Residual Malaria Transmission in Two Districts in South-Eastern Tanzania-Implications for Improved Control. *J Infect Dis.* 2021;223:S143-54.

16. Fillinger U, Kannady K, William G, Vanek MJ, Dongus S, Nyika D, et al. A tool box for operational mosquito larval control: Preliminary results and early lessons from the Urban Malaria Control Programme in Dar es Salaam, Tanzania. *Malar J.* 2008;7:1-25.

17. Killeen GF, Chitnis N, Moore SJ, Okumu FO. Target product profile choices for intra-domiciliary malaria vector control pesticide products: Repel or kill? *Malar J.* 2011;10.

18. Griffin JT, Hollingsworth TD, Okell LC, Churcher TS, White M, Hinsley W, et al. Reducing *Plasmodium falciparum* malaria transmission in Africa: A model-based evaluation of intervention strategies. *PLoS Med.* 2010;7.

19. Ngowo HS, Hape EE, Matthiopoulos J, Okumu FO. Fitness Characteristics of the Malaria Vector, *Anopheles Funestus*, During an Attempted Laboratory Colonization. *Malar J. BioMed Central*; 2020;20:148.

20. Nambunga IH, Ngowo HS, Mapua SA, Hape EE, Msugupakulya BJ, Msaky DS, et al. Aquatic habitats of the malaria vector *Anopheles funestus* in rural south-eastern Tanzania. *Malar J. BioMed Central*; 2020;19:1-11.

21. Gimnig JE, Ombok M, Kamau L, Hawley WA. Characteristics of Larval Anopheline (Diptera: Culicidae) Habitats in Western Kenya. *Source J Med Entomol J Med Entomol.* 2001;38:282-8.

22. Ngowo HS, Kaindoa EW, Matthiopoulos J, Ferguson HM, Okumu FO. Variations in household microclimate affect outdoor-biting behaviour of malaria vectors. *Wellcome Open Res.* 2017;2:102.

23. Minakawa N, Sonye G, Mogi M, Yan G, Communication S. Habitat characteristics of *Anopheles gambiae* s . s . larvae in a Kenyan highland. *Med Vet Entomol.* 2004;18:301-5.

24. Kabbale FG, Akol AM, Kaddu JB, Onapa AW. Biting patterns and seasonality of *Anopheles gambiae sensu lato* and *Anopheles funestus* mosquitoes in Kamuli District,

Uganda. *Parasit Vectors*. 2013;6:340.

25. Russell TL, Lwetoijera DW, Knols BGJ, Takken W, Killeen GF, Ferguson HM. Linking individual phenotype to density-dependent population growth: the influence of body size on the population dynamics of malaria vectors. *Proc R Soc B Biol Sci*. 2011;278:3142-51.

26. Nowicki P, Bonelli S, Barbero F, Balletto E. Relative importance of density-dependent regulation and environmental stochasticity for butterfly population dynamics. *Oecologia*. 2009;161:227-39.

27. Gimnig JE, Ombok M, Otieno S, Michael G, Vulule JM, Walker ED, et al. Density-Dependent Development of *Anopheles gambiae* (Diptera : Culicidae) Larvae in Artificial Habitats. *J Med Entomol*. 2002;

28. Ng'habi, John B, Nkwengulila G, Knols BGJ, Killeen GF, Ferguson HM. Effect of larval crowding on mating competitiveness of *Anopheles gambiae* mosquitoes. *Malar J*. 2005;4:1-9.

29. Koenraadt CJM, Majambere S, Hemerik L, Takken W. The effects of food and space on the occurrence of cannibalism and predation among larvae of *Anopheles gambiae* s.l. *Entomol Exp Appl*. John Wiley & Sons, Ltd; 2004;112:125-34.

30. Charlwood JD. May the force be with you: measuring mosquito fitness in the field. *Ecol Asp Appl Genet Modif Mosquitoes*. 2003;2:47-62.

31. Henderson PA, Magurran AE. Direct evidence that density-dependent regulation underpins the temporal stability of abundant species in a diverse animal community. *Proc R Soc B Biol Sci*. 2014;281.

32. Viana M, Ng'habi K, Lyimo I, Ferguson HM, Matthiopoulos J, Killeen G. Mesocosm experiments reveal the impact of mosquito control measures on malaria vector life history and population dynamics. *Sci Rep*. 2018;8:1-12.

33. Viana M, Hughes A, Matthiopoulos J, Ranson H, Ferguson HM. Delayed mortality effects cut the malaria transmission potential of insecticide-resistant mosquitoes. *Proc Natl Acad Sci U S A*. 2016;113:8975-80.

34. Lyimo IN, Haydon DT, Russell TL, Mbina KF, Daraja AA, Mbehela EM, et al. The impact of host species and vector control measures on the fitness of African malaria vectors. *Proc Biol Sci*. 2013;280:20122823.

35. Mweresa CK, Omusula P, Otieno B, van Loon JJA, Takken W, Mukabana WR. Molasses as a source of carbon dioxide for attracting the malaria mosquitoes *Anopheles gambiae* and *Anopheles funestus*. *Malar J*. 2014;13:160.

36. Lyimo IN, Haydon DT, Mbina KF, Daraja AA, Mbehela EM, Reeve R, et al. The fitness of African malaria vectors in the presence and limitation of host behaviour. *Malar J*. 2012;11:1-12.

37. Ferguson HM, Ng'habi KR, Walder T, Kadungula D, Moore SJ, Lyimo I, et al. Establishment of a large semi-field system for experimental study of African malaria vector ecology and control in Tanzania. *Malar J.* 2008;7:158.
38. Okoye PN, Brooke BD, Hunt RH, Coetzee M. Relative developmental and reproductive fitness associated with pyrethroid resistance in the major southern African malaria vector, *Anopheles funestus*. *Bull Entomol Res.* 2007;97:599-605.
39. Newman KB, Buckland ST, Lindley ST, Thomas L, Fernández C. Hidden process models for animal population dynamics. *Ecol Appl.* 2006;16:74-86.
40. Millar R. Bayesian state-space modeling of age-structured data: fitting a model is just the beginning. *Can J Fish Aquat Sci.* 2000;50:43-50.
41. Viana M, Cleaveland S, Matthiopoulos J, Halliday J, Packer C, Craft ME, et al. Dynamics of a morbillivirus at the domestic-wildlife interface: Canine distemper virus in domestic dogs and lions. *Proc Natl Acad Sci.* 2015;112:1464-9.
42. Rivot E, Prévost E, Parent E, Baglinière JL. A Bayesian state-space modelling framework for fitting a salmon stage-structured population dynamic model to multiple time series of field data. *Ecol Modell.* 2004;179:463-85.
43. Mboera LEG. Sampling techniques for adult Afrotropical malaria vectors and their reliability in the estimation of entomological inoculation rate. *Tanzan Health Res Bull.* 2005;7:117-24.
44. Gillies MT. The duration of the gonotrophic cycle in *Anopheles gambiae* and *Anopheles funestus*, with a note on the efficiency of hand catching. *East Afr Med J.* 1953;30:129-35.
45. Rúa GL, Quiñones ML, Vélez ID, Zuluaga JS, Rojas W, Poveda G, et al. Laboratory estimation of the effects of increasing temperatures on the duration of gonotrophic cycle of *Anopheles albimanus* (Diptera: Culicidae). *Mem Inst Oswaldo Cruz.* 2005;100:515-20.
46. Fillinger U, Sonye G, Killeen GF, Knols BGJ, Becker N. The practical importance of permanent and semipermanent habitats for controlling aquatic stages of *Anopheles gambiae* sensu lato mosquitoes: Operational observations from a rural town in western Kenya. *Trop Med Int Heal.* 2004;
47. Sciences B, Laboratories S, Bayoh MN, Lindsay SW. Temperature-related duration of aquatic stages of the Afrotropical malaria vector mosquito *Anopheles gambiae* in the laboratory. *Med Vet Entomol.* 2004;18:174-9.
48. Lyimo., Takken W, Koella JC. Effect of rearing temperature and larval density on larval survival, age at pupation and adult size of *Anopheles gambiae*. *Entomol Exp Appl.* 1992;63:265-71.
49. Beier JC, Copeland R, Oyaro C, Masinya a, Odago WO, Oduor S, et al. *Anopheles*

gambiae complex egg-stage survival in dry soil from larval development sites in western Kenya. *J Am Mosq Control Assoc.* 1990;6:105-9.

50. Koenraadt CJM, Majambere S, Hemerik L, Takken W. Cannibalism and predation among larvae of *Anopheles gambiae* s.l. *Entomol Exp Appl.* 2004;112:125-34.

51. Paaijmans KP, Heinig RL, Seliga RA, Blanford JI, Blanford S, Murdock CC, et al. Temperature variation makes ectotherms more sensitive to climate change. *Glob Chang Biol. John Wiley & Sons, Ltd;* 2013;19:2373-80.

52. Paaijmans KP, Imbahale SS, Thomas MB, Takken W. Relevant microclimate for determining the development rate of malaria mosquitoes and possible implications of climate change. *Malar J.* 2010;9:196.

53. Blanford JI, Blanford S, Crane RG, Mann ME, Paaijmans KP, Schreiber K V, et al. Implications of temperature variation for malaria parasite development across Africa. *Sci Rep.* 2013;3:1300.

54. Paaijmans KP, Thomas MB. The influence of mosquito resting behaviour and associated microclimate for malaria risk. *Malar J.* 2011;10:183.

55. Kirby MJ, Lindsay SW. Effect of temperature and inter-specific competition on the development and survival of *Anopheles gambiae sensu stricto* and *An. arabiensis* larvae. *Acta Trop.* 2009;109:118-23.

56. Paaijmans KP, Blanford S, Bell AS, Blanford JI, Read AF, Thomas MB. Influence of climate on malaria transmission depends on daily temperature variation. *Proc Natl Acad Sci U S A.* 2010;107:15135-9.

57. Mboera L. Sampling techniques for adult Afrotropical malaria vectors and their reliability in the estimation of entomological inoculation rate. *Tanzan J Health Res. National Institute for Medical Research;* 2006;7:117-24.

58. Msugupakulya BJ, Kaindo EW, Ngowo HS, Kihonda JM, Kahamba NF, Msaky DS, et al. Preferred resting surfaces of dominant malaria vectors inside different house types in rural south - eastern Tanzania. *Malar J. BioMed Central;* 2020;1-15.

59. Gelman A, Jakulin A, Pittau MG, Su YS. A weakly informative default prior distribution for logistic and other regression models. *Ann Appl Stat.* 2008;2:1360-83.

60. R Development Core Team. R: A language and environment for statistical computing. *R Found. Stat. Comput.* 2021.

61. Plummer M. JAGS : A Program for Analysis of Bayesian Graphical Models Using Gibbs Sampling JAGS : Just Another Gibbs Sampler. 2003;

62. Denwood MJ. runjags : An R Package Providing Interface Utilities, Model Templates, Parallel Computing Methods and Additional Distributions for MCMC Models in JAGS . *J Stat Softw.* 2016;

63. Plummer M, Best N, Cowles K, Vines K. CODA: Convergence Diagnosis and Output Analysis for MCMC. R News. 2006;
64. Gelman A, Rubin DB. Inference from iterative simulation using multiple sequences. Stat Sci. 1992;
65. Spiegelhalter DJ, Best NG, Carlin BP, Linde A van der. Bayesian Measures of Model Complexity and Fit. J R Stat Soc Ser C, Appl Stat. 2002;64:583-639.
66. Kreppel KS, Viana M, Main BJ, Johnson PCD, Govella NJ, Lee Y, et al. Emergence of behavioural avoidance strategies of malaria vectors in areas of high LLIN coverage in Tanzania. Sci Rep. Nature Publishing Group UK; 2020;10:1-11.
67. Smith T, Charlwood JD, Takken W, Tanner M, Spiegelhalter DJ. Mapping the densities of malaria vectors within a single village. Acta Trop. 1995;59:1-18.
68. Charlwood JD, Vij R, Billingsley PF. Dry season refugia of malaria-transmitting mosquitoes in a dry savannah zone of east Africa. Am J Trop Med Hyg. 2000;62:726-32.
69. Kelly-Hope, A L, Hemingway J, McKenzie FE. Environmental factors associated with the malaria vectors *Anopheles gambiae* and *Anopheles funestus* in Kenya. Malar J. 2009;8:268.
70. Lyons CL, Coetzee M, Terblanche JS, Chown SL. Thermal limits of wild and laboratory strains of two African malaria vector species, *Anopheles arabiensis* and *Anopheles funestus*. Malar J. 2012;11:226.
71. Zhou G, Munga S, Minakawa N, Githeko AK, Yan G. Spatial relationship between adult malaria vector abundance and environmental factors in western Kenya highlands. Am J Trop Med Hyg. 2007;77:29-35.
72. Lyimo EO, Takken W, Koella JC. Effect of rearing temperature and larval density on larval survival, age at pupation and adult size of *Anopheles gambiae*. Entomol Exp Appl. 1992;63:265-71.
73. Paaijmans KP, Huijben S, Githeko AK, Takken W. Competitive interactions between larvae of the malaria mosquitoes *Anopheles arabiensis* and *Anopheles gambiae* under semi-field conditions in western Kenya. Acta Trop. 2009;109:124-30.
74. Lyons CL, Coetzee M, Chown SL. Stable and fluctuating temperature effects on the development rate and survival of two malaria vectors, *Anopheles arabiensis* and *Anopheles funestus*. Parasit Vectors. 2013;6:104.
75. Muriu SM, Coulson T, Mbogo CM, Godfray HCJ. Larval density dependence in *Anopheles gambiae* s.s., the major African vector of malaria. Boots M, editor. J Anim Ecol. 2013;82:166-74.
76. Yang GJ, Brook BW, Whelan PI, Cleland S, Bradshaw CJA. Endogenous and exogenous factors controlling temporal abundance patterns of tropical mosquitoes.

Ecol Appl. John Wiley & Sons, Ltd; 2008;18:2028-40.

77. Hancock PA, White VL, Callahan AG, Godfray CHJ, Hoffmann AA, Ritchie SA. Density-dependent population dynamics in *Aedes aegypti* slow the spread of wMel Wolbachia. J Appl Ecol. John Wiley & Sons, Ltd; 2016;53:785-93.

78. Porretta D, Mastrantonio V, Crasta G, Bellini R, Comandatore F, Rossi P, et al. Intra-instar larval cannibalism in *Anopheles gambiae* (s.s.) and *Anopheles stephensi* (Diptera: Culicidae). Parasites and Vectors. BioMed Central; 2016;9:1-9.

79. Tuno N, Githeko A, Yan G, Takagi M. Interspecific variation in diving activity among *Anopheles gambiae* Giles, *An. arabiensis* Patton, and *An. funestus* Giles (Diptera: Culicidae) larvae. J Vector Ecol. 2007;32:112-7.

80. Matthews J, Bethel A, Osei G. An overview of malarial *Anopheles* mosquito survival estimates in relation to methodology. Parasites and Vectors. BioMed Central; 2020;13:1-12.

81. Takken W, Charlwood JD, Billingsley PF, Gort G. Dispersal and survival of *Anopheles funestus* and *A. gambiae* s.l. (Diptera: Culicidae) during the rainy season in southeast Tanzania. Bull Entomol Res. 1998;88:561.

Figures

Figure 1

Schematic representation of the state-space population model showing different life stages compartment (circles) and flows (arrows) of *Anopheles funestus*. Abundance data were only available for unfed, blood-fed and gravid stages. The model assumes that once a gravid mosquito has laid eggs, they return to the unfed stage. The annotations are described in Table 1. The model incorporates six life stages (eggs, larvae, pupae, unfed, bloodfed and gravid) of *An. funestus*.

Figure 2

Reconstruction of the abundance trajectories for all the six life-stages. The red line indicates the mean posterior values and the respective 95% confidence intervals are shown in “sky-blue”. Left column (a,c,e,g,i,k) is data collected from June 2018 to May 2019 and right column (b,d,f,h,j,l) is data collected from Jan-Dec 2015. The grey area indicates the period with rainfall.

Figure 3

Observed vs. model estimated values for the three adult stages with data collected using CDC light trap both in May 2018 – June 2019 (left column- a,c,d) and Jan-Dec 2015 (right column- b,d,e). Red lines are the model estimated trajectories with “sky-blue” showing their 95% credible intervals. The blue circles are the observed values from the Light trap catches. Grey areas are the periods with rainfalls episodes.

Figure 4

Reconstruction of the survival trajectories for all the four stages (larvae, unfed, bloodfed and gravid) which were affected by the environmental covariates with the two bottom row showing the larval development period and fecundity trends. Left column (a,c,e,g,i,k) is trajectories from June 2018 to May 2019 and right column (b,d,f,h,j,l) is from Jan-Dec 2015. Grey area is the period with rainfall. Y-axis shows the survival rates of different life stages and the bottom row (k&l) shows per-capita fecundity

Figure 5

Relationship between environmental covariates and fitness parameters as estimated from the SSMs of population dynamic of *Anopheles funestus*.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [NgowoetalSupplementaryInformation.pdf](#)