

Relationship Between Parents' Health Literacy and Children's Sleep Problems: A Multicenter Cross-Sectional Study in Japan

Sae Ono

Kobe University

Hiroto Ogi

Kobe University

Masato Ogawa

Kobe University

Daisuke Nakamura

Fudousan Technologies Corporation

Teruhiko Nakamura

Educational Corporation Tsukushi Gakuen

Kazuhiro Izawa (✉ izawapk@harbor.kobe-u.ac.jp)

Kobe University

Research Article

Keywords: parents, health literacy, children, sleep disorder, preschoolers

Posted Date: December 10th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-121155/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Public Health on April 24th, 2021. See the published version at <https://doi.org/10.1186/s12889-021-10864-z>.

Abstract

Background: Sleep disorder in preschool children can stunt their health and growth. However, the factors that cause sleep disorders in children are not well understood. The aim of this study was to determine the relationship between parents' health literacy (HL) and children's sleep problems. The study was conducted at two kindergartens, two nursery schools, and a center for early childhood education in Chitose-city, Hokkaido, Japan.

Method: This study used a multicenter cross-sectional design. The sample comprised 354 preschoolers (aged 3–6 years) and their parents. In families with two or more children attending the same facility, only the oldest child was asked to participate in the study. Exclusion criteria included participants whose completed questionnaires had missing values. Children's sleep disorder was assessed using the Japanese version of the Children's Sleep Habits Questionnaire (CSHQ-J). Parents' HL was assessed using the 14-item Health Literacy Scale (HLS-14). The parents were classified into two groups (high HL group and low HL group). Multiple regression modelling was used to determine the association between HLS-14 and CSHQ-J scores.

Results: Of the 354 parents, 255 (72%) were in the high HL group and 99 (28%) in the low HL group. The mean CSHQ-J score was significantly lower in the high HL group than in the low HL group (45.3 ± 6.0 points vs. 46.8 ± 5.9 points, $p = 0.043$). In multiple regression analyses, CSHQ-J was independently associated with parents' HL after adjusting for all confounding factors ($R^2 = 0.41$, $\beta = -0.11$; $p = 0.0043$).

Conclusions: Parents' HL appears to affect their children's sleep disorder. This finding suggests that parents' HL may be a target for intervention to improve children's sleep disorders.

Introduction

Given that the preschool age is a period of rapid functional and cognitive development, sleep in preschoolers is of critical importance.[1, 2] It has been reported that preschoolers in Asian countries have significantly later bedtime and shorter total sleep duration compared with other Western countries. Particularly among the Asian countries, Mindell et al. revealed that preschoolers in Japan have the shortest sleep time.[3] Sleep is important not only in its quantity, but also in terms of quality and habits among all generations, including preschoolers. Sleep deficiency is an all-encompassing definition that includes sleep quality, sleep quantity, and sleep habits.[4] Sleep disorders among preschoolers aged 4–5 years are associated with anxiety,[5] and among children aged 4–11 years, are associated with behavior problems and anxious/depressed mood.[6] Additionally, sleep problems among children aged 4–5 years predict poor health-related quality of life, behavior, language, and learning scores at age 6–7 years.[7] Of note, in a previous study, sleep disorders at age 4 predicted adolescent depression and anxiety, and the strength of the relationship between sleep disorders and anxiety or depression increased with age.[8] In summary, sleep disorders in preschoolers cause poor health outcomes in both the short-term and the long-term.

Health literacy (HL) is an important aspect of the provision of modern healthcare.[9] HL is defined as “the personal skills that enables individuals to obtain, understand, and use information to make decisions and take actions that will have an impact on their health”.[10] Low HL in adults has been associated with poor health outcomes such as obesity,[11] underweight,[12] and depression.[13] Further, low HL among parents has been reported to be associated with poor health outcomes for their children, such as poor body mass index (BMI),[14] low sleep duration,[15] and oral health deterioration.[16] In summary, parents’ HL is not only associated with their own health outcomes, but also their children’s health outcomes. Previous studies have suggested that parents play an important role in preschoolers’ sleep.[17, 18] Many previous studies have investigated the relationship between parents’ HL and children’s sleep duration; however, very few studies have focused on all aspects of sleep, including the regularity and quality of sleep.[18] In addition, the relationship between parents’ HL and sleep disorders among children is not clear. Therefore, we hypothesized that if parents have low HL, their children are more likely to have sleep disorders.

Clarifying these relationships will serve as a breakthrough for improving children's sleep disorders and, ultimately, their long-term health outcomes through interventions designed for their parents. The purpose of the present study was to assess the relationship between parents’ HL and sleep problems in their children.

Method

Participants

This was a cross-sectional multicenter study of children aged 3–6 years at two kindergartens, two nursery schools, and a Center for Early Childhood Education and Care in Chitose-city, Hokkaido, Japan. Questionnaires were distributed in February 2020, inviting preschool-aged children and their parents to participate in the study. Participation was completely anonymous and voluntary, and the final dataset was fixed after consultation with the heads of the multicenter facilities in October 2020.

In families with two or more children attending the same facility, only the oldest child was asked to participate in the study. Exclusion criteria included participants whose completed questionnaires had missing values.

The study procedures were conducted in accordance with the Declaration of Helsinki and the Good Clinical Practice guidelines. This study was approved in advance by the Research Ethics Committee of Kobe University (approval number: 498-2). All participants were notified about their participation in this study and it was explained to them that they could withdraw their participation at any time.

Demographic data

Data were self-reported by the parents for themselves and their children. The following is a list of the questionnaire items used for children: age (in months); sex (boy/girl); tooth decay (without decay, with decay); birth weight; number of siblings; breakfast (not everyday, everyday); lessons; sleep duration; and gaming and TV hours. BMI and the International Obesity Task Force (IOTF)[19] criteria were used to

determine children's weight status. BMI was calculated as the weight in kilograms divided by the height in meters squared. The IOTF classifies BMI as the norm to assess the prevalence of overweight and underweight conditions in children.

The following is a list of the questionnaire items used for parents: age (in years); sex (male, female); BMI; smoking and drinking habit; marital status (not married, married); education level (years); household income (≤ 6 million yen, > 6 million yen); sleep duration; PSQI (Pittsburgh Sleep Quality Index).[20]

Health literacy in parents

The 14-item Health Literacy Scale (HLS-14)[21] developed in Japan was used to assess parents' HL. A previous study has shown that the HLS-14 is an adequate instrument to assess comprehensive HL.[21] Each item is answered on a 5-point scale. The total score is calculated by adding the scores of all items. Higher scores indicate more comprehensive HL. The validity and reliability of this tool has already been demonstrated. Participants were divided into two groups: high HL group (HLS-14 score > 50) and low HL group (HLS-14 score ≤ 50).[21]

Sleep problems in children

Sleep problems among children were assessed using the Japanese version of the Children's Sleep Habits Questionnaire (CSHQ-J).[22] It is one of the most widely used tools used to comprehensively screen for children's sleep disorders. The CSHQ-J requires parents to retrospectively report their children's sleep habits and behaviors. The scale includes 33 distinct items grouped into eight subscales with the following sleep domains: bedtime resistance, sleep onset delay, sleep duration, sleep anxiety, night wakings, parasomnias, sleep disordered breathing, and daytime sleepiness. Parents are asked to recall sleep behavior during a typical week and respond using the following options: 'Usually' (5–7 times/week), 'Sometimes' (2–4 times/week), and 'Rarely' (0–1 times/week). Higher scores indicate more sleep disorders.

Statistical analysis

As mentioned above, we classified the study participants into two groups: a high HL group and a low HL group; this has been described in detail elsewhere.[21] Statistical analyses were conducted after confirming that the data were normally distributed using the Shapiro-Wilk test. The differences in the clinical characteristics between the two groups were determined using an unpaired *t*-test or chi-square test. Correlations between parents' HL and the CSHQ-J scores were assessed using Pearson's correlation coefficient. Multiple linear regression analysis was carried out to determine whether parents' HL was associated with the children's sleep states. The dependent variable was parents' HL, whereas the independent variable was CSHQ-J or sleep duration of children. Multiple linear regression analyses were adjusted for characteristics of children and parents. All statistical analyses were performed using EZR (Saitama Medical Center, Jichi Medical University, Saitama, Japan), which is a graphical user interface for R (The R Foundation for Statistical Computing, Vienna, Austria). Differences and correlations were considered significant when $p < 0.05$.

Results

Figure 1 shows the participant flow during this study. We distributed the questionnaires to 537 parents, of whom 362 (67.4%) agreed to participate in this study. Parents whose questionnaire responses had missing values ($n = 8$) were excluded. After the exclusions, the final sample for the analysis consisted of 354 parents and their children.

Table 1 shows the demographic differences between the low and high HL groups. The number of participants in the high HL group was 255 (72.0%) and that in the low HL group was 99 (28.0%). The average CSHQ-J score was significantly lower in the high HL group (45.3 ± 6.0 points) than in the low HL group (46.8 ± 5.9 points) ($p = 0.043$). In addition, parents' BMI ($p = 0.032$) and years of education ($p = 0.032$) were higher in the high HL group than in the low HL group.

Table 1
Differences in Characteristics between High Health Literacy and Low Health Literacy Groups

Characteristic	High HL (n = 255)	Low HL (n = 99)	<i>t</i> or χ^2	<i>P</i> value
Child				
Age (months)	64.6 ± 10.3	63.3 ± 10.2	1.07	0.287
Sex (boys (%))	133 (52.2)	56 (56.6)	0.35	0.553
Teeth decay (non (%))	203 (79.6)	80 (80.8)	1.95	0.136
Birth weight (g)	2987.4 ± 445.1	3065.9 ± 399.9	-1.53	0.127
Number of siblings (n)	2.3 ± 0.8	2.2 ± 0.8	1.33	0.186
Breakfast (every day (%))	240 (94.1)	91 (91.9)	0.26	0.474
Lessons (%)	111 (43.5)	43 (43.4)	0	1.00
Sleep duration (hours)	10 ± 1.0	9.9 ± 1.1	1.21	0.228
Game (hours)	0.5 ± 0.8	0.5 ± 0.7	0.84	0.401
TV (hours)	2.1 ± 1.2	2.1 ± 1.2	0.22	0.826
BMI (kg/m ²)	15.6 ± 1.5	15.6 ± 1.5	-0.13	0.896
IOTF (problem (%))	50 (19.6)	20 (20.2)	0	0.882
CSHQ-J (point)	45.3 ± 6.0	46.8 ± 5.9	-2.04	0.043
Parent				
Age (years)	36.1 ± 5.1	35.9 ± 5.8	0.41	0.685
Sex (men (%))	17 (6.7)	5 (5.1)	0.10	0.806
BMI (kg/m ²)	21.1 ± 2.9	22.0 ± 4.0	-2.15	0.032
Smoking (%)	36 (14.1)	10 (10.1)	0.69	0.380
Drinking (everyday (%))	34 (13.3)	7 (7.1)	2.15	0.137
Material status (married (%))	238 (93.3)	96 (97.0)	2.01	0.699
Education level (years)	13.6 ± 1.7	13.1 ± 1.5	2.15	0.032
Income (> 6 million (%))	97 (38.0)	28 (28.3)	2.45	0.106

Data are expressed as mean ± SD or number (percentage)

HL: health literacy; BMI: body mass index; IOTF: International Obesity Task Force; CSHQ-J: The Japanese version of Children's Sleep Habits Questionnaire; PSQI: Pittsburgh Sleep Quality Index.

Characteristic	High HL (n = 255)	Low HL (n = 99)	<i>t</i> or χ^2	<i>P</i> value
Sleep duration (hours)	6.8 ± 1.2	6.9 ± 1.2	-0.48	0.633
PSQI (alright (%))	165 (64.7)	53 (53.5)	3.26	0.066
Data are expressed as mean ± SD or number (percentage)				
HL: health literacy; BMI: body mass index; IOTF: International Obesity Task Force; CSHQ-J: The Japanese version of Children's Sleep Habits Questionnaire; PSQI: Pittsburgh Sleep Quality Index.				

The correlation between parents' HL and children's sleeping habits is illustrated in Fig. 2. There was a significantly negative correlation between parents' HL and CSHQ-J ($p = 0.0378$, $r = -0.11$).

Table 2 shows the results of the regression analyses. In the univariate regression analyses, CSHQ-J scores were statistically significantly associated with parents' HL; nevertheless, children's sleep duration did not show a statistically significant relationship with parents' HL. In the multiple regression analysis, CSHQ-J remained statistically significant in predicting parents' HL after adjustment for all other confounding factors such as children's age, sex, IOTF, and number of siblings ($R^2 = 0.21$, $\beta = -0.12$; $p = 0.03$), but not the sleep duration of children ($R^2 = 0.16$, $\beta = 0.10$; $p = 0.08$) (Model 1).

Table 2. Regression analysis of sleep behavior and parents' health literacy

	Univariate model			Model 1 ^a			Model 2 ^b		
	R^2	β	<i>p</i> value	R^2	β	<i>p</i> value	R^2	β	<i>p</i> value
CSHQ-J	0.12	-0.11	0.038	0.21	-0.12	0.03	0.41	-0.11	0.043
Sleep duration of children	0.06	0.08	0.16	0.16	0.10	0.08	0.28	0.05	0.45

CSHQ-J: The Japanese version of Children's Sleep Habits Questionnaire; R^2 : Coefficient of determination; β : Regression coefficient

^a Model 1 was adjusted for children's age, sex, IOTF, and number of siblings

^b Model 2 was adjusted for Model 1 + parent's age, sex, BMI, educational status, household income, and smoking status

In addition to Model 2, CSHQ-J remained statistically significant for predicting parents' HL after adjustment for parents' age, sex, BMI, years of education, household income, and smoking status ($R^2 = 0.41$, $\beta = -0.11$; $p = 0.043$), but not the sleep duration of children ($R^2 = 0.28$, $\beta = 0.05$; $p = 0.45$).

Discussion

To our knowledge, this is the first study to assess the relationship between parents' HL and children's sleep disorders. Consistent with our hypothesis, we found a statistically significant relationship between parents' HL and sleep disorders in children.

There are two possible reasons for the association between parents' HL and sleep disorder in children. First, parents with low HL may have mental health issues and parenting-related stress. For example, a study suggested that people with low HL were more likely to be depressed.[23] Another study suggested that parents of children with sleep disorders were more likely to have parenting stress, mental health issues, and depression.[24] Therefore, issues related to the mental health of parents with low HL may affect their children's sleep disorders.

Second, parents with low HL may be likely to create inappropriate living conditions and engage in parenting behaviors unfavorable to children's sleep. A previous study suggested that low HL in parents is associated with suboptimal parenting practices and inadequate parenting behavior, such as putting a TV in the child's bedroom.[18]

Another study suggested that parents' bedtime and the frequency of parents' presence at the children's bedtime are linked to children's sleep problems.[25] Therefore, the behaviors of parents with low HL may play a role in their children's sleep disorders.

The results of the present study differed from those of previous studies in some respects. Our results revealed that the BMI of parents was significantly lower in the low HL group than in the high HL group; however, there was no significant difference in the children's BMI between the two groups. Nakamura et al. [14] showed that the HL of parents appears to affect the BMI – overweight and underweight – of their children independently of the parents' own BMI. A possible explanation is that the present study treated children's BMI as a continuous variable, which may have failed to adequately treat the poor BMI group. Another possibility could be that we did not measure other factors that affect children's BMI, such as exercise, nutrition, and chronic disease. A previous study showed that children's physical activity and eating behavior affect BMI.[26] In the present study, parents' HL was independently associated with children's sleep disorders, but not with their sleep duration. Ogi et al.[15] showed that parents' HL was correlated with children's sleep duration only in parents with low HL. A possible explanation is that the measurement of sleep duration in the present study was not robust.

In the present study, sleep duration was not measured directly but was reported by the parents. Previous studies have shown that children's parent-measured sleep duration overestimated actual sleep duration by approximately 30 minutes.[27] Therefore, children's sleep duration reported by their parents may not always be accurate and the results may be highly variable depending on the individual reporting. Another possible explanation is that unlike CSHQ-J, which includes several items, sleep duration is a single item. Therefore, it is likely that the CSHQ-J revealed differences between the high HL and low HL parent groups but did not expose the differences between the two groups in terms of sleep duration.

On the other hand, parents' BMI and years of education were associated with parents' HL in the present study. The association between parents' BMI and HL is likely to be due to the parents' own health behaviors. Previous studies have shown that people with high HL are less likely to have unhealthy habits. [28] Additionally, a number of studies have shown a link between HL and educational levels.[29] It is thought that this is because education may improve the ability to read, analyze, and take actions based on information. Higher education may lead to better access to health-related information.

According to Fig. 2, the parents' HL scores were significantly but negatively correlated with the CSHQ-J scores, albeit the correlation was weak ($r = -0.11$; weak: $r < 0.4$, moderate: $0.4 \leq r < 0.6$, strong: $0.6 \leq r < 0.8$). Although the correlation between the parents' HL scores and the CSHQ-J scores was not strong, the relationship was statistically significant. This suggests that improving parents' HL will lead to improvements in children's sleep disorders. In fact, there are many reports of HL interventions causing improvements in HL.[30] HL education might improve sleep quality but not sleep duration. More studies are required to examine the effectiveness of intervention for parents' HL in the future. Further, although the present study dealt with the total CSHQ-J scores, investigating each item on the CSHQ-J may provide an opportunity to identify the causes of sleep disorders in children.

There are several limitations to the present study. First, as this is a cross-sectional study, a causal relationship between parents' HL and other factors cannot be concluded. Second, a sampling bias may have occurred due to the small sample size and the fact that the data were collected in one particular area, that is, Chitose City, Hokkaido. Finally, it is possible that we did not collect enough information relevant to parents' HL. This is evident from Table 2, which shows $R^2 = 0.41$, which means that when we fed in all the factors, only 41% of the parents' HL was predicted by the factors we studied.

Conclusion

The findings of this study suggest that parents' HL independently affects the sleep disorders of their children after adjusting for all other confounding factors. These results suggest that interventions targeting parents' HL are needed to improve children's sleep disorders. Additionally, children's sleep disorders need to be examined in more detail.

List Of Abbreviations

CSHQ-J: The Japanese version of Children's Sleep Habits Questionnaire

HLS-14: The 14-item Health Literacy Scale

Declarations

Ethics approval and consent to participate:

The study procedures were conducted in accordance with the Declaration of Helsinki and the Good Clinical Practice guidelines. This study was approved in advance by the Research Ethics Committee of Kobe University (approval number: 498-2). Written informed consent was obtained from all participants. All participants were notified about their participation in this study and it was explained to them that they could withdraw their participation at any time.

Consent for publication:

Not Applicable.

Availability of Data and information:

All relevant data are present within the paper. Furthermore, if additional data or permission is needed, for example, for use in a meta-analysis, it can be made available from corresponding author for researchers who meet the criteria for access to confidential data.

Competing interests' statement:

The authors declare that they have no competing interests

Funding

This work was supported by Mayekawa Foundation, Tokyo, Japan (<https://www.mayekawafoundation.org/en/>).

Authors' contributions

Sae Ono wrote the first draft of the manuscript. Hiroto Ogi and Masato Ogawa developed the methodological approach, interpreted the data and wrote the discussion. Kazuhiro P. Izawa revised the article. Daisuke Nakamura and Teruhiko Nakamura helped to develop the study design and all study materials including the study protocol. All authors conceived the study and revised and approved the final manuscript.

Acknowledgments

We express our gratitude to the parents, children, and school staff at the Tsukushi Gakuen in Hokkaido who agreed to participate and cooperate in this study. This study benefited from the support and encouragement of Mr. Kodai Ishihara, Dr. Masashi Kanai, Dr. Masahiro Kitamura, Ms. Akemi Ogura, Mr. Ikko Kubo, Mr. Yuji Kanejima, and Dr. Shinichi Shimada, Graduate School of Health Sciences, Kobe University.

References

1. Reynaud E, Vecchierini MF, Heude B, Charles MA, Plancoulaine S. Sleep and its relation to cognition and behaviour in preschool-aged children of the general population: a systematic review. *J Sleep Res.* 2018;27(3):e12636.
2. Black MM, Walker SP, Fernald LCH, Anderson CT, DiGirolamo AM, Lu, C, Mc Coy, DC, et al. Early childhood development coming of age: science through the life course. *Lancet* 2017;389(10064):77–90.
3. Mindell JA, Sadeh A, Wiegand B, How TH, Goh DY. Cross-cultural differences in infant and toddler sleep. *Sleep Med.* 2010;11(3):274–80.
4. Czeisler CA. Impact of sleepiness and sleep deficiency on public health—utility of biomarkers. *J Clin Sleep Med.* 2011;7(5 Suppl):S6–8.
5. Brown WJ, Wilkerson AK, Boyd SJ, Dewey D, Mesa F, Bunnell BE. A review of sleep disturbance in children and adolescents with anxiety. *J Sleep Res.* 2018;27(3):e12635.
6. Coulombe JA, Reid GJ, Boyle MH, Racine Y. Concurrent associations among sleep problems, indicators of inadequate sleep, psychopathology, and shared risk factors in a population-based sample of healthy Ontario children. *J Pediatr Psychol.* 2010;35(7):790–9.
7. Quach J, Hiscock H, Canterford L, Wake M. Outcomes of child sleep problems over the school-transition period: Australian population longitudinal study. *Pediatrics.* 2009;123(5):1287–92.
8. Gregory AM, O'Connor TG. Sleep problems in childhood: a longitudinal study of developmental change and association with behavioral problems. *J Am Acad Child Adolesc Psychiatry.* 2002;41(8):964–71.
9. Bitzer EM, Sørensen K. [Health Literacy]. *Gesundheitswesen.* 2018;80(8–09):754–66.
10. Nutbeam D, Levin-Zamir D, Rowlands G. Health Literacy in Context. *Int J Environ Res Public Health.* 2018;15(12), 2657.
11. Friis K, Vind BD, Simmons RK, Maindal HT. The relationship between health literacy and health behaviour in people with diabetes: a Danish population-based study. *J Diabetes Res.* 2016;2016:7823130.
12. Paasche-Orlow MK, Wolf MS. Promoting health literacy research to reduce health disparities. *J Health Commun.* 2010;15 Suppl 2:34–41.
13. Berkman ND, Sheridan SL, Donahue KE, Halpern DJ, Vierra A, Krotty K, et al. Health literacy interventions and outcomes: an updated systematic review. *Evid Rep Technol Assess (Full Rep).* 2011;199(1):941.
14. Nakamura D, Ogawa M, Nakamura T, Izawa KP. Impact of parents' comprehensive health literacy on BMI in children: a multicenter cross-sectional study in Japan. *J Sch Health.* 2018;88(12):910–6.
15. Ogi H, Nakamura D, Ogawa M, Nakamura T, Izawa KP. Associations between parents' health literacy and sleeping hours in children: a cross-sectional study. *Healthcare (Basel).* 2018;6(2):32.
16. Miller E, Lee JY, DeWalt DA, Vann WF. Impact of caregiver literacy on children's oral health outcomes. *Pediatrics.* 2010;126(1):107–14.

17. Durduran Y, Pekcan S, Çolpan B. Sleep habits and related factors in kindergarten children. *Niger J Clin Pract.* 2019;22(9):1218–23.
18. Bathory E, Tomopoulos S, Rothman R, Sanders L, Perrin EM, Mendelsohn A, et al. Infant sleep and parent health literacy. *Acad Pediatr.* 2016;16(6):550–7.
19. Cole TJ, Lobstein T. Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes.* 2012;7(4):284–94.
20. Buysse DJ, Reynolds CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res.* 1989;28(2):193–213.
21. Suka M, Odajima T, Kasai M, Igarasi A, Ishiwara H, Kusama M, et al. The 14-item health literacy scale for Japanese adults (HLS-14). *Environ Health Prev Med.* 2013;18(5):407–15.
22. Owens JA, Spirito A, McGuinn M. The Children's Sleep Habits Questionnaire (CSHQ): psychometric properties of a survey instrument for school-aged children. *Sleep.* 2000;23(8):1043–52.
23. Cheng ER, Bauer NS, Downs SM, Sanders LM. Parent health literacy, depression, and risk for pediatric injury. *Pediatrics.* 2016;138(1):e20160025.
24. Byars KC, Yeomans-Maldonado G, Noll JG. Parental functioning and pediatric sleep disturbance: an examination of factors associated with parenting stress in children clinically referred for evaluation of insomnia. *Sleep Med.* 2011;12(9):898–905.
25. Bonuck KA, Schwartz B, Schechter C. Sleep health literacy in head start families and staff: exploratory study of knowledge, motivation, and competencies to promote healthy sleep. *Sleep Health.* 2016;2(1):19–24.
26. Verjans-Janssen SRB, van de Kolk I, Van Kann DHH, Kremers SPJ, Gerards SMPL. Effectiveness of school-based physical activity and nutrition interventions with direct parental involvement on children's BMI and energy balance-related behaviors— A systematic review. *PLoS One.* 2018;13(9):e0204560.
27. Nelson TD, Lundahl A, Molfese DL, Waford RN, Roman A, Gozal D, et al. Estimating child sleep from parent report of time in bed: development and evaluation of adjustment approaches. *J Pediatr Psychol.* 2014;39(6):624–32.
28. Suka M, Odajima T, Okamoto M, Sumitani M, Igarashi A, Ishikawa H, et al. Relationship between health literacy, health information access, health behavior, and health status in Japanese people. *Patient Educ Couns.* 2015;98(5):660–8.
29. Mantwill S, Monestel-Umaña S, Schulz PJ. The relationship between health literacy and health disparities: a systematic review. *PLoS One.* 2015;10(12):e0145455.
30. Nutbeam D, McGill B, Premkumar P. Improving health literacy in community populations: a review of progress. *Health Promot Int.* 2018;33(5):901–11.

Figures




Figure 1

Participant flow during the study




Figure 1

Participant flow during the study




Figure 2

Correlation between parents' health literacy and children's sleep disorder




Figure 2

Correlation between parents' health literacy and children's sleep disorder