Bacterial Sensitivity To Chlorhexidine and Povidone-Iodine Antiseptics Over Time: A Systematic Review and Meta-Analysis of Human-Derived Data

Raiyyan Aftab

Portsmouth Hospitals NHS Trust

Vikash H Dodhia

Royal Hampshire County Hospital

Christopher Jeanes

Norfolk and Norwich University Hospital

Ryckie G Wade (✉ ryckiewade@gmail.com)

University of Leeds

Research Article

Keywords:

Posted Date: January 12th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1214575/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Surgical site infection (SSI) is the most common complication of surgery, increasing healthcare costs and hospital stay. Chlorhexidine (CHX) and povidone-iodine (PVI) are used for skin antisepsis, minimising SSIs. There is increasing concern of developing resistance to topical biocides, though clinical implications remain unclear.

Outcomes:

Determining whether the Minimum Bactericidal Concentration (MBC) for topical preparations of CHX or PVI have changed over time, in microbes relevant to SSI.

Methods

We included studies reporting the mean bactericidal concentration (MBC) of laboratory and clinical isolates of common microbes to CHX and PVI. Excluding non-human samples and studies using antimicrobial solvents or mixtures with other active substances. MBC was pooled in random effects meta-analyses. Change in MBC over time was explored using meta-regression.

Results

79 studies were including, analysing 6218 microbes over 45 years. Most used CHX (93%), with insufficient data for meta-analysis of PVI. There was no change in MBC of CHX to *Staphylococci* (β 0.12 [-1.13, 1.37]; I^2 99%) or *Streptococci* (β 0.13 [-0.35, 0.62]; I^2 97%).

Conclusions

There is no evidence of reduced susceptibility of common SSI-causing microbes to CHX over time. Providing reassurance that the worldwide guidance that CHX should remain the first-choice agent for surgical skin antisepsis.

Introduction

Surgical site infection (SSI) is the most common and costly complication of surgery^{1,2}, occurring in approximately 5% of all surgical interventions³. They represent an important economic burden across all surgical specialties⁴, increasing hospital inpatient stay time and adversely affecting patient's mental and

physical health.⁴ The use of skin antisepsis prior to surgery significantly reduces the risk of SSI and consequently, post-operative morbidity and mortality^{5–7}.

Staphylococcus aureus and *Streptococci* spp. are commonly implicated microbes in SSI, along with *Enterococcus* spp. and *Escherichia coli*⁸. To reduce the risk of SSI, the World Health Organisation (WHO)⁷, United States of America Centre for Disease Control (CDC)⁹ and United Kingdom National Institute for Health and Care Excellence (NICE)¹⁰ recommend the application of topical chlorhexidine (CHX) in alcohol to the planned operative site, for skin antisepsis. CHX in an alcoholic solvent has been shown to halve the risk of SSI following clean¹¹, contaminated^{12,13} and dirty surgery when compared to other antiseptics such as povidone-iodine (PVI).

CHX is a biguanide compound, utilised both as a broad spectrum antimicrobial and topical antiseptic¹⁴. By binding to the cell membrane and cell wall of bacteria, at lower concentrations it has a bacteriostatic effect by displacing the cations and destabilising the cell wall. At higher concentrations there is a complete loss of cellular structural integrity, having a bactericidal effect.¹⁵ PVI is an iodophor; a chemical complex between a water soluble povidone polymer and iodine¹⁶. When dissolved in water, iodine is released, penetrating microorganisms and oxidising proteins, nucleotides and fatty acids¹⁷ causing cell death. Both PVI and CHX are active against gram positive and negative bacteria, fungi and viruses^{16,18}.

There are growing fears that as antiseptic use increases, sensitivity may reduce and resistance emerge¹⁹. This is particularly concerning given the accelerating global antibiotic resistance crisis²⁰. Multiple bacteria have shown reduced sensitivity (perhaps even resistance) to CHX, particularly through multidrug resistance efflux protein *qacA*¹⁹. Methicillin resistant *Staphylococcus aureus* (MRSA) samples with *qacA/B* genes showed persistent MRSA carriage despite de-colonisation therapy²¹. Not only is the presence of reduced sensitivity concerning, but the presence of resistance conferring genes seems to be increasing annually²². PVI resistance has been less commonly reported²³ although this might be secondary to the multimodal effect of iodine on microbes²³ or otherwise. Overall, the current state of microbial sensitivity to CHX and PVI remains unclear.

The aim of this review is to summarise the sensitivity profiles of skin microbes (relevant to surgical site infection) to CHX and PVI, and explore how these have changed over time

Methods

This review was designed and conducted in accordance with the Cochrane Handbook of Systematic Reviews²⁴, the protocol was published in the PROSPERO database (CRD42021241089) and the report has been authored in accordance with the PRISMA checklist²⁵.

Types of Studies

We included all studies which reported the resistance of microbes to CHX or PVI based topical biocides derived from human samples. There were no language restrictions. We excluded case reports and studies which used antimicrobial solvents (e.g., alcohol) or mixtures of antiseptics (e.g. chlorhexidine mixed with cetrimide).

Search strategy

The NICE Healthcare Databases (hdas.nice.org.uk) was searched according to appendix 1 (supplementary materials). The medRxiv and bioRxiv preprint archives were searched with the same strategy using the R package `medrxivr`²⁶. This yielded 582 hits in PubMed, 720 in Embase, 993 in Web of Science and 896 in CINAHL. After de-duplication, there were 2318 unique citations which were screened (Fig 1). A further 3 articles were found by manual searching of these articles.

Study selection

Two review authors (RA and VHD) independently screened titles and abstracts for relevance, in accordance with the eligibility criteria. The full texts of potentially eligible articles were obtained and again independently assessed by the same authors. Disagreements were resolved by discussion with RGW.

Data extraction

Two review authors (RA and VHD) independently double extracted data. The colony was the unit of analysis. Where data was missing or unclear, the corresponding author was contacted by email and if no reply was received, these values were estimated from the available data²⁷.

Outcomes

The outcome of interest was the Minimum Bactericidal Concentration (MBC). MBC was chosen over the minimum inhibitory concentration (MIC) as a measure of susceptibility because it is generally felt to be a more appropriate for topical antiseptics^{15,28}. We were interested in deriving a pooled estimate of the MBC for different species to understand how this may have changed over time.

Methodological quality assessment

The risk of bias was not assessed because there are no validated tools available for studies of this nature and the study selection process ensured that only the highest quality research was included.

Missing data

After estimating some parameters, the overall rate of missing data (in the predictor variables) was 14.5%. Importantly, the variance of the mean MBC was missing at random in 36 observations (61%) and because this was required for the primary analyses, we imputed this data using chained equations^{29,30}.

Statistical analysis

The raw data are available open-source at <https://osf.io/khnb2>. Using metafor^{31,32}, 5 studies^{33–37} reporting the MBC of CHX for *Staphylococci* were identified as outliers (based on externally standardised residuals) or influential studies (based on the Cook's distance and leave-one-out values of the test statistics for heterogeneity), and their 95% confidence intervals (CIs) were far outside the 95% CI of the pooled estimate, so they were excluded from the meta-analyses. Data were then analysed in Stata/MP v16 (StataCorp LLC, Texas) using the meta suite. To synthesise a pooled MBC, we used a random-effects meta-analyses. We sub-grouped by the microbial family. To estimate change in MBC over time, we performed meta-regression for *Staphylococci* and *Streptococci*, separately. A sensitivity multivariable meta-regression for *Staphylococci* was performed, controlling for methicillin-resistance as a binary co-variate. The REML estimator was used throughout. To align with calls for the abolition of p-values, we minimise their use and avoid the term “statistical significance”^{38,39}, instead focusing on how our findings may be clinically applicable and what might explain uncertainty in the estimates.

Results

Ultimately, 79 studies^{33–37,40–113} were included (Fig 1).

Study characteristics

The details of the included studies are summarised in S1 Table; readers who wish to know more detail should refer to the raw data (<https://osf.io/khnb2/>). The included studies originated from 24 countries. Articles were published between 1976 and 2021, although the majority (95%) were published this century. The antiseptics used included five different CHX salts (digluconate^{33,35,37,40,47,50,55–58,61,63,67,69,74,75,81,83,88,90,94,97,99,101,106,107,111,113}, gluconate^{34,36,42,42,53,91,102,103,108,108,109,112}, dichlorohydrate^{85–87}, diacetate^{71,99} and dihydrochloride⁹⁶) and povidone-iodine^{37,46,49,70,76,91}. In total, MBC data from 6218 microbes were extracted. The microbes tested are shown in S2 Table. Most samples were laboratory isolates (61%) and not multi-drug resistant (88%). The reporting standards used to establish the MBC were the according to the Clinical Laboratory Standards Institute (CLSI, 67%), European Committee on Antimicrobial Testing (EUCAST, 6%), German Institute for Standardisation (DIN, 5%), British Society for Antimicrobial Chemotherapy (BSAC, 3%) or International Organisation for Standardisation (ISO, 1%).

Evidence Synthesis

The MBC of CHX differed significantly between the families of microbes (Fig 2). Enterobacteriales had the highest MBC for CHX (20 mg/L [95% CI 14, 25]; I^2 95%) whilst MRSA had the lowest (3 mg/L [95% CI 1, 2]; I^2 93%).

Overall, 23 studies reported the mean MBC for *Staphylococci*; observations based on MSSA were more common^{41,44,51,59,61,67,68,74,88,93–95,102,106,107,110–113} than MRSA^{34,35,37,50,67,68,80,90,93,106,110,113}. The pooled mean MBC of CHX for *Staphylococci* was 5.93 mg/L (95% CI 3.09, 8.77; I^2 99%). Meta-regression showed no change in the MBC of CHX for *Staphylococci* over time (β 0.12 [-1.13, 1.37]; I^2 99%; Fig 3). When controlling for resistance to methicillin (MRSA vs MSSA), there was still no evidence of a change in the MBC over time (β 0.26 [-0.87, 1.34]; I^2 99%). Study level estimates for MSSA, MRSA and coagulase-negative *Staphylococci* are shown in S1 Fig.

Overall, 25 studies reported the MBC of CHX for *Streptococci* species; observations of viridans *Streptococci* were most common^{44,45,47,48,52–54,58,66,69,75,77–79,81,82,84–86,89,99,100,103,105,106} and 1 study¹⁰⁶ provided an estimate for *Streptococcus pyogenes* (Lancefield group A). The pooled mean MBC of CHX for *Streptococci* was 8.54 mg/L (95% CI 4.75, 12.3; I^2 99%). Meta-regression showed that the MBC of CHX for *Streptococci* had not changed over time (β 0.13 [-0.35, 0.62]; I^2 97%; Fig 4).

There were insufficient data for meta-analysis of the MBC of PVI. Also, the majority of MBC data for PVI was derived from studies of Enterobacteriales, which is not a common cause of surgical site infection and so equally, not clinically relevant.

Discussion

This review summarise the evidence to-date and suggests that there has been no increase in MBC of CHX for the main SSI-causing skin microbes in recent decades. The stability of CHX susceptibility is reassuring for clinicians and policy makers alike, as it endorses current surgical guidance worldwide advocates topical alcoholic chlorhexidine for skin asepsis prior to surgery.

The definition of bacterial susceptibility and resistance to biocides is still a matter of debate. Most clinically relevant bacteria have defined susceptibility and resistance to systemic antibacterials based upon the MIC relative to an epidemiologically derived clinical breakpoint. However, the use of MICs is less useful in determining the efficacy of topical antiseptics biocides given the desire to induce death of specific microbes rather than inhibition. Understanding the lethality of a biocide (hence MBC) is therefore a potentially more attractive measure than MIC for topical antiseptics¹¹⁴. Additionally, the relevance of both MICs and MBCs with respect to biocides has been questioned.¹⁵ Chlorhexidine for skin prep is used in concentrations of 5000 to 50000 $\mu\text{g/mL}$ and with MBC values ranging from 0 to 30 $\mu\text{g/mL}$, this is one thousand times greater than the apparent required concentration¹¹⁵.

MIC and MBC rely on attaining a steady concentration in bodily fluids as seen by the pharmacodynamics of antibiotics¹¹⁶. The EUCAST definition of a susceptible organism is “a microorganism is defined as susceptible by a level of antimicrobial activity associated with a high likelihood of therapeutic success”¹¹⁷. Therapeutic success in the context of topical CHX prior to surgery is disinfection; complete elimination of all relevant micro-organisms, except certain bacterial spores¹¹⁸. Measurement of MIC and

MBC are gained from *in vitro* susceptibility testing of microorganisms to topical biocides and therefore, provide little information as to the mechanism of resistance or likely clinical outcome. Therefore, our recommendation is the utilisation of epidemiological cut-off values (ECOFF) based on MBC distributions to better understand the response of bacteria to CHX in clinical practice. An analysis of ECOFF values of CHX to common bacterium, including those commonly causing SSI did not reveal a bimodal distribution, concluding that resistance is uncommon to CHX in natural populations of clinically relevant organisms⁸⁸. While values above a certain cut off may be defined as a breakpoint and hence resistant, this needs to be correlated with the clinical picture. Does CHX still achieve adequate disinfection in a population of bacterium with a MBC value greater than the 95% ECOFF? Without this information, our understanding of how MBC values beyond the normal distribution impacts clinical use remains poor. However this is contrary to the application of biocides in clinical settings prior to surgery, particularly during skin preparation of sensitive areas (e.g. the face) and in the presence of open wounds⁶. We speculate that real-time methods of quantifying bacterial colonisation (e.g. free bacterial DNA/RNA sequencing tools such as MinION), taken before and serially after topical application of CHX, may be more clinically useful in understanding the true bactericidal effect of CHX, what microbes it acts upon preferentially, how long it exerts its action and how this is achieved.

Limitations

Most of the included studies reported MIC rather than MBC, which meant that much data could not be synthesised. This might explain why our meta-data (for *Staphylococci* and *Streptococci* at least) disagrees with individual articles^{18,19,21,22}. Papers also often failed to report the type and concentration of chlorhexidine used which hindered meta-regression. Some papers clearly stated the year that the microbial strain was isolated, although this was often unclear and therefore the date of the paper was taken as the year of the isolate which may not adequately represent the change in MBC over time. This was also the case with location, so where possible the location of the laboratory or hospital was used as a surrogate. Isolates from a clinical setting are exposed to different selection pressures. As 61% of the microbes used were laboratory isolates as opposed to clinical isolates, it is unclear how our data can be generalised to clinical environments.

Conclusion

There has been no demonstrable change in the susceptibility of surgical site infection causing pathogens to chlorhexidine over time. A clear definition of reduced susceptibility and resistance of pathogens to biocides is needed, alongside consensus on the methods for measuring these phenomena.

Declarations

Contributions

RA and VHD contributed equally to this paper and are joint first authors. They co-designed the protocol, extracted data and co-authored the manuscript.

CJ provided clinical oversight for this study, supporting the design, advising on data extraction, analysis techniques and co-authored the manuscript.

RGW conceived the idea of this study, co-designed the protocol, supervised all elements of the delivery, extracted, and checked data, performed the statistical analysis and co-authored the manuscript

Funding

Ryckie Wade is a Doctoral Research Fellow funded by the National Institute for Health Research (NIHR, DRF-2018-11-ST2-028). The views expressed are those of the author(s) and not necessarily those of the United Kingdom's National Health Service, NIHR or Department of Health.

Conflicting Interests

None

Availability of data

The raw data are available via the Open Science Framework (<https://osf.io/khnb2/>). The statistical syntax are available from the senior author (RGW) upon request.

References

1. Gibson, A., Tevis, S. & Kennedy, G. Readmission after delayed diagnosis of surgical site infection: a focus on prevention using the American College of Surgeons National Surgical Quality Improvement Program. *Am. J. Surg.* **207**, 832–839 (2014).
2. Zimlichman, E. *et al.* Health Care–Associated Infections. *JAMA Intern. Med.* **173**, 2039 (2013).
3. Garner, B. H. & Anderson, D. J. Surgical Site Infections. *Infect. Dis. Clin. North Am.* **30**, 909–929 (2016).
4. Badia, J. M. *et al.* Impact of surgical site infection on healthcare costs and patient outcomes: a systematic review in six European countries. *J. Hosp. Infect.* **96**, 1–15 (2017).
5. Hirsch, T. *et al.* Antiseptics in surgery. *Eplasty* **10**, e39 (2010).
6. Wade, R. G., Burr, N. E., McCauley, G., Bourke, G. & Efthimiou, O. The Comparative Efficacy of Chlorhexidine Gluconate and Povidone-iodine Antiseptics for the Prevention of Infection in Clean Surgery: A Systematic Review and Network Meta-analysis. *Ann. Surg.* **Publish Ahead of Print**, (2020).
7. World Health Organization. *Global guidelines for the prevention of surgical site infection.* (2016).
8. Owens, C. D. & Stoessel, K. Surgical site infections: epidemiology, microbiology and prevention. *J. Hosp. Infect.* **70**, 3–10 (2008).

9. Berríos-Torres, S. I. *et al.* Centers for Disease Control and Prevention Guideline for the Prevention of Surgical Site Infection, 2017. *JAMA Surg.* **152**, 784 (2017).
10. National Institute for Health and Care Excellence (Great Britain) & National Guideline Centre (Great Britain). *Surgical site infections: prevention and treatment.* (2019).
11. Wade, R. G., Burr, N. E., McCauley, G., Bourke, G. & Efthimiou, O. The Comparative Efficacy of Chlorhexidine Gluconate and Povidone-iodine Antiseptics for the Prevention of Infection in Clean Surgery. *Ann. Surg.* **Publish Ah**, In press (2020).
12. Chen, S., Chen, J. W., Guo, B. & Xu, C. C. Preoperative Antisepsis with Chlorhexidine Versus Povidone-Iodine for the Prevention of Surgical Site Infection: a Systematic Review and Meta-analysis. *World J. Surg.* **44**, 1412–1424 (2020).
13. Ayoub, F., Quirke, M., Conroy, R. & Hill, A. Chlorhexidine-alcohol versus povidone-iodine for pre-operative skin preparation: A systematic review and meta-analysis. *Int. J. Surg. Open* **1**, 41–46 (2015).
14. Gilbert, P. & Moore, L. E. Cationic antiseptics: diversity of action under a common epithet. *J. Appl. Microbiol.* **99**, 703–715 (2005).
15. Horner C, Mawer D, & Wilcox M. Reduced susceptibility to chlorhexidine in staphylococci: is it increasing and does it matter? *J. Antimicrob. Chemother.* **67**, 2547–2559 (2012).
16. Lepelletier, D., Maillard, J. Y., Pozzetto, B. & Simon, A. Povidone Iodine: Properties, Mechanisms of Action, and Role in Infection Control and *Staphylococcus aureus* Decolonization. *Antimicrob. Agents Chemother.* **64**, (2020).
17. Lachapelle, J.-M. *et al.* Antiseptics in the era of bacterial resistance: a focus on povidone iodine. *Clin. Pract.* **10**, 579–592 (2013).
18. Cieplik, F. *et al.* Resistance Toward Chlorhexidine in Oral Bacteria – Is There Cause for Concern? *Front. Microbiol.* **10**, 587 (2019).
19. Kampf, G. Acquired resistance to chlorhexidine – is it time to establish an ‘antiseptic stewardship’ initiative? *J. Hosp. Infect.* **94**, 213–227 (2016).
20. O’Neill, J. Tackling Drug-Resistant Infections Globally: Final Report and Recommendations the Review on Antimicrobial Resistance. *Rev. Antimicrob. Resist.* (2016) doi:10.1016/j.jpha.2015.11.005.
21. Lee, A. S. *et al.* Impact of Combined Low-Level Mupirocin and Genotypic Chlorhexidine Resistance on Persistent Methicillin-Resistant *Staphylococcus aureus* Carriage After Decolonization Therapy: A Case-control Study. *Clin. Infect. Dis.* **52**, 1422–1430 (2011).
22. McNeil, J. C., Hulten, K. G., Kaplan, S. L., Mahoney, D. H. & Mason, E. O. *Staphylococcus aureus* Infections in Pediatric Oncology Patients: High Rates of Antimicrobial Resistance, Antiseptic Tolerance and Complications. *Pediatr. Infect. Dis. J.* **32**, 124–128 (2013).
23. Eggers, M. Infectious Disease Management and Control with Povidone Iodine. *Infect. Dis. Ther.* **8**, 581–593 (2019).

24. Higgins, J. P. T. & Green, S. (editors). Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0 [updated March 2011]. *Cochrane Collab.* (2011).
25. Moher, D., Liberati, A., Tetzlaff, J. & Altman, D. G. Systematic Reviews and Meta-Analyses: The PRISMA Statement. *Annu. Intern. Med.* **151**, 264–269 (2009).
26. McGuinness, L. & Schmidt, L. medrxiv: Accessing and searching medRxiv and bioRxiv preprint data in R. *J. Open Source Softw.* **5**, 2651 (2020).
27. Wan, X., Wang, W., Liu, J. & Tong, T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med. Res. Methodol.* **14**, 135 (2014).
28. Walker, E. M. & Lowes, J. A. An investigation into in vitro methods for the detection of chlorhexidine resistance. *J. Hosp. Infect.* **6**, 389–397 (1985).
29. Jakobsen, J. C., Gluud, C., Wetterslev, J. & Winkel, P. When and how should multiple imputation be used for handling missing data in randomised clinical trials – a practical guide with flowcharts. *BMC Med. Res. Methodol.* **17**, 162 (2017).
30. Kontopantelis, E., White, I. R., Sperrin, M. & Buchan, I. Outcome-sensitive multiple imputation: a simulation study. *BMC Med. Res. Methodol.* **17**, 2 (2017).
31. Viechtbauer, W. & Cheung, M. W.-L. Outlier and influence diagnostics for meta-analysis. *Res. Synth. Methods* **1**, 112–125 (2010).
32. Viechtbauer, W. Conducting Meta-Analyses in R with the metafor Package. *J. Stat. Softw.* **36**, (2010).
33. Biagi, M., Giachetti, D., Miraldi, E. & Figura, N. New non-alcoholic formulation for hand disinfection. *J. Chemother.* **26**, 86–91 (2014).
34. McGann, P. *et al.* Detection of qacA/B in clinical isolates of methicillin-resistant staphylococcus aureus from a regional healthcare network in the eastern United States. *Infect. Control Hosp. Epidemiol.* **32**, 1116–1119 (2011).
35. Eick, S., Radakovic, S., Pfister, W., Nietzsche, S. & Sculean, A. Efficacy of taurolidine against periodontopathic species—an in vitro study. *Clin. Oral Investig.* **16**, 735–744 (2012).
36. Knapp, L. *et al.* The effect of cationic microbicide exposure against Burkholderia cepacia complex (Bcc); the use of Burkholderia lata strain 383 as a model bacterium. *J. Appl. Microbiol.* **115**, 1117–1126 (2013).
37. Koburger, T., Hübner, N. O., Braun, M., Siebert, J. & Kramer, A. Standardized comparison of antiseptic efficacy of triclosan, PVP-iodine, octenidine dihydrochloride, polyhexanide and chlorhexidine digluconate. *J. Antimicrob. Chemother.* **65**, 1712–1719 (2010).
38. Wasserstein, R. L. & Lazar, N. A. The ASA Statement on p-Values: Context, Process, and Purpose. *Am. Stat.* **70**, 129–133 (2016).
39. Amrhein, V., Greenland, S. & McShane, B. Scientists rise up against statistical significance. *Nature* **567**, 305–307 (2019).
40. Alotaibi, S. M. I. *et al.* Susceptibility of vancomycin-resistant and -sensitive Enterococcus faecium obtained from Danish hospitals to benzalkonium chloride, chlorhexidine and hydrogen peroxide

- biocides. *J. Med. Microbiol.* **66**, 1744–1751 (2017).
41. Antonelli, A. *et al.* In vitro antimicrobial activity of the decontaminant hybenx compared to chlorhexidine and sodium hypochlorite against common bacterial and yeast pathogens. *Antibiotics* **8**, (2019).
 42. Arslan, S., Er, O., Ozbilge, H. & Kaya, E. G. In vitro antimicrobial activity of propolis, BioPure MTAD, sodium hypochlorite, and chlorhexidine on *Enterococcus faecalis* and *Candida albicans*. *Saudi Med. J.* **32**, 479–483 (2011).
 43. Azad, A., Rostamifar, S., Bazrafkan, A., Modaresi, F. & Rezaie, Z. Assessment of the antibacterial effects of bismuth nanoparticles against *enterococcus faecalis*. *BioMed Res. Int.* 2020, (2020).
 44. Baena-Santillan, E. S. *et al.* Comparison of the Antimicrobial Activity of Hibiscus sabdariffa Calyx Extracts, Six Commercial Types of Mouthwashes, and Chlorhexidine on Oral Pathogenic Bacteria, and the Effect of Hibiscus sabdariffa Extracts and Chlorhexidine on Permeability of the Bacterial Membrane. *J. Med. Food* (2020) doi:10.1089/jmf.2019.0273.
 45. Batubara, I., Wahyuni, W. T. & Susanta, M. Antibacterial activity of zingiberaceae leaves Essential oils against *streptococcus mutans* And teeth-biofilm degradation. *Int. J. Pharma Bio Sci.* **7**, (2016).
 46. Bhatia, M., Mishra, B., Thakur, A., Dogra, V. & Loomba, P. S. Evaluation of Susceptibility of Glycopeptide-resistant and Glycopeptide-sensitive *Enterococci* to Commonly Used Biocides in a Super-speciality Hospital: A Pilot Study. *J. Nat. Sci. Biol. Med.* **8**, 199–202 (2017).
 47. Caiaffa, K. S. *et al.* KR-12-a5 is a non-cytotoxic agent with potent antimicrobial effects against oral pathogens. *Biofouling* **33**, 807–818 (2017).
 48. Chen, H. *et al.* Effects of *S. mutans* gene-modification and antibacterial monomer dimethylaminohexadecyl methacrylate on biofilm growth and acid production. *Dent. Mater. Off. Publ. Acad. Dent. Mater.* **36**, 296–309 (2020).
 49. Chen, Y. *et al.* Determining the susceptibility of carbapenem resistant *Klebsiella pneumoniae* and *Escherichia coli* strains against common disinfectants at a tertiary hospital in China. *BMC Infect. Dis.* **20**, (2020).
 50. Conceicao, T., de Lencastre, H. & Aires-de-Sousa, M. Prevalence of biocide resistance genes, chlorhexidine and mupirocin non-susceptibility in Portuguese hospitals during a 31-years period (1985-2016). *J. Glob. Antimicrob. Resist.* (2020) doi:10.1016/j.jgar.2020.12.010.
 51. Cowley, N. L. *et al.* Effects of Formulation on Microbicide Potency and Mitigation of the Development of Bacterial Insusceptibility. *Appl. Environ. Microbiol.* **81**, 7330–7338 (2015).
 52. da Silva, K. R. *et al.* Antibacterial and cytotoxic activities of *Pinus tropicalis* and *Pinus elliottii* resins and of the diterpene dehydroabietic acid against bacteria that cause dental caries. *Front. Microbiol.* **10**, (2019).
 53. Dadpe, M. *et al.* Evaluation of antimicrobial efficacy of (Ajwain) oil and chlorhexidine against oral bacteria: An study. *J. Indian Soc. Pedod. Prev. Dent.* **36**, 357–363 (2018).
 54. Dong L. *et al.* Effects of sub-minimum inhibitory concentrations of antimicrobial agents on *Streptococcus mutans* biofilm formation. *Int. J. Antimicrob. Agents* **35**, 390–5 (2012).

55. Duarte B *et al.* 2CS-CHXT Operon Signature of Chlorhexidine Tolerance among *Enterococcus faecium* Isolates. *Appl. Environ. Microbiol.* **85**, e01589-19 (2019).
56. Fernandez-Cuenca, F. *et al.* Effect of Sub-MIC concentrations of biocides on the expression of genes coding for efflux pumps and porins in *acinetobacter baumannii* ATCC 19606. *Clin. Microbiol. Infect.* **17**, (2011).
57. Ferreira, C. M., da Silva Rosa, O. P., Torres, S. A., Ferreira, F. B. & Bernardinelli, N. Activity of endodontic antibacterial agents against selected anaerobic bacteria. *Braz. Dent. J.* **13**, 118–122 (2002).
58. Filho, J. G. *et al.* Genetic and physiological effects of subinhibitory concentrations of oral antimicrobial agents on *Streptococcus mutans* biofilms. *Microb. Pathog.* **150**, 104669 (2020).
59. Forbes, S., Humphreys, G. J., McBain, A. J. & Dobson, C. B. Transient and sustained bacterial adaptation following repeated sublethal exposure to microbicides and a novel human antimicrobial peptide. *Antimicrob. Agents Chemother.* **58**, 5809–5817 (2014).
60. Freire, I. C. M. *et al.* Atividade antibacteriana de oleos essenciais sobre *Streptococcus mutans* e *Staphylococcus aureus* Antibacterial activity of essential oils against strains of *Streptococcus* and *Staphylococcus*. *Rev. Bras. Plantas Med.* **16**, 372–377 (2014).
61. Furi, L. *et al.* Evaluation of reduced susceptibility to quaternary ammonium compounds and bisbiguanides in clinical isolates and laboratory-generated mutants of *staphylococcus aureus*. *Antimicrob. Agents Chemother.* **57**, 3488–3497 (2013).
62. Ghahramani, Y., Yaghoobi, F., Motamedi, R., Jamshidzadeh, A. & Abbaszadegan, A. Effect of Endodontic Irrigants and Medicaments Mixed with Silver Nanoparticles against Biofilm Formation of *Enterococcus faecalis*. *Iran. Endod. J.* **13**, 559–564 (2018).
63. Graziano, T. S. *et al.* In vitro effects of *Melaleuca alternifolia* essential oil on growth and production of volatile sulphur compounds by oral bacteria. *J. Appl. Oral Sci. Rev. FOB* **24**, 582–589 (2016).
64. Guerin, F. *et al.* The Transcriptional Repressor SmvR Is Important for Decreased Chlorhexidine Susceptibility in *Enterobacter cloacae* Complex. *Antimicrob. Agents Chemother.* **64**, (2020).
65. Guo, J. & Li, C. Molecular epidemiology and decreased susceptibility to disinfectants in carbapenem-resistant *Acinetobacter baumannii* isolated from intensive care unit patients in central China. *J. Infect. Public Health* **12**, 890–896 (2019).
66. Hajifattahi, F., Moravej-Salehi, E., Taheri, M., Mahboubi, A. & Kamalinejad, M. Antibacterial Effect of Hydroalcoholic Extract of *Punica granatum* Linn. Petal on Common Oral Microorganisms. *Int. J. Biomater.* 2016, (2016).
67. Hardy, K. *et al.* Increased usage of antiseptics is associated with reduced susceptibility in clinical isolates of *Staphylococcus aureus*. *mBio* **9**, (2018).
68. Hendry, E. R., Worthington, T., Conway, B. R. & Lambert, P. A. Antimicrobial efficacy of eucalyptus oil and 1,8-cineole alone and in combination with chlorhexidine digluconate against microorganisms grown in planktonic and biofilm cultures. *J. Antimicrob. Chemother. JAC* **64**, 1219–1225 (2009).
69. Hirose, N. *et al.* Development of a Cavity Disinfectant Containing Antibacterial Monomer MDPB. *J. Dent. Res.* **95**, 1487–1493 (2016).

70. Houang, E. T., Gilmore, O. J. A., Reid, C. & Shaw, E. J. ABSENCE OF BACTERIAL RESISTANCE TO POVIDONE IODINE. *J. Clin. Pathol.* **29**, 752–755 (1976).
71. Izutani, N., Imazato, S., Noiri, Y. & Ebisu, S. Antibacterial effects of MDPB against anaerobes associated with endodontic infections. *Int. Endod. J.* **43**, 637–645 (2010).
72. Joy Sinha, D. *et al.* Antibacterial Effect of Azadirachta indica (Neem) or Curcuma longa (Turmeric) against Enterococcus faecalis Compared with That of 5% Sodium Hypochlorite or 2% Chlorhexidine in vitro. *Bull. Tokyo Dent. Coll.* **58**, 103–109 (2017).
73. Karpanen, T. J., Worthington, T., Hendry, E. R., Conway, B. R. & Lambert, P. A. Antimicrobial efficacy of chlorhexidine digluconate alone and in combination with eucalyptus oil, tea tree oil and thymol against planktonic and biofilm cultures of Staphylococcus epidermidis. *J. Antimicrob. Chemother.* **62**, 1031–1036 (2008).
74. Koljalg, S., Naaber, P. & Mikelsaar, M. Antibiotic resistance as an indicator of bacterial chlorhexidine susceptibility. *J. Hosp. Infect.* **51**, 106–113 (2002).
75. Kreling, P. F. *et al.* Cytotoxicity and the effect of cationic peptide fragments against cariogenic bacteria under planktonic and biofilm conditions. *Biofouling* **32**, 995–1006 (2016).
76. Lacey, R. W. Antibacterial activity of povidone iodine towards non-sporing bacteria. *J. Appl. Bacteriol.* **46**, 443–449 (1979).
77. Lavaee, F., Ghapanchi, J., Motamedifar, M. & Sharifzade Javidi, M. Experimental Evaluation of the Effect of Zinc Salt on Inhibition of Streptococcus mutans. *J. Dent. Shiraz Iran* **19**, 168–173 (2018).
78. Lee, S. Y. Susceptibility of oral streptococci to chlorhexidine and cetylpyridinium chloride. *Biocontrol Sci.* **24**, 13–21 (2019).
79. Li, Y. F. *et al.* Inhibited biofilm formation and improved antibacterial activity of a novel nanoemulsion against cariogenic Streptococcus mutans in vitro and in vivo. *Int. J. Nanomedicine* **10**, 447–462 (2015).
80. Machuca, J., Lopez-Rojas, R., Fernandez-Cuenca, F. & Pascual, A. Comparative activity of a polyhexanide-betaine solution against biofilms produced by multidrug-resistant bacteria belonging to high-risk clones. *J. Hosp. Infect.* **103**, (2019).
81. Marcoux, E., Lagha, A. B., Gauthier, P. & Grenier, D. Antimicrobial activities of natural plant compounds against endodontic pathogens and biocompatibility with human gingival fibroblasts. *Arch. Oral Biol.* **116**, 104734 (2020).
82. Martins, M. L. *et al.* Antibacterial and cytotoxic potential of a brazilian red propolis. *Pesqui. Bras. Em Odontopediatria E Clin. Integrada* **19**, (2019).
83. Massunari, L. *et al.* Antimicrobial Activity and Biocompatibility of the Psidium cattleianum Extracts for Endodontic Purposes. *Braz. Dent. J.* **28**, 372–379 (2017).
84. McBain, A. J., Ledger, R. G., Gilbert, P. & Sreenivasan, P. Selection for high-level resistance by chronic triclosan exposure is not universal. *J. Antimicrob. Chemother.* **53**, 772–777 (2004).

85. Moraes, T. S. *et al.* In vitro evaluation of copaifera oblongifolia oleoresin against bacteria causing oral infections and assessment of its cytotoxic potential. *Curr. Pharm. Biotechnol.* **17**, 894–904 (2016).
86. Moreira, M. R. *et al.* Ent-Kaurenoic acid-rich extract from Mikania glomerata: In vitro activity against bacteria responsible for dental caries. *Fitoterapia* **112**, 211–216 (2016).
87. Moreti, D. L. C. *et al.* Mikania glomerata Sprengel extract and its major compound ent-kaurenoic acid display activity against bacteria present in endodontic infections. *Anaerobe* **47**, 201–208 (2017).
88. Morrissey, I. *et al.* Evaluation of epidemiological cut-off values indicates that biocide resistant subpopulations are uncommon in natural isolates of clinically-relevant microorganisms. *PLoS ONE* **9**, (2014).
89. Motamedifar, M., Khosropanah, H. & Dabiri, S. Antimicrobial Activity of Peganum Harmala L. on Streptococcus mutans Compared to 0.2% Chlorhexidine. *J. Dent. Shiraz Iran* **17**, 213–218 (2016).
90. Munoz-Gallego, I., Infiesta, L., Viedma, E., Perez-Montarelo, D. & Chaves, F. Chlorhexidine and mupirocin susceptibilities in methicillin-resistant Staphylococcus aureus isolates from bacteraemia and nasal colonisation. *J. Glob. Antimicrob. Resist.* **4**, 65–69 (2016).
91. Nakase, K. *et al.* Propionibacterium acnes has low susceptibility to chlorhexidine digluconate. *Surg. Infect.* **19**, 298–302 (2018).
92. Nazemisalman, B. *et al.* Comparison of antimicrobial effect of Ziziphora tenuior, Dracocephalum moldavica, Ferula gummosa, and Prangos ferulacea essential oil with chlorhexidine on Enterococcus faecalis: An in vitro study. *Dent. Res. J.* **15**, 111–116 (2018).
93. Nicolae Dopcea, G., Diguta, C. F., Matei, F., Dopcea, I. & Nanu, A. E. Resistance and cross-resistance in Staphylococcus spp. strains following prolonged exposure to different antiseptics. *J. Glob. Antimicrob. Resist.* **21**, 399–404 (2020).
94. O'Driscoll, N. H., Labovitiadi, O., Matthews, K. H., Lamb, A. J. & Cushnie, T. P. T. Potassium loss from chlorhexidine-treated bacterial pathogens is time- and concentration-dependent and variable between species. *Curr. Microbiol.* **68**, 6–11 (2014).
95. Pal, S., Yoon, E. J., Park, S. H., Choi, E. C. & Song, J. M. Metallopharmaceuticals based on silver(I) and silver(II) polydiguanide complexes: Activity against burn wound pathogens. *J. Antimicrob. Chemother.* **65**, 2134–2140 (2010).
96. Roedel, A. *et al.* Evaluation of a Newly Developed Vacuum Dried Microtiter Plate for Rapid Biocide Susceptibility Testing of Clinical Enterococcus Faecium Isolates. *Microorganisms* **8**, (2020).
97. Rosa, O. P., Torres, S. A., Ferreira, C. M. & Ferreira, F. B. In vitro effect of intracanal medicaments on strict anaerobes by means of the broth dilution method. *Pesqui. Odontol. Bras. Braz. Oral Res.* **16**, 31–36 (2002).
98. Rose, H., Mahenthiralingam, E., Baldwin, A. & Dowson, C. G. Biocide susceptibility of the Burkholderia cepacia complex. *J. Antimicrob. Chemother.* **63**, 502–510 (2009).
99. Shani, S., Friedman, M. & Steinberg, D. Relation between surface activity and antibacterial activity of amine-fluorides. *Int. J. Pharm.* **131**, 33–39 (1996).

100. Sherry, L. *et al.* Investigating the biological properties of carbohydrate derived fulvic acid (CHD-FA) as a potential novel therapy for the management of oral biofilm infections. *BMC Oral Health* **13**, 47 (2013).
101. Skovgaard, S. *et al.* Recently introduced qacA/B genes in *Staphylococcus epidermidis* do not increase chlorhexidine MIC/MBC. *J. Antimicrob. Chemother.* **68**, 2226–2233 (2013).
102. Suwantararat, N. *et al.* High Prevalence of Reduced Chlorhexidine Susceptibility in Organisms Causing Central Line- Associated Bloodstream Infections. *Infect. Control Hosp. Epidemiol.* **35**, 1183–1186 (2014).
103. Suzuki, Y. *et al.* Effects of a sub-minimum inhibitory concentration of chlorhexidine gluconate on the development of in vitro multi-species biofilms. *Biofouling* **36**, 146–158 (2020).
104. Tambe, S. M., Sampath, L. & Modak, S. M. In vitro evaluation of the risk of developing bacterial resistance to antiseptics and antibiotics used in medical devices. *J. Antimicrob. Chemother.* **47**, 589–598 (2001).
105. Targino, A. G. R. *et al.* An innovative approach to treating dental decay in children. A new anti-caries agent. *J. Mater. Sci. Mater. Med.* **25**, 2041–2047 (2014).
106. Tetz, G. & Tetz, V. In vitro antimicrobial activity of a novel compound, Mul-1867, against clinically important bacteria. *Antimicrob. Resist. Infect. Control* **4**, (2015).
107. Touzel, R. E., Sutton, J. M. & Wand, M. E. Establishment of a multi-species biofilm model to evaluate chlorhexidine efficacy. *J. Hosp. Infect.* **92**, 154–160 (2016).
108. Uzunbayir-Akel, N. *et al.* Effects of disinfectants and ciprofloxacin on quorum sensing genes and biofilm of clinical *Pseudomonas aeruginosa* isolates. *J. Infect. Public Health* **13**, 1932–1938 (2020).
109. Vahabi, S., Hakemi-Vala, M. & Gholami, S. In vitro Antibacterial Effect of Hydroalcoholic Extract of *Lawsonia inermis*, *Malva sylvestris*, and *Boswellia serrata* on *Aggregatibacter actinomycetemcomitans*. *Adv. Biomed. Res.* **8**, 22 (2019).
110. Weaver Jr, A. J. *et al.* Antibacterial activity of THAM trisphenylguanide against methicillin-resistant *Staphylococcus aureus*. *PLoS ONE* **9**, (2014).
111. Xing, M. *et al.* Antimicrobial efficacy of the alkaloid harmaline alone and in combination with chlorhexidine digluconate against clinical isolates of *Staphylococcus aureus* grown in planktonic and biofilm cultures. *Lett. Appl. Microbiol.* **54**, 475–482 (2012).
112. Zhang, M., Boost, M. V., O'Donoghue, M. M., Ito, T. & Hiramatsu, K. Prevalence of antiseptic-resistance genes in *Staphylococcus aureus* and coagulase-negative staphylococci colonising nurses and the general population in Hong Kong. *J. Hosp. Infect.* **78**, 113–7 (2011).
113. Zhou, Z., Lu, Y. & Wei, D. Polyhexamethylene guanidine hydrochloride shows bactericidal advantages over chlorhexidine digluconate against ESKAPE bacteria. *Biotechnol. Appl. Biochem.* **62**, 268–274 (2015).
114. Russell, A. D. Bacterial resistance to disinfectants: present knowledge and future problems. *J. Hosp. Infect.* **43**, S57–S68 (1999).

115. Platt, J. H. & Bucknall, R. A. MIC tests are not suitable for assessing antiseptic handwashes. *J. Hosp. Infect.* **11**, 396–397 (1988).

116. Russell, A. D. Do Biocides Select for Antibiotic Resistance? *J. Pharm. Pharmacol.* **52**, 227–233 (2010).

117. EUCAST. On recent changes in clinical microbiology susceptibility reports - new interpretation of susceptibility categories S, I and R. *EUCAST.org* <https://www.eucast.org/newsiandr/> (2021).

118. *Essentials of neuroanesthesia*. (Academic Press, an imprint of Elsevier, 2017).

Figures

Figure 1

Figure 2

Forest plot of the mean MBC for different species and families of bacteria

Figure 3

A scatterplot of study-level estimates of mean MBC over time, for *Staphylococci*. The size of the points corresponds to the precision (inverse variance) of the study.

Mean Bacteriocidal Concentration of Chlorhexidine for Streptococci over time

Figure 4

A scatterplot of study-level estimates of mean MBC over time, for *Streptococci*. The size of the points corresponds to the precision (inverse variance) of the study.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [S1Fig.jpg](#)
- [S1Table.docx](#)
- [S2Table.docx](#)