Bioinformatics analysis of expression profiling by high throughput sequencing for identification of potential key genes among SARS-CoV-2/COVID 19

Basavaraj Mallikarjunayya Vastrad

Department of Pharmaceutics, SET'S College Of Pharmacy, SR Nagar, Dharwad

Chanabasayya Mallikarjunayya Vastrad (✉ channu.vastrad@gmail.com)

Chanabasava Nilaya <https://orcid.org/0000-0003-3615-4450>

Research Article

Keywords: SARS-CoV-2 infection, bioinformatics analysis, differentially expressed genes, novel biomarkers, hub genes

Posted Date: January 4th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-122015/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Severe acute respiratory syndrome corona virus 2 (SARS-CoV-2) is pandemic recently emerged and is rapidly spreading in humans. However, the precise molecular mechanisms of the advancement and progression of SARS-CoV-2 infection remain unclear. The current investigation attempted to identify and functionally analyze the differentially expressed genes (DEGs) between SARS-CoV-2 infection and mock by using comprehensive bioinformatics analyses. The GSE148729 expression profiling by high throughput sequencing was downloaded from the Gene Expression Omnibus (GEO) and analyzed using the limma package in R software to identify DEGs. Pathway and gene ontology (GO) enrichment analysis of the up and down regulated genes were performed in ToppGene. The HIPPIE database was used to evaluate the interactions of up and down regulated genes and to construct a protein-protein interaction (PPI) network using Cytoscape software. Hub genes were selected using the Network Analyzer plugin. Subsequently, extensive target prediction and network analyses methods were used to assess, target gene - miRNA regulatory network and target gene - TF regulatory network. Receiver operating characteristic (ROC) analysis was utilized for validation. A total of 928 DEGs (461 up regulated genes and 467 down regulated genes) were identified between SARS-CoV-2 infection and mock samples. The Pathway enrichment analysis results showed that these up and down regulated genes were significantly enriched in cytokine-cytokine receptor interaction, and ascorbate and aldarate metabolism. Several significant GO terms, including the response to biotic stimulus and oxoacid metabolic process, were identified as being closely associated with these up and down regulated genes. The top hub genes and target genes were screened and included JUN, FBXO6, PCLAF, CFTR, TXNIP, PMAIP1, BRI3BP, FAHD1, PROX1, CXCL11, SERHL2 and CFI. ROC curve analysis showed that messenger RNA levels of these ten genes (DDX58, IFITM2, IRF1, PML, SAMHD1, ACSS1, CYP2U1, DDC, PNMT and UGT2A3) exhibited better diagnostic efficiency for SARS-CoV-2 infection and mock. The current investigation identified a series of key genes and pathways that may be involved in the progression of SARS-CoV-2 infection, providing a new understanding of the underlying molecular mechanisms of SARS-CoV-2 infection.

Introduction

Severe acute respiratory syndrome corona virus 2 (SARS-CoV-2) is key causes of respiratory disease in pandemic form [1] and named as COVID - 19. In 2020, more than 406,207 people die of SARS-CoV-2 complications in the entire world. Restriction of the spread of SARS-CoV-2 virus has been difficult, but transmissions of this virus develop to require close contact through large-particle aerosols [2]. SARS-CoV-2 infections in humans cause a severe pneumonia that is disastrous in about 11% of infected individuals [3].

Intensive investigation has been achieved on the virulence and evolution of the SARS-CoV-2 virus [4]. However, very little is known about the influence of specific genes or biomarkers in humans that add to the susceptibility to SARS-CoV-2 infections. Consequently, it needs more effort to clarify the molecular mechanism underlying SARS-CoV-2 infections advancement and progression, holding promise for finding potential drug targets and diagnostic biomarkers of SARS-CoV-2 infections.

Expression profiling by high throughput sequencing analysis based on microarray technology is a widely used, which can simultaneously detect expression change of thousands of genes on the mRNA level. By gene expression profiling analysis with microarray technology, some investigations have found many differently expressed genes which played a key role in SARS-CoV-2 infections initiation and development and could be assessed as possible molecular targets and diagnostic markers. ACE2 and TMPRSS2 could function as a key mediator of SARS-CoV-2 infections by means of gene expression analysis [5]. In the present investigation, we identified SARS-CoV-2 infection linked with differential expressed genes (DEGs) between SARS-CoV-2 infected samples and mock samples, and successively performed pathway and GO term enrichment analysis, PPI network analysis, modules analysis, target gene - miRNA regulatory network and target gene - TF regulatory network analysis to discover the hub and target genes and pathways closely related to SARS-CoV-2 infections. Finally, hub genes were validated by receiver operating characteristic (ROC) curve analysis.

Material And Methods

Microarray data

The expression profiling by high throughput sequencing dataset GSE148729 was downloaded from the National Center for Biotechnology Information Gene Expression Omnibus Dataset (<https://www.ncbi.nlm.nih.gov/gds>) [6] and was based on the GPL18573 Illumina NextSeq 500 (Homo sapiens) array. The GSE148729 dataset, which is expression profiling by high throughput sequencing dataset, contains 22 samples, including 14 SARS-CoV-2 infections samples and 8 mock samples.

Data pre-processing and identification of DEG

Limma package in R bioconductor was used for the pre-processing for GSE148729 dataset, which included background correcting, normalizing and calculating expression [7]. The DEGs in SARS-CoV-2 infections samples compared with mock samples were determined using limma package. DEGs were screened with a adj. P value corrected $P < 0.05$ and $|\log \text{fold-change (FC)}| > 1$. The adj.PVal for up and down regulated genes corrected $P < 0.05$ and $|\log \text{FC}| > 1$ were used as the screening thresholds.

Pathway enrichment analysis of DEGs

The route of gene cluster and related functions was determined using the BIOCYC (<https://biocyc.org/>) [8], Kyoto Encyclopedia of Genes and Genomes (KEGG) (<http://www.genome.jp/kegg/pathway.html>) [9], Pathway Interaction Database (PID) (<https://wiki.nci.nih.gov/pages/viewpage.action?pagelD=315491760>) [10], REACTOME (<https://reactome.org/>) [11], GenMAPP (<http://www.genmapp.org/>) [12], MSigDB C2 BIOCARTEA (<http://software.broadinstitute.org/gsea/msigdb/collections.jsp>) [13], PantherDB (<http://www.pantherdb.org/>) [14], Pathway Ontology (<http://www.obofoundry.org/ontology/pw.html>) [15] and Small Molecule Pathway Database (SMPDB) (<http://smpdb.ca/>) [16]. This investigation used the ToppGene (ToppFun)

(<https://toppgene.cchmc.org/enrichment.jsp>) [17] to perform the pathway enrichment analysis. A P-value <0.05 was considered to be statistically significant.

Gene ontology (GO) enrichment analysis

Gene ontology (GO) (<http://www.geneontology.org>) enrichment analysis defines approach used to characterize gene function and relationships [18]. It classifies functions such as biological process (BP), cellular component (CC) and molecular function (MF). ToppGene (ToppFun) (<https://toppgene.cchmc.org/enrichment.jsp>) [17] was used to classify significant DEGs by their BP, CC and MF through GO and the significant transcripts (Benjamini–Hochberg false discovery rate <0.05) were identified using the Functional Annotation clustering tool.

Protein-protein interaction network construction and module analysis

The online Human Integrated Protein-Protein Interaction rEference (HIPPIE) (<http://cbdm.uni-mainz.de/hippie/>) [19] database was used to identify potential interaction among the up and down regulated genes and associated with various PPI data bases such as IntAct (<https://www.ebi.ac.uk/intact/>) [20], BioGRID (<https://thebiogrid.org/>) [21], HPRD (<http://www.hprd.org/>) [22], MINT (<https://mint.bio.uniroma2.it/>) [23], BIND (<http://download.baderlab.org/BINDTranslation/>) [24], MIPS (<http://mips.helmholtz-muenchen.de/proj/ppi/>) [25] and DIP (<http://dip.doe-mbi.ucla.edu/dip/Main.cgi>) [26]. Cytoscape 3.8.0 (<http://www.cytoscape.org/>) [27] was used to visualize the results from the PPI network. The Network Analyzer Cytoscape plug-in was used to explore hub genes were generated using highest node degree [28], betweenness centrality [29], stress centrality [30], closeness centrality [31] and lowest clustering coefficient [Wang et al 2012] methods and hub genes were further analyzed for pathway and GO enrichment analysis. Module analysis on the PPI network results was performed using the PEWCC1 (<http://apps.cytoscape.org/apps/PEWCC1>) [32] clustering algorithm that comes with Cytoscape. Module analysis was applied to analyze more connected gene groups. In addition, the module analysis results were further analyzed for pathway and GO enrichment analysis.

Construction of target genes - miRNA regulatory network

miRNet database (<https://www.mirnet.ca/>) [33] is a bioinformatics platform for predicting target genes (up and down regulated genes) and miRNA pairs. In the current investigation, the targets genes were predicted using ten databases: TarBase (<http://diana.imis.athena-innovation.gr/DianaTools/index.php?r=tarbase/index>) [34], miRTarBase (<http://mirtarbase.mbc.nctu.edu.tw/php/download.php>) [35], miRecords (<http://miRecords.umn.edu/miRecords>) [36], miR2Disease (<http://www.mir2disease.org/>) [37], HMDD (<http://www.cuilab.cn/hmdd>) [38], PhenomiR (<http://mips.helmholtz-muenchen.de/phenomir/>) [39], SM2miR (<http://bioinfo.hrbmu.edu.cn/SM2miR/>) [40], PharmacomiR (<http://www.pharmacomir.org/>) [41], EpimiR (<http://bioinfo.hrbmu.edu.cn/EpimiR/>) [42] and starBase (<http://starbase.sysu.edu.cn/>) [43]. The screening criterion was that the miRNA target exists in the ten miRNA databases concurrently. The target genes - miRNA regulatory network was depicted and visualized using Cytoscape software.

Construction of target genes - TF regulatory network

NetworkAnalyst database (<https://www.networkanalyst.ca/>) [44] is a bioinformatics platform for predicting target genes (up and down regulated genes) and TF pairs. In the current investigation, the targets genes were predicted using JASPAR (<http://jaspar.genereg.net/>) [45]. The screening criterion was that the TF target exists in the TF database concurrently. The target genes - TF regulatory network was depicted and visualized using Cytoscape software.

ROC analysis

Receiver operating characteristic (ROC) curve analysis was implemented to calculate the sensitivity and specificity of the up and down regulated genes for SARS-CoV-2 infections diagnosis using the R “pROC” package [46]. An area under the curve (AUC) value was determined and used to label the ROC effect.

Results

Data pre-processing and identification of DEG

Expression profiling by high throughput sequencing dataset GSE148729 was downloaded from GEO database. After quality detection of microarray raw data, microarray based on GPL18573 platform, including 14 SARS-CoV-2 infections samples and 8 mock samples. Limma in R package was used for preprocessing and gene differential expression analysis of microarray data (Fig. 1A and Fig. 1B). Total of 928 DEGs (fold change >1, Fig. 2) consisting 461 significantly up regulated genes and 467 significantly down regulated genes, for the subsequent bioinformatics analysis. The hierarchical clustering heat map of up and down regulated genes shows the differences between SARS-CoV-2 infections and mock (Fig. 3 and Fig. 4).

Pathway enrichment analysis of DEGs

To analyze the biological classification of up and down regulated genes, functional and pathway enrichment analyses were performed using ToppGene. Pathway analysis revealed that the up regulated genes were mainly enriched in superpathway of steroid hormone biosynthesis, aspirin triggered resolvins biosynthesis, cytokine-cytokine receptor interaction, TNF signaling pathway, calcineurin-regulated NFAT-dependent transcription in lymphocytes, ATF-2 transcription factor network, cytokine signaling in immune system, interferon signaling, steroid hormone metabolism, androgen and estrogen metabolism, genes encoding secreted soluble factors, cytokines and inflammatory response, toll receptor signaling pathway, interleukin signaling pathway, nuclear factor kappa B signaling, c-Jun N-terminal kinases MAPK signaling, 17-beta hydroxysteroid dehydrogenase III deficiency and adrenal hyperplasia type 3 or congenital adrenal hyperplasia due to 21-hydroxylase deficiency are listed in Table 2, while down regulated genes were mainly enriched in catecholamine biosynthesis, noradrenaline and adrenaline degradation, ascorbate and aldarate metabolism, steroid hormone biosynthesis, signaling mediated by p38-gamma and p38-delta, aurora A signaling, biological oxidations, the citric acid (TCA) cycle and

respiratory electron transport, bile acid biosynthesis, valineleucine and isoleucine degradation, angiotensin-converting enzyme 2 regulates heart function, apoptotic DNA fragmentation and tissue homeostasis, nicotine degradation, serine glycine biosynthesis, renin-angiotensin system signaling, histidine metabolic, fatty Acid elongation in mitochondria and gluconeogenesis are listed in Table 3.

Gene ontology (GO) enrichment analysis

GO term enrichment analysis results varied from GO classification and expression change of up and down regulated genes are listed in Table 4 and Table 5. As to BP, the up regulated genes significantly enriched in response to biotic stimulus and defense response, and the down regulated genes significantly enriched in oxoacid metabolic process, and monocarboxylic acid metabolic process. For CC, the up regulated genes significantly enriched in cell surface and synapse, and the down regulated genes significantly enriched in mitochondrion and polymeric cytoskeletal fiber. About MF, the up regulated genes significantly enriched in cytokine activity and signaling receptor binding and the down regulated genes significantly enriched in cofactor binding and oxidoreductase activity.

Protein-protein interaction network construction and module analysis

PPI network of up regulated genes, consisting of 5698 nodes and 11384 edges, was constructed by Cytoscape software, based on HIPPIE database (Fig. 5). The up and down regulated genes with highest node degree, betweenness centrality, stress centrality, closeness centrality and lowest clustering coefficient were selected as the hub genes of SARS-CoV-2 infections and are listed in Table 6. The scatter plots of node degree, betweenness centrality, stress centrality, closeness centrality and clustering coefficient for up regulated genes are shown in Fig. 6. The up regulated hub genes (JUN, FBXO6, CYLD, IFI16, ADRB2, PML, ZC3HAV1, SYNPR, CH25H, LSMEM1, CRTAM and RAET1L) in this PPI network were significantly enriched in TNF signaling pathway, RIG-I/MDA5 mediated induction of IFN-alpha/beta pathways, NOD-like receptor signaling pathway, cAMP signaling pathway, influenza A, response to biotic stimulus, IL-17 signaling pathway, cytokine-cytokine receptor interaction, positive regulation of RNA metabolic process, transition metal ion binding and response to endogenous stimulus. Similarly, PPI network of down regulated genes, consisting of 5712 nodes and 11002 edges (Fig. 7). The scatter plots of node degree, betweenness centrality, stress centrality, closeness centrality and clustering coefficient for down regulated genes are shown in Fig. 8. The down regulated hub genes (PCLAF, CFTR, TUBA1A, AURKA, ARRB1, KPNA2, PRDX2, PGK1, GALM, ACYP2, HSD17B2, CEACAM7 and SEMA3E) in this PPI network were significantly enriched in microtubule cytoskeleton, peptidase activity, metabolism of proteins, aurora A signaling, morphine addiction, regulation of cytoplasmic and nuclear SMAD2/3 signaling, oxidation-reduction process, metabolic pathways, protein homodimerization activity, identical protein binding, steroid hormone biosynthesis and genes encoding proteins affiliated structurally or functionally to extracellular matrix proteins.

PEWCC1 was used to identified the total 1503 modules in the PPI network of up regulated genes and the four significant modules were selected (Fig. 9). Pathway and GO enrichment analysis, module 25 (nodes 92 and edges 173), module 33 (nodes 75 and edges 153), module 37 (nodes 63 and edges 109) and

module 70 (nodes 20 and edges 41) revealed that the four significant modules were mainly enriched in cytokine-cytokine receptor interaction, influenza A, Jak-STAT signaling pathway, inflammation mediated by chemokine and cytokine signaling pathway, response to cytokine, response to biotic stimulus, response to virus and response to endogenous stimulus. Similarly, PEWCC1 was used to identify the total 1405 modules in the PPI network of down regulated genes and the four significant modules were selected (Fig. 10). Pathway and GO enrichment analysis, module 14 (nodes 111 and edges 135), module 15 (nodes 107 and edges 121), module 27 (nodes 81 and edges 97) and module 45 (nodes 54 and edges 75) revealed that the four significant modules were mainly enriched in hedgehog signaling events mediated by Gli proteins, metabolism of proteins, metabolism of lipids and lipoproteins, the citric acid (TCA) cycle and respiratory electron transport, mitochondrion, growth, intrinsic component of plasma membrane and cellular lipid metabolic process.

Construction of target genes - miRNA regulatory network

To further understand the regulatory network between miRNA and target genes, various miRNA databases such as TarBase, miRTarBase, miRecords, miR2Disease, HMDD, PhenomiR, SM2miR, PharmacomiR, EpmiR and starBase through miRNet were constructed by Cytoscape. As shown in Fig. 11, the target genes - miRNA regulatory network for up regulated genes with 5667 edges and 2144 nodes was obtained for target genes. Different up regulated target genes regulated by miRNA are shown in Table 7, which top five miRNAs such as hsa-mir-4511, hsa-mir-3924, hsa-mir-4478, hsa-mir-3650 and hsa-mir-4252 had been predicted to regulate target genes such as TXNIP, PMAIP1, APOL6, CHAC1 and KLF2. Pathway and GO enrichment analysis revealed that target genes in this network were mainly involved in NOD-like receptor signaling pathway, Direct p53 effectors, signaling receptor binding and response to cytokine. Similarly, as shown in Fig. 12, the target genes - miRNA regulatory network for down regulated genes with 6679 edges and 2236 nodes was obtained for target genes. Different down regulated target genes regulated by miRNA are shown in Table 7, which top five miRNAs such as hsa-mir-4312, hsa-mir-4527, hsa-mir-4673, hsa-mir-4496 and hsa-mir-3153 had been predicted to regulate target genes such as BRI3BP, FAHD1, CPM, HOXA13 and TMBIM6. Pathway and GO enrichment analysis revealed that target genes in this network were mainly involved in mitochondrion, metabolic pathways, metabolism of proteins and ion transport.

Construction of target genes - TF regulatory network

To further understand the regulatory network between TFs and target genes, JASPAR through NetworkAnalyst were constructed by Cytoscape. As shown in Fig. 13, the target genes - TF regulatory network for up regulated genes with 2844 edges and 439 nodes was obtained for target genes. Different up regulated target genes regulated by TF are shown in Table 8, which top five TFs such as FOXC1, GATA2, YY1, FOXL1 and NFkB1 had been predicted to regulate target genes such as PROX1, CXCL11, C4A, JAK2 and IL15RA. Pathway and GO enrichment analysis revealed that target genes in this network were mainly involved in viral genome replication, cytokine-cytokine receptor interaction, innate immune system, influenza A and Jak-STAT signaling pathway. Similarly, as shown in Fig. 14, the target genes - TF regulatory network for down regulated genes with 3191 edges and 505 nodes was obtained for target

genes. Different down regulated target genes regulated by TF are shown in Table 8, which top five TFs such as FOXC1, GATA2, YY1, NFIC and FOXL1 had been predicted to regulate target genes such as SERHL2, CFI, TCN1, PGP and PGK1. Pathway and GO enrichment analysis revealed that target genes in this network were mainly involved in mitochondrion, peptidase activity, drug metabolic process, metabolic pathways and carbon metabolism.

ROC analysis

As these five up and five down regulated genes are prominently expressed in SARS-CoV-2 infection, we performed a ROC curve analysis to evaluate their sensitivity and specificity for the diagnosis of SARS-CoV-2 infection. As shown in Fig. 15, DDX58, IFITM2, IRF1, PML, SAMHD1, ACSS1, CYP2U1, DDC, PNMT and UGT2A3 achieved an AUC value of 0.777, 0.857, 0.929, 0.804, 0.759, 0.714, 0.728, 0.754, 0.714 and 0.763, demonstrating that these genes have high sensitivity and specificity for SARS-CoV-2 infection diagnosis. The results suggested that DDX58, IFITM2, IRF1, PML, SAMHD1, ACSS1, CYP2U1, DDC, PNMT and UGT2A3 can be used as biomarkers for the diagnosis of SARS-CoV-2 infection.

Discussion

SARS-CoV-2 infection transmits worldwide and its mortality rate has increased in recent years. This research represents the first complete study of a genes linked with SARS-CoV-2 virus infection, using bioinformatics analysis. However, the molecular mechanisms of SARS-CoV-2 infection remain poorly understood. Thus, probable biomarkers for diagnosis and treatment with high competence are crucially demanded. Microarray technology has been proved to be a needful approach to diagnose novel biomarkers in SARS-CoV-2 infection.

In the current investigation, expression profiling by high throughput sequencing dataset was analyzed to obtain DEGs between SARS-CoV-2 infections samples and mock samples. A total of 928 DEGs were identified, including 461 up regulated genes and 467 down regulated genes. Genes such as SOCS3 [47] and IL1A [48] were liable for progression of influenza virus infection, but these genes may be linked with progression of SARS-CoV-2 infection. Genes such as KLF6 [49], DUSP1 [50] and OLFM4 [51] were associated with development of respiratory syncytial virus infection, but these genes may be responsible for advancement of SARS-CoV-2 infection. IL6 was involved in progression of SARS-CoV-2 infection [52]. Genes such as MAGT1 [53], CD24 [54] and UGT1A7 [55] were involved in progression of various viral infections, but these genes may be associated with development of SARS-CoV-2 infection. AQP1 was linked with development of porcine reproductive and respiratory syndrome virus infection [56], but this gene may be responsible for progression of SARS-CoV-2 infection.

In the pathway enrichment analysis, many up and down regulated genes may play a very key role in progression of SARS-CoV-2 infection. Enriched genes such as CXCL9 [57], CX3CL1 [58], IFNB1 [59], IL7 [60], TNFSF10 [61], IFNL3 [62], EGR1 [63], USP18 [64], DDX58 [65], UBA7 [66], IFI35 [67], IFIT2 [68], IFIT1 [69], TRIM5 [70], RSAD2 [71], XAF1 [72], MX2 [73], GBP1 [74], TRIM21 [75], FGF19 [76], PELI1 [77], OAS1 [78], DUSP10 [79], TLR6 [80], TICAM1 [81], DDC (dopa decarboxylase) [82], UGT1A1 [83], STMN1 [84],

CYP27A1 [85] and ACE (angiotensin I converting enzyme) [86] were linked with advancement of various viral infections, but these genes may be responsible for progression of SARS-CoV-2 infection. Enriched genes such as CCL2 [87], FLT3 [88], CXCL8 [89], IL11 [90], CXCL10 [91], TNF (tumor necrosis factor) [92], EGR2 [93], TRIM22 [94], PML (promyelocytic leukemia) [95], ICAM1 [96], IFITM1 [97], IFITM2 [98], ISG15 [99], SOCS1 [100], IRF1 [101], IRF7 [102], ISG20 [103], STAT2 [104], OAS2 [105], HERC5 [106] and TNFAIP3 [107] were associated with advancement of influenza virus infection, but these genes may be liable for development of SARS-CoV-2 infection. Enriched genes such as CCL20 [108], TSLP (thymic stromal lymphopoietin) [109], IL7R [110], IFIT3 [111] and OASL (2'-5'-oligoadenylate synthetase like) [112] were involved in the development of respiratory syncytial virus infection, but these genes may be associated with development of SARS-CoV-2 infection. SAMHD1 was involved in progression of porcine reproductive and respiratory syndrome virus infection [113], but this gene may be liable for progression of SARS-CoV-2 infection. Enriched genes such as JAK2 [114] and ACE2 [115] were responsible for progression of SARS-CoV-2 infection. As given in the pathway enrichment analysis for up and down regulated genes, HSD17B3, CYP21A2, TNFRSF4, CXCL11, FLT3LG, CSF2, TNFSF13B, IL12A, IL15RA, INHBA (inhibin subunit beta A), IL17C, IL22RA1, BMP2, CXCL2, CXCL3, LTB (lymphotoxin beta), IFNL2, IFNL1, FOS (Fos proto-oncogene, AP-1 transcription factor subunit), GATA3, PTGS2, JUN (Jun proto-oncogene, AP-1 transcription factor subunit), EGR3, EGR4, UBE2L6, HLA-F, SPTBN5, RASGRP3, TRIM38, KLB (klotho beta), MAP3K8, MT2A, PSMB9, FGF18, DUSP8, RELB (RELB proto-oncogene, NF-kB subunit), LGALS9, RASGRP1, FST (follistatin), WIF1, NFKBIA (NFKB inhibitor alpha), PNMT (phenylethanolamine N-methyltransferase), ALDH2, UGT1A10, UGT1A8, UGT1A6, UGT1A5, UGT1A9, UGT1A4, UGT1A3, UGT2A3, SNTA1, EPHX1, ADH6, FDX1, ACSS1, GSTO2, GGT1, NR1H4, GLYATL2, GSTA4, CYP2U1, CYP4F3, HADHB (hydroxyacyl-CoA dehydrogenase trifunctional multienzyme complex subunit beta), AGT (angiotensinogen), MME (membrane metalloendopeptidase), PPT1 and PRDX3 are novel biomarkers for SARS-CoV-2 infection.

Similarly, GO enrichment analysis, many up and down regulated genes may play a very key role in progression of SARS-CoV-2 infection. Enriched genes such as IFNL4 [116], ZFP36 [117], CD274 [118], MIR21 [119], CYP1A1 [120], CMPK2 [121], C4A [122], SERPINE1 [123], CCRL2 [124], CD69 [125], FAH (fumarylacetoacetate hydrolase) [126], ERF (erythroferrone) [127], SERINC5 [128], ADI1 [129], AP2M1 [130], PRIMPOL (primase and DNA directed polymerase) [131], BIRC5 [132] and TF (transferrin) [133] were responsible for advancement of various viral infections, but these genes may be involved in progression of SARS-CoV-2 infection. Enriched genes such as APOBEC3G [134], NLRC5 [135], PTX3 [136], FAP (fibroblast activation protein alpha) [137] and CAT (catalase) [138] were linked with development of influenza viral infection, but these genes may be associated with progression of SARS-CoV-2 infection. Enriched genes such as IFI16 [139], TAP1 [140] and PIN1 [141] were involved in development of porcine reproductive and respiratory syndrome virus infection, but these genes may be associated with progression of SARS-CoV-2 infection. NLRP3 was key for advancement of SARS-CoV-2 infection [142]. PDK1 was liable for development of SARS corona virus infection [143], but this gene may be linked with advancement of SARS-CoV-2 infection. As given in the GO enrichment analysis for up and down regulated genes, PDE2A, PDE4B, NT5C3A, PGLYRP2, DTX3L, PLAC8, NOCT (nocturnin), PMAIP1,

ZC3H12A, DHX58, TENT5A, IFI44, BATF2, TXNIP (thioredoxin interacting protein), SH2D1B, MIR181B1, RAET1L, STX11, ZC3HAV1, CYLD (CYLD lysine 63 deubiquitinase), TAGAP (T cell activation RhoGTPase activating protein), BCL3, PTGER4, RTP4, GFI1, IFIH1, CLEC4A, GLI2, C4B, GPER1, CRTAM (cytotoxic and regulatory T cell molecule), PCK1, acetylcholinesterase (Cartwright blood group), TRPC4, DSCAML1, HEG1, PLAUR (plasminogen activator, urokinase receptor), SLITRK6, ACKR4, LAYN (layilin), GRIA2, MMP25, EPHA4, SECTM1, VSTM1, RIMKLA (ribosomal modification protein rimK like family member A), ACO1, ETFB (electron transfer flavoprotein subunit beta), ETFDH (electron transfer flavoprotein dehydrogenase), MDH1, HNMT (histamine N-methyltransferase), PGK1, FABP6, ACSM3, FBP1, HOGA1, SCP2, ECI2, ALDH6A1, PCYOX1, MPC2, HACD3, FOLR1, SHMT1, ACSF2, SLC3A1, CRYM (crystallin mu), CSPG5, PPA2, AKR1C3, DHFRP1, SPR (sepiapterinreductase), BDH2, GATM (glycine amidinotransferase), BCKDHB (branched chain keto acid dehydrogenase E1 subunit beta), PSMD10, ITIH2, PHGDH (phosphoglycerate dehydrogenase), BTD (biotinidase), SLC25A21, OAT (ornithine aminotransferase), OAZ1, HACD2, ECH1, BCO2, BNIPL (BCL2 interacting protein like), PAH (phenylalanine hydroxylase), LTC4S, DSEL (dermatansulfateepimerase like), ATPSCKMT (ATP synthase c subunit lysine N-methyltransferase), COX16, ACYP2, TRMT2B, TSTD1, ASB9, UCP2, AK4, CEBPZOS (CEBPZ opposite strand), SERHL2, HSPE1, SLC25A27, COA3, AGR2, MACROD1, COX11, NUBPL (nucleotide binding protein like), CRLS1, BRI3BP, DNLZ (DNL-type zinc finger), CYBA (cytochrome b-245 alpha chain), KLK6, NDUFS7, PSAP (prosaposin), NDUFAF8, ACOT13, SRI (sorcin), TOMM20, NDUFA2, SGPP1, TMEM160, HEBP1, AKR1B10, DPYSL2, ATP5IF1, NRG1 (neurogranin), DTYMK (deoxythymidylate kinase), NTHL1, PRDX2, TMBIM6, GSR (glutathione-disulfidereductase), TFRC (transferrin receptor), ANXA10, SAMD13, FAHD1, ENDOG (endonuclease G), COX14, FMO5, CRYZ (crystallin zeta), ACBD7, KCNAB1, CREG2 and TCN1 are novel biomarkers for SARS-CoV-2 infection.

PPI network was constructed and modules extracted to explore the molecular mechanism SARS-CoV-2 infection. In this investigation, the expression of hub genes in SARS-CoV-2 infection. Hub gene CH25H was liable for progression of porcine reproductive and respiratory syndrome virus infection [144], but this gene may be involved in advancement of SARS-CoV-2 infection. Hub genes such as TRAF1 [145], IFIT5 [146], CFTR (CF transmembrane conductance regulator) [147] and SMARCA2 [148] were key for progression of influenza viral infection, but these genes may be linked with progression of SARS-CoV-2 infection. Hub gene SAP30 was responsible for advancement of Rift Valley fever virus infection [149], but this gene may be key for advancement of SARS-CoV-2 infection. Hub gene APOE (apolipoprotein E) was responsible for development of SARS-CoV-2 infection [150]. As given in the PPI network and module analysis for up and down regulated genes, FBXO6, ADRB2, SYNPR (synaptoporin), LSMEM1, ARRDC3, ACTA1, LMO2, ATF3, KLF4, PCLAF (PCNA clamp associated factor), TUBA1A, AURKA (aurora kinase A), ARRB1, KPNA2, GALM (galactosemutarotase), HSD17B2, CEACAM7, SEMA3E, EPN3, ROM1, SORL1 and PDCD4 are novel biomarkers for SARS-CoV-2 infection.

Target gene - miRNA regulatory network and target gene - TF regulatory network analysis (up and down regulated genes) can be regarded as key to the understanding of SARS-CoV-2 infection and might also lead to new therapeutic approaches. Hub genes such as PROX1 [151] and HOXA13 [152] were responsible for development of hepatitis B virus infection, but these genes may be essential for advancement of for

SARS-CoV-2 infection. As given in the target gene - miRNA regulatory network and target gene - TF regulatory network for up and down regulated genes, APOL6, CHAC1, KLF2, CPM (carboxypeptidase M), CFI (complement factor I) and PGP (phosphoglycolate phosphatase) are novel biomarkers for SARS-CoV-2 infection.

Conclusion

The current investigation provided a comprehensive bioinformatics analysis of up and down regulated genes in SARS-CoV-2 infection. Analysis of these genes provided information regarding the molecular mechanisms of SARS-CoV-2 infection and key biomarkers or targets for the diagnosis and treatment of SARS-CoV-2 infection. However, further molecular biological experiments are required to confirm the action of the up and down regulated genes and pathways in different types of SARS-CoV-2 infection.

Declarations

Acknowledgements

We also thank Emanuel Wyler, Max Delbrück Center for Molecular Medicine, Berlin Institute for Medical Systems Biology, Robert Roessle Str 10, Berlin, Germany., very much, the author who deposited their expression profiling by high throughput sequencing dataset, GSE148729, into the public GEO database.

Conflict of interest

The authors declare that they have no conflict of interest.

Ethical approval

This article does not contain any studies with human participants or animals performed by any of the authors.

Informed consent

No informed consent because this study does not contain human or animals participants.

Author Contributions

B. V - Writing original draft, and review and editing

C. V - Software and investigation

Availability of data and materials

The datasets supporting the conclusions of this article are available in the GEO (Gene Expression Omnibus) (<https://www.ncbi.nlm.nih.gov/geo/>) repository. [(GSE148729)]

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Authors

Basavaraj Vastrad ORCID ID: [0000-0003-2202-7637](https://orcid.org/0000-0003-2202-7637)

Chanabasayya Vastrad ORCID ID: [0000-0003-3615-4450](https://orcid.org/0000-0003-3615-4450)

References

1. Andersen, G.; Rambaut, A.; Lipkin, W.I.; Holmes, E.C.; Garry, R.F. The proximal origin of SARS-CoV-2. *Nat. Med.* **2020**,26,450-452. doi:10.1038/s41591-020-0820-9
2. Wang, W.; Xu, Y.; Gao, R.; Lu, R.; Han, K.; Wu, G.; Tan, W. Detection of SARS-CoV-2 in Different Types of Clinical Specimens. **2020**,323,1843-1844. doi:10.1001/jama.2020.3786
3. Kissler, S.M.; Tedijanto, C.; Goldstein, E.; Grad, Y.H.; Lipsitch, M. Projecting the transmission dynamics of SARS-CoV-2 through the postpandemic period. **2020**,368,860-868. doi:10.1126/science.abb5793
4. Ou, X.; Liu, Y.; Lei, X.; Li, P.; Mi, D.; Ren, L.; Guo, L.; Guo, R.; Chen, T.; Hu, J.; et al. Characterization of spike glycoprotein of SARS-CoV-2 on virus entry and its immune cross-reactivity with SARS-CoV. *Commun.* **2020**,11,1620. doi:10.1038/s41467-020-15562-9
5. Hoffmann, M.; Kleine-Weber, H.; Schroeder, S.; Krüger, N.; Herrler, T.; Erichsen, S.; Schiergens, T.S.; Herrler, G.; Wu, N.H.; Nitsche, A.; et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. **2020**,181,271-280.e8. doi:10.1016/j.cell.2020.02.052
6. Clough, E.; Barrett, T. The Gene Expression Omnibus Database. *Mol. Biol.* **2016**,1418,93-110. doi:10.1007/978-1-4939-3578-9_5
7. Ritchie, M.E.; Phipson, B.; Wu, D.I.; Hu, Y.; Law, C.W.; Shi, W.; Smyth, G.K. limma powers differential expression analyses for RNA-sequencing and microarray studies. *Acids. Res.* **2015**,43,e47. doi:10.1093/nar/gkv007
8. Caspi, R.; Billington, R.; Ferrer, L.; Foerster, H.; Fulcher, C.A.; Keseler, I.M.; Kothari, A.; Krummenacker, M.; Latendresse, M.; Mueller, L.A.; et al The MetaCyc database of metabolic pathways and enzymes and the BioCyc collection of pathway/genome databases. *Acids. Res.* **2016**,44,D471–D480. doi:10.1093/nar/gkv1164

9. Kanehisa, M.; Sato, Y.; Furumichi, M.; Morishima, K.; Tanabe, M. New approach for understanding genome variations in KEGG. *Acids. Res.* **2019**,*47*,D590–D595. doi:10.1093/nar/gky962
10. Schaefer, C.F.; Anthony, K.; Krupa, S.; Buchoff, J.; Day, M.; Hannay, T.; Buetow, K.H. PID: the Pathway Interaction Database. *Acids. Res.* **2009**,*37*,D674–D679. doi:10.1093/nar/gkn653
11. Fabregat, A.; Jupe, S.; Matthews, L.; Sidiropoulos, K.; Gillespie, M.; Garapati, P.; Haw, R.; Jassal, B.; K€orninger, F.; May, B.; et al. The Reactome Pathway Knowledgebase. *Acids. Res.* **2018**,*46*,D649–D655. doi:10.1093/nar/gkx1132
12. Dahlquist, K.D.; Salomonis, N.; Vranizan, K.; Lawlor, S.C.; Conklin, B.R. GenMAPP, a new tool for viewing and analyzing microarray data on biological pathways. *Genet.* **2002**,*31*,19–20. doi:10.1038/ng0502-19
13. Subramanian, A.; Tamayo, P.; Mootha, V.K.; Mukherjee, S.; Ebert, B.L.; Gillette, M.A.; Paulovich, A.; Pomeroy, S.L.; Golub, T.R.; Lander, E.S.; et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Natl. Acad. Sci. U. S. A.* **2005**,*102*,15545–15550. doi:10.1073/pnas.0506580102
14. Mi, H.; Huang, X.; Muruganujan, A.; Tang, H.; Mills, C.; Kang, D.; Thomas, P.D. PANTHER version 11: expanded annotation data from Gene Ontology and Reactome pathways, and data analysis tool enhancements. *Acids. Res.* **2017**,*45*,D183–D189. doi:10.1093/nar/gkw1138
15. Petri, V.; Jayaraman, P.; Tutaj, M.; Hayman, G.T.; Smith, J.R.; De Pons, J.; Lauderkind, S.J.; Lowry, T.F.; Nigam, R.; Wang, S.J.; et al. The pathway ontology - updates and applications. *Biomed. Semantics.* **2014**,*5*,7. doi:10.1186/2041-1480-5-7
16. Jewison, T.; Su, Y.; Disfany, F.M.; Liang, Y.; Knox, C.; Maciejewski, A.; Poelzer, J.; Huynh, J.; Zhou, Y.; Arndt, D.; et al. SMPDB 2.0: big improvements to the Small Molecule Pathway Database. *Acids. Res.* **2014**,*42*,D478–D484. doi:10.1093/nar/gkt1067
17. Chen, J.; Bardes, E.E.; Aronow, B.J.; Jegga, A.G. ToppGene Suite for gene list enrichment analysis and candidate gene prioritization. *Acids. Res.* **2009**,*37*,W305-W311. doi:10.1093/nar/gkp427
18. Thomas, P.D. The Gene Ontology and the Meaning of Biological Function. *Mol. Biol.* **2017**,*1446*,15-24. doi:10.1007/978-1-4939-3743-1_2
19. Alanis-Lobato, G.; Andrade-Navarro, M.A.; Schaefer, M.H. HIPPIE v2.0: enhancing meaningfulness and reliability of protein-protein interaction networks. *Acids. Res.* **2017**,*45*,D408-D414. doi:10.1093/nar/gkw985
20. Orchard, S.; Ammari, M.; Aranda, B.; Breuza, L.; Briganti, L.; Broackes-Carter, F.; Campbell, N.H.; Chavali, G.; Chen, C.; del-Toro, N.; et al. The MIntAct project–IntAct as a common curation platform for 11 molecular interaction databases. *Acids. Res.* **2014**,*42*,D358-D363. doi:10.1093/nar/gkt1115
21. Chatr-Aryamontri, A.; Oughtred, R.; Boucher, L.; Rust, J.; Chang, C.; Kolas, N.K.; O'Donnell, L.; Oster, S.; Theesfeld, C.; Sellam, A.; et al. The BioGRID interaction database: 2017 update. *Acids. Res.* **2017**,*45*,D369-D379. doi:10.1093/nar/gkw1102
22. Keshava Prasad, T.S.; Goel, R.; Kandasamy, K.; Keerthikumar, S.; Kumar, S.; Mathivanan, S.; Telikicherla, D.; Raju, R.; Shafreen, B.; Venugopal, A.; et al. Human Protein Reference Database–2009

- update. *Acids.Res.* **2009**,37,D767-D772. doi:10.1093/nar/gkn892
23. Licata, L.; Briganti, L.; Peluso, D.; Perfetto, L.; Iannuccelli, M.; Galeota, E.; Sacco, F.; Palma, A.; Nardozza, A.P.; Santonico, E.; et al. MINT, the molecular interaction database: 2012 update. *Acids. Res.* **2012**,40,D857-D861. doi:10.1093/nar/gkr930
 24. Isserlin, R.; El-Badrawi, R.A.; Bader, G.D. The Biomolecular Interaction Network Database in PSI-MI 2.5. *Database (Oxford)*. **2011**,2011.;baq037. doi:1093/database/baq037
 25. Pagel, P.; Kovac, S.; Oesterheld, M.; Brauner, B.; Dunger-Kaltenbach, I.; Frishman, G.; Montrone, C.; Mark, P.; Stümpflen, V.; Mewes, H.W.; et al. The MIPS mammalian protein-protein interaction database. **2005**,21,832-834. doi:10.1093/bioinformatics/bti115
 26. Salwinski, L.; Miller, C.S.; Smith, A.J.; Pettit, F.K.; Bowie, J.U.; Eisenberg, D. The Database of Interacting Proteins: 2004 update. *Acids. Res.* **2004**,32,D449-D451. doi:10.1093/nar/gkh086
 27. Shannon, P.; Markiel, A.; Ozier, O.; Baliga, N.S.; Wang, J.T.; Ramage, D.; Amin, N.; Schwikowski, B.; Ideker, T. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Res.* **2003**,13,2498-2504. doi:10.1101/gr.1239303
 28. Przulj, N.; Wigle, D.A.; Jurisica, I. Functional topology in a network of protein interactions. **2004**,20,340–348. doi:10.1093/bioinformatics/btg415
 29. Nguyen, T.P.; Liu, W.C.; Jordán, F. Inferring pleiotropy by network analysis: linked diseases in the human PPI network. *Syst. Biol.* **2011**,5,179. doi:10.1186/1752-0509-5-179
 30. Shi, Z.; Zhang, B. Fast network centrality analysis using GPUs. *Bioinformatics.* **2011**,12,149. doi:10.1186/1471-2105-12-149
 31. Fadhal, E.; Gamiieldien, J.; Mwambene, E.C. Protein interaction networks as metric spaces: a novel perspective on distribution of hubs. *Syst. Biol.* **2014**,8,6. doi:10.1186/1752-0509-8-6
 32. Wang, J.; Li, M.; Wang, H.; Pan, Y. Identification of essential proteins based on edge clustering coefficient. *IEEE/ACM. Trans. Comput. Biol. Bioinform.* **2012**,9,1070–1080. doi:1109/TCBB.2011.147
 33. Zaki, N.; Efimov, D.; Berenguères, J. Protein complex detection using interaction reliability assessment and weighted clustering coefficient. *Bioinformatics.* **2013**,14,163. doi:10.1186/1471-2105-14-163
 34. Vlachos, I.S.; Paraskevopoulou, M.D.; Karagkouni, D.; Georgakilas, G.; Vergoulis, T.; Kanellos, I.; Anastasopoulos, I.L.; Maniou, S.; Karathanou, K.; Kalfakakou, D.; et al. DIANA-TarBase v7.0: indexing more than half a million experimentally supported miRNA:mRNA interactions. *Acids. Res.* **2015**,43,D153-D159. doi:10.1093/nar/gku1215
 35. Chou, C.H.; Shrestha, S.; Yang, C.D.; Chang, N.W.; Lin, Y.L.; Liao, K.W.; Huang, W.C.; Sun, T.H.; Tu, S.J.; Lee, W.H.; et al. miRTarBase update 2018: a resource for experimentally validated microRNA-target interactions. *Acids. Res.* **2018**,46,D296-D302. doi:10.1093/nar/gkx1067
 36. Xiao, F.; Zuo, Z.; Cai, G.; Kang, S.; Gao, X.; Li, T. miRecords: an integrated resource for microRNA-target interactions. *Acids. Res.* **2009**,37,D105-D110. doi:10.1093/nar/gkn851
 37. Jiang, Q.; Wang, Y.; Hao, Y.; Juan, L.; Teng, M.; Zhang, X.; Li, M.; Wang, G.; Liu, Y. miR2Disease: a manually curated database for microRNA deregulation in human disease. *Acids. Res.* **2009**,37,D98-

104. doi:10.1093/nar/gkn714
38. Huang, Z.; Shi, J.; Gao, Y.; Cui, C.; Zhang, S.; Li, J.; Zhou, Y.; Cui, Q. HMDD v3.0: a database for experimentally supported human microRNA-disease associations. *Acids. Res.* **2019**,47,D1013-D1017. doi:10.1093/nar/gky1010Z
 39. Ruepp, A.; Kowarsch, A.; Schmidl, D.; Buggenthin, F.; Brauner, B.; Dunger, I.; Fobo, G.; Frishman, G.; Montrone, C.; Theis, F.J. PhenomiR: a knowledgebase for microRNA expression in diseases and biological processes. *Biol.* **2010**,11,R6. doi:10.1186/gb-2010-11-1-r6
 40. Liu, X.; Wang, S.; Meng, F.; Wang, J.; Zhang, Y.; Dai, E.; Yu, X.; Li, X.; Jiang, W. SM2miR: a database of the experimentally validated small molecules' effects on microRNA expression. **2013**,29,409-411. doi:10.1093/bioinformatics/bts698
 41. Rukov, J.L.; Wilentzik, R.; Jaffe, I.; Vinther, J.; Shomron, N. PharmacomiR: linking microRNAs and drug effects. *Bioinform.* **2014**,15,648-659. doi:10.1093/bib/bbs082
 42. Dai, E.; Yu, X.; Zhang, Y.; Meng, F.; Wang, S.; Liu, X.; Liu, D.; Wang, J.; Li, X.; Jiang, W. EpimiR: a database of curated mutual regulation between miRNAs and epigenetic modifications. (*Oxford*). **2014**,2014,bau023. doi:10.1093/database/bau023
 43. Li, J.H.; Liu, S.; Zhou, H.; Qu, L.H.; Yang, J.H. starBase v2.0: decoding miRNA-ceRNA, miRNA-ncRNA and protein-RNA interaction networks from large-scale CLIP-Seq data. *Acids. Res.* **2014**,42,D92-D97. doi:10.1093/nar/gkt1248
 44. Zhou, G.; Soufan, O.; Ewald, J.; Hancock, R.E.W.; Basu, N.; Xia, J. NetworkAnalyst 3.0: a visual analytics platform for comprehensive gene expression profiling and meta-analysis. *Acids. Res.* **2019**,47,W234-W241. doi:10.1093/nar/gkz240
 45. Khan, A.; Fornes, O.; Stigliani, A.; Gheorghe, M.; Castro-Mondragon, J.A.; van der Lee, R.; Bessy, A.; Chèneby, J.; Kulkarni, S.R.; Tan, G.; et al. JASPAR 2018: update of the open-access database of transcription factor binding profiles and its web framework. *Acids. Res.* **2018**,46,D260-D266. doi:10.1093/nar/gkx1126
 46. Robin, X.; Turck, N.; Hainard, A.; Tiberti, N.; Lisacek, F.; Sanchez J.C.; Müller M. pROC: an open-source package for R and S+ to analyze and compare ROC curves. *Bioinformatics.* **2011**,12,77. doi:10.1186/1471-2105-12-77
 47. Lin, X.; Yu, S.; Ren, P.; Sun, X.; Jin, M. Human microRNA-30 inhibits influenza virus infection by suppressing the expression of SOCS1, SOCS3, and NEDD4. *Microbiol.* **2020**,22,e13150. doi:10.1111/cmi.13150
 48. Liu, Y.; Li, S.; Zhang, G.; Nie, G.; Meng, Z.; Mao, D.; Chen, C.; Chen, X.; Zhou, B.; Zeng, G. Genetic variants in IL1A and IL1B contribute to the susceptibility to 2009 pandemic H1N1 influenza A virus. *Immunol.* **2013**,14,37. doi:10.1186/1471-2172-14-37
 49. Mgbemena, V.; Segovia, J.; Chang, T.H.; Bose, S. KLF6 and iNOS regulates apoptosis during respiratory syncytial virus infection. *Cell Immunol.* **2013**,283,1-7. doi:10.1016/j.cellimm.2013.06.002
 50. Robitaille, A.C.; Caron, E.; Zucchini, N.; Mukawera, E.; Adam, D.; Mariani, M.K.; Gélinas, A.; Fortin, A.; Brochiero, E.; Grandvaux, N. DUSP1 regulates apoptosis and cell migration, but not the JIP1-protected

- cytokine response, during Respiratory Syncytial Virus and Sendai Virus infection. *Rep.* **2017**,7,17388. doi:10.1038/s41598-017-17689-0
51. Brand, H.K.; Ahout, I.M.; De Ridder, D.; van Diepen, A.; Li, Y.; Zaalberg, M.; Andeweg, A.; Roeleveld, N.; De Groot, R.; Warris, A.; et al. Olfactomedin 4 Serves as a Marker for Disease Severity in Pediatric Respiratory Syncytial Virus (RSV) Infection. *One.* **2015**,10,e0131927. doi:10.1371/journal.pone.0131927
 52. Lagunas-Rangel, F.A.; Chávez-Valencia, V. High IL-6/IFN- γ ratio could be associated with severe disease in COVID-19 patients. *Med. Virol.* **2020**,10.1002/jmv.25900. doi:10.1002/jmv.25900
 53. Diao, B.; Huang, X.; Guo, S.; Yang, C.; Liu, G.; Chen, Y.; Wu, Y. MAGT1-mediated disturbance of Mg²⁺ homeostasis lead to exhausted of HBV-infected NK and CD8⁺ T cells. *Rep.* **2017**,7,13594. doi:10.1038/s41598-017-11522-4
 54. Mazar, J.; Li, Y.; Rosado, A.; Phelan, P.; Kedarinath, K.; Parks, G.D.; Alexander, K.A.; Westmoreland, T. J. Zika virus as an oncolytic treatment of human neuroblastoma cells requires CD24. *One.* **2018**,13,e0200358. doi:10.1371/journal.pone.0200358
 55. Stücker, I.; Lorient, M.A.; N'Koutchou, G.; Cénée, S.; Bodin, L.; Mulot, C.; Gelu-Simeon, M.; Pelletier, L.; Bronowicki, J.P.; Degos, F.; et al. UDP-glucuronosyltransferase UGT1A7 genetic polymorphisms in hepatocellular carcinoma: a differential impact according to seropositivity of HBV or HCV markers?. *Cancer.* **2007**,7,214. doi:10.1186/1471-2407-7-214
 56. Zhang, J.; Yan, M.; Gu, W.; Chen, A.; Liu, J.; Li, L.; Zhang, S.; Liu, G. Downregulation of Aquaporins (AQP1 and AQP5) and Na,K-ATPase in Porcine Reproductive and Respiratory Syndrome Virus-Infected Pig Lungs. **2018**,41,1104-1114. doi:10.1007/s10753-018-0762-2
 57. Huang, W.; Hu, K.; Luo, S.; Zhang, M.; Li, C.; Jin, W.; Liu, Y.; Griffin, G.E.; Shattock, R.J.; Hu, Q. Herpes simplex virus type 2 infection of human epithelial cells induces CXCL9 expression and CD4⁺ T cell migration via activation of p38-CCAAT/enhancer-binding protein- β pathway. *Immunol.* **2012**,188,6247-6257. doi:10.4049/jimmunol.1103706
 58. Bertin, J.; Jalaguier, P.; Barat, C.; Roy, M.A.; Tremblay, M.J.; Exposure of human astrocytes to leukotriene C4 promotes a CX3CL1/fractalkine-mediated transmigration of HIV-1-infected CD4⁺ T cells across an in vitro blood-brain barrier model. **2014**,454-455,128-138. doi:10.1016/j.virol.2014.02.007
 59. Gagné, B.; Tremblay, N.; Park, A.Y.; Baril, M.; Lamarre, D. Importin β 1 targeting by hepatitis C virus NS3/4A protein restricts IRF3 and NF- κ B signaling of IFNB1 antiviral response. **2017**,18,362-377. doi:10.1111/tra.12480
 60. Levy, Y.; Lacabaratz, C.; Weiss, L.; Viard, J.P.; Goujard, C.; Lelièvre, J.D.; Boué, F.; Molina, J.M.; Rouzioux, C.; Avettand-Fénoël, V.; et al. Enhanced T cell recovery in HIV-1-infected adults through IL-7 treatment. *Clin. Invest.* **2009**,119,997-1007. doi:10.1172/JCI38052
 61. Shin, G.C.; Kang, H.S.; Lee, A.R.; Kim, K.H. Hepatitis B virus-triggered autophagy targets TNFRSF10B/death receptor 5 for degradation to limit TNFSF10/TRAIL response. **2016**,12,2451-2466. doi:10.1080/15548627.2016.1239002

62. Lu, Y.F.; Mauger, D.M.; Goldstein, D.B.; Urban, T.J.; Weeks, K.M.; Bradrick, S.S. IFNL3 mRNA structure is remodeled by a functional non-coding polymorphism associated with hepatitis C virus clearance. *Rep.* **2015**,*5*,16037. Published 2015 Nov 4. doi:10.1038/srep16037
63. Lee, S.; Park, U.; Lee, Y.I. Hepatitis C virus core protein transactivates insulin-like growth factor II gene transcription through acting concurrently on Egr1 and Sp1 sites. **2001**,283,167-177. doi:10.1006/viro.2001.0892
64. Chen, L.; Li, S.; McGilvray, I. The ISG15/USP18 ubiquitin-like pathway (ISGylation system) in hepatitis C virus infection and resistance to interferon therapy. *J Biochem. Cell. Biol.* **2011**,*43*,1427-1431. doi:10.1016/j.biocel.2011.06.006
65. Xia, M.; Gonzalez, P.; Li, C.; Meng, G.; Jiang, A.; Wang, H.; Gao, Q.; Debatin, K.M.; Beltinger, C.; Wei, J. Mitophagy enhances oncolytic measles virus replication by mitigating DDX58/RIG-I-like receptor signaling. *Virology*. **2014**,*88*,5152-5164. doi:10.1128/JVI.03851-13
66. Zhao, P.; Jiang, T.; Zhong, Z.; Zhao, L.; Yang, S.; Xia, X. Inhibition of rabies virus replication by interferon-stimulated gene 15 and its activating enzyme UBA7. *Genet. Evol.* **2017**,*56*,44-53. doi:10.1016/j.meegid.2017.10.016
67. Das, A.; Dinh, P.X.; Panda, D.; Pattnaik, A.K. Interferon-inducible protein IFI35 negatively regulates RIG-I antiviral signaling and supports vesicular stomatitis virus replication. *Virology*. **2014**,*88*,3103-3113. doi:10.1128/JVI.03202-13
68. Fensterl, V.; Wetzel J.L.; Sen G.C. Interferon-induced protein Ifit2 protects mice from infection of the peripheral nervous system by vesicular stomatitis virus. *Virology*. **2014**,*88*,10303-10311. doi:10.1128/JVI.01341-14
69. Buggele, W.A.; Horvath, C.M. MicroRNA profiling of Sendai virus-infected A549 cells identifies miR-203 as an interferon-inducible regulator of IFIT1/ISG56. *Virology*. **2013**,*87*,9260-9270. doi:10.1128/JVI.01064-13
70. van Manen, D.; Rits, M.A.; Beugeling, C.; van Dort, K.; Schuitemaker, H.; Kootstra, N.A. The effect of Trim5 polymorphisms on the clinical course of HIV-1 infection. *Pathog.* **2008**,*4*,e18. doi:10.1371/journal.ppat.0040018
71. Kurokawa, C.; Iankov, I.D.; Galanis, E. A key anti-viral protein, RSAD2/VIPERIN, restricts the release of measles virus from infected cells. *Res.* **2019**,*263*,145-150. doi:10.1016/j.virusres.2019.01.014
72. Long, X.; Li, Y.; Qi, Y.; Xu, J.; Wang, Z.; Zhang, X.; Zhang, D.; Zhang, L.; Huang, J. XAF1 contributes to dengue virus-induced apoptosis in vascular endothelial cells. *J.* **2013**,*27*,1062-1073. doi:10.1096/fj.12-213967
73. Wang, H.; Bai, J.; Fan, B. Li.; Y, Zhang.; Q, Jiang, P. The Interferon-Induced Mx2 Inhibits Porcine Reproductive and Respiratory Syndrome Virus Replication. *Interferon. Cytokine. Res.* **2016**,*36*,129-139. doi:10.1089/jir.2015.0077
74. Niu, P.; Shabir, N.; Khatun, A.; Seo, B.J.; Gu, S.; Lee, S.M.; Lim, S.K.; Kim, K.S.; Kim, W.I. Effect of polymorphisms in the GBP1, Mx1 and CD163 genes on host responses to PRRSV infection in pigs. *Microbiol.* **2016**,*182*,187-195. doi:10.1016/j.vetmic.2015.11.010

75. Manocha, G.D, Mishra, R, Sharma, N, Kumawat, K.L, Basu, A, Singh, S.K. Regulatory role of TRIM21 in the type-I interferon pathway in Japanese encephalitis virus-infected human microglial cells. *Neuroinflammation*. **2014**,11,24. doi:10.1186/1742-2094-11-24
76. Gallego-Escuredo, J.M.; Domingo, P.; del Mar Gutiérrez, M.; Mateo, M.G.; Cabeza, M.C.; Fontanet, A.; Vidal, F.; Domingo, J.C.; Giralt, M.; Villarroya, F. Reduced levels of serum FGF19 and impaired expression of receptors for endocrine FGFs in adipose tissue from HIV-infected patients. *Acquir. Immune. Defic. Syndr*. **2012**,61,527-534. doi:10.1097/QAI.0b013e318271c2c7
77. Luo, H.; Winkelmann, E.R.; Zhu, S.; Ru, W.; Mays, E.; Silvas, J.A.; Vollmer, L.L.; Gao, J.; Peng, B.H.; Bopp, N.E.; et al. Peli1 facilitates virus replication and promotes neuroinflammation during West Nile virus infection. *Clin. Invest*. **2018**,128,4980-4991. doi:10.1172/JCI99902
78. Knapp, S.; Yee, L.J.; Frodsham, A.J.; Hennig, B.J.; Hellier, S.; Zhang, L.; Wright, M.; Chiaramonte, M.; Graves, M.; Thomas, H.C.; et al. Polymorphisms in interferon-induced genes and the outcome of hepatitis C virus infection: roles of MxA, OAS-1 and PKR. *Immun*. **2003**,4,411-419. doi:10.1038/sj.gene.6363984
79. Manley, G.C.A, Stokes, C.A, Marsh, E.K, Sabroe, I, Parker, L.C. DUSP10 Negatively Regulates the Inflammatory Response to Rhinovirus through Interleukin-1 β Signaling. *Virology*. **2019**,93,e01659-18. doi:10.1128/JVI.01659-18
80. Chen, J.; Ng, M.M.; Chu, J.J. Activation of TLR2 and TLR6 by Dengue NS1 Protein and Its Implications in the Immunopathogenesis of Dengue Virus Infection. *PLoS Pathog*. **2015**,11,e1005053. doi:1371/journal.ppat.1005053
81. Leong, C.R.; Oshiumi, H.; Okamoto, M.; Azuma, M.; Takaki, H.; Matsumoto, M.; Chayama, K.; Seya, T. A MAVS/TICAM-1-independent interferon-inducing pathway contributes to regulation of hepatitis B virus replication in the mouse hydrodynamic injection model. *Innate. Immun*. **2015**,7,47-58. doi:10.1159/000365113
82. Frakolaki, E.; Kalliampakou, K.I.; Kaimou, P.; Moraiti, M.; Kolaitis, N.; Boleti, H.; Koskinas, J.; Vassilacopoulou, D.; Vassilaki, N. Emerging Role of L-Dopa Decarboxylase in Flaviviridae Virus Infections. **2019**,8,837. doi:10.3390/cells8080837
83. Panagopoulos, P.; Maltezos, E.; Hatzakis, A.; Paraskevis, D. Hyperbilirubinemia in atazanavir treated HIV-infected patients: the impact of the UGT1A1*28 allele. *Pers. Med*. **2017**,10,205-208. doi:10.2147/PGPM.S107152
84. Wu, N.; Gao, N.; Fan, D.; Wei, J.; Zhang, J.; An, J. miR-223 inhibits dengue virus replication by negatively regulating the microtubule-destabilizing protein STMN1 in EAhy926 cells. *Infect*. **2014**,16,911-922. doi:10.1016/j.micinf.2014.08.011
85. Yang, J.J.; Fan, H.Z.; Tian, T.; Wu, M.P.; Xie, C.N.; Huang, P.; Yu, R.B.; Yi, H.G.; Zhang, Y.; Wang, J. Impact of CYP2R1, CYP27A1 and CYP27B1 genetic polymorphisms controlling vitamin D metabolism on susceptibility to hepatitis C virus infection in a high-risk Chinese population. *Virology*. **2019**,164,2909-2918. doi:10.1007/s00705-019-04378-8

86. Kozlowski, S.; Corr, M.; Takeshita, T.; Boyd, L.F.; Pendleton, C.D.; Germain, R.N.; Berzofsky, J.A.; Margulies, D.H. Serum angiotensin-1 converting enzyme activity processes a human immunodeficiency virus 1 gp160 peptide for presentation by major histocompatibility complex class I molecules. *Exp. Med.* **1992**,175,1417-1422. doi:10.1084/jem.175.6.1417
87. Lai, C.; Wang, K.; Zhao, Z.; Zhang, L.; Gu, H.; Yang, P.; Wang, X. C-C Motif Chemokine Ligand 2 (CCL2) Mediates Acute Lung Injury Induced by Lethal Influenza H7N9 Virus. *Microbiol.* **2017**,8,587. doi:10.3389/fmicb.2017.00587
88. Beshara, R.; Sencio, V.; Soulard, D.; Barthélémy, A.; Fontaine, J.; Pinteau, T.; Deruyter, L.; Ismail, M.B.; Paget, C.; Sirard, J.C.; et al. Alteration of Flt3-Ligand-dependent de novo generation of conventional dendritic cells during influenza infection contributes to respiratory bacterial superinfection. *Pathog.* **2018**,14,e1007360. doi:10.1371/journal.ppat.1007360
89. Huipao, N.; Borwornpinyo, S.; Wiboon-Ut, S.; Campbell, C.R.; Lee, I.H.; Hiranyachattada, S.; Sukasem, C.; Thitithanyanont, A.; Pholpramool, C.; Cook, D.I.; et al. P2Y6 receptors are involved in mediating the effect of inactivated avian influenza virus H5N1 on IL-6 & CXCL8 mRNA expression in respiratory epithelium. *One.* **2017**,12,e0176974. doi:10.1371/journal.pone.0176974
90. Oh, J.W.; Lee, H.B.; Park, I.K.; Kang, J.O. Interleukin-6, interleukin-8, interleukin-11, and interferon-gamma levels in nasopharyngeal aspirates from wheezing children with respiratory syncytial virus or influenza A virus infection. *Allergy. Immunol.* **2002**,13,350-356. doi:10.1034/j.1399-3038.2002.02018.x
91. Law, A.H.; Lee, D.C.; Yuen, K.Y.; Peiris, M.; Lau, A.S. Cellular response to influenza virus infection: a potential role for autophagy in CXCL10 and interferon-alpha induction. *Mol. Immunol.* **2010**,7,263-270. doi:10.1038/cmi.2010.25
92. Lee, D.C.; Cheung, C.Y.; Law, A.H.; Mok, C.K.; Peiris, M.; Lau, A.S. p38 mitogen-activated protein kinase-dependent hyperinduction of tumor necrosis factor alpha expression in response to avian influenza virus H5N1. *Viol.* **2005**,79,10147-10154. doi:10.1128/JVI.79.16.10147-10154.2005
93. Du, N.; Kwon, H.; Li, P.; West, E.E.; Oh, J.; Liao, W.; Yu, Z.; Ren, M.; Leonard, W.J. EGR2 is critical for peripheral naïve T-cell differentiation and the T-cell response to influenza. *Natl. Acad. Sci. U S A.* **2014**,111,16484-16489. doi:10.1073/pnas.1417215111
94. Pagani, I.; Di Pietro, A.; Oteiza, A.; Ghitti, M.; Mechti, N.; Naffakh, N.; Vicenzi, E. Mutations Conferring Increased Sensitivity to Tripartite Motif 22 Restriction Accumulated Progressively in the Nucleoprotein of Seasonal Influenza A (H1N1) Viruses between 1918 and 2009. **2018**,3,e00110-18.doi:10.1128/mSphere.00110-18
95. Li, W.; Wang, G.; Zhang, H.; Zhang, D.; Zeng, J.; Chen, X.; Xu, Y.; Li, K. Differential suppressive effect of promyelocytic leukemia protein on the replication of different subtypes/strains of influenza A virus. *Biophys. Res. Commun.* **2009**,389,84-89. doi:10.1016/j.bbrc.2009.08.091
96. Jiang, H.; Shen, S.M.; Yin, J.; Zhang, P.P.; Shi, Y. Sphingosine 1-phosphate receptor 1 (S1PR1) agonist CYM5442 inhibits expression of intracellular adhesion molecule 1 (ICAM1) in endothelial cells infected with influenza A viruses. *One.* **2017**,12,e0175188. doi:10.1371/journal.pone.0175188

97. Yu, M.; Qi, W.; Huang, Z.; Zhang, K.; Ye, J.; Liu, R.; Wang, H.; Ma, Y.; Liao, M.; Ning, Z. Expression profile and histological distribution of IFITM1 and IFITM3 during H9N2 avian influenza virus infection in BALB/c mice. *Microbiol. Immunol.* **2015**,204,505-514. doi:10.1007/s00430-014-0361-2
98. Wang, H.; Chen, L.; Luo, J.; He, H. NP and NS1 proteins of H5N1 virus significantly upregulated IFITM1, IFITM2, and IFITM3 in A549 cells. *Health. Sci.* **2019**,19,1402-1410. doi:10.4314/ahs.v19i1.13
99. Sanyal, S.; Ashour, J.; Maruyama, T.; Altenburg, A.F.; Cragnolini, J.J.; Bilate, A.; Avalos, A.M.; Kundrat, L.; García-Sastre, A.; Ploegh, H.L. Type I interferon imposes a TSG101/ISG15 checkpoint at the Golgi for glycoprotein trafficking during influenza virus infection. *Host. Microbe.* **2013**,14,510-521. doi:10.1016/j.chom.2013.10.011
100. Ye, S.; Lowther, S.; Stambas, J. Inhibition of reactive oxygen species production ameliorates inflammation induced by influenza A viruses via upregulation of SOCS1 and SOCS3. *Viol.* **2015**,89,2672-2683. doi:10.1128/JVI.03529-14
101. Kuriakose, T.; Zheng, M.; Neale, G.; Kanneganti, T.D. IRF1 Is a Transcriptional Regulator of ZBP1 Promoting NLRP3 Inflammasome Activation and Cell Death during Influenza Virus Infection. *Immunol.* **2018**,200,1489-1495. doi:10.4049/jimmunol.1701538
102. Rosenberger, C.M.; Podyminogin, R.L.; Diercks, A.H.; Treuting, P.M.; Peschon, J.J.; Rodriguez, D.; Gundapuneni, M.; Weiss, M.J.; Aderem, A. miR-144 attenuates the host response to influenza virus by targeting the TRAF6-IRF7 signaling axis. *Pathog.* 2017;13(4):e1006305. doi:10.1371/journal.ppat.1006305
103. Chai, W.; Li J.; Shanguan, Q.; Liu, Q.; Li, X.; Qi, D.; Tong, X.; Liu, W.; Ye, X. Lnc-ISG20 Inhibits Influenza A Virus Replication by Enhancing ISG20 Expression. *Viol.* **2018**,92,e00539-18. doi:10.1128/JVI.00539-18
104. Warnking, K.; Klemm, C.; Löffler, B.; Niemann, S.; van Krüchten, A.; Peters, G.; Ludwig, S.; Ehrhardt, C. Super-infection with *Staphylococcus aureus* inhibits influenza virus-induced type I IFN signalling through impaired STAT1-STAT2 dimerization. *Microbiol.* **2015**,17,303-317. doi:10.1111/cmi.12375
105. Zhao, L.; Zhang, X.; Wu, Z.; Huang, K.; Sun, X.; Chen, H.; Jin, M. The Downregulation of MicroRNA hsa-miR-340-5p in IAV-Infected A549 Cells Suppresses Viral Replication by Targeting RIG-I and OAS2. *Ther. Nucleic. Acids.* **2019**,14,509-519. doi:10.1016/j.omtn.2018.12.014
106. Tang, Y.; Zhong, G.; Zhu, L.; Liu, X.; Shan, Y.; Feng, H.; Bu, Z.; Chen, H.; Wang, C. Herc5 attenuates influenza A virus by catalyzing ISGylation of viral NS1 protein. *J. Immunol.* **2010**,184,5777-5790. doi:10.4049/jimmunol.0903588
107. Maelfait, J.; Roose, K.; Bogaert, P.; Sze, M.; Saelens, X.; Pasparakis, M.; Carpentier, I.; van Loo, G.; Beyaert, R. A20 (Tnfrsf25) deficiency in myeloid cells protects against influenza A virus infection. *Pathog.* **2012**,8,e1002570. doi:10.1371/journal.ppat.1002570
108. Shi, T.; He, Y.; Sun, W.; Wu, Y.; Li, L.; Jie, Z.; Su, X. Respiratory Syncytial virus infection compromises asthma tolerance by recruiting interleukin-17A-producing cells via CCR6-CCL20 signaling. *Immunol.* **2017**,88,45-57. doi:10.1016/j.molimm.2017.05.017

109. Stier, M.T.; Bloodworth, M.H.; Toki, S.; Newcomb, D.C.; Goleniewska, K.; Boyd, K.L.; Quitalig, M.; Hotard, A.L.; Moore, M.L.; Hartert, T.V.; et al. Respiratory syncytial virus infection activates IL-13-producing group 2 innate lymphoid cells through thymic stromal lymphopoietin. *Allergy. Clin. Immunol.* **2016**,138,814-824.e11. doi:10.1016/j.jaci.2016.01.050
110. Inchley, C.S.; Osterholt, H.C.; Sonerud, T.; Fjærli, H.O.; Nakstad, B. Downregulation of IL7R, CCR7, and TLR4 in the cord blood of children with respiratory syncytial virus disease. *Infect. Dis.* **2013**,208,1431-1435. doi:10.1093/infdis/jit336
111. Ternette, N.; Wright, C.; Kramer, H.B.; Altun, M.; Kessler, B.M. Label-free quantitative proteomics reveals regulation of interferon-induced protein with tetratricopeptide repeats 3 (IFIT3) and 5'-3'-exoribonuclease 2 (XRN2) during respiratory syncytial virus infection. *J.* **2011**,8,442. doi:10.1186/1743-422X-8-442
112. Dhar, J.; Cuevas, R.A.; Goswami, R.; Zhu, J.; Sarkar, S.N.; Barik S. 2'-5'-Oligoadenylate Synthetase-Like Protein Inhibits Respiratory Syncytial Virus Replication and Is Targeted by the Viral Nonstructural Protein 1. *Virology*. **2015**,89,10115-10119. doi:10.1128/JVI.01076-15
113. Yang, S.; Shan, T.; Zhou, Y.; Jiang, Y.; Tong, W.; Liu, F.; Wen, F.; Zhang, Q.; Tong, G. Molecular cloning and characterizations of porcine SAMHD1 and its roles in replication of highly pathogenic porcine reproductive and respiratory syndrome virus. *Comp. Immunol.* **2014**,47,234-246. doi:10.1016/j.dci.2014.07.024
114. Wu, D.; Yang, X.O. TH17 responses in cytokine storm of COVID-19: An emerging target of JAK2 inhibitor Fedratinib. *Microbiol. Immunol. Infect.* **2020**,53,368-370. doi:10.1016/j.jmii.2020.03.005
115. Monteil, V.; Kwon, H.; Prado, P.; Hagelkrüys, A.; Wimmer, R.A.; Stahl, M.; Leopoldi, A.; Garreta, E.; Del Pozo, C.H.; Prosper, F.; et al. Inhibition of SARS-CoV-2 Infections in Engineered Human Tissues Using Clinical-Grade Soluble Human ACE2. **2020**,181,905-913.e7. doi:10.1016/j.cell.2020.04.004
116. O'Brien, T.R.; Pfeiffer, R.M.; Paquin, A.; Kuhs, K.A.; Chen, S.; Bonkovsky, H.L.; Edlin, B.R.; Howell, C.D.; Kirk, G.D.; Kuniholm, M.H.; et al. Comparison of functional variants in IFNL4 and IFNL3 for association with HCV clearance. *Hepatology*. **2015**,63,1103-1110. doi:10.1016/j.jhep.2015.06.035
117. Jin, W.J.; Chen, C.F.; Liao, H.Y.; Gong, L.L.; Yuan, X.H.; Zhao, B.B.; Zhang, D.; Feng, X.; Liu, J.J.; Wang, Y.; et al. Downregulation of the AU-rich RNA-binding protein ZFP36 in chronic HBV patients: implications for anti-inflammatory therapy. *One.* **2012**,7,e33356. doi:10.1371/journal.pone.0033356
118. Jeong, H.Y.; Lee, Y.J.; Seo, S.K.; Lee, S.W.; Park, S.J.; Lee, J.N.; Sohn, H.S.; Yao, S.; Chen, L.; Choi, I. Blocking of monocyte-associated B7-H1 (CD274) enhances HCV-specific T cell immunity in chronic hepatitis C infection. *Leukoc. Biol.* **2008**,83,755-764. doi:10.1189/jlb.0307168
119. Kanokudom, S.; Vilaivan, T.; Wikan, N.; Thepparit, C.; Smith, D.R.; Assavalapsakul, W. miR-21 promotes dengue virus serotype 2 replication in HepG2 cells. *Res.* **2017**,142,169-177. doi:10.1016/j.antiviral.2017.03.020
120. Anderson, G.R.; Hasan, A.; Yin, H.; Qadri, I.; Quattrochi, L.C. Regulation of the CYP1A1 gene by 2,3,7,8-tetrachlorodibenzo-p-dioxin but not by beta-naphthoflavone or 3-methylcholanthrene is altered in

- hepatitis C virus replicon-expressing cells. *Pharmacol.* **2006**,70,1062-1070.
doi:10.1124/mol.106.024125
121. El-Diwany, R.; Soliman, M.; Sugawara, S.; Breitwieser, F.; Skaist, A.; Coggiano, C.; Sangal, N.; Chattergoon, M.; Bailey, J.R.; Siliciano, R.F.; et al. CMPK2 and BCL-G are associated with type 1 interferon-induced HIV restriction in humans. *Adv.* **2018**,4,eaat0843. doi:10.1126/sciadv.aat0843
 122. Imakiire, K.; Uto, H.; Sato, Y.; Sasaki, F.; Mawatari, S.; Ido, A.; Shimoda, K.; Hayashi, K.; Stuver, S.O.; Ito, Y.; et al. Difference in serum complement component C4a levels between hepatitis C virus carriers with persistently normal alanine aminotransferase levels or chronic hepatitis C. *Med. Rep.* **2012**,6,259-264. doi:10.3892/mmr.2012.924
 123. Yang, C.H.; Li, H.C.; Ku, T.S.; Wu, P.C.; Yeh, Y.J.; Cheng, J.C.; Lin, T.Y.; Lo, S.Y. Hepatitis C virus down-regulates SERPINE1/PAI-1 expression to facilitate its replication. *Gen. Virol.* **2017**,98,2274-2286. doi:10.1099/jgv.0.000901
 124. An, P.; Li, R.; Wang, J.M.; Yoshimura, T.; Takahashi, M.; Samudralal, R.; O'Brien, S.J.; Phair, J.; Goedert, J.J.; Kirk, G.D.; et al. Role of exonic variation in chemokine receptor genes on AIDS: CCRL2 F167Y association with pneumocystis pneumonia. *Genet.* **2011**,7,e1002328. doi:10.1371/journal.pgen.1002328
 125. Yong, Y.K.; Tan, H.Y.; Saeidi, A.; Rosmawati, M.; Atiya, N.; Ansari, A.W.; Rajarajeswaran, J.; Vadivelu, J.; Velu, V.; Larsson, M.; et al. Decrease of CD69 levels on TCR V α 7.2+CD4+ innate-like lymphocytes is associated with impaired cytotoxic functions in chronic hepatitis B virus-infected patients. *Immun.* **2017**,23,459-467. doi:10.1177/1753425917714854
 126. Li, F.; Nio, K.; Yasui, F.; Murphy, C.M.; Su, L. Studying HBV Infection and Therapy in Immune-Deficient NOD-Rag1^{-/-}IL2RgammaC-null (NRG) Fumarylacetoacetate Hydrolase (Fah) Knockout Mice Transplanted with Human Hepatocytes. *Mol. Biol.* **2017**,1540,267-276. doi:10.1007/978-1-4939-6700-1_23
 127. Inomata, S.; Anan, A.; Yamauchi, E.; Yamauchi, R.; Kunimoto, H.; Takata, K.; Tanaka, T.; Yokoyama, K.; Morihara, D.; Takeyama, Y.; et al. Changes in the Serum Hepcidin-to-ferritin Ratio with Erythroferrone after Hepatitis C Virus Eradication Using Direct-acting Antiviral Agents. *Med.* **2019**,58,2915-2922. doi:10.2169/internalmedicine.2909-19
 128. Dai, W.; Usami, Y.; Wu, Y.; Göttlinger, H. A Long Cytoplasmic Loop Governs the Sensitivity of the Anti-viral Host Protein SERINC5 to HIV-1 Nef. *Rep.* **2018**,22,869-875. doi:10.1016/j.celrep.2017.12.082
 129. Cheng J.C.; Yeh Y.J.; Pai L.M.; Chang M.L.; Yeh C.T. 293 cells over-expressing human ADI1 and CD81 are permissive for serum-derived hepatitis C virus infection. *Med. Virol.* **2009**,81,1560-1568. doi:10.1002/jmv.21495
 130. Neveu, G.; Barouch-Bentov, R.; Ziv-Av, A.; Gerber, D.; Jacob, Y.; Einav, S. Identification and targeting of an interaction between a tyrosine motif within hepatitis C virus core protein and AP2M1 essential for viral assembly. *Pathog.* **2012**,8,e1002845. doi:10.1371/journal.ppat.1002845
 131. Duong, V.N.; Zhou, L.; Martínez-Jiménez, M.I.; He, L.; Cosme, M.; Blanco, L.; Paintsil, E.; Anderson, K.S. Identifying the role of PrimPol in TDF-induced toxicity and implications of its loss of function

- mutation in an HIV+ patient. *Rep.* **2020**,10,9343. doi:10.1038/s41598-020-66153-z
132. Moudi, B.; Heidari, Z.; Mahmoudzadeh-Sagheb, H. Polymorphisms of BIRC5 Gene is Associated with Chronic HBV Infection in Iranian Population. *J. Clin. Biochem.* **2020**,35,158-168. doi:10.1007/s12291-018-0805-3
 133. Masaisa, F.; Gahutu, J.B.; Mukiibi, J.; Delanghe, J.; Philippé, J. Transferrin polymorphism and opportunistic infections in HIV-infected women in Rwanda. *Haematol.* **2012**,128,100-106. doi:10.1159/000338262
 134. Pauli, E.K.; Schmolke, M.; Hofmann, H.; Ehrhardt, C.; Flory, E.; Münk, C.; Ludwig, S. High level expression of the anti-retroviral protein APOBEC3G is induced by influenza A virus but does not confer antiviral activity. **2009**,6,38. doi:10.1186/1742-4690-6-38
 135. Ranjan, P.; Singh, N.; Kumar, A.; Neerincx, A.; Kremmer, E.; Cao, W.; Davis, W.G.; Katz, J.M.; Gangappa, S.; Lin, R.; et al. NLRC5 interacts with RIG-I to induce a robust antiviral response against influenza virus infection. *J. Immunol.* **2015**,45,758-772. doi:10.1002/eji.201344412
 136. Job E.R.; Bottazzi B.; Short K.R.; Deng Y.M.; Mantovani A.; Brooks A.G.; Reading P.C. A single amino acid substitution in the hemagglutinin of H3N2 subtype influenza A viruses is associated with resistance to the long pentraxin PTX3 and enhanced virulence in mice. *Immunol.* **2014**,192,271-281. doi:10.4049/jimmunol.1301814
 137. Tan S.Y.; Chowdhury S.; Polak N.; Gorrell M.D.; Weninger W. Fibroblast activation protein is dispensable in the anti-influenza immune response in mice. *PLoS One.* **2017**,12,e0171194. doi:1371/journal.pone.0171194
 138. Shi, X.; Shi, Z.; Huang, H.; Zhu, H.; Zhou, P.; Zhu, H.; Ju, D. Ability of recombinant human catalase to suppress inflammation of the murine lung induced by influenza A. **2014**,37,809-817. doi:10.1007/s10753-013-9800-2
 139. Chang, X.; Shi, X.; Zhang, X.; Wang, L.; Li, X.; Wang, A.; Deng, R.; Zhou, E.; Zhang, G. IFI16 Inhibits Porcine Reproductive and Respiratory Syndrome Virus 2 Replication in a MAVS-Dependent Manner in MARC-145 Cells. *Viruses.* **2019**,11,1160. doi:3390/v11121160
 140. Sun, N.; Liu, D.; Chen, H.; Liu X.; Meng F.; Zhang X.; Chen H.; Xie S.; Li X.; Wu Z. Localization, expression change in PRRSV infection and association analysis of the porcine TAP1 gene. *J. Biol. Sci.* **2012**,8,49-58. doi:10.7150/ijbs.8.49
 141. Shen, Z.J.; Malter J.S. Eosinophils, Pin1 and the Response to Respiratory Viral Infection and Allergic Stimuli. *Rev. Immunol.* **2019**,39,135-149. doi:10.1615/CritRevImmunol.2019031697
 142. Ratajczak, M.Z.; Kucia, M. SARS-CoV-2 infection and overactivation of Nlrp3 inflammasome as a trigger of cytokine "storm" and risk factor for damage of hematopoietic stem cells. *Leukemia.* **2020**,1-4. doi:1038/s41375-020-0887-9
 143. Tsoi, H.; Li, L.; Chen, Z.S.; Lau, K.F.; Tsui, S.K.; Chan, H.Y. The SARS-coronavirus membrane protein induces apoptosis via interfering with PDK1-PKB/Akt signalling. *J.* **2014**,464,439-447. doi:10.1042/BJ20131461

144. Ke, W.; Fang, L.; Tao, R.; Li, Y.; Jing, H.; Wang, D.; Xiao, S. Porcine Reproductive and Respiratory Syndrome Virus E Protein Degrades Porcine Cholesterol 25-Hydroxylase via the Ubiquitin-Proteasome Pathway. *Virology*. **2019**,93,e00767-19. doi:10.1128/JVI.00767-19
145. Zhou, A.C.; Batista, N.V.; Watts T.H. 4-1BB Regulates Effector CD8 T Cell Accumulation in the Lung Tissue through a TRAF1-, mTOR-, and Antigen-Dependent Mechanism to Enhance Tissue-Resident Memory T Cell Formation during Respiratory Influenza Infection. *Immunology*. **2019**,202,2482-2492. doi:10.4049/jimmunol.1800795
146. Rohaim, M.A.; Santhakumar, D.; Naggar, R.F.E.; Iqbal, M.; Hussein, H.A.; Munir, M. Chickens Expressing IFIT5 Ameliorate Clinical Outcome and Pathology of Highly Pathogenic Avian Influenza and Velogenic Newcastle Disease Viruses. *Immunology*. **2018**,9,2025. doi:10.3389/fimmu.2018.02025
147. Londino J.D.; Lazrak A.; Noah J.W.; Aggarwal S.; Bali V.; Woodworth B.A.; Bebok Z.; Matalon S. Influenza virus M2 targets cystic fibrosis transmembrane conductance regulator for lysosomal degradation during viral infection. *J. Virol.* **2015**,29,2712-2725. doi:10.1096/fj.14-268755
148. Dornfeld, D.; Dudek, A.H.; Vausselin, T.; Günther, S.C.; Hultquist, J.F.; Giese, S.; Khokhlova-Cubberley, D.; Chew, Y.C.; Pache, L.; Krogan, N.J.; et al. SMARCA2-regulated host cell factors are required for MxA restriction of influenza A viruses. *Rep.* **2018**,8,2092. doi:10.1038/s41598-018-20458-2
149. Le May, N.; Mansuroglu, Z.; Léger, P.; Josse, T.; Blot, G.; Billecocq, A.; Flick, R.; Jacob, Y.; Bonnefoy, E.; Bouloy, M. A SAP30 complex inhibits IFN-beta expression in Rift Valley fever virus infected cells. *Pathog.* 2008;4(1):e13. doi:10.1371/journal.ppat.0040013
150. Goldstein, M.R.; Poland, G.A.; Graeber, C.W. Does apolipoprotein E genotype predict COVID-19 severity? **2020**,hcaa142. doi:10.1093/qjmed/hcaa142
151. Qin, J.; Zhai, J.; Hong, R.; Shan, S.; Kong, Y.; Wen, Y.; Wang, Y.; Liu, J.; Xie Y. Prospero-related homeobox protein (Prox1) inhibits hepatitis B virus replication through repressing multiple cis regulatory elements. *J. Gen. Virol.* **2009**,90,1246-1255. doi:1099/vir.0.006007-0
152. Zhao, X.; Fan, H.; Chen, X.; Zhao X.; Wang X.; Feng Y.; Liu M.; Li S.; Tang H. HBV DNA polymerase restrains viral replication via the CREB1-HOTTIP-HOXA13 axis. **2020**,10.1002/hep.31284. doi:10.1002/hep.31284

Tables

Tables 1-8 are available in the Supplemental Files section

Figures

regulated significant genes.

Figure 3

Heat map of up regulated differentially expressed genes. Legend on the top left indicate log fold change of genes. (A1 – A8 = Mock samples; B1 – B14 = SARS-CoV-2 infection samples)

Figure 4

Heat map of downregulated differentially expressed genes. Legend on the top left indicate log fold change of genes. (A1 – A8 = Mock samples; B1 – B14 = SARS-CoV-2 infection samples)

Figure 5

Protein–protein interaction network of up regulated genes. Green nodes denotes up regulated genes.

Figure 6

Scatter plot for up regulated genes. (A- Node degree; B- Betweenness centrality; C- Stress centrality ; D- Closeness centrality; E- Clustering coefficient)

Figure 7

Protein–protein interaction network of down regulated genes. Red nodes denotes down regulated genes

Figure 8

Scatter plot for down regulated genes. (A- Node degree; B- Betweenness centrality; C- Stress centrality ; D- Closeness centrality; E- Clustering coefficient)

Module 25

Module 33

Module 37

Module 70

Figure 9

Modules in PPI network. The green nodes denote the up regulated genes

Module 14

Module 15

Module 27

Module 45

Figure 10

Modules in PPI network. The red nodes denote the down regulated genes

Figure 11

The network of up regulated genes and their related miRNAs. The green circles nodes are the up regulated genes, and purple diamond nodes are the miRNAs

Figure 12

The network of down regulated genes and their related miRNAs. The red circles nodes are the down regulated genes, and blue diamond nodes are the miRNAs

Figure 13

The network of up regulated genes and their related TFs. The green circles nodes are the up regulated genes, and purple triangle nodes are the TFs

Figure 14

The network of down regulated genes and their related TFs. The green circles nodes are the down regulated genes, and blue triangle nodes are the TFs.

Figure 15

ROC curve validated the sensitivity, specificity of hub genes as a predictive biomarker for SARS-CoV-2 infection. A) DDX58 B) IFITM2 C) IRF1 D) PML E) SAMHD1 F) ACSS1 G) CYP2U1 H) DDC I) PNMT J) UGT2A3

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Tables.docx](#)