Cardiac complications and pertaining mortality rate in COVID-19 patients; a systematic review and meta-analysis

Amirmohammad Toloui

Physiology Research Center, Iran University of Medical Sciences, Tehran

Donya Moshrefiaraghi

Physiology Research Center, Iran University of Medical Sciences, Tehran

Arian Madani Neishaboori

Physiology Research Center, Iran University of Medical Sciences, Tehran

Saeed Safari

Emergency Department, Shohadye Tajrish Hospital, Shahid Beheshti University of Medical Sciences, Tehran

Mahmoud Yousefifard ([✉ yousefifard.m@iums.ac.ir](mailto:yousefifard.m@iums.ac.ir))

Physiology Research Center, Iran University of Medical Sciences, Tehran

Mohammad Haji Aghajani

Prevention of Cardiovascular Disease Research Center, Shahid Beheshti University of Medical Sciences, Tehran

Research Article

Keywords: COVID-19, Cardiovascular, Cardiac Complication, In-hospital Mortality

Posted Date: December 21st, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-122109/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background

Raising knowledge over cardiac complications and managing them can play a key role in their recovery. In this study, we aim to investigate the evidence regarding the prevalence of cardiac complications and the resulting mortality rate in COVID-19 patients.

Results

The initial search resulted in 853 records, from which 40 articles were included. Overall analysis showed the prevalence of acute cardiac injury, heart failure and cardiac arrest were 19.46% (95% CI: 18.23–20.72), 19.07% (95% CI: 15.38–23.04) and 3.44% (95% CI: 3.08–3.82), respectively. Moreover, abnormal serum troponin level was observed in 22.86% (95% CI: 21.19–24.56) of the COVID-19 patients. Further analysis revealed that the overall odds of mortality is 14.24 [odds ratio (OR) = 14.24; 95% CI: 8.67–23.38] times higher, when patients develop acute cardiac injury. The pooled odds ratio of mortality when the analysis was limited to abnormal serum troponin level was 19.03 (OR = 19.03; 95% CI: 11.85–30.56).

Conclusion

Acute cardiac injury and abnormal serum troponin level were the most prevalent cardiac complications/abnormalities in COVID-19 patients. The importance of cardiac complications becomes crucial due to the higher mortality rate among patients with these complications. Thus, troponin screenings and cardiac evaluations are recommended to be performed in routine patient assessments.

Background

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is a novel coronavirus, emerged in Wuhan, China in December 2019 and has spread to over 200 countries in the world, ever since (1). Coronavirus disease 2019 (COVID-19) mostly causes lower respiratory tract infection symptoms such as fever, being the most prevalent, cough and dyspnea (2). Although people with severe type of the disease form the minority of the patients, it is illustrated that mortality rate is the highest among patients who have comorbidities such as cardiovascular diseases (3).

The angiotensin converting enzyme 2 (ACE2) has been shown to be the receptor of the virus in human cells. This enzyme being expressed in many organs such as lungs, heart, kidneys and brain, extrapulmonary manifestations and complications are being studied rigorously, all around the world (4). Cardiac involvement is one of the most common causes of death in COVID-19 patients (5). Cardiovascular complications, whether being chronic or acute, can lead to critically imbalanced homeostasis and could be a serious strike to patient's recovery from the disease (6). Therefore, raising awareness over cardiac complications caused by the disease and managing them as early as possible can play a key role in recovery of the patients. While, there is no comprehensive evaluation in this issue. Therefore, in this study, we aim to investigate the evidence regarding the prevalence of cardiac complication in COVID-19 patients. Moreover, the risk of mortality after COVID-19 related cardiac involvements has also been assessed.

Methods

Study design

The present systematic review and meta-analysis aims to investigate the evidence regarding cardiac complications in patients, confirmed with COVID-19, and to see whether there is any potential association between the prevalence of cardiac complications and their mortality rate. As a result, PECO in the current study is defined as: P (patients): Patients with confirmed COVID-19 disease, E (exposure): SARS-CoV-2 infection, O (outcome): prevalence of the cardiac complication and mortality after COVID-19 related cardiac involvement. As for comparison (C in PECO), mortality rate was evaluated between COVID-19 patients with cardiac complications and patients without any cardiac involvements.

Search strategy

The searching process was initiated, selecting keywords with the help of experts in the field and by screening titles of similar articles. Then, using MeSH and Emtree, equivalent and related synonyms were identified. As a result, a search strategy was constructed based on the instructions of electronic databases Medline, Embase, Scopus and Web of Science, and an extensive search was performed in each of the mentioned databases for articles published from the February 1st of 2019 until April 30th, 2020. The search strategy in Medline database through PubMed search engine is presented in Appendix 1. In addition, a manual search was performed in preprint databases, Google and Google scholar to obtain preprinted and possibly missing manuscripts. Moreover, references of the obtained review articles were screened to find additional articles.

Selection criteria

The inclusion criteria in the present meta-analysis was reporting cardiac complications in patients with confirmed COVID-19. Furthermore, the exclusion criteria were case report studies, review articles, studies which did not report a cardiac involvement and studies whose entire target population was patients with cardiac comorbidities as underlying disease.

Data collection

Two independent reviewers screened titles and abstracts of the gathered articles. Then, full texts of the related articles were obtained and included articles were selected and entered to the present systematic review and meta-analysis. Finally, a summary of the included studies was recorded using a checklist, consisting of following variables: first author's name, publication year, country in which the study was conducted in, number of patients, study design, number of patients in which the cardiac complication was assessed in, mean/median age of the patients, number of males among the patients, type of cardiac complication along with its diagnostic method, number of the deceased in patients presenting with cardiac complication (if reported) and number of the deceased among patients without the cardiac complication of the included articles. If several types of cardiac complications were reported in the studies, the number of each cardiac involvement was recorded separately. Any disagreements within the process was resolved using a third reviewer's opinion.

Outcome

The primary outcome of the present meta-analysis was the prevalence of cardiac complications in COVID-19 patients. Secondary outcome was the risk of mortality after COVID-19 related cardiac involvement. Our screening showed that most studies reported abnormal serum troponin level and acute cardiac injury, separately, as cardiac assessments/complications. Other reported cardiac complications were heart failure, cardiomyopathy, cardiac arrest, myocarditis, pericardial effusion, cardiac insufficiency and myocardial infarction.

Quality assessment

Two independent reviewers scored the quality of the studies according to National Heart, Lung, and Blood Institute (NHLBI) quality assessment tools (7). Any disagreement was resolved by discussion with third reviewers.

Statistical analysis

All analyses were performed using STATA 14.0 statistical program. Analyses were performed in two parts. Initially, the cardiac complications reported in the articles were categorized into 9 subgroups: acute cardiac injury, abnormal serum troponin level, heart failure, cardiac arrest, myocarditis, cardiac insufficiency, pericardial effusion, myocardial infarction and cardiomyopathy. The prevalence of each cardiac complication in the COVID-19 patients were calculated with a 95% confidence interval (95% CI) using the "metaprop_one" command performed on the total sample size and the number of patients presenting each complication in the included articles.

In the second part of the analysis, the association between the appearance of cardiac complications/abnormalities and mortality rate was calculated and presented as an overall odds ratio (OR) with a 95% CI using the "metan" command performed on four groups of data: number of the deceased in patients with a cardiac complication, number of the alive in patients with the cardiac complication, number of the deceased in patients without the cardiac complication, number of the alive in patients without the cardiac complication. I^2 test was performed to assess heterogeneity, and if the heterogeneity among data was considerable, random effect model was used to calculate 95% CI. Egger's test was used to evaluate publication bias.

Results

Study characteristics

The systematic search resulted in 853 records, and after eliminating duplicates, 557 articles remained. Then, after screening titles and abstracts of the remaining articles, 181 studies were selected to be potentially eligible. Afterwards, based on the mentioned exclusion and inclusion criteria, 40 articles were included in the present systematic review and meta-analysis (Fig. 1) (5, 8–46). These 40 articles were treated as 60 different experiments, as some studies reported more than one cardiac complication/abnormality. Two studies were taken place in the United States (41, 45), one study was conducted in Spain (9), one study was in Italy (42) and the rest of the included studies were conducted in China. Regarding the study design of the included articles, one study was ambispective (22), four studies were conducted prospectively (9, 14, 19, 39) and the other 35 studies were retrospective. Overall, 15616 patients confirmed with COVID-19 were enrolled in the included studies, with 2985 males among them [patients' gender was not reported in two of the studies (42, 46)]; however, not all of the patients were assessed for each cardiac complication, so the total number of patients tested for each manifestation is presented for each complication in Table 1. In general, 9 different cardiac complications/abnormalities were reported in the studies including acute cardiac injury, which was reported in 26 studies (5, 8, 11, 13, 16, 18–20, 22–24, 26–30, 32, 33, 35, 37–41, 43, 46), abnormal serum troponin level, reported in 19 studies (8, 10–12, 14, 15, 17, 19, 25, 30, 32, 36–38, 40, 41, 43–45), heart failure, reported in four studies (8, 11, 32, 41), cardiac arrest, reported in three studies (5, 34, 42), myocarditis, reported in two studies (9, 44), cardiac insufficiency, reported in two studies (29, 30), pericardial effusion, reported in two studies (21, 31) and cardiomyopathy (45) and myocardial infarction (32), each reported in one study. Mortality rate was reported in 18 studies (8, 11, 13, 14, 16–18, 20, 26, 29, 30, 33, 34, 37, 39, 40, 44, 46). These numbers were further used to evaluate the odds ratio (OR) between the appearance of each cardiac complication and the mortality rate in COVID-19 patients. Table 1 demonstrates a summary of the characteristics of the included studies.

Table 1
Summary of the included studies

Author; Year; Country	Study design	Total study population	Mean age	Number of males	Diagnostic methods	Type of cardiac complication	No. of patients tested for the complication	No. of the complication	No. of deceased in patients with the complication	No. of decease patients without complic
Aggarwal S; 2020; USA	Retrospective	16	67	12	Blood Sample, Echo	Acute cardiac injury	16	3	NR	NR
					NR	Heart failure	16	2	NR	NR
					Blood Sample	Abnormal Troponin	16	10	NR	NR
Arentz M; 2020; USA	Retrospective	21	70	11	Blood Sample, Echo, Clinical	Cardiomyopathy	21	7	NR	NR
					Blood Sample	Abnormal Troponin	21	3	NR	NR
Baldi E; 2020; Italy	Retrospective	9806	NR	NR	Hospital report	Cardiac arrest	9806	362	NR	NR
Barrasa H; 2020; Spain	Prospective	48	63.2	27	NR	Myocarditis	48	1	NR	NR
Chen C; 2020; China	Retrospective	150	59	84	Blood Sample	Abnormal Troponin	150	22	NR	NR
Chen T; 2020; China	Retrospective	274	62	171	NR	Acute cardiac injury	203	89	72	22
					NR	Heart failure	176	43	41	42
					Blood Sample	Abnormal Troponin	203	83	68	26
Deng Q; 2020; China	Retrospective	112	65	57	Blood Sample	Abnormal Troponin	112	42	NR	NR
Deng Y; 2020; China	Retrospective	225	54	124	Blood Sample	Acute cardiac injury	225	66	65	44
Du Ra; 2020; China	Prospective	179	57.6	97	Blood Sample	Abnormal Troponin	179	31	13	8
Du Rb; 2020; China	Retrospective	109	70.7	75	Blood Sample	Abnormal Troponin	109	52	NR	NR
Du Y; 2020; China	Retrospective	85	65.8	62	Blood Sample	Acute cardiac injury	85	38	NR	NR
					Clinical	Cardiac arrest	85	7	NR	NR
Guo T; 2020; China	Retrospective	187	58.5	91	Blood Sample	Acute cardiac injury	187	52	31	12
Han H; 2020; China	Retrospective	273	NR	97	Blood Sample	Abnormal Troponin	273	27	13	8
He X; 2020; China	Retrospective	54	68	34	Blood Sample	Acute cardiac injury	54	24	18	8
Hu L; 2020; China	Retrospective	323	61	166	Blood Sample	Acute cardiac injury	323	24	NR	NR

CT: Computed tomography scan; Echo: Echocardiography; ECG: Electrocardiography; NR: Not reported

Author, Year, Country	Study design	Total study population	Mean age	Number of males	Diagnostic methods	Type of cardiac complication	No. of patients tested for the complication	No. of the complication	No. of deceased in patients with the complication	No. of decease patients without complic
					Blood Sample	Abnormal Troponin	323	68	NR	NR
Huang C; 2020; China	Prospective	41	49	30	Blood Sample	Abnormal Troponin	41	5	NR	NR
						Acute cardiac injury	41	5	NR	NR
Lei S; 2020; China	Retrospective	34	55	14	Blood Sample	Acute cardiac injury	34	5	4	3
Li K; 2020; China	Retrospective	83	45.5	44	CT	Pericardial effusion	83	4	NR	NR
Li X; 2020; China	Ambispective	548	60	279	Blood Sample	Acute cardiac injury	548	119	NR	NR
Li Y; 2020; China	Retrospective	54	61.8	34	Blood Sample	Acute cardiac injury	41	23	NR	NR
Liu M; 2020; China	Retrospective	30	35	10	Blood Sample	Acute cardiac injury	30	5	NR	NR
Liu Y; 2020; China	Retrospective	76	45	49	Blood Sample	Abnormal Troponin	76	14	NR	NR
Ma K; 2020; China	Retrospective	84	48	48	Blood Sample	Abnormal Troponin	84	36	0	0
					Blood Sample, Clinical Symptom	Myocarditis	84	4	0	0
Ruan Q; 2020; China	Retrospective	150	NR	NR	NR	Acute cardiac injury	68	5	5	63
Shi S; 2020; China	Retrospective	416	64	205	Blood Sample	Acute cardiac injury	416	82	42	15
Wan S; 2020; China	Retrospective	135	47	72	Blood Sample	Acute cardiac injury	135	10	NR	NR
Wang Da; 2020; China	Retrospective	138	56	75	Blood Sample, ECG, Echo	Acute cardiac injury	138	10	NR	NR
Wang Db; 2020; China	Retrospective	107	51	57	Blood Sample	Abnormal Troponin	107	6	5	14
					Blood Sample, ECG, Echo	Acute cardiac injury		12	8	19
Wang La; 2020; China	Retrospective	339	69	166	Blood Sample	Acute cardiac injury	339	70	39	26
					Blood Sample, Clinical Symptom	Cardiac insufficiency	339	58	25	52
Wang Lb; 2020; China	Retrospective	202	63	88	Blood Sample	Acute cardiac injury	202	27	17	14

CT: Computed tomography scan; Echo: Echocardiography; ECG: Electrocardiography; NR: Not reported

Author, Year, Country	Study design	Total study population	Mean age	Number of males	Diagnostic methods	Type of cardiac complication	No. of patients tested for the complication	No. of the complication	No. of deceased in patients with the complication	No. of deceased patients without complic
Wei J; 2020; China	Prospective	101	49	54	Blood Sample, ECG, Echo	Cardiac insufficiency	202	24	14	19
					Blood Sample	Abnormal Troponin	202	27	NR	NR
					Blood Sample	Acute cardiac injury	101	16	3	0
Xu X; 2020; China	Retrospective	90	50	39	CT	Pericardial effusion	90	1	NR	NR
Yang F; 2020; China	Retrospective	92	69.8	49	Blood Sample	Acute cardiac injury	92	31	NR	NR
					NR	Myocardial infarction	92	6	NR	NR
					NR	Heart failure	92	2	NR	NR
					Blood Sample	Abnormal Troponin	92	31	NR	NR
Yang X; 2020; China	Retrospective	52	59.7	35	Blood Sample	Acute cardiac injury	52	12	9	23
Yao W; 2020; China	Retrospective	202	63.4	136	Clinical	Cardiac arrest	202	4	0	21
Zhang G; 2020; China	Retrospective	221	55	108	NR	Acute cardiac injury	221	17	NR	NR
Zhao X; 2020; China	Retrospective	91	46	49	Blood Sample	Acute cardiac injury	91	14	NR	NR
Zheng Y; 2020; China	Retrospective	99	49.4	51	Blood Sample	Abnormal Troponin	99	88	NR	NR
					Blood Sample	Abnormal Troponin	99	88	NR	NR
					Blood Sample	Abnormal Troponin	99	88	NR	NR
Zhou F; 2020; China	Retrospective	191	56	119	Blood Sample	Abnormal Troponin	145	24	23	31
						Heart failure	145	44	28	26
						Acute cardiac injury	145	33	32	22
						Acute cardiac injury	145	33	32	22
Zou X; 2020; China	Retrospective	178	60.68	67	Blood Sample, ECG, Echo	Acute cardiac injury	154	45	34	18
					Blood Sample	Abnormal Troponin	154	33	28	24
					Blood Sample	Abnormal Troponin	154	33	28	24

CT: Computed tomography scan; Echo: Echocardiography; ECG: Electrocardiography; NR: Not reported

Risk of bias assessment

No study was provided a sample size justification, power description, or variance and effect estimates. In addition, 39 studies were not measured to have key potential confounders in their assessment of outcomes. 10 studies did not report the details of their inclusion and exclusion criteria. Moreover, the participation rate of eligible persons was not reported in 6 studies (Table 2).

Table 2
Risk of bias assessment of included studies

Author; Year	Item 1	Item 2	Item 3	Item 4	Item 5	Item 6	Item 7	Item 8	Item 9	Item 10	Item 11	Item 12	Item 13	Item 14
Aggarwal S; 2020	Yes	Yes	No	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Arentz M; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Baldi E; 2020	Yes	Yes	Yes	No	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Barrasa H; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Chen C; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Chen T; 2020	Yes	Yes	No	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Deng Q; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Deng Y; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Du Ra; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Du Rb; 2020	Yes	Yes	No	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Du Y; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Guo T; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Han H; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
He X; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Huang C; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Hu L; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	No	NA	Yes	NA	Yes	No
Lei S; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Li K; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Liu M; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Liu Y; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Li X; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Li Y; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Ma K; 2020	Yes	Yes	Yes	No	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Ruan Q; 2020	Yes	Yes	Yes	No	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Shi S; 2020	Yes	Yes	No	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Wang Da; 2020	Yes	Yes	Yes	No	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Wang Db; 2020	Yes	Yes	No	No	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Wang La; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Wang Lb; 2020	Yes	Yes	Yes	Yes	No	Yes	No	NA	Yes	NA	Yes	NA	Yes	No
Wan S; 2020	Yes	Yes	Yes	Yes	No	Yes	No	NA	Yes	NA	Yes	NA	Yes	No
Wei J; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Xu X; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Yang F; 2020	Yes	Yes	Yes	No	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	Yes
Yang X; 2020	Yes	Yes	No	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Yao W; 2020	Yes	Yes	Yes	No	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Zhang G; 2020	Yes	Yes	Yes	No	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Zhao X; 2020	Yes	Yes	Yes	No	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Zheng Y; 2020	Yes	Yes	Yes	No	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Zhou F; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No
Zou X; 2020	Yes	Yes	Yes	Yes	No	Yes	Yes	NA	Yes	NA	Yes	NA	Yes	No

Author, Year	Item 1	Item 2	Item 3	Item 4	Item 5	Item 6	Item 7	Item 8	Item 9	Item 10	Item 11	Item 12	Item 13	Item 14
NA: Not applicable; NR: Not reported														
Items:														
1. Was the research question or objective in this paper clearly stated?														
2. Was the study population clearly specified and defined?														
3. Was the participation rate of eligible persons at least 50%?														
4. Were all the subjects selected or recruited from the same or similar populations (including the same time period)? Were inclusion and exclusion criteria for being in the study prespecified and applied uniformly to all participants?														
5. Was a sample size justification, power description, or variance and effect estimates provided?														
6. For the analyses in this paper, were the exposure(s) of interest measured prior to the outcome(s) being measured?														
7. Was the timeframe sufficient so that one could reasonably expect to see an association between exposure and outcome if it existed?														
8. For exposures that can vary in amount or level, did the study examine different levels of the exposure as related to the outcome (e.g., categories of exposure, or exposure measured as continuous variable)?														
9. Were the exposure measures (independent variables) clearly defined, valid, reliable, and implemented consistently across all study participants?														
10. Was the exposure(s) assessed more than once over time?														
11. Were the outcome measures (dependent variables) clearly defined, valid, reliable, and implemented consistently across all study participants?														
12. Were the outcome assessors blinded to the exposure status of participants?														
13. Was loss to follow-up after baseline 20% or less?														
14. Were key potential confounding variables measured and adjusted statistically for their impact on the relationship between exposure(s) and outcome(s)?														

Publication bias

There are evidences of publication bias in the assessment of the prevalence of acute cardiac injury ($p = 0.004$) and abnormal serum troponin level ($p < 0.001$) among the included studies. However, no evidence of publication bias was observed in the assessment of cardiac related mortality and the prevalence of other cardiac complications (Fig. 2).

Meta-analysis

In the beginning, the prevalence of cardiac complications following SARS-CoV-2 infection was evaluated, and the results are depicted in Figs. 3, 4, 5 and 6 and Table 3. The abnormal serum level of troponin was observed in 22.86% (95% CI: 21.19 to 24.56) of the patients (Fig. 4). Moreover, the prevalence of acute cardiac injury, heart failure and cardiac arrest were 19.46% (95% CI: 18.23 to 20.72), 19.07% (95% CI: 15.38 to 23.04) and 3.44% (95% CI: 3.08 to 3.82), respectively (Fig. 3 and Table 3). Furthermore, the I^2 test revealed no heterogeneity regarding the prevalence of abnormal troponin levels and acute cardiac injury. The prevalence of other cardiac complications including myocarditis, cardiac insufficiency, pericardial effusion, myocardial infarction and cardiomyopathy are depicted in Table 3.

Table 3
Summary of finding regarding the cardiac complications in COVID-19

Complication	Number of studies	Prevalence	95% CI	Number of studies	Odds ratio	95% CI	P
Acute cardiac injury	26	19.46	18.23, 20.72	14	14.24	8.67, 23.38	< 0.001
Abnormal troponin	19	22.86	21.19, 24.56	6	19.03	11.85, 30.56	< 0.001
Heart failure	4	19.07	15.38, 23.04	2	10.66	5.69, 19.97	< 0.001
Cardiac arrest	3	3.44	3.08, 3.82	1	0.04	0.00, > 999.0	0.651
Myocarditis	2	3.66	0.88, 7.82	1	1.00	0.00, > 999.0	> 0.99
Pericardial effusion	2	2.62	0.58, 5.73	—	—	—	—
Cardiac insufficiency	2	15.06	12.15, 18.22	2	4.65	2.82, 7.66	< 0.001
Cardiomyopathy	1	33.33	17.19, 54.63	—	—	—	—
Myocardial infarction	1	6.52	3.02, 13.51	—	—	—	—
—: No data; CI: Confidence interval							

Further analysis revealed that the odds of mortality in COVID-19 patients with acute cardiac injury is 14.24 (OR = 14.24, 95% CI: 8.67 to 23.38) times higher than the COVID-19 patients without acute cardiac injury. However, I^2 test showed some degrees of heterogeneity regarding the relationship of cardiac complication and mortality of COVID-19 patients (Fig. 5). Moreover, the odds of mortality in a COVID-19 patient presenting with abnormal serum troponin level in his/her blood sample was 19.03 (OR = 19.03; 95% CI: 11.85 to 30.56) times higher than the patients not having this manifestation. Interestingly, no heterogeneity was observed when calculating the OR for the mortality in COVID-19 patients with abnormal serum troponin level (Fig. 6). The odds of mortality of patients having other cardiac complications are presented in Table 3.

Discussion

In this systematic review and meta-analysis, we investigated the prevalence of 9 cardiac complications in COVID-19 patients and their subsequent mortality rates. Abnormal serum level of troponin with 22.86% prevalence, detected by laboratory tests, and acute cardiac injury with 19.46% prevalence, defined with laboratory test results and other diagnostic techniques were among the most prevalent complications/abnormalities observed in COVID-19 patients. Although a high number of studies reported the prevalence of the two mentioned cardiac complications, no heterogeneity was observed in this section. To contemplate even more on the matter, the prevalence of abnormal troponin level and acute cardiac injury are rather close numbers, which might mean that by close observation of troponin changes in a COVID-19 patient, we could possibly anticipate acute cardiac injuries in them and take appropriate measures. However, serum troponin levels also increase due to damage to other tissues, such as the kidneys, which may make it impossible to use troponin level alone to detect cardiac damage. Other cardiac complications such as heart failure and cardiomyopathy were also prevalent among the patients. However, with the few number of the studies observing them, more data is needed on the matter.

Regarding the limitations, each individual article used different reference range for troponin or other injury indicators to be accepted as abnormal, and each one used different diagnostic and laboratory test results to define acute cardiac injury, which could cause the slight differences in the reported prevalence of acute cardiac injury and abnormal serum level of troponin. For example, Huang et al. defined acute cardiac injury as cardiac troponin rising to 3 or more times than normal or appearance of new abnormalities in echocardiography or ECG (19); While many others defined it only with the appearance of abnormal troponin levels in blood samples. Moreover, some articles' study population consisted of only critically ill patients or deceased ones, which could shift prevalence statistics, causing inevitable heterogeneity in reported numbers (5, 15, 32). Additionally, it is worth mentioning that the mechanism of myocardial injury, whether being done by direct viral invasion to the host tissue or due to imbalanced homeostasis, was of no concern for the writers at the time. Although more research should be conducted to identify the exact causes of this damage, the effects it carries are the most important subject needed to be studied at these critical times. Even though limitations were present in data reporting the prevalence of cardiac outcomes, it is important to pay enough attention to these complications, since many of them are directly related to patients' general situation and illness severity.

On the other hand, our study shows intriguing data regarding mortality rate in patients presenting with cardiac complications, especially acute cardiac injury and abnormal serum level of troponin. Abnormal troponin levels are associated with about 19 times more mortality chance in COVID-19 patients, which is of great importance in disease management for health care providers around the world. Considering the fact that no heterogeneity was observed regarding the risk of mortality in abnormal troponin level, laboratory screenings in routine patient assessments can be critical in determining early cardiac injury and intervening accordingly. In addition, we suggest that blood levels of troponin can be counted as a prognostic factor in patients with cardiac involvement.

Furthermore, our data indicated that acute cardiac injury could raise the mortality to about 14 times more in COVID-19 patients. However, this number was associated with an overall heterogeneity, which could be attributed to different definitions of cardiac injury in studies, discussed above. Nonetheless, although one can conclude that there is a relationship between acute cardiac injury following COVID-19 and increased risk of mortality, the threat of acute cardiac injury to patients' health status in future is inevitable and demands careful monitoring and managements when confronting this situation. Moreover, it is noteworthy that articles reporting mortality rates of cardiac complications were mostly focused on abnormal troponin level and myocardial injury and the number of the articles reporting correlations between other cardiac complications such as heart failure, and mortality rate were extremely low; so it is inevitable that we cannot present an exact estimate of what other cardiac outcomes such as heart failure and myocarditis would contribute to patients' fatality. In addition, inability of blinding of the outcome assessors in the studies, not presenting the cardiac complications with respect to prior cardiac comorbidities of the patients and not presenting explicit inclusion and exclusion criteria in some articles were further limitations detected in the studies.

To be concluded, our findings thoroughly approve other published articles investigating associations between COVID-19 cardiac outcomes and related mortalities (47, 48) confirming that cardiac injury regardless of the mechanism of establishment is a factor determining disease severity and patient prognosis in a large and reliable scale and should be considered and monitored from early stages of disease.

Conclusion

Cardiac complications/abnormalities can be prevalent in COVID-19 patients in the forms of acute cardiac injury, serum troponin levels abnormalities, heart failure, cardiac arrest and other types. The importance of cardiac involvements becomes crucial when observing the higher mortality rate among COVID-19 patients presenting with cardiac involvements. Thus, careful monitoring of heart involvements should be performed in COVID-19 patients.

Abbreviations

SARS-CoV-2

Severe Acute Respiratory Syndrome Coronavirus 2

COVID-19

Coronavirus disease 2019

ACE2
Angiotensin Converting Enzyme 2
PECO
Problem, Exposure, Comparison, Outcome
NHLBI
National Heart, Lung, and Blood Institute
CI
Confidence Interval
OR
Odds Ratio
ECG
Electro Cardio Graphy

Declarations

Ethics approval and consent to participate

This study received ethics approval from Ethics committee of Iran University medical Sciences.

Consent for publication

Not applicable.

Availability of data and materials

All data generated or analyzed during this study are included in this published article [and its supplementary information files].

Conflict of Interest

There is no conflict of interest.

Funding

This study was supported by Prevention of Cardiovascular Disease Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Authors' contribution

Study design: MY, SS, MHA; Data gathering: AT, DM, AMN. MY; Analysis: MY; Interpreting the results: MY, SS, MHA; Drafting: AT, DM, AMN; Critically revised the paper: All authors.

Acknowledgments

None.

References

1. Organization WH. Coronavirus disease 2019 (COVID-19): situation report, 121. 2020.
2. Guan W-j, Ni Z-y, Hu Y, Liang W-h, Ou C-q, He J-x, et al. Clinical characteristics of coronavirus disease 2019 in China. *New England journal of medicine*. 2020;382(18):1708-20.
3. Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW, et al. Presenting characteristics, comorbidities, and outcomes among 5700 patients hospitalized with COVID-19 in the New York City area. *Jama*. 2020.
4. Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, et al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *The Lancet*. 2020;395(10224):565-74.
5. Du Y, Tu L, Zhu P, Mu M, Wang R, Yang P, et al. Clinical Features of 85 Fatal Cases of COVID-19 from Wuhan. *A Retrospective Observational Study*. *American journal of respiratory and critical care medicine*. 2020;201(11):1372-9.
6. Hu H, Ma F, Wei X, Fang Y. Coronavirus fulminant myocarditis treated with glucocorticoid and human immunoglobulin. *European heart journal*. 2020.
7. Health Nlo. Study Quality Assessment Tools| National Heart, Lung, and Blood Institute (NHLBI). National Institutes of Health. 2014.
8. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The lancet*. 2020.
9. Barrasa H, Rello J, Tejada S, Martín A, Balziskueta G, Vinuesa C, et al. SARS-CoV-2 in Spanish intensive care: early experience with 15-day survival in Vitoria. *Anaesthesia Critical Care & Pain Medicine*. 2020.
10. Chen C, Yan J, Zhou N, Zhao J, Wang D. Analysis of myocardial injury in patients with COVID-19 and association between concomitant cardiovascular diseases and severity of COVID-19. *Zhonghua xin xue guan bing za zhi*. 2020;48:E008-E.

11. Chen T, Wu D, Chen H, Yan W, Yang D, Chen G, et al. Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *Bmj*. 2020;368.
12. Deng Q, Hu B, Zhang Y, Wang H, Zhou X, Hu W, et al. Suspected myocardial injury in patients with COVID-19: Evidence from front-line clinical observation in Wuhan, China. *International journal of cardiology*. 2020.
13. Deng Y, Liu W, Liu K, Fang Y-Y, Shang J, Zhou L, et al. Clinical characteristics of fatal and recovered cases of coronavirus disease 2019 in Wuhan, China: a retrospective study. *Chinese medical journal*. 2020;133(11):1261-7.
14. Du R-H, Liang L-R, Yang C-Q, Wang W, Cao T-Z, Li M, et al. Predictors of mortality for patients with COVID-19 pneumonia caused by SARS-CoV-2: a prospective cohort study. *European Respiratory Journal*. 2020;55(5).
15. Du R-H, Liu L-M, Yin W, Wang W, Guan L-L, Yuan M-L, et al. Hospitalization and critical care of 109 decedents with COVID-19 pneumonia in Wuhan, China. *Annals of the American Thoracic Society*. 2020(ja).
16. Guo T, Fan Y, Chen M, Wu X, Zhang L, He T, et al. Cardiovascular implications of fatal outcomes of patients with coronavirus disease 2019 (COVID-19). *JAMA cardiology*. 2020.17. Han H, Xie L, Liu R, Yang J, Liu F, Wu K, et al. Analysis of heart injury laboratory parameters in 273 COVID-19 patients in one hospital in Wuhan, China. *Journal of Medical Virology*. 2020.
17. He X, Lai J, Cheng J, Wang M, Liu Y, Xiao Z, et al. Impact of complicated myocardial injury on the clinical outcome of severe or critically ill COVID-19 patients. *Zhonghua xin xue guan bing za zhi*. 2020;48:E011-E.
18. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The lancet*. 2020;395(10223):497-506.
19. Lei S, Jiang F, Su W, Chen C, Chen J, Mei W, et al. Clinical characteristics and outcomes of patients undergoing surgeries during the incubation period of COVID-19 infection. *EClinicalMedicine*. 2020:100331.
20. Li K, Wu J, Wu F, Guo D, Chen L, Fang Z, et al. The clinical and chest CT features associated with severe and critical COVID-19 pneumonia. *Investigative radiology*. 2020.
21. Li X, Xu S, Yu M, Wang K, Tao Y, Zhou Y, et al. Risk factors for severity and mortality in adult COVID-19 inpatients in Wuhan. *Journal of Allergy and Clinical Immunology*. 2020.
22. Li Y, Hu Y, Yu J, Ma T. Retrospective analysis of laboratory testing in 54 patients with severe-or critical-type 2019 novel coronavirus pneumonia. *Laboratory Investigation*. 2020:1-7.
23. Liu M, He P, Liu H, Wang X, Li F, Chen S, et al. Clinical characteristics of 30 medical workers infected with new coronavirus pneumonia. *Zhonghua jie he hu xi za zhi= Zhonghua jiehe he huxi zazhi= Chinese journal of tuberculosis and respiratory diseases*. 2020;43:E016-E.
24. Liu Y, Liao W, Wan L, Xiang T, Zhang W. Correlation between relative nasopharyngeal virus RNA load and lymphocyte count disease severity in patients with COVID-19. *Viral immunology*. 2020.
25. Shi S, Qin M, Shen B, Cai Y, Liu T, Yang F, et al. Association of cardiac injury with mortality in hospitalized patients with COVID-19 in Wuhan, China. *JAMA cardiology*. 2020.
26. Wan S, Xiang Y, Fang W, Zheng Y, Li B, Hu Y, et al. Clinical features and treatment of COVID-19 patients in northeast Chongqing. *Journal of medical virology*. 2020.
27. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus–infected pneumonia in Wuhan, China. *Jama*. 2020;323(11):1061-9.
28. Wang L, He W, Yu X, Hu D, Bao M, Liu H, et al. Coronavirus disease 2019 in elderly patients: Characteristics and prognostic factors based on 4-week follow-up. *Journal of Infection*. 2020.
29. Wang L, He W, Yu X, Liu H, Zhou W, Jiang H. Prognostic value of myocardial injury in patients with COVID-19. [*Zhonghua yan ke za Zhi*] *Chinese Journal of Ophthalmology*. 2020;56:E009-E.
30. Xu X, Yu C, Qu J, Zhang L, Jiang S, Huang D, et al. Imaging and clinical features of patients with 2019 novel coronavirus SARS-CoV-2. *European journal of nuclear medicine and molecular imaging*. 2020:1-6.
31. Yang F, Shi S, Zhu J, Shi J, Dai K, Chen X. Analysis of 92 deceased patients with COVID-19. *Journal of medical virology*. 2020.
32. Yang X, Yu Y, Xu J, Shu H, Liu H, Wu Y, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *The Lancet Respiratory Medicine*. 2020.
33. Yao W, Wang T, Jiang B, Gao F, Wang L, Zheng H, et al. Emergency tracheal intubation in 202 patients with COVID-19 in Wuhan, China: lessons learnt and international expert recommendations. *British Journal of Anaesthesia*. 2020.
34. Zhang G, Hu C, Luo L, Fang F, Chen Y, Li J, et al. Clinical features and short-term outcomes of 221 patients with COVID-19 in Wuhan, China. *Journal of Clinical Virology*. 2020:104364.
35. Zheng Y, Xu H, Yang M, Zeng Y, Chen H, Liu R, et al. Epidemiological characteristics and clinical features of 32 critical and 67 noncritical cases of COVID-19 in Chengdu. *Journal of Clinical Virology*. 2020:104366.
36. Zou X, Li S, Fang M, Hu M, Bian Y, Ling J, et al. Acute Physiology and Chronic Health Evaluation II Score as a Predictor of Hospital Mortality in Patients of Coronavirus Disease 2019. *Critical care medicine*. 2020.
37. Hu L, Chen S, Fu Y, Gao Z, Long H, Wang J-m, et al. Risk factors associated with clinical outcomes in 323 COVID-19 hospitalized patients in Wuhan, China. *Clinical infectious diseases*.
38. Wei J-F, Huang F-Y, Xiong T-Y, Liu Q, Chen H, Wang H, et al. Acute myocardial injury is common in patients with covid-19 and impairs their prognosis. *Heart*. 2020.

39. Wang D, Yin Y, Hu C, Liu X, Zhang X, Zhou S, et al. Clinical course and outcome of 107 patients infected with the novel coronavirus, SARS-CoV-2, discharged from two hospitals in Wuhan, China. *Critical Care*. 2020;24:1-9.
40. Aggarwal S, Garcia-Telles N, Aggarwal G, Lavie C, Lippi G, Henry BM. Clinical features, laboratory characteristics, and outcomes of patients hospitalized with coronavirus disease 2019 (COVID-19): Early report from the United States. *Diagnosis*. 2020;7(2):91-6.
41. Baldi E, Sechi GM, Mare C, Canevari F, Brancaglione A, Primi R, et al. Out-of-hospital cardiac arrest during the Covid-19 outbreak in Italy. *New England Journal of Medicine*. 2020.
42. Zhao X-Y, Xu X-X, Yin H-S, Hu Q-M, Xiong T, Tang Y-Y, et al. Clinical characteristics of patients with 2019 coronavirus disease in a non-Wuhan area of Hubei Province, China: a retrospective study. *BMC Infectious Diseases*. 2020;20:1-8.
43. Ma K-L, Liu Z-H, Cao C-f, Liu M-K, Liao J, Zou J-B, et al. COVID-19 myocarditis and severity factors: an adult cohort study. *medRxiv*. 2020.
44. Arentz M, Yim E, Klaff L, Lokhandwala S, Riedo FX, Chong M, et al. Characteristics and outcomes of 21 critically ill patients with COVID-19 in Washington State. *Jama*. 2020;323(16):1612-4.
45. Ruan Q, Yang K, Wang W, Jiang L, Song J. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive care medicine*. 2020;46(5):846-8.
46. Zhu H, Rhee J-W, Cheng P, Waliany S, Chang A, Witteles RM, et al. Cardiovascular complications in patients with COVID-19: consequences of viral toxicities and host immune response. *Current cardiology reports*. 2020;22:1-9.
47. Santoso A, Pranata R, Wibowo A, Al-Farabi MJ, Huang I, Antariksa B. Cardiac injury is associated with mortality and critically ill pneumonia in COVID-19: a meta-analysis. *The American Journal of Emergency Medicine*. 2020.

Figures

Figure 1

Flow diagram of the present meta-analysis

Figure 2

Funnel plot for publication bias in prevalence of cardiac complication in COVID-19 patients and mortality rate after incidence of cardiac manifestation. There are evidences of publication bias among the studies, which reported prevalence of cardiac complications. While, there no publication bias was observed regarding the relationship of cardiac complication and mortality in COVID-19 patients.

Figure 3

Forest plot for the assessment of the prevalence of acute cardiac injury in COVID-19 patients; CI: confidence interval.

Figure 4

Forest plot for the assessment of the prevalence of abnormal troponin levels in COVID-19 patients; CI: confidence interval.

Figure 5

Forrest plot for the assessment of risk of mortality in COVID-19 patients with acute cardiac injury; OR: odds ratio; CI: confidence interval.

Figure 6

Forrest plot for the assessment of risk of mortality in COVID-19 patients with abnormal troponin level; OR: odds ratio; CI: confidence interval.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Appendix1.docx](#)