

# Prehospital Handling of Postpartum Haemorrhage- Healthcare Personnel's Experiences and Perspectives

Ann-Chatrin Linqvist Leonardsen (✉ [ann.c.leonardsen@hiof.no](mailto:ann.c.leonardsen@hiof.no))

Østfold University College: Hogskolen i Ostfold <https://orcid.org/0000-0001-5608-9401>

Ann Karin Helgesen

Østfold University College

Linn Eik Ulvøy

Østfold Hospital Trust

Vigdis Abrahamsen Grøndahl

Østfold University College

---

## Original research

**Keywords:** postpartum hemorrhage, manual aortic compression, prehospital, ambulance, questionnaire, delphi technique, self-efficacy, collective efficacy

**Posted Date:** December 10th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-122155/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

**Background:** Postpartum hemorrhage (PPH) is a serious obstetric emergency, and one of the top five causes of maternal mortality globally. The most common causes of PPH include uterine atony, placental disorders, birth trauma and coagulation defects. Timely diagnosis and early management are critical to reduce morbidity, the need for blood transfusion or even mortality. External, manual aortic compression (AC) has been suggested as an intervention that reduce PPH and extend time for control of bleeding or resuscitation. This procedure is not commonly utilized by healthcare personnel. The incidence of home-births is increasing, and competence in PPH handling is essential in prehospital personnel. The objective was to explore prehospital personnel's competence in PPH and AC, utilizing different tools.

**Methods:** The study was conducted in a county in South-eastern Norway, including five ambulance stations. All prehospital personnel (n=250) were invited to participate in a questionnaire study. The questionnaire included the PPH self-efficacy (PPHSE) and PPH collective efficacy (PPHCE) tools, as well as a developed tool. Descriptive statistics were used to analyse the quantitative data, while quantitative content analysis was used to analyse free-text responses.

**Results:** A total of 87 prehospital personnel responded to the questionnaire, 57.5 percent male, mean age 37.9 years. 80.4 percent were ambulance workers and/or paramedics, and 96.6 and 97.7 percent respectively reported to need more education or training in PPH. Results show that prehospital personnel lack knowledge about what PPH is, how to estimate and handle PPH, and how to perform AC. 82.8 percent reported having experienced PPH, but only 2.9 percent had performed AC. Mean self-efficacy varied from 3.3 to 5.6, while collective efficacy varied from 1.9 to 3.8. Educational background did not seem to have any impact on participants' responses.

**Conclusions and implications:** This study indicates that prehospital personnel lack knowledge about PPH and AC, and also report to need more education and training. Hence, PPH should be included in educational programs at all levels of the education, as well as in quality improvement initiatives. After the conduction of this study, prehospital personnel attended a digital course in PPH, as well as a practical course in AC.

## Background

Postpartum hemorrhage (PPH) is a serious obstetric emergency, and one of the top five causes of maternal mortality globally [1]. Internationally, the prevalence of severe PPH appears to be rising, with increasing morbidity and need for transfusion therapy, and the mortality rates are high in low-income countries [2-6]. Maternal mortality rates (MMR) in the Nordic countries are among the lowest in the world, but women still die from complications of pregnancy or birth. In Norway, 168 maternal deaths were identified between 2005 and 2013 (Maternal Mortality Rate, MMR, 6.6 per 100 000), of whom 14 died due to severe PPH [7].

The incidence of PPH has been reported to vary between one to five percent, depending upon the diagnostic criteria applied [2]. Although PPH is traditionally defined by the volume of blood loss observed, bleeding may not be visible externally or the blood may be mixed with amniotic fluid [4]. In 2017, the American College of Obstetricians and Gynecologists revised their definition of PPH to consist of the following criteria:

1. Cumulative blood loss  $\geq 1000$  ml or
2. Bleeding associated with signs or symptoms of hypovolemia within 24 hours of the birth process, regardless of the route of delivery [8]

The most common causes of PPH include uterine atony, placental disorders, birth trauma and coagulation defects [9].

Timely diagnosis, appropriate resources and early management are critical to prevent death [9]. According to the World Health Organization, immediate solutions are needed to prevent women from dying of PPH [1]. Maternal mortality reviews have demonstrated that deaths caused by PPH are most likely to be preventable [10, 11]. The consistent application of a comprehensive protocol for management of PPH have been demonstrated to result in improved outcomes [12, 13]. Other important interventions include fluid resuscitation, removal of the placenta, bimanual uterine compression, uterotonics, tranexamic acid, suturing of lower genital tract injury, blood product replacement, balloon tamponade, laparotomy, stepwise uterine de-vascularization, uterine compression sutures and hysterectomy. Emergency temporizing measures include application of the non-pneumatic anti-shock garment, uterine tourniquet application and aortic compression [14].

External, manual aortic compression (AC) is an emergency manoeuvre proposed to reduce postpartum haemorrhage and extend time for resuscitation and control of bleeding. The technique can be applied immediately to reduce bleeding from the uterus by reducing the blood supply. This again may prevent cardiac arrest from hypovolemia, and allow transfer to definitive care in-hospital. The Swedish obstetrician Bergstrom has been teaching this life-saving technique for many years in African countries, with great effect on maternal morbidity and mortality [15]. Nevertheless, in many countries, including Norway, manual AC is not actively used by healthcare personnel [16].

There is an increasing incidence of home births in Norway, as in Nordic countries [17, 18]. Hence, knowledge of PPH on competence in handling this condition is essential in prehospital personnel. Consequently, as researchers, healthcare personnel educators, nurse anesthetist and paramedic, we wanted to explore prehospital personnel's knowledge about and self-assessed competence in PPH and AC, their experience with this condition, perceived need for more education and/or training, as well as their perceived self-efficacy and collective efficacy in PPH handling.

## Methods

The study had a cross-sectional design, utilizing a questionnaire to assess prehospital personnel's knowledge, self-assessed competence, self-efficacy and team-efficacy.

### *Setting*

The study was conducted in a county with approximately 317.000 inhabitants, within one hospital catchment area. There are five ambulance stations in this area. Prehospital personnel include ambulance assistants, ambulance workers (upper high school), bachelor in paramedicine or – nursing (180 ECTs), paramedics (further education, 60 ECTs).

### *Questionnaire*

The questionnaire consisted of three parts:

1. a validated questionnaire in PPH self-efficacy (PPHSE), including eight items, using an eight-point Likert scale that range from 1 (never) to 8 (always) [19]. The items on self-efficacy focus on individual perception of control.
2. a validated questionnaire in PPH collective efficacy (PPHCE), including 13 items (same scoring format as the PPHSE) (Egenberg, Øian, Eggebø, Arsenovic, & Bru, 2016). In the current study, collective efficacy was defined as team-efficacy in the prehospital team, most commonly consisting of two healthcare personnel.
3. a questionnaire developed as part of this study

### *Development of the questionnaire*

Since no validated questionnaire to measure knowledge and self-assessed competence in PPH handling could be identified, we developed a questionnaire. Here, we used recommendations from the Delphi technique, which is suitable to obtain expert opinions in a systematic manner, and includes four steps: 1) expert input, 2) interaction with feedback, 3) statistical group responses, and 4) confidentiality [20, 21].

Experts were defined as specialists in their field, and included six anesthesiologists, three obstetricians and two midwives, knowledgeable in the field of obstetrics and obstetric anesthesia, and recommended by other experts [22]. The expert group consisted of four males, seven females, mean age was 53 years, and mean years of experience 15. In step 1 and 2, the expert group participated in the development of questions, and gave constructive inputs on clarity, wording, and contents of the whole questionnaire, as suggested by Streiner & Norman [23]. In these steps, experts received the questionnaire in two or three rounds depending on their inputs. In step 3, experts were asked to score the questionnaire regarding relevance, clarity and logic, on a scale from 1=strongly disagree, to 5=strongly agree. Table 1 presents the mean and range of responses to these scorings.

The experts were involved in several rounds until consensus was reached. The final version of the questionnaire consisted of a) 13 knowledge questions with free-text answers, and two questions with

alternatives yes/no/undecided, b) two questions about perceived need for more education and/or simulation, and c) five questions about experience with PPH and the use of AC.

### *Analysis*

Data were analyzed using the Statistical Package for the Social Sciences (SPSS 26.0)[24]. The descriptive statistics frequency, mean and SD were used to analyse data. Internal consistency for the scales was tested by Cronbach's alpha. There were no missing items in the validated questionnaires. The free-text responses were analyzed through a quantitative content analysis, reading through the responses and searching for similarities and code-words repeated throughout [25].

## **Results**

A total of 87 prehospital healthcare personnel (34.8 percent) responded to the questionnaire. Table 2 gives an overview of respondents' gender, age, educational background and years of experience.

In this study, the Cronbach's alpha was 0.89 on the PPHSE scale, and 0.96 on the PPHCE, which is assumed excellent.

### *Responses to the developed questionnaire*

#### *Knowledge about PPH*

On the question "How much is normal hemorrhage during birth, and when is it defined as postpartum hemorrhage?" most of the respondents assumed a hemorrhage of up to 500 ml as normal per-partum. Hemorrhage above 500 ml was interpreted as postpartum hemorrhage by 37 of the respondents. Other answers were 'above 1 litre' (n=17), '2 litres' (n=3), '1.5 litre' (n=1) and '3-4 litres' (n=1). The rest were undecided.

Regarding the question "How do you estimate the amount of hemorrhage during/after birth?" 39 respondents found this 'difficult'. A total of 18 respondents reported to assess the sheets or diapers, how often they needed to be changed, or even to weigh them. In addition, 21 of the respondents reported to assess the patients' vital parameters or level of consciousness.

#### *Knowledge about interventions*

On the question "Which interventions should be initiated in postpartum hemorrhage?", the responses varied. Reported from most to least frequent response; uterus massage (n=29), fluid resuscitation (n=22), put the baby to the breast (n=14), abdominal massage (n=10), establish intravenous access (n=9), add pressure on the abdominal aorta (n=9), elevate legs (n=9), quick transport to hospital (n=8), oxygen treatment (n=7), put pressure to the abdomen (n=5), shock-treatment (n=5), put the fist into the woman and add pressure from the inside (n=5), areola massage (n=4), hemorrhage control (n=4), and oxytocin (n=3). Other suggestions were 'add pressure on the inguinal aorta', 'analgesia', 'tranexamacid', 'comfort

the mother', 'compression', 'early warning to the hospital'. The questionnaire also included the question "When you observe a life threatening hemorrhage, what do you do first?" Answers here were similar to those above regarding interventions.

Other "Clinical situations than postpartum hemorrhage where manual aortic compression can be lifesaving" reported were 'hemorrhage in the lower extremities' (n=7), 'abdominal aorta aneurism' (AAA) (n=7), 'other vaginal hemorrhage' (n=6), amputations (n=4), other causes of massive hemorrhage such as extrauterine pregnancy (n=2), and open wounds (n=2).

Whether the ambulance had any drugs for use in situations of postpartum hemorrhage, most respondents reported 'no' (82.8 percent), while 11.5 percent were undecided, and 5.7 percent of the respondents reported 'yes'. Drug reported accessible was oxytocin, and side-effects of this drug was reported to be 'high blood-pressure' (n=1), and 'nausea and vomiting' (n=1).

To the question "When is manual aortic compression (using a fist on aorta) appropriate?", responses were 'in massive hemorrhage' (n=32), 'in PPH' (n=13), 'in life-threatening hemorrhage' (n=5), 'when the child is delivered'(n=2), 'when uterus massage does not have an effect' (n=2), and 'AAA'(n=1) (non-response, n=32).

Contra-indications to AC reported were 'limited hemorrhage' (n=28), 'child not delivered' (n=6), 'pain'(n=1), and 'patient awake' (n=1) (non-response, n=51).

### *Knowledge about performance of AC*

When asked «How would you provide manual aortic compression?» 12 of the respondents reported 'establish pulse in arteria femoralis, add pressure above the uterus until absence of pulse'. And 20 respondents reported to 'add pressure on the abdomen', but location of pressure varied from 'under the diaphragm', 'umbilical area', or 'in the middle'. Ten respondents answered 'add pressure on the aorta', four respondents reported 'add pressure both from the inside and outside', and three 'add vaginal pressure'.

When asked what the purpose of AC is, 71 respondents reported 'to stop the hemorrhage'. On the question "How do you assess whether the maneuver is conducted correct?", 35 responded 'when the hemorrhage stops', and 18 responded 'when the pulse in arteria femoralis is absent'. Regarding considerations during drug administration and ongoing AC, five respondents reported 'side-effects', and one reported 'that drugs are not transported beyond the location of pressure'. A total of 82.8 percent of the respondents reported 'no', 11.5 percent reported 'undecided' and 5.7 percent reported 'yes', to the question about whether there are potential complications related to AC. Suggested complications were 'damage due to ischemia' (n=4), 'reduced blood pressure'(n=3), 'damage to inner organs'(n=1), and 'pain' (n=1).

No relation between educational background and level of knowledge could be identified.

### *Need for education and/or training*

When asked “Do you want more education in handling postpartum hemorrhage?”, 96.6 percent responded ‘yes’, 1.1 percent responded ‘no’, and 2.3 percent responded ‘undecided’.

Among the respondents, 97.7 percent answered ‘yes’ that they want more training/simulation in handling postpartum hemorrhage.

Participants that responded ‘no’ or ‘undecided’ that they needed more education or training were all assistants or ambulance workers.

### *Experience*

Prehospital personnel’s experience with PPH and AC is shown in table 3.

Reasons for not using AC were ‘lack of education’ (74.7 percent), ‘lack of training’ (10.3 percent), ‘feel unsecure on the procedure’ (10.3 percent), and ‘difficult to cause the patient pain’ (4.6 percent) (fixed response alternatives).

### *Self-efficacy in PPH*

## **Discussion**

Our findings indicate that prehospital personnel lack knowledge about postpartum hemorrhage (PPH) and manual aortic compression (AC). As much as 82.8 percent had experienced PPH, but only 2.3 percent had utilized AC. Participants scored lower on self-efficacy than on collective efficacy in PPH handling. 96.6 and 97.7 percent respectively reported a need for more education or training in PPH/PPH handling.

Our findings enlighten the knowledge gap in prehospital personnel regarding PPH and AC. This indicates a need to include this topic in educational programs regardless of educational level. Studies have indicated a need to develop and implement robust clinical research regarding treatment of PPH, to establish an international knowledge platform [26]. Senthiles et al. [27] emphasize a need to reach a broad consensus about the most efficient interventions to prevent and treat PPH. This include PPH prevention initiatives, estimation of blood loss, when and which uterotonica to administer, and use of blood products. Nevertheless, consensus has been reached regarding the main steps for the initial management of PPH: manual exploration of the uterus, visual assessment of genital tract, bladder indwelling catheter, measures to maintain maternal temperature, supplemental oxygen, uterine massage, maintenance of venous access, infusion of crystalloids rather than colloids, continuous monitoring of pulse, blood pressure and respiratory rate, and use of a uterotonic [27].

The participants’ self-efficacy was scored lowest on the items «I have experienced being able to act in situations with PPH” (mean score 2.6, SD=1.5) and “I am confident in how to treat PPH” (mean score 3.5, SD=2.6), while the highest score was on the item “I remain calm when handling PPH” (mean score 5.5, SD=2.1). This may be because prehospital personnel is used to handling acute and unpredictable situations. The lowest score regarding collective efficacy was on the item “I think that every

member of the team will express themselves clearly during PPH” (mean score 4.8, SD=2.0), and the highest score on the item “As a team, we help each other prevent excessive PPH” (mean score 6.9, SD=1.4). This may mostly reflect the nature of prehospital personnel’s work, rather than PPH itself [28]. In contrast, a recent study including pre- and post-simulation scores of self- and collective efficacy in PPH in intra-hospital personnel, participants scored 5.9 (SD=1.1) on self-efficacy before the simulations, and 6.5 (SD=0.9) after. Scores on collective efficacy in PPH handling increased from 5.8 (SD=0.9) to 6.3 (SD=0.8) [19]. This may indicate that personnel working in obstetric/resuscitation teams in hospital are more trained in this obstetric emergency, which is also natural.

Aronson and Bergström’s research [15] indicates that AC may effectively reduce need for blood transfusion, morbidity and mortality. Nevertheless, our participants rarely used this technique. This study, as well as earlier research enlighten the challenges in distributing knowledge to larger groups of personnel, and from high-income to low-income countries. Hence, a PPH digital learning program was developed, in addition to a simulator that enables participants to evaluate the pressure, placement and effect of AC. After the conduction of this study, prehospital personnel attended the course and were given a demonstration and opportunity to apply AC. In addition, a new PPH handling guideline has been implemented in hospital, resulting in that midwives and obstetricians now often arrive to the operating room in the patient’s bed, adding AC. In Tanzania, a structured inter-professional simulation program on PPH handling lead to a significant reduction in the use of  $\geq 5$  units of blood products related to severe bleeding after birth [19]. Moreover, training that included all levels of maternity staff, repeated sessions with realistic scenarios, and debriefing contributed to reduced blood transfusion rates in this high-risk maternity settings [29]. The same research team conducted a study on the effect of inter-professional simulations to reduce PPH, in a Norwegian university hospital. The researchers emphasized the importance of team training as a learning feature, and that inter-professional simulation enhanced self-efficacy and reduced perception of stress. Here, personnel also experienced an improved competence to provide efficient PPH management [30].

## Limitations

Due to the small sample size, findings here may not be generalizable to other countries or settings. Nevertheless, research support our findings, stating a need for more knowledge and competence in PPH and PPH handling internationally. Two validated tools were used to measure self-efficacy and collective efficacy, which increase the reliability of the study. Moreover, the new questionnaire was developed in-line with recommendations from the Delphi technique, and face- and content validity were high.

## Conclusion

Postpartum haemorrhage is a critical medical emergency, that may occur prehospital due to the increasing amount of home-births. External, manual aortic compression may efficiently reduce morbidity and mortality. This study indicates a need for more education and training in PPH and PPH handling in prehospital personnel, to be able to identify and intervene in PPH.

### *Implications for further research*

Responses on the free-text questionnaire in this study will be used to develop a tool for assessment of healthcare personnel's knowledge about PPH and AC that can be used in other settings to assess knowledge gaps and what to focus on in quality improvement initiatives.

## **List Of Abbreviations**

AAA - abdominal aorta aneurism

AC - aortic compression

PPH - postpartum haemorrhage

PPHCE - postpartum haemorrhage collective efficacy

PPHSE - postpartum haemorrhage self-efficacy

SD - standard deviation

## **Declarations**

### *Ethics approval and consent to participate*

The study was approved by the Norwegian Centre for Research Data (NSD) (project no.740937). The study was based on the research ethical principles in the Declaration of Helsinki [31], on willing participation, consent to participate implicated by completion and submission of questionnaire, anonymity and confidentiality.

### *Consent for publication*

Not applicable.

### *Availability of data and materials*

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

### *Competing interests*

The authors declare that they have no competing interests

### *Funding*

The study was funded by Østfold University College. The funding body did not take part in design of the study, collection, analysis, and interpretation of data or in writing the manuscript.

### *Authors' contributions*

ACL participated in planning the study, facilitated the expert-group inputs, development of the questionnaire, datacollection, analysis, and writing and approval of the whole draft of the manuscript.

LEU participated in planning the study, facilitating the intervention, datacollection, and critical revisions of the manuscript, and approval of the final version of the manuscript.

AKH participated in planning the study, development of the questionnaire, datacollection, analysis, and writing and approval of the whole draft of the manuscript.

VAG participated in planning the study, development of the questionnaire, datacollection, analysis, and writing and approval of the whole draft of the manuscript.

### *Acknowledgements*

Participants in the expert group are acknowledged for their thorough work with the development of the questionnaire. Grethe Heitmann is acknowledged for taking the initiative to this quality improvement initiative both intra- and prehospital, development of a local guideline and the digital course in PPH, as well as training in AC. Heitmann was not involved in the research project.

## **References**

1. WHO Recommendations on prevention and treatment of postpartum haemorrhage and the WOMAN trial. [https://www.who.int/reproductivehealth/topics/maternal\\_perinatal/pph-woman-trail/en/](https://www.who.int/reproductivehealth/topics/maternal_perinatal/pph-woman-trail/en/). Accessed Nov 20 2020.
2. Kramer M, Berg C, Abenhaim H. Incidence, risk factors, and temporal trends in severe postpartum hemorrhage. *Am J Obstet Gynecol*. 2013;209:441-447. doi: 10.1016/j.ajog.2013.07.007.
3. Bateman B, Berman M, Riley L, Leffert L. The epidemiology of postpartum hemorrhage in a large, nationwide sample of deliveries. *Anesth Analg*. 2010; 110:1368-1373. doi: 10.1213/ANE.0b013e3181d74898.
4. Oyolese Y, Ananth C. Postpartum hemorrhage: epidemiology, risk factors and causes. *Clin Obstet Gynecol*. 2010;53:147-156. doi: 10.1097/GRF.0b013e3181cc406d..
5. Fleischer A, Meirowitz N. Care bundles for management of obstetrical hemorrhage. *Seminars in Perinatology*. 2016;40:99-108. doi: 10.1053/j.semperi.2015.11.015.
6. Merriam A, Bateman B, Ananth C, Wright J, Siddiq Z, D'Alton M, Friedman A. Risk for postpartum hemorrhage, transfusion and hemorrhage-related morbidity at low, moderate and high volume hospitals. *J Matern Fetal Neonatal Med*. 2018;31(8):1025-1034. doi: 10.1080/14767058.2017.1306050.

7. Vangen S, Bødker B, Ellingsen L, Saltvedt S, Gissler M, Gerisson R, Nyfløt L. Maternal deaths in the Nordic countries. *Acta Obstet Gynecol Scand.* 2017; 96(9):1112-1119. doi: 10.1111/aogs.13172.
8. Committee on Practice Bulletins-Obstetrics. Postpartum Hemorrhage. Practice Bulletin No. 183. *Obstet Gynecol.* 2017;130(4):168. doi: 10.1097/AOG.0000000000002351.
9. Belfort M. Overview of postpartum hemorrhage. UptoDate 2019. <https://www.uptodate.com/contents/overview-of-postpartum-hemorrhage>. Accessed Nov 18 2020.
10. Clark S, Belfor M, Dildy G, Herbst M, Meyers J, Hankins G. Maternal death in the 21st century: causes, prevention and relationship to cesarean delivery. *Am J Obstet Gynecol.* 2008;199:e1-5. doi:10.1016/j.ajog.2008.05.009.
11. Berg C, Harler H, Atkinson S. Preventability of pregnancy-related deaths: results of a state-wide review. *Obstetrics and Gynecology.* 2005;106:1228-1234. doi: 10.1097/01.AOG.0000187894.71913.e8.
12. Shields L, Wiesner S, Fulton J, Pelletreau B. Comprehensive maternal hemorrhage protocols reduce the use of blood products and improve patient safety. *Am J Obstet Gynecol.* 2015;212(3):272-280. doi: 10.1016/j.ajog.2014.07.012.
13. Skupski D, Brady D, Lowenwirt I, Sample J, Lin S, Lohana R, Eglinton G. Improvement in outcomes of major obstetric hemorrhage through systematic change. *Obstet Gynecol.* 2017;130(4):770. doi: 10.1097/AOG.0000000000002207.
14. Hofmeyr G, Quereshi Z. Preventing deaths due to hameorrhage. *Best Pract Res Clin Obst Gynaecol.* 2016;36:68-82. doi: 10.1016/j.bpobgyn.2016.05.004.
15. Aronsson A, Bergstrom S. Abdominal compression of the aorta effective in severe postpartum hemorrhage. *Läkartidningen.* 2012;109(34-35):1478-1480.
16. Mousa H, Blum J, El Senoun G, Shakur H, Alfirevic Z. Treatment for primary postpartum hemorrhage. *Cochrane Database Syst Rev.* 2014;2(CD0003249). doi:https://doi.org/10.1002/14651858.CD003249.pub3.
17. Medical birth registry. 2019. <https://www.fhi.no/en/hn/health-registries/medical-birth-registry-of-norway/>. Accessed Nov 01 2020.
18. Blix E, Kumle M, Ingversen K, Huitfeldt A, Hegaard H, Olafsdottir O, et al. Transfers to hospital in planned home birth in four Nordic countries- a prospective cohort study. *Acta Anesthesiol Gynecol Scand.* 2016;95(4):420-8. doi: 10.1111/aogs.12858.
19. Egenberg S, Øian P, Eggebø T, Arsenovic M, Bru L. Changes in self-efficacy, collective efficacy and patint outcome following interprofessional simulation training on postpartum haemorrhage. *J Clin Nurs.* 2016;26:3174-3187. doi: 10.1111/jocn.13666.
20. Jirwe M, Gerrish K, Keeney S. Identifying the core components of cultural competence: findings from a Delphi study. *J Clin Nurs.* 2009;18:2622-2634. doi: 10.1111/j.1365-2702.2008.02734.x.
21. Keeney S, Hasson F, McKenna H. *The Delphi technique in nursing and health research.* West Sussex: Wiley-Blackwell; 2011.

22. Bing-Jonsson P, Bjørk I, Hofoss D, Kirkevold M, Foss C. Competence in advanced older people nursing: development of 'Nursing older people- Competence evaluation tool'. *Int J Older People Nursing*. 2014;10:59-72. doi: 10.1111/opn.12057.
23. Streiner D, Norman G. *Health measurement scales: A practical guide to their development and use*. Oxford: Oxford Scholarship; 2008.
24. IBM Corporation. *IBM SPSS Statistics for Windows, Version 25.0*. Armonk, New York; 2017.
25. White M, Marsh E. Content analysis: A flexible methodology. *Library Trends*. 2006; 55(1):22-45. doi: [10.1353/lib.2006.0053](https://doi.org/10.1353/lib.2006.0053).
26. Dahlke J, Mendez-Figueroa H, Maggio L, Hauspurg A, Sperling J, Chauhan S, et al. Prevention and management of postpartum hemorrhage: a comparison of 4 national guidelines. *Am J Obstet Gynecol*. 2015;213:1-10. doi: 10.1016/j.ajog.2015.02.023.
27. Senthiles L, Goffinet F, Vayssiere C, Deneux-Tharaux C. Comparison of postpartum haemorrhage guidelines: discrepancies underline our lack of knowledge. *Int J Obstet Gynaecol*. 2016;124(5):718-722. doi: 10.1111/1471-0528.14305.
28. Patterson P, Arnold R, Abebe K, Lave J, Krackhardt D, Carr M, et al. Variation in emergency medical technician partner familiarity. *Health Serv Res*. 2011;46(4):1319-1331. doi: [10.1111/j.1475-6773.2011.01241.x](https://doi.org/10.1111/j.1475-6773.2011.01241.x).
29. Egenberg S, Masenga G, Bru L, Eggebø T, Mushi C, Massay D, et al. Impact of multi-professional, scenario-based training on postpartum hemorrhage in Tanzania: a quasi-experimental, pre- vs. post-intervention study. *BMC Pregnancy and Childbirth*. 2017; 17(287). doi: <https://doi.org/10.1186/s12884-017-1478-2>.
30. Egenberg S, Karlsen B, Massay D, Kimaro H, Bru L. "No patient should die of PPH just for the lack of training!" Experiences from multi-professional simulation training on postpartum hemorrhage in northern Tanzania: a qualitative study. *BMC Medical Education*. 2017;17(119). doi: 10.1186/s12909-017-0957-5.
31. Declaration of Helsinki- Ethical Principles for Medical Research Involving Human Subjects. <http://www.wma.net/en/30publications/10policies/b3>. Accessed Nov 20 2020.

## Tables

<b>Table 1. Expert group scorings on the questionnaire</b>						
	a)	b)	c)	d)	e)	f)
Mean (range) I	3.3 (2-5)	3.6 (2-4)	4.6 (4-5)	4.1 (2-5)	3.9 (3-4)	3.9 (34)
Mean (range) II	4.3 (4-5)	4.3 (4-5)	4.8 (4-5)	4.6 (4-5)	4.6 (4-5)	4.5 (4-5)
Abbreviations: a) relevance to assess knowledge about PPH, b) relevance to assess competence in PPH handling, c) relevance to assess knowledge about manual aortic compression (AC), d) relevance to assess competence in performing AC, e) whether questions were clear, relevant and understandable, and f) whether the questionnaire was logic. 1=strongly disagree, 2=disagree, 3=neither agree nor disagree, 4=agree, 5=strongly agree. I=first round, II=last round.						

<b>Table 2. Descriptives of the respondents</b>	
<b>Gender (male)</b>	57.5
<b>Age, mean</b>	37.9 (14.6)
<b>Age, range</b>	22-62
<b>Educational background</b>	
Assistant	6.9
Ambulance worker	47.1
Bachelor paramedicine	2.3
Bachelor nursing	4.6
Paramedic	33.3
Other	5.7
<b>Full time employe</b>	81.6
<b>Years of experience, mean</b>	11.1 (14)
Abbreviations; Gender in proportion of male, percent. Age in years. Educational background; ambulance worker=upper grade school, Bachelor= 180 ECT (European <i>Credit</i> Transfer and Accumulation System), three years full-time, Paramedic=30 ECT further education. Numbers in percent. Standard deviation (SD) in parenthesis.	

<b>Table 3. Experiences with PPH and AC</b>	
Have experience with PPH	82.8
Have used AC	2.3
Have considered using AC	5.7
Had patients where AC may have been appropriate	
Yes	
No	6.9
Undecided	79.3
	13.8
AC= external, manual, aortic compression.	

<b>Table 4. Self-efficacy in PPH</b>	
I remain calm when handling PPH	5.5(2.1)
I have experienced being able to act in situations with PPH	2.6(1.5)
I can handle PPH whenever it happens	4.4(2.1)
I can carry out the necessary actions to handle PPH	4.7(2.1)
I am confident in how to treat PPH	3.5(2.6)
I am able to stay calm in emergency situations	4.1(2.2)
I am able to identify PPH at an early stage	3.6(2.0)
PPH will make me feel paralyzed/unable to act	4.2(2.2)
Total score	4.0(1.6)
PPH= postpartum hemorrhage. Scored on a Likert scale from 1=never to 8= always. Mean scores, standard deviation in parenthesis.	

<b>Table 5. Collective efficacy in PPH</b>	
As a team, we help each other prevent excessive PPH	6.9(1.4)
As a team, we are able to carry out the necessary actions to treat PPH	5.7(2.4)
I think the team will share tasks in an appropriate way during PPH	6.5(1.7)
The team can handle PPH	5.0(2.1)
I think that every member of the team will express themselves clearly during PPH	4.8(2.0)
As a team we can cope with PPH	5.6(2.1)
The team usually has clear leadership in emergency situations like PPH	6.1(1.7)
When PPH arises, our team is able to take action	5.1(2.2)
As a team we communicate clearly and efficiently whenever PPH arises	5.5(2.1)
Everyone knows what to do during an ongoing PPH situation	5.9(1.9)
We are able to identify PPH at an early stage	5.9(2.0)
We as a team remain calm during situations involving PPH	5.3(2.0)
We are supportive of each other when we are in high-pressure situations	5.6(2.0)
Total score	5.6(1.6)
PPH= postpartum hemorrhage. Scored on a Likert scale from 1=never, to 8= always.	