Complementary Omics Strategies to Dissect p53 Signaling Networks Under Nutrient Stress

Markus Galhuber

Medical University of Graz: Medizinische Universität Graz

Helene Michenthaler

Medical University of Graz: Medizinische Universität Graz

Christoph Heininger

Medical University of Graz: Medizinische Universität Graz

Isabel Reinisch

Medical University of Graz: Medizinische Universität Graz

Christoph Nössing

Beatson Institute for Cancer Research

Jelena Krstic

Medical University of Graz: Medizinische Universität Graz

Nadja Kupper

Medical University of Graz: Medizinische Universität Graz

Elisabeth Moyschewitz

Medical University of Graz: Medizinische Universität Graz

Martina Auer

Medical University of Graz: Medizinische Universität Graz

Ellen Heitzer

Medical University of Graz: Medizinische Universität Graz

Peter Ulz

Medical University of Graz: Medizinische Universität Graz

Ruth Birner-Gruenberger

Technical University of Vienna: Technische Universität Wien

Laura Liesinger

Medical University of Graz: Medizinische Universität Graz

Georgia Ngawai Lenihan-Geels

German Institute of Human Nutrition: Deutsches Institut für Ernährungsforschung

Moritz Oster

Charité Universitätsmedizin Berlin: Charité Universitätsmedizin Berlin

Emil Spreitzer

Medical University of Graz: Medizinische Universität Graz

Riccardo Zenezini Chiozzi

University of Utrecht: Universiteit Utrecht

Tim J. Schulz

German Institute of Human Nutrition: Deutsches Institut für Ernährungsforschung

Michael Schupp

Charité Universitätsmedizin Berlin: Charité Universitätsmedizin Berlin

Tobias Madl

Medical University of Graz: Medizinische Universität Graz

Albert J.R. Heck

University of Utrecht: Universiteit Utrecht

Andreas Prokesch (✉ andreas.prokesch@medunigraz.at)

Medical University of Graz <https://orcid.org/0000-0002-8487-7103>

Research Article

Keywords: p53 signaling, nutrient stress, starvation, interactome, p53 targets

Posted Date: January 12th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1224764/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Signaling through p53 is a major cellular stress response mechanism and increases upon nutrient stresses such as starvation. Here, we show in a human hepatoma cell line that starvation leads to robust nuclear p53 stabilization. Using BioID, we determine the cytoplasmic p53 interaction network within the immediate-early starvation response and show that p53 is dissociated from several metabolic enzymes and the kinase PAK2 for which direct binding with the p53 DNA-binding domain was confirmed with NMR studies. Furthermore, proteomics after p53 immunoprecipitation (RIME) uncovered the nuclear interactome under prolonged starvation, where we confirmed the novel p53 interactors SORBS1 (insulin receptor signaling) and UGP2 (glycogen synthesis). Finally, transcriptomics after p53 re-expression revealed a distinct starvation-specific transcriptome response and suggested previously unknown nutrient-dependent p53 target genes. Together, our complementary approaches delineate several nodes of the p53 signaling cascade upon starvation, shedding new light on the mechanisms of p53 as nutrient stress sensor. Given the central role of p53 in cancer biology and the beneficial effects of fasting in cancer treatment, the identified interaction partners and networks could pinpoint novel pharmacologic targets to fine-tune p53 activity.

Introduction

The tumor suppressor p53 is a multifunctional protein involved in various cellular stress responses [1]. Acting either as transcription factor or through protein-protein interactions [2], p53 can coordinate downstream cell fate decisions to maintain genomic stability, control cell cycle progression, induce cell death or senescence, modulate autophagy, and determine cell differentiation and pluripotency [3, 4]. Coinciding with inclusion of inflammation and metabolism as central hallmarks of cancer [5], recent reports implicate p53 in the regulation of immune responses [6] and of various metabolic pathways involved in the management of all major classes of macromolecules (i.e. glucose, amino acids, and lipids; [7, 8]). To coordinate these fine-tuned activities, cellular p53 levels need to be regulated in defined margins, as loss-of-function is often detrimental (tumorigenic) and hyperactivation can lead to excessive cell death [9, 10].

A key determinant of p53 activation in cells is the interplay between p53 and MDM2. In fact, MDM2 can act as both inhibitor and activator of p53. Functioning as a p53 inhibitor, MDM2 directly binds to the N-terminal region of p53, causing p53 ubiquitination followed by p53 nuclear export, proteasomal degradation, or transactivation potential inhibition [11, 12]. Certain post-translational modifications of MDM2, however, stimulate binding to and stabilization of p53 mRNA, which cumulates into p53 activation [13, 14]. Adding to the regulatory complexity, MDM2 is a major transcriptional target of p53, forming a negative feedback loop to constrain cellular p53 protein abundance under conditions favoring MDM2-mediated p53 inhibition [15, 16]. Of note, in addition to MDM2, p53 stability can be influenced through interaction with many other proteins [17] and numerous post-translational modifications [18].

Once activated, p53 can regulate cell fate through protein-protein interactions (e.g. regulating glucose flux by binding to G6PDH in cancer cells [19]) or through acting as transcription factor to directly activate hundreds of target genes in various pathways [20, 21]. Determinants of p53 selectivity and affinity on genomic binding sites include cellular p53 levels and expression dynamics, p53 post-translational modifications, chromatin state, and interaction with nuclear cofactors (elegantly reviewed in [22]). The context-dependent selection of a specific ensemble of target genes by p53 as well as the upstream cues and specific stresses leading to p53 activation are ongoing subjects of intense research.

While fasting regimens have shown health benefits in many pathologies [23, 24] and are currently undergoing clinical testing as cancer adjuvant therapy [25–27]), from the viewpoint of a single cell, starvation constitutes a severe stress. Consistent with its role as stress response factor, we recently reported that p53 is stabilized in starved HepG2 cells and in mouse and human primary hepatocytes [28], and that p53 is necessary and sufficient for the therapy-enhancing effect of fasting on sorafenib treatment of hepatocellular carcinoma (Preprint at BioRxiv [29]). Moreover, hepatic p53 was recently shown to be stabilized by O-GlcNAcylation in fasted mice, which was shown to be essential for the regulation of gluconeogenesis in the liver [30]. Here we investigate the upstream regulatory events that activate p53 signaling upon starvation, as well as the chromatin-level interactions and transcriptional programs that define the downstream activities of p53 as a starvation response hub. We show that immediate-early dissociation of MDM2 and p53 leads to sustained stabilization of nuclear p53. A proximity-based labeling proteomics approach (BioID) identified p53 interactors early upon nutrient withdrawal and suggests the kinase PAK2 as directly interacting p53 regulator. Downstream, nuclear p53 is associated with a network of chromatin modifiers and other nuclear factors to mount a defined transcriptional starvation response, regulating canonical as well as yet unknown p53 target genes. Hence, using complementary mass spectrometry (MS)- and sequencing-based omics technologies we reveal p53 as a central node in the cellular response to nutrient withdrawal.

Results

Starvation dissociates p53-MDM2 regulatory feedback loop leading to robust nuclear p53 stabilization

p53 is key in integrating extracellular cues, such as nutrient stress, to regulate cell fate either through transcriptional mechanisms or through protein-protein interactions [3, 4]. A central determinant of p53-mediated cellular outcome is its interaction with its main endogenous inhibitor MDM2 [31]. Extending our previously published results that indicated stabilized p53 protein in response to starvation in HepG2 cells [28], we conducted a time series experiment to probe p53 and MDM2 protein dynamics after exposure to starvation medium [28, 32]. While p53 protein was immediately (after 1 hour of starvation medium) and increasingly induced, MDM2 protein abundance declined reciprocally to very low levels at 24 hours of starvation (Fig. 1A), whereas mRNA levels of both p53 and MDM2 did not reflect protein abundances (Fig S1A). As both a transcriptional target and major negative regulator of p53, MDM2 is described to elicit a

negative feedback loop to constrain p53 induction, as very high p53 levels lead to apoptosis in many cell systems [33]. In our system, however, sustained p53 levels after incubation in starvation medium for 24 hours did not lead to reduced cell viability (Fig S1B). The continuous reduction of MDM2 suggests disruption of the p53/MDM2-feedback loop by starvation as an underlying mechanism of sustained p53 stabilization. To test this, we reasoned that the effect of nutlin-3a should be dampened in starvation conditions. The small molecule nutlin-3a is a highly specific and potent inhibitor of the MDM2/p53-interaction, leading to p53 protein stabilization and target gene activation [34]. Accordingly, nutlin-3a under starvation conditions led to diminished induction of MDM2 protein (Fig. 1B and S1C) and to blunted mRNA fold-changes of known p53 target genes (Fig. 1C), when compared to nutlin-3a responses in full growth medium.

p53 transactivation potential is regulated largely through shuttling between the cytoplasm and nucleus [35]. To investigate p53/MDM2 dynamics in cellular compartments we performed fractionation experiments following three time points of nutlin-3a treatment after keeping cells either in growth medium or starvation medium for 24 hours. While in growth medium nuclear p53 protein as well as p53 targets MDM2 and p21 increase with time, these nutlin-3a responses were severely blunted in starvation medium (Fig. 1D), again indicating disruption of the p53/MDM2 feedback loop under starvation.

To further investigate prolonged nuclear stabilization of p53 in starvation medium we performed a protein degradation assay with the translation inhibitor cycloheximide (CHX) after keeping the cells in growth or starvation medium for 24 hours. As shown in Fig. 1E, in growth medium p53 is subject to rapid degradation following 1 hour of CHX treatment, commensurate with earlier reports [36]. After 24 hours in starvation medium, when p53 is mainly localized in the nucleus, it is markedly stabilized, as no degradation could be detected after 5 hours of CHX. This is likely owed to the nuclear exclusion (Fig. 1E) and proteasomal degradation (Fig S1D) of MDM2 in starvation conditions.

Together, these data indicate that starvation conditions disrupt the p53/MDM2 regulatory cycle, which leads to robust and sustained nuclear stabilization of the p53 protein.

Proximity-based labeling proteomics reveals p53 interactome under starvation

To gain further insight into the p53-regulating mechanisms in starved cells we established a BioID-based proteomics approach to determine the cytoplasmic p53 interactome early upon nutrient withdrawal. Therefore, we cloned the p53 coding sequence in an expression vector coding for an N-terminal V5-tag and a biotin ligase with a nuclear export sequence (NES) fused to the p53 C-terminus. The miniTurbo (mT) biotin ligase is optimized for minimal steric hindrance and short labeling time [37]. As a background labeling control, we cloned the coding sequence for EGFP instead of p53 into the fusion vector. To perform the overexpression on a background with non-detectable endogenous p53, we used CRISPR/Cas9 to knock out p53 in HepG2 cells (Fig S2A). Pilot experiments established efficient biotin labeling of cellular proteins with 250 μ M exogenous biotin supplemented to the medium (Fig S2B) within a 2-hour period, before p53 is enriched in nuclei (Fig S2C). Immunofluorescence microscopy after

transfection in these p53KO cells validated cytoplasmic location of both fusion proteins (Fig S2D). Importantly, when performing subcellular fractionation followed by Western blot, we observed a clear enrichment of cytoplasmic p53 protein under starvation (Fig. 2A). Hence, we overexpressed either p53-mT or EGFP-mT fusion protein in HepG2 p53KO cells and subjected the cells to either growth medium or starvation medium for 2 hours, both supplemented with biotin (Fig. 2B). As proteins in close proximity to p53-mT are biotin labelled in a cumulative manner, we reasoned that the time frame of 2 hours should facilitate identification of immediate-early cytoplasmic interactions (Fig S2C) upon change to starvation medium. Subsequently, the cell lysate was enriched for biotin labeled proteins through streptavidin pull-down and the enriched material was analyzed with MS (Fig. 2B). After peptide mapping, hierarchical clustering identified three clusters with proteins specifically enriched in the p53 group over the EGFP control (Fig. 2C, green marked in Fig S2E). p53 is found in growth and starvation medium samples with a high coverage of 36 peptides (Table S1), as would be expected as a consequence of auto-biotinylation of the fusion protein (Fig S2B marked with asterisks). Performing statistical analysis on these three clusters defines 115 and 75 proteins as potential p53 interactors in growth medium and starvation medium, respectively (FDR=0.01; Fig. 2D). The abundance of proteins in non-transfected negative controls does not significantly change between growth medium or starvation conditions (Fig S2F). Examining the list of significantly enriched proteins (Table S1), we noticed a number of known p53 interactors. Firstly, several members of the high-mobility protein group (HMG) family are detected specifically in starvation medium. In particular, HMGB1 known to directly bind to p53 [38] shows a high peptide abundance (27 peptides) in our MS data, indicating intricate interaction with p53 under starvation conditions. Secondly, FKBP prolyl isomerase 3 (FKBP3), found enriched upon starvation (Fig. 2D), was described to interact with MDM2 to promote its proteasomal degradation (as shown in Fig S1D under starvation) and p53 stabilization [39]. Additionally, Peroxiredoxin-2 (PRDX2), recently reported as promoting p53 ubiquitination and degradation [40], was found de-enriched upon starvation (Fig. 2D). Moreover, p53 and its negative regulators MDM2/MDMX have been shown to be regulated by the 14-3-3 protein interaction network [41] and we found several members of the adapter protein family 14-3-3 (gene names: YWHAB, YWHAE, YWHAG, YWHAQ, YWHAZ) with high sequence coverage under growth medium as compared to starvation medium conditions, suggesting dissociation of the 14-3-3/p53 complex upon nutrient withdrawal.

Performing KEGG pathway over-representation analysis on p53 interactors under growth medium conditions yields an enrichment of proteins of metabolic pathways such as the pentose phosphate pathway (PPP), glycolysis, amino acid biosynthesis, and carbon metabolism (Fig. 2E, proteins in significant metabolic pathways marked in Fig. 2D, blue circle). In fact, mapping EC numbers to proteins in all three clusters shows a significant enrichment of enzymes in the growth medium group (Fisher's exact test, $p=0.0101$; Fig S2E). Our data set identified several p53-associated enzymes, including key metabolic enzymes such as Phosphoglycerate kinase 1 (PGK1), Citrate synthase (CS), Malate dehydrogenase (MDH2), L-lactate dehydrogenase A chain (LDHA), Glucose-6-phosphate 1-dehydrogenase (G6PD), Ribose-phosphate pyrophosphokinase 1 (PRPS1), and Pyruvate kinase (PKM). Intriguingly, we found two members of the very small group of metabolic enzymes that are altered in cancers in the list of potential p53 interactors in growth medium: D-3-phosphoglycerate dehydrogenase (PHGDH) [42, 43] and Isocitrate

dehydrogenase [NADP] cytoplasmic (IDH1), the latter of which is involved in the regulation of the oncometabolite D-2-hydroxyglutarate [44]. Furthermore, our approach identified a number of mitochondrial proteins, such as components of the electron transport chain, as putative p53 interactors under starvation. Most noticeably, transcription factor A (TFAM) [45], the master regulator of mitochondrial DNA replication, is significantly enriched under starvation conditions with a high peptide coverage (47%). To identify more stringently controlled true positive p53 interactions, we utilized the Significance Analysis of Interactome (SAINT) algorithm (Fig. 2F) [46–48]. Overall, 16 high-confidence p53 interactors surpassed a stringent SAINT probability (SP) cut-off ≥ 0.9 . Strikingly, we found PHGDH robustly scored as interactor in growth medium conditions, confirming our finding with hierarchical clustering. PHGDH (SP = 0.96), which controls glycolytic flux into serine and glycine synthesis has been described as a direct p53 target gene [43], but we are the first to identify direct protein-protein interaction. Furthermore, another enzyme mediating *de novo* serine synthesis, Phosphoserine aminotransferase (PSAT1) (SP = 0.93), is enriched in the growth medium group.

While these novel, putative p53 interaction partners need verification with complementary methods, they harbor the potential to extend the functional spectrum of p53 in the cellular response to nutrient changes and as a metabolic regulator [10, 49, 50] of serine biosynthesis and oncometabolite production.

Differential peptide mapping reveals serine/threonine-protein kinase PAK2 as upstream regulator directly binding to p53

Through mapping several MS-detected peptides to protein amino acid sequences, information of single peptide occurrences and abundances, and therefore domain-specific interactions, might be obscured. Thus, we performed a complementary analysis on the level of unique peptides that are present exclusively in p53 overexpressing samples and identified 1050 peptides (Table S2). Additionally, a BioID approach allows for detection of differentially biotinylated peptides within the generated MS spectra. This analysis revealed 17 biotinylated peptides exclusively present in the p53 samples (Table S2). Focusing on proteins that can modify protein activity by posttranslational mechanisms, among the peptides differentially biotinylated under growth but not starvation medium we noticed one peptide mapping to the serine/protein kinase PAK2 (Fig. 3A, B; Table S2). Strikingly, also among the 1050 unique peptides found in the p53 groups, 9 peptides correspond to PAK2 or its activator, Cell division control protein 42 homolog (CDC42, [51]) (Table S2). Interestingly, overlapping the BioGRID [52] interactome of PAK2 with the list of growth medium-enriched p53 interactors (Fig. 2D) yields four 14-3-3 proteins and the Ras-related C3 botulinum toxin substrate 1 (RAC1), another well-described PAK2 activator, as common interactors (Fig S3A) [53]. Thus, we provide evidence for a close association of p53, PAK2 and its activators, and 14-3-3 proteins under growth medium conditions, which is corroborated by prediction of 14-3-3 protein binding sites in both PAK2 and p53 (Fig S3B) [54].

To address the question whether PAK2 and p53 indeed physically interact and to determine which domains mediate these interactions we employed nuclear magnetic resonance (NMR) spectroscopy.

Since the differentially biotinylated PAK2 peptide is localized within the N-terminus (Fig. 3B, Table S2), we titrated a recombinant PAK2 N-terminal domain (amino acids (aa) 1-212) to a solution containing either ¹⁵N- labeled p53 DNA binding domain (DBD; aa 94-312) or p53 transactivation domain (TAD; aa 1-94). This titration revealed a progressive chemical shift perturbation (CSP) and progressive loss of signal intensity of ¹H, ¹⁵N HSQC cross peaks with the p53 DBD (Fig. 3C), while no CSPs were observed upon testing ¹⁵N- labeled p53 TAD (Fig S3C). Inversely, titration of the p53 DBD to ¹⁵N labeled PAK2 N-terminal domain induced complementary CSPs (Fig. 3C). Together, these results indicate that the PAK2 N-terminal domain binds specifically to the p53 DBD in a concentration-dependent manner.

Since PAK2 is a potential upstream regulator of p53, we reasoned that PAK2 kinase inhibition would represent a stimulus mimicking nutrient deprivation and directly translate into p53 stabilization. Indeed, treating wild-type HepG2 cells with FRAX597 [55], a group I PAK-specific kinase inhibitor, led to p53 accumulation and gradual p21 upregulation over 6 hours of treatment (Fig. 3E). Based on our findings and previous reports [56], we propose a model (Fig. 3F) where PAK2/p53 interaction (maybe through PAK2-dependent p53 phosphorylation) renders p53 prone to degradation (p53 “off state”), while starvation-induced disruption of this complex (p53 “on state”) leads to p53 stabilization followed by nuclear translocation (Fig. 1E) and p53 activation.

Hence, our peptide-based analyses in conjunction with robust NMR results identified PAK2 as a novel p53 interactor and as an immediate-early regulator of cellular p53 levels upon nutrient withdrawal.

Nuclear p53 interactome in starved cells

Our data identify novel p53 interaction partners during the immediate-early response to starvation and show that, as a consequence, p53 is stabilized in the nucleus. Hence, we next sought to investigate nuclear p53 interactions upon prolonged starvation. First evidence of starvation-mediated nuclear p53 interactions was derived from cross-linking HepG2 cells followed by Western blot with α-p53 antibody. This approach showed p53-containing high-molecular weight complexes in cells under growth medium (Fig. 4A). When incubated in starvation medium for 24 hours - when p53 is mainly nuclear (Fig. 1D and 1E) - this complex formation increased in abundance (Fig. 4A). Importantly, this signal is p53-specific, as it was almost absent in p53KO cells (Fig. 4A). To further define the nuclear p53 interactome under starvation we employed rapid immunoprecipitation (IP) MS of endogenous proteins [57]. For this, p53-proficient HepG2 cells were kept in either growth or starvation medium for 24 hours, before they were cross-linked with formaldehyde. Nuclei were isolated, lysed, and chromatin was immunoprecipitated with p53 (with an IP-validated DO1 antibody (Fig S4A)) or with isotype-matched IgG control antibody. Enriched eluates were analyzed with liquid chromatography-MS/MS (Fig. 4B). High Pearson correlation values between replicate p53 pull-downs (>0.9; Fig S4B) and clear principal component (PCA) separation of p53-pull down groups from IgG controls (Fig S4C) indicates the robustness of the protocol. From the LC-MS/MS-identified 1,008 proteins, stringent filtering against the respective IgG groups (SAINT score = 0.85) [58] yielded 82 p53 IP-specific proteins (Fig. 4B, Table S3). Validating efficient pull-down, six p53 peptides were detected in both p53-pulldown groups (but not in the IgG samples; Fig S4D) with an

average sequence coverage of 16% (Fig. 4C). A lack of peptide detection in the N-terminus and the central DBD of p53 suggests that these domains were not amenable to trypsin digestion, likely because they were obscured by molecular interactions in the cross-linked material. The majority of the 82 p53 co-enriched proteins are differentially overabundant in the starvation medium group (Fig. 4D) confirming p53 nuclear translocation (Fig. 1D and 1E) and increase in high molecular weight complexes upon 24 hours of starvation (Fig. 4A). Depicting these in an interaction network, reveals a list of known (red edges in Fig. 4E) and novel potential p53 interactors, including several chromatin-related proteins, such as PPP1CC which has been previously shown to modify RNA polymerase 2 phosphorylation state [59], and mediator complex family member MED12. These interactions indicate that p53 is intricately involved in transcriptional complexes upon starvation and is conserved when using a more stringent SAINT score (top 22 with SAINT score ≥ 0.95 in Fig. 4F). Based on our prior work on p53 as a coordinator of the response to fasting [28, 60], we were intrigued by the number of metabolic enzymes mapping to the REACTOME pathway “carbohydrate metabolism” (circled green in Fig. 4E). In particular, UGP2, an enzyme involved in glycogen synthesis, raised our attention as we reported previously that hepatic p53 loss leads to deregulation of glycogen storage [28]. We observed that UGP2 peptides were absent in RIME GM samples, while abundantly present in SM conditions (three out of three replicates). This observation was also reflected in overexpression experiments, where UGP2-FLAG overexpression was pronounced in SM compared to GM conditions (Fig. 4G). Importantly, p53 was co-immunoprecipitated specifically under starvation upon UGP2-FLAG pull-down (Fig. 4G), confirming the RIME result by reverse pull-down. Furthermore, with Sorbin and SH3 domain-containing protein 1 (SORBS1) we confirmed another previously unreported p53 interactor (as judged by the prePPI prediction server <http://honig.c2b2.columbia.edu/>) [61, 62] as increasingly interacting with p53 under SM conditions (Fig. 4H). SORBS1 has been reported to be involved in glucose metabolism via insulin receptor regulation [63]. These findings will spur further investigations to delineate the functional significance of the nuclear interaction of p53 and SORBS1 or UGP2, as well as other candidates identified by the RIME approach.

p53-specific starvation-response transcriptional landscape

Nuclear localization and interactions with chromatin regulators upon 24 hours of starvation suggests that p53 functions to transactivate a starvation-specific set of target genes. Therefore, using our model with CRISPR/Cas9-mediated p53 knock out, we performed RNA-seq analysis with and without re-expression of full-length wild-type FLAG-tagged p53 both in growth medium and after prolonged starvation (workflow shown in Figure 5A, validation of overexpression in Fig. 5B, group clustering in Fig S5A, and expression data in Table S4).

As p53 was previously suggested to solely act as transcriptional activator, while repressive actions were ascribed to secondary effects [20, 22], we limited our analysis to upregulated transcripts (min. 2-fold change, adjusted p-value after BH correction <0.05). Selecting transcripts that are specifically upregulated by p53 overexpression, as well as being starvation-responsive mainly in the presence of p53 (Fig. 5C and S5B), revealed a distinct set of 88 transcripts (Fig. 5C and 5D). Of these, 40 were known, validated direct p53 targets [20] and 48 are putative novel p53 targets under starvation conditions (Fig. 5C, Table S4).

Expression patterns of *bona fide* p53 targets [20] are exemplified for the four transcripts *BTG2*, *PHLDA3*, *CYFIP2*, and *ZNF385A* in Fig. 5E. Confirming our approach and discerning known from potentially novel p53 target genes, mapping of the list of novel p53 targets to gene ontology biological processes or to REACTOME pathways did not yield any significant hits (FDR = 0.01). The list of known p53 target genes returned several highly significantly enriched p53-specific terms and p53-related processes/pathways such as cell cycle control and DNA damage (Fig S5C). Next, we performed *in silico* mapping of p53 binding sites (consensus: GGACATGCCCGGGCATGTCY, Table S5) in the proximity of transcription start sites (TSSs) and compared TSS-centered p53 binding site distribution density from the 48 novel p53 target candidates, with those of the 40 known transcripts, and 48 transcripts that are starvation regulated in a p53-independent manner in our data set. This analysis showed that, similar to known targets, p53 binding sites in the identified target candidates are enriched around TSSs, whereas p53-unregulated transcripts showed no such enrichment peak (Fig. 5F and Table S5). Expression of four novel p53 target gene candidates with p53 binding sites within 1,500 base pairs of TSSs was confirmed via qPCR (Fig. 5G).

Hence, our transcriptome analyses with p53 re-expression on a knock-out background show that p53 coordinates very specific transcriptome changes in response to prolonged starvation. Bioinformatics analyses suggest a number of novel starvation-specific p53 target genes, extending the current knowledge base of p53's transactivation landscape.

Discussion

The specifics of p53 action are highly dependent on the cellular context, the nature of the upstream stressor, and the intensity and duration of the stressor. Activated p53 signaling significantly influences cell fate decisions that range from survival to cell death. p53 activation through various nutrient deprivation conditions has been reported in several cancer model systems [64, 65]. Recent reports showed that p53 protein abundance is increased in cultured hepatocytes by starvation [28] and that p53 signaling is necessary for the physiological response to fasting in the mouse liver [28, 30]. Furthermore, our most recent data provide evidence for the requirement of p53 for the effects of fasting in enhancing hepatocellular carcinoma therapy (Preprint at BioRxiv [29]). Thus, our findings add to a flourish of recent reports that establish p53 as central and potent metabolic regulator in transformed as well as in non-transformed cells [49, 65]. Here, we focused on p53's role as a starvation-response factor and aimed to systematically delineate the entire signaling cascade in response to a starvation stimulus. Using tailored omics methods combined with in-depth bioinformatics analyses and biochemical assays, we define the p53 interaction network leading to nuclear p53 stabilization, the chromatin interactions under prolonged starvation, and the corresponding p53 "targetome" at the transcriptional level.

The interaction of p53 and MDM2 is instrumental for determining cellular p53 levels. Our data suggest a dissociation of MDM2 and p53 as a primary mechanism of p53 stabilization under starvation. MDM2 is regulated by many posttranslational modifications [18]. Our approach, performing p53-BioID to identify cytoplasmatic proteins vicinal to p53 within two hours of nutrient withdrawal, identified increased

association with FKBP3. FKBP3 was shown to interfere with the p53/MDM2 feed-back loop by directly binding to and inducing auto-ubiquitination and degradation of MDM2, subsequently leading to p53 stabilization [39]. Given our observations that MDM2 levels continuously decline during the course of starvation (Fig. 1A) and that the FKBP3/p53 association is increased early after nutrient withdrawal, FKBP3-mediated MDM2 degradation is a plausible mechanism to explain p53 stabilization under starvation. Moreover, we observed a starvation-mediated dissociation of p53 and several members of the 14-3-3 protein family in the BioID data set. From the seven known human isoforms [41], our BioID data suggest p53 associates with five, each showing high sequence coverage and each of which are enriched in growth medium when compared to starvation medium conditions. Dissociation from 14-3-3 proteins under starvation could constitute a part of the mechanism of p53 stabilization and nuclear translocation, as all members of the 14-3-3 family were shown to directly or indirectly contribute to regulation of p53 activity [41] and a commonly described function of 14-3-3 proteins is sequestering proteins in the cytoplasm [66].

Further insights from the BioID-derived p53 interactome include the starvation-enriched association with the known interactor HGMB1 [67, 68] that was shown to prime p53 for DNA interaction [69], and an unexpectedly large number of metabolic enzymes that are preferentially enriched in growth medium conditions. For example, G6PDH was previously described to directly interact with p53 to regulate glycolytic flow through the PPP, supporting biosynthetic processes during growth of HCT116 colon cancer cells [19]. Finding G6PDH in our BioID data set validates our approach and suggests that among the other detected metabolic enzymes physical interactions with p53 are likely and may lead to respective functional consequences. In support of involvement of p53 with the PPP, we also found PRPS1 as a candidate p53 interactor that not only controls purine biosynthesis but also PPP [70], therefore playing a central role in metabolic and oncogenic pathologies [71]. Along these lines, we also find association of p53 with two enzymes involved in oncometabolite production, IDH1 and PGHDH. A number of tumor entities harbor PHGDH overexpression or IDH1 mutations and the subsequent changes in cancer metabolism can be targeted [44]. In this context it might be essential to take a potential interaction with p53 into account.

AMP-activated protein kinase (AMPK) is a central energy sensor with multiple outputs aimed to regenerate ATP levels when cellular energy levels run low [72]. In cancer cells, AMPK was shown to phosphorylate p53 to initiate a mitotic checkpoint upon glucose starvation [64]. The kinase PAK2 has been detected in an AMPK substrate screen and its phosphorylation at Ser20 was shown to increase with rising AMPK activation during nutrient deprivation [73]. Together with our novel finding of a direct p53/PAK2 interaction, which was found to decrease upon starvation, one could surmise that AMPK-mediated PAK2 phosphorylation releases p53 from the p53/PAK2 complex. Thus, our data add another regulatory level to the nutrient sensing AMPK/p53 axis and warrants further investigation.

Further confirming p53 translocation under prolonged starvation, we found exclusive enrichment of potential p53 interactors in starvation medium over growth medium conditions (Fig. 4D) using RIME, a protocol that starts with nuclei isolation of cross-linked cells followed by MS identification of proteins co-

immunoprecipitated with p53. Using IgG pull-down to control for background contamination [47] yielded a *bona fide* starvation-mediated, nuclear p53 interactome. Verifying the validity of this approach, we identified several p53 peptides mostly from its C-terminal domains (Fig. 4C). This suggests that the N-terminal domain and the DBD are obscured in chromatin complexes and therefore not amenable for tryptic digestion and that p53 may be largely bound to DNA. Further (structural) studies are needed to resolve the mode of p53/DNA interaction under starvation conditions. Our RIME approach also identified some known p53 interactors (e.g. MAPK1 [74, 75]) and a number of proteins not known to directly associate with p53. Among these, we focused on UGP2, which we confirmed by reverse co-IP (Fig. 4G). UGP2 is an essential enzyme that synthesizes UDP-glucose from glucose-1-phosphate as a precursor for glycogen synthesis [76, 77]. A plausible scenario is that nuclear translocation of p53 in starvation conditions facilitates UGP2 nuclear transport (or retention) to quench cytoplasmic glycogen synthesis under prolonged starvation. This would make glucose, derived from glycogenolysis or gluconeogenesis, available as an energy substrate during times of starvation, a mechanism that would promote both growth in cancer cells and effective energy partitioning during fasting in normal hepatocytes. Alternatively, a recent report described a surprising role of nuclear glycogen synthesis involving nuclear UGP2 in non-small cell lung cancer progression [78], which could also be affected by p53 under starvation. Another finding from our RIME screen we could confirm with co-IP experiments, was the association of p53 with SORBS1, that has been shown to be involved in insulin receptor signaling [63]. In the light of recent findings, showing nuclear translocation and transcriptional activity of the insulin receptor [79], it would be intriguing to scrutinize the role of p53 in this process under starvation. However, such scenarios need to be further tested, but we view the UGP2/p53 and SORBS1/p53 association as examples of how biological relevant hypotheses can be generated from data obtained with our approaches.

Finally, we devised a tailored transcriptome study to delineate the p53-specific transcriptional response under starvation. CRISPR/Cas9-mediated knock out of p53 in our cell model followed by long-term starvation, yielded transcripts that are regulated by starvation in a p53-independent manner. Subtracting this set from starvation-responsive transcripts upon isogenic re-expression of full-length p53 delivers a starvation-specific targetome, containing known and potential novel p53 target genes. To further increase confidence in the putative novel targets, we used *in silico* binding site prediction and found an enrichment of p53 binding sites in the set of novel p53 target candidate genes when compared to transcripts that are starvation-regulated in a p53-independent manner.

Taken together, our study systematically dissects upstream mechanisms that lead to nuclear p53 stabilization upon starvation, as well as downstream interaction networks and the transcriptome response (Fig. 6). Verification experiments for selective findings prove the validity of the stringent experimental designs and suggest novel starvation-specific mechanisms, such as PAK2-mediated p53 stabilization, nuclear interaction between p53 and UGP2 or SORBS1, and novel high-confidence downstream target genes. These results are indicative for the utility of our complementary omics approaches to extend the already vast landscape of regulatory and functional p53 elements (summarized in Fig. 6).

Our study provides unprecedented details of the p53 signaling cascade upon nutrient withdrawal, thereby expanding the universe of p53 action and interaction in the context of cellular stress. However, we note that global approaches such as the ones used here might produce a considerable number of indirect interactions (MS-based approaches) or indirect regulatory targets (transcriptomics), and may produce false-positive candidates. The latter problem is particularly pertinent to affinity-based proteomics methods that are known to enrich for unspecific proteins merely based on their over-abundance in the specific cellular context. To compensate for this complication, we spent considerable efforts to estimate background noise in the experimental contexts investigated. To this end, we used the same number of background control samples in both the BioID (i.e. EGFP fusion vector) and RIME (i.e. IgG pulldown) experiments and analyzed all our samples in triplicates. In addition, stringent cut-offs in the bioinformatics analyses using the SAINT score [47, 58] reduced the space of potential interactors and increased the degree of confidence in our final data sets. Finally, we note that our findings are specific to HepG2 cells that are transformed and far removed from *in vivo* conditions. To infer general biological relevance of the identified p53 interactions and targets under starvation, key findings would need to be confirmed in physiological models such as primary hepatocytes or liver (cancer) organoids.

Materials And Methods

Cell culture

Hepatocellular carcinoma-derived cells (HepG2) were purchased from ATCC (HB-8065). Cells were maintained in DMEM (gibco, 41966-029) containing 4.5 g/l glucose supplemented with 10% (v/v) heat inactivated fetal bovine serum (FBS; HyCloneTM, SV30160.03) and 1% Penicilin/Streptomycin (gibco, 15140-122). Cells were cultured in a humidified atmosphere of 5% CO₂, 95% air at 37°C. From those cells, HepG2 p53 knockout cell lines (HepG2 p53KO) were generated using p53 CRISPR/Cas9 KO plasmid (h) (sc416469, Santa Cruz Biotechnology). Hepatocellular carcinoma-derived cells (HepG2) were maintained in DMEM (gibco, 41966-029) ("growth medium", GM) containing 4.5 g/l glucose supplemented with 1 % (v/v) heat inactivated fetal bovine serum (FBS; HyCloneTM, SV30160.03) and % Penicilin/Streptomycin (gibco, 15140-122). Cells were cultured in a humidified atmosphere of % CO₂, 9 % air at 37°C. For starvation of cells, medium was removed, cells were washed with phosphate buffered saline (PBS; gibco, 10010-015), and treated in HBSS (gibco, 14175-053) ("starvation medium", SM) containing 1 g/l glucose supplemented with 10 mM HEPES (gibco, 1560-080).

Compound treatments

For treatments with different small molecules, the following concentrations were used:

10 µM nutlin-3a (Biomol, Plymouth Meeting, PA, USA),

10 mM cycloheximide (CHX; Caymen Chem 14126).

10 µM MG132 (Calbiochem 474790)

20 nM FRAX597 (Selleckchem, S7271)

Cell lysis

Lysis buffers were supplemented with 1x protease inhibitor cocktail (PIC) (cOmplete Tablets EASYpack, Roche, 04693116001) and 1x PhosSTOP (Roche, 04906837001). Total protein lysates were obtained after washing the cells with PBS, and application of the respective buffers provided in each protocol segment.

Western Blot (WB)

Whole cell lysates: Cultured cells were scraped and collected in radioimmunoprecipitation assay (RIPA) buffer (50 mM Tris-HCl, 150 mM NaCl, 2 mM EDTA, 50 mM NaF, 0.1% SDS, 0.5% Na-deoxycholate, 1% NP-40, adjusted to pH 7.2 – 7.4) mixed with PhosStop and PIC. Samples were sonicated for 5 min (30 secs on/off) in a cooled water bath ultrasound sonicator (Bioruptor, Diagenode). Cell lysates were centrifuged (14500 rpm 15 min 4°C) and clear supernatants were used to measure protein concentrations with a bicinchoninic acid assay (BCA; Thermo Fisher Scientific). Immunoblotting was performed as described elsewhere (Prokesch et al. 2016), using 50 µg protein of each sample.

Subcellular fractionation: Cytoplasmic and nuclear fractions were separated using a method previously described (Baldwin, Rockland). Briefly, cells were scraped in hypotonic cytoplasmic lysis buffer (10 mM HEPES, 60 mM KCl, 1 mM EDTA, 1 mM, adjusted to pH 7.6) and incubated for 10 min on ice. After addition of IGEPAL 10% to a final concentration of 0.5%, cells were vortexed, incubated on ice for 1 min, vortexed and centrifuged for 4 min at 1500 rpm. The supernatant was collected as the cytoplasmic fraction. Crude nuclei cell pellet was washed once with cytoplasmic lysis buffer (without PIC/PhosSTOP) and resuspended in nuclear extraction buffer (usually one fifth of cytoplasmic buffer volume used; 20 mM Tris-HCl, 100 mM NaCl, 1.5 mM MgCl₂, 0.2 mM EDTA, 0.05% SDS and 25% (v/v) glycerol, adjusted to pH 8.0). The nuclear fraction was sonicated for 5 min (30 secs on/off) in a cooled water bath ultrasound sonicator (Bioruptor, Diagenode). After sonication, 100 U/ml benzonase (Chem Cruz, sc-202391) was added to the nuclear lysates and incubated on ice for 1 – 2 hours, vortexing the mixture periodically (15 min) to resuspend the pellet. Digested sample was centrifuged at 14500 rpm for 15 min and 4°C to precipitate any remaining cell debris. The supernatant was used as the nuclear fraction.

RNA isolation and reverse transcription

After cell lysis with RNA-Lysis-Puffer T (peqGOLD VWR, 12-TRK-88) RNA was isolated with PeqGOLD Total RNA Kit (C-Line) (peqGOLD VWR, 12-6634-02) according to the manufacturer's manual. Sample purification and concentrations was measured with NanoDrop® ND-1000 (peqlab Biotechnologie GmbH). Reverse transcription was performed with High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems by Thermo Fischer Scientific, 4368814). RT Puffer (10x), 0,8 µl dNTP (100 mM), 2 µl RT Random Primer (10x), nuclease free water and MultiScribe Reverse Transcriptase (50 U/µl) were added to defined RNA concentrations. The thermo cycler program was set to 10 min at 25°C, 120 min at 37°C, 5

min at 85°C and subsequent cooling at 4°C. cDNA was diluted afterwards to a concentration of 1 - 20 ng/μl and stored at -80°C.

Real time quantitative PCR (qPCR)

qPCR was performed in 96-well plate or 384 well plates (BioRad, HSP3805) with a cDNA concentration between 1 - 20 ng/μl. A master mix of Blue SybrGreen qPCR (Biozym, 331416) and cDNA (2:1) were prepared and added to each well to pre-pipetted primer pairs (800 nM). After performing the qPCR in a C1000TM Thermal Cycler (BIO –RAD, CFX96TM Real-Time System) (Program: 1 cycle (10 min) at 95°C; 40 cycles: 15 sec at 95°C, 1 min at 60°C, 1 min at 72°C; 1 cycle: 30 sec at 95°C, 30 sec at 60°C, 30 sec at 95°C) data was analyzed with Bio-Rad CFX Manager 3.1. The list of primers is provided in Table S6.

Viability assay

For the viability assay, 2×10^5 cells/well were seeded in a 96-well plate (Thermo Fisher Scientific) in 200 μl of GM to reach full confluency. Cells were allowed to attach for 24 hours at 37°C in 5% CO₂, after which the medium was discarded, and cells were washed with PBS and treated with either GM or SM for 24 hours. Cell viability was analyzed using an EZ4U assay (Biomedica Immunoassays) according to the manufacturer's instructions. Briefly, at the end of treatment the media was replaced with fresh GM (200 μl/well) and 20 μl/well EZ4U working solution. After 2 hours of incubation at 37°C, the absorbance was measured at 492 nm with a reference-wavelength of 620 nm (Spark® 10M multimode microplate reader).

Molecular cloning

Inserts (p53 wt sequence or EGFP wt sequence) for V5-p53-miniTurbo-NES and V5-EGFP-miniTurbo-NES were amplified from Flag-p53 (Addgene #10838; Primer: F: 5'-ACAGCGCTAGCGAGGAGCCGAGTCAGATC-3' R: 5'-ACAGCGCTAGCGGATCCGTCTGAGTCAGGCCCTTCTGT-3') or EGFP (Addgene #13031; Primer: F: 5'-ACAGCGCTAGCGTGAGCAAGGGCGAGGAG-3' R: 5'-ACAGCGGCTAGCGGATCCCTTGTACAGCTCGTCCATGCCG AGAG-3'), cloned into the NheI site in V5-miniTurbo-NES (Addgene #107170) and validated by Sanger sequencing.

Proximity-biotinylation (BioID)

HepG2 p53KO cells were cultured in collagen-coated T75 flasks (approx. 1.5×10^8 cells) and transiently transfected (VIAfect transfection reagent, E4981, Promega) with either V5-p53-miniTurbo-NES or V5-EGFP-miniTurbo-NES. After 24 hours of transfection, intracellular biotinylation was induced by changing cell culture media to either GM or SM (medium composition described in subsection cell culture) both containing 250 μM biotin. Each group was analyzed in 3 biological replicates (GM_p53, SM_p53, GM_EGFP, SM_EGFP; 12 samples total). After 2 hours of biotin labeling, samples were harvested and processed as previously published (Roux, 2018). In brief, cells were lysed in lysis buffer (8 M urea, 50 mM Tris-HCl, pH 7.4) containing protease inhibitor and 1 mM dithiothreitol (DTT). After addition of 20% Triton

X-100 to a final concentration of 1% and sonication for 10 min (30 secs on/off) in a cooled water bath ultrasound sonicator (Bioruptor, Diagenode), samples were subjected to affinity purification with streptavidin beads (Invitrogen, Cat.#65001) on a rotator o/n at 4°C. Afterwards, beads with captured biotinylated proteins were thoroughly washed (10 times) with washing buffer (50 mM Tris-HCl, pH 7.4, 8 M urea) and final washing steps were performed with 50 mM Tris-HCl pH 7.4 and 50 mM AMBIC containing 1 mM biotin to block remaining streptavidin on the beads. The beads including the captured proteins were reconstituted in 100 mM Tris-HCl pH 8.5, 2% sodium deoxycholate (SDC) and reduced/alkylated with 5 mM TCEP/30 mM chloroacetamide at 56°C for 10 min. Next, the proteins were quantified (using a BCA assay) and digested with 1:100 Lys-C and 1:50 trypsin overnight at 37°C with continuous shaking. Digestion was stopped by adding 1% trifluoroacetic acid (TFA) to a final pH of 2, in which SDC was completely precipitated. SDC and the beads were removed by centrifugation at 14000 rpm for 10 min and the supernatant, containing peptides was desalted on an Oasis HLB plate (Waters). Peptides were dried and dissolved in 2% formic acid before liquid chromatography-tandem mass spectrometry analysis.

Mass spectrometry run: One thousand ng of each sample was analyzed using an Ultimate3000 high-performance liquid chromatography system (Thermo Fisher Scientific) coupled to an EXPLORIS 480 mass spectrometer (Thermo Fisher Scientific). Buffer A consisted of water acidified with 0.1% formic acid, while Buffer B was 80% acetonitrile (ACN) and 20% water with 0.1% FA. The peptides were first trapped for 1 min at 30 μ l/min with 100% Buffer A on a trap (0.3 $\times$ 5 mm with PepMap C18, 5 μ m – 100 Å Thermo Fisher Scientific); after the trapping peptides were separated by a 50 cm analytical column packed with C18 beads (Poroshell 120 EC-C18, 2.7 μ m, Agilent Technologies). The gradient was 9 – 40% B in 40 min at 400 nL/min. Buffer B was then raised to 55% in 5 min and increased to 99% for the cleaning step. Peptides were ionized using a spray voltage of 2 kV and a capillary heated at 275°C. The mass spectrometer was set to acquire full-scan MS spectra (350–1400 m/z) for a maximum injection time of 120 ms at a mass resolution of 60,000 and an automated gain control (AGC) target value of 300%. For a total cycle of 1 second the most intense precursor ions were selected for tandem MS (MS/MS). HCD fragmentation was performed in the HCD cell, with the readout in the Orbitrap mass analyzer at a resolution of 15,000 (isolation window of 1.4 Th) and an AGC target value of 200% with a maximum injection time of 25 ms and a normalized collision energy of 28%.

BioID data analysis: All raw files were analyzed by MaxQuant v1.6.17 software using the integrated Andromeda Search engine and searched against the Human UniProt Reference Proteome (October 2020 release with 75,088 protein sequences). MaxQuant was used with the standard parameters (“Label-Free Quantification”, “iBAQ” and “Match between runs” were selected with automatic values) with only the addition of Deamidation (N) as variable modification. Data analysis was performed with Perseus v1.6.14: proteins reported in the file “proteinGroups.txt” were filtered for reverse, potential contaminants and identified by site. For the quantitation, we used the iBAQ values calculated by MaxQuant and we kept only proteins found in at least 2 biological replicates in each group. At this point intensities were log10 transformed and missing values were imputed by Perseus with the automatic settings (width: 0.3, down shift: 1.8, mode: separately for each column) leading to 2,500 proteins left for statistical analysis with

ANOVA testing (Benjamini-Hochberg FDR 0.05), z-score (mean per row) and hierarchical clustering (distance: Euclidian). Clusters were selected if enriched over EGFP background control under both nutrient conditions. Volcano plots were generated with Perseus 1.4.16 using the setting t-test, GM_p53 vs SM_p53, FDR0.05. KEGG overrepresentation analysis was performed in webgestalt.org [80] functional database 'pathway – KEGG' against 'genome protein-coding' reference list with 272 high confidence p53 interactors originating from the hierarchical clustering analysis (3 clusters enriched over EGFP background, Fig. 2C). In a second MaxQuant screen with the annotated proteins from the initial search only, biotinylated peptides were identified, by adding the modification Biotin(K)="C(10) H(14) N(2) O(2) S" (+226.077598) possible on any K or protein N-term as search criteria. The resulting biotinylated peptides were filtered for occurring only in the p53 samples (Table S2). Biotinylated peptides are added in the search output (PRIDE) with file name "Biotin (K)Sites.txt" together with the other submitted files. Unique peptides found only in p53 samples were analyzed in Perseus v1.6.14: peptides from "peptideGroups.txt" were first filtered for reverse, potential contaminant, identified by site and afterwards only for occurring in p53 samples (Table S2). For SAINTscore analysis, the peptide counts for each sample were retrieved from MaxQuant output file "proteinGroups.txt" and prepared as input file for analysis in the Contaminant Repository for Affinity Purification (CRAPome, lit). Analysis in CRAPome was conducted with experiment type: Proximity Dependent Biotinylation; Quantitation type: SPC without selecting CRAPome internal controls, instead only experimental controls as background (GM_p53, SM_p53, and CONTROL) EGFP control groups were merged to a single group, as we specifically look for proteins enriched over background control in both conditions (GM and SM), similar to the hierarchical clustering analysis. Interaction scoring was conducted with standard analysis options. For generating dot plots, the CRAPome output file was downloaded via the External Tools section and used as input file in ProHits-viz [48].

Fluorescence microscopy

Cells were seeded onto collagen-coated glass Chamber Slides and transfected with ViaFECT (Promega) according to the manufacturer's protocol. After 18 hours, cells were washed two times with PBS and fixed with 1% paraformaldehyde (PFA; Thermo scientific) for 15 min at room temperature. PFA was removed and cells were washed two times with PBS. After washing, cells were blocked using Ultravision Hydrogen Peroxide Block (Thermo scientific, Rockford, USA) for 5 min at RT. Then, cells were incubated with primary antibodies (listed in Table 11) diluted in Antibody diluent (Dako, California, USA) for 45 min in a dark environment. Cells were then washed two times with PBS, followed by incubation with secondary antibody Anti-rabbit IgG (H+L) Alexa Fluor 555 (Cell Signaling) for 30 min in a dark environment. Secondary antibodies were diluted 1:1000 in Antibody diluent. After another two times PBS washing, nuclei were stained using DAPI 1:2000 (Thermo scientific) diluted in PBS for 5 min at RT. Finally, chambers were removed from the slides and the slides were washed twice with 1x PBS before mounting with ProLong™ Gold antifade reagent (Invitrogen) and applying a cover slip. Slides were stored in the dark until visualizing on the PALM Microbeam Laser Microdissection microscope (Zeiss, New York, USA) using 40x magnification. Images were analyzed using Zen software 2.3. Brightness and contrast were modified for enhanced visualization.

Expression and purification of recombinant proteins

Expression constructs for the fragments of p53 (Uniprot ID P04637) corresponding to amino acid 1 to 94 (p53^{TAD}) and 94 to 312, as well as for PAK2 (Uniprot ID Q13177) corresponding to amino acid 1 to 212 were generated by synthesis of the corresponding optimized cDNA-constructs of p53 or PAK2 respectively and insertion of these cDNA into a pETM11-ZZ-His₆ vector via NcoI/BamHI restriction digest.

For expression of recombinant unlabeled or ¹⁵N labelled ZZ-His₆ proteins, the bacterial expression vectors were transformed into Escherichia Coli BL21-DE3 Star strain cells. Cells were either grown in lysogeny broth medium for unlabeled proteins or minimum medium supplemented with 6g ¹²C₆H₁₂O₆ and 1 g ¹⁵NH₄Cl at 37°C until they reached an OD600 of 0.8, when protein expression was induced by addition of 0.5 mM IPTG. After proteins were expressed at 20°C for 16 hours, cells expressing disordered fragments (p53¹⁻⁹⁴ and PAK2¹⁻²¹²) were harvested in denaturing buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 2 mM Imidazol, 6 M Urea) and cells expressing folded fragments (p53⁹⁴⁻³¹²) were harvested in non-denaturing harvesting buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 2 mM Imidazol, 2 mM tris(2-carboxyethyl)phosphine)), followed by sonication and centrifugation at 6198 rcf for 45 min. Proteins were purified from the lysate using Ni-NTA agarose (Qiagen) and the ZZ-His₆ tag was cleaved by 2% (w/w) His₆-tagged TEV protease treatment for 16 hours at 4°C. After a desalting step (HiPrep 26/10, GE Healthcare) on an ÄKTA Pure system (GE Healthcare) into a low imidazole buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 2 mM Imidazol), the cleaved protein was separated from the uncleaved protein and the His₆ tagged TEV protease by Ni-NTA affinity chromatography. Finally, the proteins were purified using size exclusion chromatography (Superdex 75 Increase, GE Healthcare).

Nuclear magnetic resonance (NMR) binding assay

For binding studies all proteins were equilibrated in the same buffer containing 20 mM Hepes pH 7.0, 50 mM NaCl, 2 mM TCEP. Samples for NMR measurements contained 100 µM ¹⁵N labelled p53⁹⁴⁻³¹² in presence of 0 µM, 50 µM, 100 µM or 200 µM unlabeled PAK2¹⁻²¹² and 10% D₂O. For titrations using ¹⁵N labelled PAK2¹⁻²¹² 50 µM were used in presence 0 µM, 50 µM and 100 µM p53⁹⁴⁻³¹² and 10% D₂O. ¹H¹⁵N HSQC spectra of the aforementioned samples were recorded at 25°C on a Bruker 600 Mhz Avance Neo NMR spectrometer equipped with a TXI room temperature probe.

Rapid immunoprecipitation mass spectrometry of endogenous proteins (RIME) for analysis chromatin complexes

Wild-type (p53-proficient) HepG2 cells were grown to 100% confluency and treated with either SM or GM for 24 hours. After crosslinking with 1% FA (prepared in either SM or serum-free GM) for 8 min, the reaction was quenched with glycine at a final concentration of 0.1 M and cells washed within the flask twice with ice-cold PBS. Cells were immediately frozen at -80°C and stored in flasks until use. Cell lysates were prepared as previously described (Mohammed et al. 2016) and nuclei extracted for immune-

precipitation. IP was conducted with either DO1X antibody for p53-specific pulldown or IgG2A isotype control. After thorough washing of the bead-bound proteins with AMBIC, proteins were digested on-bead with trypsin (1:50) o/n and for an additional 4 hours the next day. Derived peptides were desalted with C18 ultra micro spin columns and analyzed with LC-MS/MS.

Mass spectrometry run: One-tenth of each sample was measured by nano-HPLC (Dionex Ultimate 3000) equipped with an Aurora Series Emitter nanocolumn with CSI fitting (C18, 1.6 μ m, 120 Å, 250 x 0.075 mm) (IonOpticks, Melbourne, Australia). Separation was carried out at 50°C at a flow rate of 300 nl/min using the following gradient. Solvent A is 0.1% formic acid in water and solvent B is acetonitrile containing 0.1% formic acid: 0-18 min: 2% B; 18-100 min: 2-25% B; 100-107 min: 25-35% B, 107-108 min: 35-95% B; 108-118 min: 95% B, 118-118 min: 95-2% B; 118-133 min: 2% B. The Bruker maXis II ETD mass spectrometer was operated with the captive source in positive mode with following settings: mass range: 150 - 2200 m/z, 4 Hz, precursor acquisition control top20 (CID), capillary 1600 V, dry gas flow 3 L/min with 150°C, nanoBooster 0.2 bar.

RIME data analysis: The MS/MS data were analyzed for protein identification and label-free quantification using MaxQuant 1.6.1.0 against the public database Swiss-Prot with taxonomy *Homo sapiens* and common contaminants (downloaded on 16.04.2019, 20482 sequences). Detailed search criteria were used as follows: trypsin, max. missed cleavage sites: 2; oxidation on Met as variable modification; search mode: MS/MS ion search with decoy database search included; precursor mass tolerance +/- 0.006 Da; product mass tolerance +/- 80 ppm; acceptance parameters for identification: 1% PSM FDR; 1% protein FDR and 1% site decoy fraction. In addition, a label free quantitation and iBAQ values including the match between runs feature of MaxQuant was performed (Cox et al. 2014) requiring a minimum of 2 ratio counts of quantified razor and unique peptides while omitting the normalization step. Data processing was performed using Perseus software version 1.6.6.0, contaminants and reverse proteins created during database search were removed. Intensities were log10 transformed in order to lower the effect of the outlier values, filtered for 3 valid values in either GM_p53 or SM_p53, or at least 4 valid values in total, allowed no values in the IgG control groups. Missing intensities were replaced with random values taken from the Gaussian distribution of values using default parameters (width: 0.3, down shift: 1.8, mode: separately for each column), in order to simulate a value for low abundant proteins. Two-sample t-tests and volcano blotting were used to identify altered proteins between GM and SM conditions. PCA analysis and multi-scatter plot were generated in Perseus. For SAINTscore analysis, the peptide counts for each sample were retrieved from MaxQuant output file "proteinGroups.txt" and prepared as input file for analysis in the Contaminant Repository for Affinity Purification (CRAPome, lit). Analysis in CRAPome was conducted with experiment type: Endogenous pull-down; Quantitation type: SPC without selecting CRAPome internal controls, instead only experimental controls as background (GM_p53, SM_p53, and IgG CONTROL) IgG control groups were merged to a single group, as we specifically look for proteins enriched over background control in both conditions (GM and SM). Interaction scoring was conducted with standard analysis options. For generating dot plots and comparison of average spectral counts, the CRAPome output file was downloaded via the External Tools section and used as input file in ProHits-viz [48].

Co-Immunoprecipitation (IP)

HA-p53 (Genscript, OHu20059C) and/or UGP2-FLAG (Genscript, OHu02074D) were transiently overexpressed in HepG2 p53KO cells with ViaFECT (Promega) according to the manufacturer's protocol. After 24 hours of transfection, cell culture media was changed to either GM or SM and kept under those conditions for another 24 hours. Next, cells were harvested by washing with ice-cold PBS and lysed in Co-IP buffer (25 mM Tris-HCl, 150 mM NaCl, 1 mM EDTA, 1% IGEPAL, 5% glycerol; adjusted to pH 7.4) including PIC and PhosSTOP. After 30 min incubation on ice (passive lysis), the suspensions were sonicated for 5 min (30 secs on/off) in a cooled water bath ultrasound sonicator (Bioruptor, Diagenode) at 4°C. The lysates were centrifuged for 5 min at 14500 rpm and 4°C. Protein concentration was determined with a BCA protein assay kit. 500 µg of protein lysates was normalized to equal volumes with Co-IP buffer and used for precipitation. 30 µl Flag-beads slurry (Sigma, M8823) was used per mg protein, equilibrated in 500 µl Co-IP buffer for 3 times and incubated with the protein lysates over night at 4°C with reciprocal shaking (BioSan MultiBio RS-24). Next day, flow-through was collected and used for control blots. Beads were washed 3 times with Co-IP buffer and proteins eluted from the beads with elution buffer (100 mM glycine, pH 3) at room temperature for 5 min.

RNA-seq experiments and data analysis

Three SEQ libraries (TruSeq® Stranded mRNA Library Prep) each of vehicle control (pcDNA4/HisMax, Invitrogen) and FLAG-p53 (pcDNA3 flag-p53, Addgene) re-expressing cells (electroporation) were prepared from purified mRNAs of HepG2 p53KO cells under untreated (GM) or treated (SM) conditions for a total of 4 x 3 (12) libraries. Raw sequencing data was generated on an Illumina NextSeq 550 and mapped to the human genome (hg18) using STAR alignment (Dobin et al. 2013). Differentially expressed genes were found using DEseq2 package in R, data was transformed (VST), Wald-Test performed on individual genes, and Benjamini-Hochberg corrected for multiple testing. Genes with an adjusted P-value ≤ 0.05 were used for further analysis.

p53 motif analysis and motif density

Analysis was conducted using HOMER (<http://homer.ucsd.edu/homer/index.html>) (Heinz et al. 2010). A motif file was generated based on p53 consensus sequence "GGACATGCCCGGGCATGTCY" (Human, hg18) derived from MotifMap (<http://motifmap.ics.uci.edu/>) (Daily et al. 2011) using HOMER's seq2profile.pl, allowing 6 mismatches (Table Sx). Motif sites were identified via HOMER's annotatePeaks.pl module in "TSS mode", searching ± 10 kb around each gene's TSS on either the sense or the antisense strand in the provided lists of genes of interest. A Gaussian density plot was generated in R (<https://R-project.org/>) (R Core Team, 2017) using the ggplot2 package (<https://ggplot2.tidyverse.org>) (Wickham H. 2016).

Declarations

DATA AND SOFTWARE AVAILABILITY

The BioID data obtained in this study has been uploaded to ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD027734.

The RIME data obtained in this study has been uploaded to ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD028341.

The RNA-Seq data obtained in this study has been uploaded to NCBI GEO datasets, under accession number GSE183127.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Conflict of Interest

The authors declare no competing interests.

Funding

A.P. was supported by the Austrian Science Fund (FWF, grants P29328, I3165, P34109) and by a MEFOgraz grant from the Medical University of Graz. This work has also been supported by EPIC-XS project (EPIC-XS-0000206) funded by the Horizon 2020 program of the European Union. I.R., M.G., K.S., N.B., N.K., H.M., and C.N. were supported by the Austrian Science Fund (FWF, grants P29328 and I3165). I.R. was funded by the PhD faculty MolMed at the Medical University of Graz. J.K. has been supported by a grant of the Oesterreichische Nationalbank (Austrian Central Bank, Anniversary Fund, project number: 18517) and by a MEFOgraz research grant. This study was also supported by Austrian Science fund (FWF) projects KLI645, Doctoral school “DK Metabolic and Cardiovascular disease” (W1226), and SFB “Lipid hydrolysis” (F73) to R.B.G. Finally, M.S. and M.O. were supported by the German Research foundation (grant SCHU 2546/4-1 to M.S).

Authors' information

Contributions: M.G. designed and performed the experiments, analyzed the data, and wrote the manuscript. A.P. designed, coordinated, and supervised the project, analyzed the data, and wrote the manuscript. I.R., J.K., G.L.G., M.O., E.M., M.A., H.M., and C.N. performed experiments and analyzed data. N.K., E.H., and P.U. performed and analyzed RNA-seq experiments. M.G., R.Z.C., R.B.G., and L.L. performed and analyzed proteomics experiments. E.S. and T.M. performed NMR interaction studies. C.H. performed in silico binding site analyses. M.S., T.J.S., G.L.G., M.O., and A.J.R.H. contributed materials and provided expertise and feedback.

References

1. Levine AJ (2020) P53: 800 Million Years of Evolution and 40 Years of Discovery. *Nat Rev Cancer* 20:471–480. <https://doi.org/10.1038/s41568-020-0262-1>
2. Kruse JP, Gu W (2009) Modes of p53 Regulation. *Cell* 137:609–622. <https://doi.org/10.1016/j.cell.2009.04.050>
3. Vousden KH, Prives C (2009) Blinded by the Light: The Growing Complexity of p53. *Cell* 137:413–431. <https://doi.org/10.1016/j.cell.2009.04.037>
4. Kasthuber ER, Lowe SW (2017) Putting p53 in Context. *Cell* 170:1062–1078. <https://doi.org/10.1016/j.cell.2017.08.028>
5. Hanahan D, Weinberg RA (2011) Hallmarks of cancer: The next generation. *Cell* 144:646–674. <https://doi.org/10.1016/j.cell.2011.02.013>
6. Levine AJ (2020) P53 and The Immune Response: 40 Years of Exploration—A Plan for the Future. *Int J Mol Sci* 21:541. <https://doi.org/10.3390/ijms21020541>
7. Lacroix M, Riscal R, Arena G et al (2020) Metabolic functions of the tumor suppressor p53: Implications in normal physiology, metabolic disorders, and cancer. *Mol Metab* 33:2–22. <https://doi.org/10.1016/j.molmet.2019.10.002>
8. Simabuco FM, Morale MG, Pavan ICB et al (2018) P53 and Metabolism: From Mechanism To Therapeutics. *Oncotarget* 9:23780–23823. <https://doi.org/10.18632/oncotarget.25267>
9. Krstic J, Galhuber M, Schulz T et al (2018) p53 as a Dichotomous Regulator of Liver Disease: The Dose Makes the Medicine. *Int J Mol Sci* 19:921. <https://doi.org/10.3390/ijms19030921>
10. Labuschagne CF, Zani F, Vousden KH (2018) BBA - Reviews on Cancer Control of metabolism by p53 – Cancer and beyond. *BBA - Rev Cancer* 1870:32–42. <https://doi.org/10.1016/j.bbcan.2018.06.001>
11. Li M, Brooks CL, Wu-Baer F et al (2003) Mono- versus polyubiquitination: differential control of p53 fate by Mdm2. *Science* 302:1972–1975. <https://doi.org/10.1126/science.1091362>
12. Honda R, Tanaka H, Yasuda H (1997) Oncoprotein MDM2 is a ubiquitin ligase E3 for tumor suppressor p53. *FEBS Lett* 420:25–27. [https://doi.org/10.1016/S0014-5793\(97\)01480-4](https://doi.org/10.1016/S0014-5793(97)01480-4)
13. Gajjar M, Candeias MM, Malbert-Colas L et al (2012) The p53 mRNA-Mdm2 interaction controls mdm2 nuclear trafficking and is required for p53 activation following dna damage. *Cancer Cell* 21:25–35. <https://doi.org/10.1016/j.ccr.2011.11.016>
14. Candeias MM, Malbert-Colas L, Powell DJ et al (2008) p53 mRNA controls p53 activity by managing Mdm2 functions. *Nat Cell Biol* 10:1098–1105. <https://doi.org/10.1038/ncb1770>
15. Barak Y, Juven T, Haffner R, Oren M (1993) mdm2 Expression is induced by wild type p53 activity. *EMBO J* 12:461–468. <https://doi.org/10.1002/j.1460-2075.1993.tb05678.x>
16. Wu X, Bayle JH, Olson D, Levine AJ (1993) The p53-mdm-2 autoregulatory feedback loop. *Cell* 73:1126–1132

17. Hock AK, Vousden KH (2014) The role of ubiquitin modification in the regulation of p53. *Biochim Biophys Acta - Mol Cell Res* 1843:137–149. <https://doi.org/10.1016/j.bbamcr.2013.05.022>
18. Liu Y, Tavana O, Gu W (2019) P53 Modifications: Exquisite Decorations of the Powerful Guardian. *J Mol Cell Biol* 11:564–577. <https://doi.org/10.1093/jmcb/mjz060>
19. Jiang P, Du W, Wang X et al (2011) p53 regulates biosynthesis through direct inactivation of glucose-6-phosphate dehydrogenase. *Nat Cell Biol* 13:310–316. <https://doi.org/10.1038/ncb2172>
20. Fischer M (2017) Census and evaluation of p53 target genes. *Oncogene* 36:3943–3956. <https://doi.org/10.1038/onc.2016.502>
21. Sullivan KD, Galbraith MD, Andrysik Z, Espinosa JM (2018) Mechanisms of transcriptional regulation by p53. *Cell Death Differ* 25:133–143. <https://doi.org/10.1038/cdd.2017.174>
22. Hafner A, Bulyk ML, Jambhekar A, Lahav G The multiple mechanisms that regulate p53 activity and cell fate. *Nat Rev Mol Cell Biol* 11:1–13. <https://doi.org/10.1038/s41580-019-0110-x>
23. Nencioni A, Caffa I, Cortellino S, Longo VD (2018) Fasting and cancer: molecular mechanisms and clinical application. *Nat Rev Cancer* 18:707–719. <https://doi.org/10.1038/s41568-018-0061-0>
24. de Cabo R, Mattson MP (2019) Effects of Intermittent Fasting on Health, Aging, and Disease. *N Engl J Med* 381:2541–2551. <https://doi.org/10.1056/NEJMr1905136>
25. Krstic J, Pieber TR, Prokesch A (2020) Stratifying nutritional restriction in cancer therapy: Next stop, personalized medicine. *International Review of Cell and Molecular Biology*. Elsevier Inc., pp 231–259
26. Lévesque S, Le Naour J, Pietrocola F et al (2019) A synergistic triad of chemotherapy, immune checkpoint inhibitors, and caloric restriction mimetics eradicates tumors in mice. *Oncoimmunology* 8:e1657375. <https://doi.org/10.1080/2162402X.2019.1657375>
27. Deligiorgi MV, Liapi C, Trafalis DT (2020) How Far Are We from Prescribing Fasting as Anticancer Medicine? *Int J Mol Sci* 21:9175. <https://doi.org/10.3390/ijms21239175>
28. Prokesch A, Graef FA, Madl T et al (2017) Liver p53 is stabilized upon starvation and required for amino acid catabolism and gluconeogenesis. *FASEB J* 31:732–742. <https://doi.org/10.1096/fj.201600845R>
29. Krstic J, Reinisch I, Schindlmaier K et al (2021) Fasting reverses drug-resistance in hepatocellular carcinoma through p53-dependent metabolic synergism. *bioRxiv* 2021.02.10.430545
30. Gonzalez-Rellan MJ, Fondevila MF, Fernandez U et al (2021) O-GlcNAcylated p53 in the liver modulates hepatic glucose production. *Nat Commun* 12:5068. <https://doi.org/10.1038/s41467-021-25390-0>
31. Wade M, Wang YV, Wahl GM (2010) The p53 orchestra: Mdm2 and Mdmx set the tone. *Trends Cell Biol* 20:299–309. <https://doi.org/10.1016/j.tcb.2010.01.009>
32. Lee JM, Wagner M, Xiao R et al (2014) Nutrient-sensing nuclear receptors coordinate autophagy. *Nature* 516:112–115. <https://doi.org/10.1038/nature13961>
33. Aubrey BJ, Kelly GL, Janic A et al (2018) How does p53 induce apoptosis and how does this relate to p53-mediated tumour suppression? *Cell Death Differ* 25:104–113.

<https://doi.org/10.1038/cdd.2017.169>

34. Vassilev LT, Vu BT, Graves B et al (2004) In Vivo Activation of the p53 Pathway by Small-Molecule Antagonists of MDM2. *Science* (80-) 303:844–848. <https://doi.org/10.1126/science.1092472>
35. O’Keefe K, Li H, Zhang Y (2003) Nucleocytoplasmic shuttling of p53 is essential for MDM2-mediated cytoplasmic degradation but not ubiquitination. *Mol Cell Biol* 23:6396–6405. <https://doi.org/10.1128/MCB.23.18.6396-6405.2003>
36. Anwar A, Norris DA, Fujita M (2011) Ubiquitin proteasomal pathway mediated degradation of p53 in melanoma. *Arch Biochem Biophys* 508:198–203. <https://doi.org/10.1016/j.abb.2010.12.012>
37. Branon TC, Bosch JA, Sanchez AD et al (2018) Efficient proximity labeling in living cells and organisms with TurboID. *Nat Biotechnol* 36:880–898. <https://doi.org/10.1038/nbt.4201>
38. Livesey KM, Kang R, Vernon P et al (2012) p53/HMGB1 complexes regulate autophagy and apoptosis. *Cancer Res* 72:1996–2005. <https://doi.org/10.1158/0008-5472.CAN-11-2291>
39. Ochocka AM, Kampanis P, Nicol S et al (2009) FKBP25, a novel regulator of the p53 pathway, induces the degradation of MDM2 and activation of p53. *FEBS Lett* 583:621–626. <https://doi.org/10.1016/j.febslet.2009.01.009>
40. Wang W, Wei J, Zhang H et al (2021) PRDX2 promotes the proliferation of colorectal cancer cells by increasing the ubiquitinated degradation of p53. *Cell Death Dis* 12:605. <https://doi.org/10.1038/s41419-021-03888-1>
41. Falcicchio M, Ward JA, Macip S, Doveston RG (2020) Regulation of p53 by the 14-3-3 protein interaction network: new opportunities for drug discovery in cancer. *Cell death Discov* 6:126. <https://doi.org/10.1038/s41420-020-00362-3>
42. Geeraerts SL, Heylen E, De Keersmaecker K, Kampen KR (2021) The ins and outs of serine and glycine metabolism in cancer. *Nat Metab* 3:131–141. <https://doi.org/10.1038/s42255-020-00329-9>
43. Ou Y, Wang SJ, Jiang L et al (2015) p53 Protein-mediated regulation of phosphoglycerate dehydrogenase (PHGDH) is crucial for the apoptotic response upon serine starvation. *J Biol Chem* 290:457–466. <https://doi.org/10.1074/jbc.M114.616359>
44. DeBerardinis RJ, Chandel NS (2016) Fundamentals of cancer metabolism. *Sci Adv* 2:e1600200. <https://doi.org/10.1126/sciadv.1600200>
45. Saleem A, Hood DA (2013) Acute exercise induces tumour suppressor protein p53 translocation to the mitochondria and promotes a p53-Tfam-mitochondrial DNA complex in skeletal muscle. *J Physiol* 591:3625–3636. <https://doi.org/10.1113/jphysiol.2013.252791>
46. Breitkreutz A, Choi H, Sharom JR et al (2010) A Global Protein Kinase and Phosphatase Interaction Network in Yeast. *Science* (80-) 328:1043–1046. <https://doi.org/10.1126/science.1176495>
47. Mellacheruvu D, Wright Z, Couzens AL et al (2013) The CRAPome: A contaminant repository for affinity purification-mass spectrometry data. *Nat Methods* 10:730–736. <https://doi.org/10.1038/nmeth.2557>

48. Knight JDR, Liu G, Zhang JP et al (2015) A web-tool for visualizing quantitative protein-protein interaction data. *Proteomics* 15:1432–1436. <https://doi.org/10.1002/pmic.201400429>
49. Flöter J, Kaymak I, Schulze A (2017) Regulation of Metabolic Activity by p53. *Metabolites* 7:21. <https://doi.org/10.3390/metabo7020021>
50. Bensaad K, Tsuruta A, Selak MA et al (2006) TIGAR, a p53-inducible regulator of glycolysis and apoptosis. *Cell* 126:107–120. <https://doi.org/10.1016/j.cell.2006.05.036>
51. Johnson DI (1999) Cdc42: An Essential Rho-Type GTPase Controlling Eukaryotic Cell Polarity. *Microbiol Mol Biol Rev* 63:54–105. <https://doi.org/10.1128/membr.63.1.54-105.1999>
52. Stark C, Breitkreutz BJ, Reguly T et al (2006) BioGRID: a general repository for interaction datasets. *Nucleic Acids Res* 34:535–539. <https://doi.org/10.1093/nar/gkj109>
53. Renkema GH, Pulkkinen K, Saksela K (2002) Cdc42/Rac1-Mediated Activation Primes PAK2 for Superactivation by Tyrosine Phosphorylation. *Mol Cell Biol* 22:6719–6725. <https://doi.org/10.1128/mcb.22.19.6719-6725.2002>
54. Madeira F, Tinti M, Murugesan G et al (2015) 14-3-3-Pred: Improved methods to predict 14-3-3-binding phosphopeptides. *Bioinformatics* 31:2276–2283. <https://doi.org/10.1093/bioinformatics/btv133>
55. Yeo D, He H, Patel O et al (2016) FRAX597, a PAK1 inhibitor, synergistically reduces pancreatic cancer growth when combined with gemcitabine. *BMC Cancer* 16:24. <https://doi.org/10.1186/s12885-016-2057-z>
56. Cai Q, Xiao B, Si H et al (2012) Kaposi's sarcoma herpesvirus upregulates Aurora A expression to promote p53 phosphorylation and ubiquitylation. *PLoS Pathog* 8:e1002566. <https://doi.org/10.1371/journal.ppat.1002566>
57. Mohammed H, Taylor C, Brown GD et al (2016) Rapid immunoprecipitation mass spectrometry of endogenous proteins (RIME) for analysis of chromatin complexes. *Nat Protoc* 11:316–326. <https://doi.org/10.1038/nprot.2016.020>
58. Choi H, Larsen B, Lin ZY et al (2011) SAINT: Probabilistic scoring of affinity purification-mass spectrometry data. *Nat Methods* 8:70–73. <https://doi.org/10.1038/nmeth.1541>
59. Bharucha DC, Zhou M, Nekhai S et al (2002) A Protein Phosphatase from Human T Cells Augments Tat Transactivation of the Human Immunodeficiency Virus Type 1 Long-Terminal Repeat. *Virology* 296:6–16. <https://doi.org/10.1006/viro.2002.1438>
60. Schupp M, Chen F, Briggs ER et al (2013) Metabolite and transcriptome analysis during fasting suggest a role for the p53-Ddit4 axis in major metabolic tissues. *BMC Genomics* 14:758. <https://doi.org/10.1186/1471-2164-14-758>
61. Zhang QC, Petrey D, Garzón JI et al (2013) PrePPI: A structure-informed database of protein-protein interactions. *Nucleic Acids Res* 41:828–833. <https://doi.org/10.1093/nar/gks1231>
62. Zhang QC, Petrey D, Deng L et al (2012) Structure-based prediction of protein-protein interactions on a genome-wide scale. *Nature* 490:556–560. <https://doi.org/10.1038/nature11503>

63. Baumann CA, Ribon V, Kanzaki M et al (2000) CAP defines a second signalling pathway required for insulin-stimulated glucose transport. *Nature* 407:202–207. <https://doi.org/10.1038/35025089>
64. Jones RG, Plas DR, Kubek S et al (2005) AMP-activated protein kinase induces a p53-dependent metabolic checkpoint. *Mol Cell* 18:283–293. <https://doi.org/10.1016/j.molcel.2005.03.027>
65. Labuschagne CF, Zani F, Vousden KH (2018) Control of metabolism by p53 - Cancer and beyond. *Biochim Biophys Acta* 56:1511–1519. <https://doi.org/10.1016/j.bbcan.2018.06.001>
66. Mhawech P (2005) 14-3-3 proteins—an update. *Cell Res* 15:228–236. <https://doi.org/10.1038/sj.cr.7290291>
67. Livesey KM, Kang R, Zeh HJ et al (2012) Direct molecular interactions between HMGB1 and TP53 in colorectal cancer. *Autophagy* 8:846–848. <https://doi.org/10.4161/auto.19891>
68. Štros M, Muselikova-Polanska E, Pospisilova S, Strauss F (2004) High-affinity binding of tumor-suppressor protein p53 and HMGB1 to hemicatenated DNA loops. *Biochemistry* 43:7215–7225. <https://doi.org/10.1021/bi049928k>
69. Rowell JP, Simpson KL, Stott K et al (2012) HMGB1-facilitated p53 DNA binding occurs via HMG-Box/p53 transactivation domain interaction, regulated by the acidic tail. *Structure* 20:2014–2024. <https://doi.org/10.1016/j.str.2012.09.004>
70. Jing X, Wang XJ, Zhang T et al (2019) Cell-cycle-dependent phosphorylation of PrPS1 fuels nucleotide synthesis and promotes tumorigenesis. *Cancer Res* 79:4650–4664. <https://doi.org/10.1158/0008-5472.CAN-18-2486>
71. Ge T, Yang J, Zhou S et al (2020) The Role of the Pentose Phosphate Pathway in Diabetes and Cancer. *Front Endocrinol (Lausanne)* 11:365. <https://doi.org/10.3389/fendo.2020.00365>
72. Hardie DG (2015) AMPK: Positive and negative regulation, and its role in whole-body energy homeostasis. *Curr Opin Cell Biol* 33:1–7. <https://doi.org/10.1016/j.ceb.2014.09.004>
73. Banko MR, Allen JJ, Schaffer BE et al (2011) Chemical Genetic Screen for AMPK α 2 Substrates Uncovers a Network of Proteins Involved in Mitosis. *Mol Cell* 44:878–892. <https://doi.org/10.1016/j.molcel.2011.11.005>
74. Yeh PY, Chuang SE, Yeh KH et al (2004) Phosphorylation of p53 on Thr55 by ERK2 is necessary for doxorubicin-induced p53 activation and cell death. *Oncogene* 23:3580–3588. <https://doi.org/10.1038/sj.onc.1207426>
75. Sablina AA, Chumakov PM, Levine AJ, Kopnin BP (2001) p53 activation in response to microtubule disruption is mediated by integrin-Erk signaling. *Oncogene* 20:899–909. <https://doi.org/10.1038/sj.onc.1204156>
76. Heyen CA, Tagliabracci VS, Zhai L, Roach PJ (2009) Characterization of mouse UDP-glucose pyrophosphatase, a Nudix hydrolase encoded by the Nudt14 gene. *Biochem Biophys Res Commun* 390:1414–1418. <https://doi.org/10.1016/j.bbrc.2009.11.007>
77. Roach PJ, Depaoli-Roach AA, Hurley TD, Tagliabracci VS (2012) Glycogen and its metabolism: some new developments and old themes. *Biochem J* 441:763–787. <https://doi.org/10.1042/BJ20111416>

78. Sun RC, Dukhande VV, Zhou Z et al (2019) Nuclear Glycogenolysis Modulates Histone Acetylation in Human Non-Small Cell Lung Cancers. *Cell Metab* 30:903–916. <https://doi.org/10.1016/j.cmet.2019.08.014>. e7
79. Hancock ML, Meyer RC, Mistry M et al (2019) Insulin Receptor Associates with Promoters Genome-wide and Regulates Gene Expression. *Cell* 177:722. <https://doi.org/10.1016/j.cell.2019.02.030>. 736.e22
80. Liao Y, Wang J, Jaehnig EJ et al (2019) WebGestalt 2019: gene set analysis toolkit with revamped UIs and APIs. *Nucleic Acids Res* 47:W199–W205. <https://doi.org/10.1093/nar/gkz401>

Tables

Table 11 and Table Sx are not available with this version

Figures

Figure 1

Starvation dissociates p53-MDM2 regulatory feedback loop leading to robust nuclear p53 stabilization

(A) Western blot from a time course experiment in HepG2 cells showing p53 and MDM2 protein abundances over 24 hours starvation. Beta-actin (ACTB) as loading control. GM...growth medium, SM...starvation medium.

(B) Western blot showing regulation of p53 and MDM2 after 24 hours growth medium or starvation medium containing pharmacological inhibitor nutlin-3a (10 μ M). Glyceraldehyde 3-phosphate dehydrogenase (GAPDH) as loading control.

(C) RT-qPCR showing fold change of p53, MDM2, p21 (CDKN1A), GADD45a, TIGAR, and SESN1 mRNA expression levels in 6h nutlin-3a treated samples over vehicle (DMSO) controls after preincubation for 24 hours growth medium or starvation medium. Shown are replicate measurements of one representative out of three independent experiments. Data is shown as mean $\pm$ SEM.

(D) Western blot of p53, MDM2, and p21. HepG2 cells were kept in growth medium (left panel) or starvation medium (right panel) for 24 hours before they were treated with nutlin-3a (10 μ M) for the indicated times. Loading controls: cytoplasm GAPDH, nuclei KI67.

(E) Western blot of protein degradation assay showing regulation of p53, MDM2, and p21. HepG2 cells were kept in growth medium (left panel) or starvation medium (right panel) for 24 hours before they were treated with cycloheximide for the indicated times. Loading controls: cytoplasm GAPDH, nuclei KI67.

Figure 2

Proximity-based biotin labeling proteomics reveals specific p53 interactome changes upon starvation

(A) Western blot showing HepG2 p53KO cells overexpressing either p53-miniTurbo or EGFP-miniTurbo fusion protein. Cells were subjected to growth medium or starvation medium (SM) and fractionated into nuclei (nuc) and cytoplasm (cyt). Overexpressed proteins were detected with a V5-tag-specific antibody. GAPDH, Histone3 as loading controls.

(B) Proximity-biotinylation and affinity purification MS: experimental and analysis workflow.

(C) Left: Hierarchical clustering of proteomics data after ANOVA testing (FDR0.05) / z-scoring showing three clusters with p53-dependently enriched signals over EGFP background control. Right: Profile blots showing protein abundances within clusters between biological replicates (n = 3 per group). Top profile: enriched p53 interactors under GM conditions (137 proteins). Middle profile: enriched p53 interactors under both, GM and SM, conditions (37 proteins). Bottom profile: enriched p53 interactors under SM conditions (98 proteins). Total 272 proteins.

(D) Volcano plot showing 272 proteins enriched over EGFP background control. Proteins with significant fold changes between GM and SM conditions are located outside the cut-off curve (red diamonds) (FDR0.01|S0=0.1; Perseus). Significantly changed proteins were subjected to KEGG pathway overrepresentation analysis (E) and proteins from overrepresented pathways (pentose phosphate pathway, biosynthesis of amino acids, glycolysis/gluconeogenesis, carbon metabolism) labelled with protein names and circled in blue. Proteins that reportedly influence p53 stability by modifying ubiquitination are marked in orange.

(E) KEGG pathway overrepresentation analysis (WebGestalt) with proteins significantly changed between GM and SM.

(F) SAINT score analysis of differential p53 interactors GM vs SM. Cut-offs SAINT probability: (SP) ≥ 0.9 ; SP ≥ 0.8 ; SP < 0.8 indicated with grey scaled circles. Relative abundance represented by circle diameter. Blue tones indicate the respective average spectral counts (avgSPC).

Figure 3

Differential peptide mapping reveals serine/threonine-protein kinase PAK2 as upstream regulator directly binding to p53

(A) Bar graph showing iBAQ intensities of PAK2 peptide 51-62 in untreated samples (control) vs biotin treated samples. Biotinylation on K52 was highly increased under GM conditions, while it significantly

declined under SM conditions. Data is shown as mean $\pm$ SEM. Student's t test, * $p < 0.05$.

(B) PAK2 protein sequence with indicated peptides uniquely identified in p53 samples (GM or SM) but absent in EGFP control samples (blue boxes). Highly significant differentially biotinylated peptide between GM and SM conditions (shown in A) (brown box).

(C) ^1H , ^{15}N HSQR NMR spectra of ^{15}N -labelled recombinant p53 DBD (94-312) incubated with increasing concentrations of recombinant PAK2 N-terminal domain (1-212). Insets show clear concentration-dependent chemical shift perturbations. Bottom: intensity ration of p53 DBD residues in bound vs free state.

(D) ^1H , ^{15}N HSQR NMR spectra of ^{15}N -labelled N-terminal PAK2 domain (1-212) incubated with increasing concentrations of p53 DBD (94-312). Insets show clear concentration-dependent chemical shift perturbations.

(E) Western blot showing time course experiment with PAK2 inhibitor FRAX597 (20 nM) over 6 hours of treatment. GAPDH as loading control.

(F) Proposed model of p53 activation mediated by dissociation from PAK2.

Figure 4

Nuclear p53 interactome under prolonged starvation: SORBS1 and UGP2 as novel interactors

(A) Western blot probed with p53 antibody (DO1) showing high molecular weight complexes (HMWC) specific for HepG2 WT cells, while p53KO cells lack p53-dependent staining in the high molecular weight area of the immunoblot. Starvation leads to a p53-dependent increase of HMWC in comparison to growth medium conditions.

(B) Experimental and analysis workflow for rapid immunoprecipitation MS of endogenous proteins (RIME, $n=3$ IPs from independent experiments per group).

(C) Scheme depicting detected p53 peptides in samples precipitated with p53-specific antibody. TAD...transactivation domain, PRD...proline rich domain, DBD...DNA binding domain, HD...hinge domain, OD...oligomerization domain, RD...regulatory domain.

(D) Scatter blot showing comparison of average spectral counts (avgSPC) in GM and SM samples for proteins precipitated with p53-specific antibody (DO1). Enrichment of proteins above $x=y$ line indicates increased p53 interactions in SM vs GM samples. Inset shows proteins with higher avgSPC values.

(E) p53 interaction network enriched under starvation over growth medium from SAINT core analysis (cut off: SAINT probability $SP \geq 0.9$). Encircled are proteins that are known p53 interactors (BioGRID, red) and/or map to the Reactome pathway “Metabolism of carbohydrates” (RSA-HAS-71387, green).

(F) Dot blot showing results of SAINT score analysis of differential nuclear p53 interactors GM vs SM. Cut-offs SAINT probability ($SP \geq 0.95$; $SP \geq 0.8$; $SP < 0.8$) indicated with grey scaled circles. Relative abundance represented by circle diameter. Blue tones indicate the respective average spectral counts (avgSPC).

(G) Western blot showing immunoprecipitation after overexpression of HA-tagged p53 and/or FLAG-tagged UGP2 in HepG2 p53KO cells. Samples were treated with either GM or SM for 24 hours and proteins precipitated with FLAG-beads. p53 co-eluted with UGP2 exclusively under starvation conditions.

(H) Western blot showing immunoprecipitation of overexpressed p53HA in HepG2 p53KO cells with co-eluted SORBS1 protein under growth medium and starvation conditions. Samples were treated with either GM or SM for 24 hours and proteins precipitated with anti-HA-beads.

Figure 5

Transcriptome analysis reveals a p53- and starvation-specific differentially expressed genes

(A) RNAseq experimental and analysis workflow in HepG2 p53KO cells re-expressing either FLAG-p53 or empty vector (EV) control (n=3 per group).

(B) Western blot showing FLAG-p53 re-expression and target gene expression (p21) in HepG2 p53KO cells treated with either growth medium or starvation medium. b-Actin as loading control.

(C) Heat map showing 88 significantly differentially expressed gene transcripts (DESeq2, Wald-test with Benjamini-Hochberg correction, $p < 0.05$) enriched in p53 overexpressing cells and sensitive to starvation. Venn diagram showing overlap with p53 target gene meta analysis (Fischer et al., 2017) indicating 48 novel, starvation-specific p53 target genes.

(D) Violin blot showing fold change ($\log_2$) distribution of 88 gene transcripts between treatment groups. EV_GM set to 1.

(E) Exemplary expression profiles of known p53 target genes found in the RNAseq data set. EV_GM set to 1.

(F) Density plot of p53 binding sites centered on transcription start sites (TSS) of gene sets of known, novel, and p53-unregulated (negative control) transcripts.

(G) RT-qPCR validation of novel p53 target genes found in the RNAseq data set. Data is normalized to PPIA. Mean values $\pm$ SEM are shown and two-way ANOVA, Tukey's multiple comparisons test was performed. **** $p < 0.001$, *** $p < 0.005$, * $p < 0.05$.

Figure 6

Model of p53 signaling and interactomes upon starvation

Model of p53 signaling and interactomes upon starvation suggested by our complementary approaches. Processes investigated are in blue italics. Proteins that are individually investigated are colored. Grey scale proteins are examples derived from affinity purification MS and described in the text. Protein shapes are schematic with no relation to native conformations. Illustration was created with BioRender.com.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementalFigure1to5Galhuberetal.pdf](#)
- [TableS1.xlsx](#)
- [TableS2.xlsx](#)
- [TableS3.xlsx](#)
- [TableS4.xlsx](#)
- [TableS5.xlsx](#)
- [TableS6oligos.xlsx](#)