

The Influence of Rhizosphere Soil Fungal Diversity and Complex Community Structure to Wheat Root Rot Disease

Xuejiang Zhang

Institute of Plant Protection and Soil & Fertilizer, Hubei Academy of Agricultural Sciences

Heyun Wang

Hubei University of Technology, China

Gavin Ash

The Centre for Crop Health, University of Southern Queensland, Toowoomba QLD, Australia

Dazhao Yu

Institute of Plant Protection and Soil & Fertilizer, Hubei Academy of Agricultural Sciences

Hua Wang (✉ wanghua4@163.com)

<https://orcid.org/0000-0003-2552-7063>

Research article

Keywords: Rhizosphere soil, Fungal diversity, Community structure, Wheat root rot disease

Posted Date: January 23rd, 2020

DOI: <https://doi.org/10.21203/rs.2.21748/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at PeerJ on December 13th, 2021. See the published version at <https://doi.org/10.7717/peerj.12601>.

The Influence of Rhizosphere Soil Fungal Diversity and Complex Community Structure to Wheat Root Rot Disease

Xuejiang Zhang¹, Heyun Wang², Gavin Ash³, Dazhao Yu¹, Hua Wang^{1,*}

The full names, email addresses and institutional addresses for all authors

¹Xuejiang Zhang

zhangxuejiang1976@126.com

Institute of Plant Protection and Soil & Fertilizer, Hubei Academy of Agricultural Sciences/Key Laboratory of Integrated Pest Management on Crops in Central China, Ministry of Agriculture. P. R. China/Hubei Key Laboratory of Crop Disease, Insect Pests and Weeds Control, Wuhan 430064, P. R. China.

²Heyun Wang

heyunwang2006@163.com

HuBei University of Technology, Wuhan 430068, P. R. China.

³Gavin Ash

Gavin.Ash@usq.edu.au

The Centre for Crop Health, University of Southern Queensland, Toowoomba QLD 4350, Australia.

¹Dazhao Yu

dazhaoyu1956@sina.com.cn

Institute of Plant Protection and Soil & Fertilizer, Hubei Academy of Agricultural Sciences/Key Laboratory of Integrated Pest Management on Crops in Central China, Ministry of

* Corresponding author: wanghua4@163.com

²Heyun Wang contributed equally to this work.

¹Institute of Plant Protection and Soil & Fertilizer, Hubei Academy of Agricultural Sciences, Nanhu Road 6, Wuhan 430064, China.

Full list of author information is available at the end of the article.

Agriculture. P. R. China/Hubei Key Laboratory of Crop Disease, Insect Pests and Weeds Control, Wuhan 430064, P. R. China.

Corresponding author:

^{1,*}Hua Wang

wanghua4@163.com

Institute of Plant Protection and Soil & Fertilizer, Hubei Academy of Agricultural Sciences/Key Laboratory of Integrated Pest Management on Crops in Central China, Ministry of Agriculture. P. R. China/Hubei Key Laboratory of Crop Disease, Insect Pests and Weeds Control, Wuhan 430064, P. R. China.

Abstract

Background: Wheat root rot disease due to soil-borne fungal pathogens leads to tremendous yield losses worth billions of dollars worldwide every year. It is very important to study the relationship between rhizosphere soil fungal diversity and wheat roots to understand the occurrence and development of wheat root rot disease.

Results: A significant difference in fungal diversity was observed in the rhizosphere soil of healthy and diseased wheat roots in the heading stage, but the trend was the opposite in the filling stage. The abundance of most genera with high richness decreased significantly from the heading to the filling stage in the diseased groups; the richness of approximately one-third of all genera remained unchanged, and only a few low-richness genera, such as *Fusarium* and *Ceratobasidium*, had a very significant increase from the heading to the filling stage. In the healthy groups, the abundance of most genera increased significantly from the heading to the filling stage; the abundance of some genera did not change markedly, or the abundance of very few genera increased significantly. Physical and chemical soil indicators showed that low soil pH and density, increases in ammonium nitrogen, nitrate nitrogen and total nitrogen contributed to the occurrence of wheat root rot disease.

Conclusions: Our results revealed that in the early stages of disease, highly diverse rhizosphere soil fungi and a complex community structure can easily cause wheat root rot disease. The existence of pathogenic fungi is a necessary condition for wheat root rot disease, but the richness of pathogenic fungi is not necessarily important. The increases in ammonium nitrogen, nitrate nitrogen and total nitrogen contributed to the occurrence of wheat root rot disease. Low soil pH and soil density are beneficial to the occurrence of wheat root rot disease.

Keywords: Rhizosphere soil, Fungal diversity, Community structure, Wheat root rot disease

Background

Soil microbial diversity is important to sustainable agriculture because microbes mediate many processes that are essential to the agricultural productivity of soil (1). However, to meet the food demand of an increasing population, intensive agricultural practices and the cultivation of exhaustive crops have destroyed soil structure and ignored the biological potential of roots or rhizospheres to efficiently mobilize and acquire soil nutrients (2-8). As such, soils have very low biological activity, and plants growing in these soils are predisposed to soil-borne pathogens (9-12).

With the destruction of soil structure, the degradation of soil and the increase in soil-borne pathogens, wheat, as one of the three major staple foods in the world, grown in Asia (China), Australia, Europe, North America, and South America, is vulnerable to attack by a complex of root pathogens, which results in tremendous yield losses (13). Annual losses in wheat industries due to soil-borne fungal pathogens amount to over billions of dollars worldwide (14-16). All cultivars of wheat are attacked by several soil-borne fungal pathogens that cause root diseases (15). The primary fungal pathogens include the following: *Fusarium culmorum*, *Fusarium pseudograminearum*, *Gaeumannomyces graminis* var. *Tritici*, *Bipolaris sorokiniana*, and *Alternaria* spp. in *Ascomycota*; *Rhizoctonia*, *Rhizopus oryzae*, *Rhizoctonia solani* and *Penicillium* spp. in *Basidiomycota*; *Pythium* spp. in *Oomycota*; and *Curvularia* spp. in *Deuteromycota* (10, 17). Root diseases are difficult to control because these soil-borne fungi are ubiquitous, the pathogens often occur as a complex (15, 18), and they can easily survive on infected plant debris or form durable chlamydospores in the soil with or in the absence of growing hosts and can outgrow or evade plant defenses (10, 19). There are no resistant varieties among adapted cultivars of wheat (20-21) and no chemical controls, although certain seed treatments can provide some early benefits to seedling health (21-22).

Therefore, it is very important to study the relationship between soil microbial diversity and plants to understand the occurrence and development of crop root rot. Plants depend on the ability of roots to communicate with rhizosphere soil microorganisms through signaling pathways, creating a connection between plants and microorganisms (23-25). The composition of the rhizosphere soil microbiota can negatively or positively influence plant traits such as stress tolerance, health, development, and productivity (26-28). The plant, in turn, cultivates the structural and functional diversity of microbial communities in the rhizosphere soil by adjusting soil pH, releasing secondary metabolites into the rhizosphere soil (25, 29-30), reducing competition for beneficial microbes, and providing an energy source, mostly in the carbon-rich rhizosphere soil (4, 21). Soil-plant-microbial health must remain in equilibrium to maintain sustainable agricultural practices (25, 31-32). Under unfavorable conditions, some fungi can cause plant diseases and sometimes even total loss of crop yields (28).

In addition to agrochemicals (33), fertilization (8, 34), soil types (35-36), tillage (1) and crop rotation (9-10), which can influence rhizosphere soil microorganisms, the structure and function of rhizosphere soil microflora may also be affected by the plant physiology of different plant genotypes (36-37) and may also fluctuate among the vegetation stages of the same plant genotype (36, 38). Plant growth stage influences root physiology and changes the quality and quantity of root exudates; consequently, these changes select for different root-associated microorganisms at different growth stages (39-41). The purpose of this study was to study the diversity of fungi, the variation in community structure and the trends in the microbial species in the rhizosphere soil of healthy and diseased wheat roots at different heading and filling stages of wheat growth. Additionally, by combining these results with the physical and chemical properties of soil, the possible causes of wheat root rot disease were revealed, providing an important theoretical basis and practice for the improved control of wheat root rot disease.

Results

Fungal diversity in the rhizosphere soil of wheat root

According to the diversity index, the community richness and diversity of the diseased groups were higher than those of the healthy groups in the heading stage, and there was a significant difference in community diversity between the diseased group and the healthy group (Shannon index: $P < 0.01$) (Supplemental table 1). The high community richness and diversity created conditions for subsequent diseases. However, the community richness and diversity of the disease group and the healthy group in the filling stage were higher than those of the disease group and the healthy group in the heading stage, and there was a significant difference between the disease group and the healthy group in the heading stage (Sobs index: $P < 0.05$; Shannon index: $P < 0.05$). The community richness and diversity of the healthy group in the filling stage were higher than those of the diseased group in the filling stage.

Community structure of fungi in the rhizosphere soil

There were 1393 OTUs distributed among 7 phyla, 346 genera and 549 species. The seven phyla were *Ascomycota*, *Basidiomycota*, *Zygomycota*, *Chytridiomycota*, *Glomeromycota*, *Blastocladiomycota* and unclassified fungi (Fig 2 A). Among these phyla, *Ascomycota* was the most abundant. The richness values were 65.48% and 67.61% in healthy groups 4 and 5 (H4 and H5, respectively), and 72.57% and 57.35% in diseased groups 4 and 5 (D4 and D5, respectively), respectively, but there was no significant difference among the populations. The other three phyla showed very interesting results. *Basidiomycota* had the highest abundance in population H4 in the heading stage (25.3%). However, in the D4, D5 and H5 populations, the abundance was significantly decreased to 12.36%, 11.16% and 14.7%, respectively. Significant differences were

found between populations H4 and D4 and populations D5 and H5 ($P < 0.01$). The abundance of *Zygomycota* in populations H4 and D4 was low in the heading stage (3.16% and 2.44%, respectively), but suddenly increased to 9.87% (H5) and 7.42% (D5) in the filling stage. However, the trends in the abundance of *Chytridiomycota* and *Zygomycota* were almost the opposite. The abundance of *Chytridiomycota* in the heading stage groups was 4.52% (H4) and 7.18% (D4), and there was a significant difference between these two groups. The abundance of *Chytridiomycota* in population D4, a diseased group, was much higher than that in the healthy groups. By May, the abundance of *Chytridiomycota* in population H5 decreased sharply to 1.37%, while that of population D5 decreased to only 4.12%.

Excluding unclassified fungi, all fungi with richness greater than 1% in a single group were counted (Table 2). There are 30 genera in the four groups belonging to *Ascomycota*, *Basidiomycota*, *Chytridiomycota* and *Zygomycota*. Among these genera, *Chytridiomycota* and *Zygomycota* had only one genus, *Olpidiaster* and *Portierella*, respectively. *Basidiomycota* had seven genera, and *Ascomycota* had the largest distribution, with 21 genera (Fig 2 B and C, Supplemental table 2). In the heading stage, the abundance of the most highly abundant genera in population D4 was significantly decreased in population D5 at the filling stage; however, the abundance of *Mortierella* significantly increased, the abundance of one-third of the genera remained almost unchanged, and the abundance of a few low abundance genera, such as *Fusarium* and *Ceratobasidium*, significantly increased in the diseased population over time. Seven genera with high richness were found in the H4 population at the heading stage, and the richness of these genera was significantly increased in the H5 population at the grain filling stage; among these genera, the variation in the richness of *Microdochium*, *Mycosphaerella*, *Scytalidium*, *Acremonium* and *Olpidiaster* from the H4 population to the H5 population was the opposite of the change in abundance from the D4 population to the D5 population. The richness of 13 genera did not markedly change, and the richness of 4 low-richness genera was significantly increased at the grain filling stage compared with the heading stage. In general, the following six genera showed no change in richness from the heading stage to the grain filling stage, between the H5 and D5 populations and the H4 and D4 populations, respectively, *Alternaria*, *Sarocladium*, *Gibberella*, *Cladosporium*, *Neosetophoma* and *Cystofilobasidium*. The genera with significant or extremely significant differences in richness among the groups were *Alternaria*, *Mortierella*, *Cryptococcus*, *Apodus*, *Epicoccum*, *Scytalidium* and *Chaetomium*.

A Venn diagram of genera with richness greater than 1% was constructed (Fig 2 D). There were 153 shared genera, 35 endemic genera that appeared most frequently in the healthy group in the filling stage and 7 and 10 genera in the heading stage and filling stage, respectively. The results showed that the root exudates of diseased plants specifically promoted the growth of microbial fungi, while the rhizosphere of healthy plants was more suitable for the growth of various fungi. An increased diversity of soil microorganisms promoted healthy plant growth.

PCoA of populations

The PCoA of fungal community structure in different samples based on OTU revealed that the populations could be divided into two groups: the healthy group and the diseased group (Figure 3 A and B). There was no overlap between the two groups, which indicated that the samples were consistent with the expectation and that the microbial community structures of the diseased group and the healthy group were quite different. The diseased group at the heading stage was significantly separated from the diseased group at the filling stage, and the microbial community structure was quite different. The healthy group at the heading stage and the healthy group at the filling stage were also clearly separated, and the microbial community structure was quite different. In addition, according to the flora classification, all rhizosphere fungi were divided into two types (Fig 3 C): type one included the diseased group, and type two included the healthy group. Only sample H5-5 was assigned to type one.

Effects of physical and chemical soil properties on rhizosphere fungi

The main physical and chemical properties of soil measured were total phosphorus (TP), ammonium nitrogen (NH_4), nitrate nitrogen (NO_3), total nitrogen (TN), pH, soil density (SD), total carbon (TC) and the soil dry-humidity ratio (DHR) (Table 3). TP and TC decreased and NH_4 increased from the heading stage to the filling stage. Moreover, the NO_3 and TN of the diseased group were higher than those of the healthy group, and the pH value and SD of the diseased group were lower than those of the healthy group. Heatmap cluster analysis based on Spearman correlation coefficients for the 30 genera with the most richness and these physical and chemical indicators was performed (Fig 4). The graph shows that pH and SD have the greatest influence on microbial abundance, and the influence of these factors is in the same direction. One-third of the genera were positively or significantly positively correlated with both pH and SD, and one-third were negatively or significantly negatively correlated with both pH and SD. The effects of NO_3 and TN on microbial richness were similar, but the impact of these factors on microbial species richness was almost the opposite of that of pH and SD. Soil DHR and TP had moderate effects on microbial richness. The physical and chemical properties of soil with the least influence on microbial abundance were NH_4 and TC.

Discussion

Microbial diversity in soil is an important factor that determines soil health and is considered one of the main contributors to soil suppressiveness (28, 42). The rhizosphere is one of the most complex environments; rhizospheres are influenced by plant roots and are an active microhabitat where plant roots and microbes interact (28, 43-44). Throughout the heading and filling stages of wheat, the fungi in the rhizosphere soil showed high diversity. The diversity of soil fungi at the

grain filling stage was significantly higher than that at the heading stage. It is already well known that most crops can significantly benefit from establishing associations with diverse soil microbes (25). Plants stimulate or inhibit the growth of specific rhizosphere microorganisms by releasing secondary metabolites into the rhizosphere (25, 29-30). In the rhizosphere soil of wheat root rot disease, the interaction between wheat and microorganisms was intense, and the diversity and richness of the fungal community in the rhizosphere soil of the diseased group were significantly higher than that of the healthy group, creating conditions for the occurrence of wheat root rot. However, at the filling stage, wheat roots were clearly diseased due to infection by pathogenic fungi. During this process, the metabolites secreted by wheat roots tend to be monotonous. As a result, the fungi in the rhizosphere of diseased plants showed lower community diversity and richness than those in the rhizosphere of healthy plants. The interaction between rhizosphere fungi and wheat roots in the healthy group reached an optimal balance during the filling stage and then became mutualistic, which was favorable for the healthy growth of the fungi and wheat. Therefore, in the filling stage, the diversity and richness of fungi in the rhizosphere were higher in the healthy group than in the diseased group.

The richness change analysis at the phylum level revealed that the phylum with the highest richness was *Ascomycota*, which consisted of more than 60% of all healthy and diseased groups, and there was no significant difference among groups. However, the main pathogenic fungi that cause wheat root rot disease, such as *F. culmorum*, *F. pseudograminearum*, *G. graminis* var. *Tritici*, *B. sorokiniana* and *Alternaria* spp., are in this phylum. There are also some fungi that cause wheat root rot in the phylum *Basidiomycota*, such as *Rhizoctonia*, *R. oryzae*, *R. solani* and *Penicillium* spp. This phylum showed high richness in the healthy group at the heading stage (25.3%), but the richness of this phylum was significantly decreased in the other groups (<14%). Analysis of the abundance of these two phyla shows that the impact of the fungi that cause wheat root rot disease is not due to phylum-level abundance. The richness of two other phyla, *Zygomycota* and *Chytridiomycota*, which contain almost no root rot fungi, showed notable differences between the heading and filling stages in the healthy group but showed no significant differences between these stages in the diseased groups. This change in abundance may contribute to the healthy growth of wheat roots or may be a cofactor in the occurrence of wheat root rot disease.

Relative to the changes at the phylum level, levels of richness at the genera level varied greatly. The abundance of most genera with high richness decreased significantly from the heading stage to the filling stage in the diseased groups; the richness of approximately one-third of all genera remained unchanged, and only very few low-richness genera, such as *Fusarium* and *Ceratobasidium*, had a very significant increase in richness over time. In the healthy group, the abundance of most genera increased significantly from the heading stage to the filling stage, except for some genera whose abundance did not markedly change or very few genera whose abundance increased significantly. This result also shows that the interactions between wheat roots

and rhizosphere fungi must achieve a balance. If this balance is lost, the wheat roots will become diseased. From the analysis of the endemic genera in each group, we also found that there were 35 endemic genera in the healthy group, which was many more than the approximately 10 endemic genera in the diseased group, indicating that the healthy growth of wheat roots can promote plant growth and suppress disease through various activities that prevent infection by pathogens. Therefore, the rhizosphere microorganisms show improved growth, and the microbial diversity and community richness are also significantly increased in the healthy plants compared with the diseased plants.

Fungi and fungus-like organisms form one of the most diverse groups of eukaryotes and represent an essential functional component of soil microbial communities (28, 45). Under unfavorable conditions, some fungi can cause plant diseases and sometimes even the total loss of crop yields. In many instances, these diseases are caused by a complex of fungal species (28). Only *Alternaria* and *Fusarium*, genera with richness values greater than one percent, have been documented to cause wheat root rot, but the abundance of these two genera is not high. Although the abundance of *Alternaria* in the diseased groups was significantly higher than that in the healthy groups, there was no significant change between the heading stage and filling stage. The abundance of *Fusarium* in the filling stage was significantly higher than that in the heading stage. Considering the abundance of these two pathogens, the richness level of a fungus does not indicate whether it can lead to wheat root rot. In addition, other pathogenic fungal genera mentioned in the literature were determined to have less than one percent abundance. Therefore, we can speculate that as long as these fungal genera exist in the rhizosphere, they can lead to wheat root rot disease, regardless of the abundance. However, whether a fungus can induce root rot disease depends on the result of the interaction between wheat root and rhizosphere microorganisms, which is also closely related to seasonal climatic conditions.

Differences in the fungal community structure among groups can clearly demonstrate the heterogeneity of each group, thus showing why there are so many differences in the rhizosphere fungi of the diseased groups and the healthy groups. PCoA clustering analysis clearly clustered the four groups into two groups: the healthy group and the diseased group. However, the healthy group and the diseased group were each clearly separated into the heading stage and grouting stage. In other words, the healthy group and the diseased group had significant differences in fungal community structure, and the difference in the fungal community structure of the heading stage and the grouting stage was significant. In addition, all rhizosphere fungi were divided into two types according to flora classification: one type included only the healthy group, and the other type included only the diseased group. When wheat roots were attacked by microorganisms and developed root disease, the structure of the rhizosphere fungal community was markedly changed. To differentiate between the fungi located very close to the epidermis in the root zone and to protect against the invasion of heterogeneous microbes, plants continuously secrete signaling

molecules, which allows for the development of pathogenic, associative, symbiotic, or naturalistic relationships between microbes and the plant (25, 46).

At the heading stage, the root is slightly diseased, and the interaction between wheat roots and rhizosphere fungi is the most intense, resulting in marked heterogeneity of the soil environment and inducing a high level of rhizosphere fungal diversity and a complex community structure. However, at the filling stage, the wheat roots are completely diseased. At this time, the soil environment is stable, and the material secreted by wheat roots is relatively limited, which leads to a decrease in the microbial diversity of the rhizosphere and a relatively simple community structure. Soil-plant-microbial health must be maintained at an equilibrium to maintain sustainable agricultural practices (25, 31-32). At the heading stage, the antagonistic interaction between healthy wheat roots and rhizosphere fungi reaches a balance and promotes the healthy growth of wheat and fungi. By the filling stage, the ecological environment, such as temperature, humidity and other factors, is improved, the rhizosphere microbial diversity naturally significantly increases, and the community structure becomes more complex.

Biotic and abiotic factors are assumed to influence the structural and functional diversity of the microbial communities in the rhizosphere (47-48). Site properties, including soil type, climatic conditions and type of agricultural management, have been shown to strongly influence the relative composition of rhizosphere microbial communities (48-54). Changes in the physical and chemical properties of soil also have a significant impact on fungal diversity and community structure in crop rhizospheres. Soil physical and chemical properties are determined not only by the nature of the soil itself but also by the physical and chemical properties of the soil after the interaction between crop roots and rhizosphere fungi. The physical and chemical soil properties we measured were determined in soil collected around the rhizosphere of wheat. Organisms that are present in the rhizosphere microbiota can have profound effects on the growth, nutrition and health of plants in agroecosystems (25, 55-57). TP and TC decreased from the heading stage to the filling stage, indicating that the later the growth period was, the less TP and TC was needed; however, these metrics seemed unrelated to the occurrence of diseases. In addition, the levels of NH_4 , NO_3 and TN in the diseased group were higher than those in the healthy group at the heading and filling stages, indicating that their increase contributed to the occurrence of wheat root rot disease. In contrast, the pH and SD of the soils around diseased plants were lower than those of the soils around healthy plants at the heading and filling stages. This finding indicated that relatively low pH and SD values were beneficial to the occurrence of wheat root rot disease. The heatmap cluster analysis of the physical and chemical soil indicators and the 30 most abundant genera also showed that soil pH and SD affected fungal abundance and diversity in the same direction.

Conclusions

In the rhizosphere, many plant-microbial interactions occur that mediate soil processes (25). The occurrence of crop root rot disease is closely related to the interaction between rhizosphere microorganisms and crop roots, as well as the physical and chemical properties of soil. At present, there are approximately 10 types of fungi that can cause wheat root rot disease alone or in a complex, according to the literature. By studying the diversity of fungi and the community structure of the rhizospheres of healthy and diseased wheat at different growth stages, the heading and filling stages, it was revealed that in the early stages of illness, the high diversity of rhizosphere fungi and a complex community structure can easily cause wheat root rot disease. Additionally, the existence of pathogenic fungi is a necessary condition for wheat root rot disease, but the richness of pathogenic fungi is not necessarily important. Based on the physical and chemical properties of the soil, an increase in NH_4 , NO_3 and TN contributes to the occurrence of wheat root rot disease. Soil pH and SD had the greatest influence on the abundance and diversity of rhizosphere fungi, and the influence was in the same direction; low soil pH and SD are beneficial to the occurrence of wheat root rot disease.

Methods

Rhizosphere soil sampling

From 2015 to 2016, we conducted a continuous follow-up survey on land planted with wheat at Xiangyang Original Farm in Hubei Province, China, and found that the wheat root rot disease at this farm was serious (Fig 1 A). In this study, the rhizosphere soil samples of healthy and diseased wheat roots were collected in early April (heading stage) and early May (filling stage) in 2017 (Fig 1 B). The sampling method consisted of first investigating the occurrence of wheat root rot, selecting weak seedlings and sampling the brown parts of the root as the diseased wheat rhizosphere. Areas with diseased wheat and healthy wheat (the diameter of each area was not more than 10 cm) were marked by inserting cards at fixed points at least 10 meters apart. Then, plants were sampled at 20 points, and 5 wheat plants were sampled at each point. The roots of the whole plant and the soil on the roots were collected. The majority of soil on the roots was shaken off, and samples were collected for the determination of physical and chemical soil properties. The roots of five wheat plants were placed in a sealed pocket and quickly stored in a dry-ice box for dry-ice preservation. After all the samples were collected, they were immediately brought to the laboratory and stored in a cryogenic refrigerator for future use. In May, the samples were collected from the areas labeled in April. The ears of diseased wheat in this area had a clear white color, and these plants had markedly brown, black or yellow roots. Five soil samples from the diseased

wheat root and five soil samples from the healthy wheat root were collected in May. The sampling method was the same as that for the previous collection. According to the cards inserted in the soil in April, the 20 samples collected in April, which were identical to the samples collected in May, were removed from the cryogenic refrigerator to analyze the diversity of fungi in the wheat rhizosphere soil.

Extraction and PCR amplification of total genomic DNA from rhizosphere soil fungi

The soil attached to the root was brushed off, a 0.1-g soil sample was accurately weighed, and the total genomic DNA from all samples was extracted according to the instructions provided with an E.Z.N.A.® Stool DNA Kit (Omega Biotek, Norcross, GA, U.S.). Total DNA was detected by 1% agarose gel electrophoresis, and the purity and concentration of DNA were determined by spectrophotometry. All DNA samples were stored in a refrigerator at -20 °C. Fungal diversity was determined by amplifying the ITS1 region using the ITS1F and ITS2R primer sets for fungi. The primer sequences were ITS1F 5'- CTTGGTCATTTAGAGGAAGTAA-3' and ITS2R 5'- GCTGCGTTCTTCATCGATGC-3', and the amplification conditions were predenaturation at 95°C for 5 min, 27 cycles of 95°C for 30 s, 55°C for 30 s, and 72°C for 45 s, and elongation at 72°C for 5 min. Three replicates of the PCR were performed, and a 20-μL reaction system (4 μL of 5× FastPfu buffer solution, 2 μL of 2.5 mM dNTPs, 0.8 μL of primer (5 μM), 0.4 μL of FastPfu polymerase and 10 ng of fungal total genomic DNA) was used.

Illumina HiSeq2500 sequencing

The PCR amplification product was recovered by 2% agarose gel electrophoresis followed by the use of an AxyPrep DNA gel Recovery Kit (Axygen Biosciences, Union City, CA, U.S.), and the quantitative analysis was carried out using a QuantiFluor®-ST (Promega, U.S.). Two-terminal sequencing was carried out using a HiSeq2500 platform after equally mixing the purified products. The original read data were deposited in the NCBI Sequence Read Archive (Accession number: SUB5829339).

Data processing

The sequencing depth was more than 30,000 original reads per library. Raw fastq files were demultiplexed, quality-filtered by Trimmomatic and merged by FLASH with the following criteria: (i) The reads were truncated at any site receiving an average quality score <20 over a 50 bp sliding window; (ii) Primers were exactly matched allowing 2 nucleotide mismatching, and reads containing ambiguous bases were removed; (iii) Sequences whose overlap longer than 10 bp were merged according to their overlap sequence.

OTU and species community analysis

Operational taxonomic units (OTUs) were clustered with 97% similarity cutoff using UPARSE (version 7.1 <http://drive5.com/uparse/>) and chimeric sequences were identified and removed using UCHIME. The taxonomy of each gene sequence was analyzed by RDP Classifier algorithm (<http://rdp.cme.msu.edu/>) against the Unite (Release 7.0 <http://unite.ut.ee/index.php>) database using confidence threshold of 70%. Finally, the effective tag data, low-frequency tag data, annotated tag data, and OTU data obtained from each sample were counted by a script. Additionally, we used R software to calculate the annotation ratio of OTUs and each taxonomic level and the relative abundance of the species in each taxonomic level.

Based on the above valid OTU data, the following evolutionary analysis was carried out: a) evolutionary relationships and relative abundance information of species systems based on OTU data in samples were determined, and species annotation results for a single sample were visualized using KRONA software (<http://sourceforge.net/projects/krona>); b) related genera were selected, a phylogenetic tree for the OTUs of these genera was constructed (QIIME software package: `make_phylogeny.py` (http://qiime.org/scripts/make_phylogeny.html)), and the systematic evolutionary relationship was displayed by combining the relative abundance of OTUs and the reliability of annotation using a Perl script; and c) local Perl scripts were used to select the dedicated OTUs for intrasample and intersample phylogenetic analysis and to compare relative abundance.

To analyze the community structure of species, relative abundance thermograms were plotted at the OTU level and the genera level by R software. Cluster analysis and principal coordinate analysis (PCoA) were also carried out to compare samples.

Availability of data and materials

Datasets generated/analyzed during this study is available in the NCBI Sequence Read Archive (Accession number: SUB5829339)

Abbreviations

OTU: Operational taxonomic unit

PCoA: principal coordinate analysis

H: healthy population

D: diseased population

TP: total phosphorus

NH₄: ammonium nitrogen

NO₃: nitrate nitrogen

TN: total nitrogen

SD: soil density

TC: total carbon

DHR: soil dry-humidity ratio

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

FUNDING

This work was financially supported by the National Special Fund for Agro-scientific Research in the Public Interest of China (201503112-8), the National Key Research and Development Program of China (2017YFD0200605), and the Key Research and Development Project (2017YFD0201600).

AUTHOR CONTRIBUTIONS

W. H., and Y.D.Z. designed research; Z.X.J., and W.H. performed research; Z.X.J., W.H.Y., and A.G. analyzed data; and Z.X.J., W.H., and W.H.Y. wrote the manuscript with input from all authors.

ACKNOWLEDGEMENTS

We thank L.J. Yang for insightful discussions; M.F. Xue and S.J. Gong for critical comments on the manuscript and J.Z. Ruan for land planted with wheat.

Authors' information

¹Institute of Plant Protection and Soil & Fertilizer, Hubei Academy of Agricultural Sciences/Key Laboratory of Integrated Pest Management on Crops in Central China, Ministry of Agriculture, P. R. China/Hubei Key Laboratory of Crop Disease, Insect Pests and Weeds Control, Wuhan 430064, P. R. China; ²HuBei University of Technology, Wuhan 430068, P. R. China; ³The Centre for Crop Health, University of Southern Queensland, Toowoomba QLD 4350, Australia.

References

1. Lupwayi NZ, Rice WA, Clayton GW. Soil microbial diversity and community structure under wheat as influenced by tillage and crop rotation. *Soil Biol Biochem.* 1998; 30(13):1733-1741.
2. Parkinson D, Coleman DC. Microbial communities, activity and biomass. *Agr Ecosyst Environ.* 1981; 34:3-33.
3. Kumar A, Kumar A, Pratush A. Molecular diversity and functional variability of environmental isolates of *Bacillus* species. *SpringerPlus.* 2014; 3:2–11.
4. Rashida MI, Mujawara LH, Shahzade T, Almeelbi TI, Iqbal MI, Oves M. Bacteria and fungi can contribute to nutrients bioavailability and aggregate formation in degraded soils. *Microbiol Res.* 2016; 183: 26–41.
5. Schreiner RP, Bethlenfalvay GJ. *Mycorrhizae*, biocides, and biocontrol. 4. Response of a mixed culture of arbuscular *mycorrhizal* fungi and host plant to three fungicides. *Biol Fert Soils.* 1996; 23:189-95.
6. Assaf TA, Turk MA, Ameen KM. Impact of olive pomace wastes and fungicide treatment on indigenous arbuscular mycorrhizal fungi associated with chickpea (*Cicerarietinum* L.) under field conditions. *Aust J Crop Sci.* 2009; 3:6-12.
7. Taheri AE, Hamel C, Gan YT. Pyrosequencing reveals the impact of foliar fungicide application to chickpea on root fungal communities of durum wheat in subsequent year. *Fungal Ecol.* 2015; 15:73-81.
8. Ai C, Liang GQ, Sun JW, Wang XB, He P, Zhou W, He XH. Reduced dependence of rhizosphere microbiome on plant-derived carbon in 32-year long-term inorganic and organic fertilized soils. *Soil Biol Biochem.* 2015; 80:70-8.
9. Kirkegaard J, Christen O, Krupinsky J, Layzell D. Break crop benefits in temperate wheat production. *Field Crops Res.* 2008; 107:185–195.
10. Wintera M, Mol FD, Tiedemann AV. Cropping systems with maize and oilseed rape for energy production may reduce the risk of stem base diseases in wheat. *Field Crops Res.* 2014; 156:249–57.
11. Sivasithamparam K. Ecology of root-infecting pathogenic fungi in Mediterranean environments. *Adv Plant Pathol.* 1993; 10:245-79.
12. Almasudy AM, You MP, Barbetti MJ. Influence of fungicidal seed treatments and soil type on severity of root disease caused by *Rhizoctonia solani* AG-8 on wheat. *Crop Prot.* 2015; 75:40-5.

13. Duffy B. Combination of *penycuron* and *pseudomonas yuorescens* strain 2-79 for integrated control of *rhizoctonia* root rot and take-all of spring wheat. *Crop Prot.* 200; 19:21-5.
14. Paulitz TC, Smiley RW, Cook RJ. Insights into the prevalence and management of soilborne cereal pathogens under direct seeding in the Pacific Northwest, USA. *Can J Plant Pathol.* 2002; 24:416–428.
15. Mavrodi OV, Walter N, Elateek S, Taylor CG, Ubara PAO. Suppression of *Rhizoctonia* and *Pythium* root rot of wheat by new strains of *Pseudomonas*. *Biol Ccontrol.* 2012; 62:93-102.
16. Okubara PA, Dickman MB, Blechl AE. Molecular and genetic aspects of controlling the soilborne necrotrophic pathogens *Rhizoctonia* and *Pythium*. *Plant Sci.* 2014; 228:61–70.
17. Mielke H., Studien zum Befall des Weizens mit *Gaeumannomyces graminis* (Sacc.) von Arx et Olivier var. *tritici* Walker unter Berücksichtigung der Sorten- und Artenanfälligkeit sowie der Bekämpfung des Erregers. *Mitteilungen aus der Biologischen Bundesanstalt für Land- und Forstwirtschaft, Berlin-Dahlem.* Heft pp. 1998; 359.
18. Paulitz TC, Adams K. Composition and distribution of *Pythium* communities in wheat fields in eastern Washington state. *Phytopathology.* 2003; 93:867–873.
19. Smith JD, Kidwell K, Evans M, Cook R, Smiley RW. Assessment of spring wheat genotypes for disease reaction to *Rhizoctonia solani* AG-8 in controlled environment and direct-seeded field evaluations. *Crop Sci.* 2003; 43:694-700.
20. Smith JD, Kidwell KK, Evans MA, Cook RJ, Smiley RW. Evaluation of spring cereal grains and wild *Triticum* relatives for resistance to *Rhizoctonia solani* AG 8. *Crop Sci.* 2002; 43:701–709.
21. Davis RA, Huggins D, Cook RJ, Paulitz TC. Can placement of seed away from relic stubble limit *Rhizoctonia* root rot in direct-seeded wheat? *Soil Till Res.* 2008; 101:37–43.
22. Paulitz TC, Scott RB. Effect of Seed Treatments for Control of *Rhizoctonia* Root Rot in Spring Wheat, 2005. *Fungicide and Nematicide Tests.* 2006; 61:ST014.
23. Meena VS, Maurya BR, Bohra JS, Verma R, Meena MD. Effect of concentrate manure and nutrient levels on enzymatic activities and microbial population under submerged rice in alluvium soil of *Varanasi*. *Crop Res.* 2013; 45 (1,2 and 3):6–12.
24. Li M, Ahammed GJ, Li C, Bao X, Yu J, Huang C, Yin H, Zhou J. *Brassinosteroid ameliorates* zinc oxide nanoparticles-induced oxidative stress by improving antioxidant potential and redox homeostasis in tomato seedling. *Front Plant Sci.* 2016; 7:615.
25. Kumar A, Maurya BR, Raghuwanshi R, Meena VS, Islam MT. Co-inoculation with *Enterobacter* and *Rhizobacteria* on yield and nutrient uptake by wheat (*Triticum aestivum* L.) in the alluvial soil under the Indo-Gangetic Plain of India. *J Plant Growth Regul.* 2017; <http://dx.doi.org/10.1007/s00344-016-9663-5>
26. Kristin A, Miranda H. The root microbiota—a fingerprint in the soil? *Plant Soil.* 2013; 370:671-686.
27. Lakshmanan V, Selvaraj G, Bais HP. Functional soil microbiome: below ground solutions to

- an aboveground problem. *Plant Physiol.* 2014; 166:689-700.
28. Miao CP, Mi QL, Qiao XG, Zheng YK, Chen YW, Xu LH, Guan HL, Zhao LX. Rhizospheric fungi of *Panax notoginseng*: diversity and antagonism to host phytopathogens. *J Ginseng Res.* 2016; 40:127-134.
29. Chakraborty U, Swarnendu R, Pratim CA, Pannalal D, Bishwanath C. Plant growth promotion and amelioration of salinity stress in crop plants by a salt-tolerant bacterium. *Recent Res Sci Technol.* 2011; 3 (11):61–70.
30. Meena SK, Rakshit A, Meena VS. Effect of seed bio-priming and N doses under varied soil type on nitrogen use efficiency (NUE) of wheat (*Triticumaestivum* L.) under greenhouse conditions. *Biocata Agric Biotechnol.* 2016; 6:68–75.
31. Narula S, Anand RC, Dudeja SS. Beneficial traits of endophytic bacteria from field pea nodules and plant growth promotion of field pea. *J Food Legumes.* 2013; 26:73–79.
32. Ramirez-Bahena MH, Tejedor C, Martín I, Velaquíez E, Peix A. *Enterobacter medicaginis* gen. nov. sp. nov., isolated from alfalfa nodules in anacidic soil. *Int J Syst Evol Microbiol.* 2013;63:1760–1765.
33. Handiseni M, Brow J, Zemetra R, Mazzol M. Effect of *Brassicaceae* seed meals with different glucosinolate profiles on *Rhizoctonia* root rot in wheat. *Crop Prot.* 2013; 48: 1-55.
34. Phillips R, Fahey T. Fertilization effects on fine root biomass, rhizosphere microbes and respiratory fluxes in hardwood forest soils. *New Phytol.* 2007; 176:655-664.
35. Buyer JS, Roberts DP, Russek-Cohen E. Microbial community structure and function in the spermosphere as affected by soil and seed type. *Can J Microbiol.* 1999; 45:138–144.
36. Rasche F, Hödl V, Poll C, Kandeler E, Gerzabek MH, van Elsas JD, Sessitsch A. Rhizosphere bacteria affected by transgenic potatoes with antibacterial activities compared with the effects soil, wild-type potatoes, vegetation stage and pathogen exposure. *FEMS Microbiol Ecol.* 2006; 56:219–235.
37. Söderberg KH, Olsson PA, Bååth E. Structure and activity of the bacterial community in the rhizosphere of different plant species and the effect of arbuscular mycorrhizal colonisation. *FEMS Microbiol Ecol.* 2002; 40:223–231.
38. Gyamfi S, Pfeifer U, Stierschneider M, Sessitsch A. Effects of transgenic glufosinate-tolerant oilseed rape (*Brassica napus*) and the associated herbicide application on eubacterial and *pseudomonas* communities in the rhizosphere. *FEMS Microbiol Ecol.* 2002; 41:181–190.
39. Dunfield KE, Germida JJ. Seasonal changes in the rhizosphere microbial communities associated with field-grown genetically modified canola (*Brassica napus*). *Appl Environ Microbiol.* 2003; 69:7310-7318.
40. Houlden A, Timms-Wilson TM, Day MJ, Bailey MJ. Influence of plant developmental stage on microbial community structure and activity in the rhizosphere of three field crops. *FEMS Microbiol Ecol* 2008; 65:193-201.
41. Li XZ, Rui JP, Mao YJ, Yannarell A, Mackie R. Dynamics of the bacterial community

- structure in the rhizosphere of a maize cultivar. *Soil Boil Biochem.* 2014; 68: 392-401.
42. Xu L, Ravnskov S, Larsen J, Nilsson RH, Nicolaisen M. Soil fungal community structure along a soil health gradient in pea fields examined using deep amplicon sequencing. *Soil Boil Biochem.* 2012; 46:26-32.
43. Mendes R, Garbeva P, Raaijmakers JM. The rhizosphere microbiome: significance of plant beneficial, plant pathogenic, and human pathogenic microorganisms. *FEMS Microbiol Rev.* 2013; 37:634-663.
44. Singh AK, Singh M, Dubey SK. Rhizospheric fungal community structure of a Bt brinjal and a near isogenic variety. *J Appl Microbiol.* 2014; 117:750-765.
45. Buée M, Reich M, Murat C, Morin E, Nilsson RH, Uroz S, Martin F. 454 Pyrosequencing analyses of forest soils reveal an unexpectedly high fungal diversity. *New Phytol.* 2009; 184:449-456.
46. Hayat R, Ali S, Amara U, Khalid R, Ahmed I. Soil beneficial bacteria and their role in plant growth promotion: a review. *Ann Microbiol.* 2010; 60:579–598.
47. Berg G, Smalla K. Plant species and soil type cooperatively shape the structure and function of microbial communities in the rhizosphere. *FEMS Microbiol Ecol.* 2009; 68:1–13.
48. Weinert N, Piceno Y, Ding GC, Meincke R, Heuer H, Berg G. PhyloChip hybridization uncovered an enormous bacterial diversity in the rhizosphere of different potato cultivars: many common and few cultivar-dependent taxa. *FEMS Microbiol Ecol.* 2011; 75:497–506.
49. Heuer H, Kroppenstedt RM, Lottmann J, Berg G, Smalla K. Effects of T4 lysozyme release from transgenic potato roots on bacterial rhizosphere communities are negligible relative to natural factors. *Appl Environ Microbiol.* 2002; 68(3):1325-1335.
50. Kowalchuk GA, Buma DS, de Boer W, Klinkhamer PGL, van Veen JA. Effects of above-ground plant species composition and diversity on the diversity of soil-borne microorganisms. *Antonie van Leeuwenhoek.* 2002; 81:509–520.
51. Berg G, Opelt K, Zachow C, Lottmann J, Götz M, Costa R, Smalla K. The rhizosphere effect on bacteria antagonistic towards the pathogenic fungus *Verticillium* differs depending on plant species and site. *FEMS Microbiol Ecol.* 2006; 56:250–261.
52. Costa R, Götz M, Mrotzek N, Lottmann J, Berg G, Smalla K. Effects of site and plant species on rhizosphere community structure as revealed by molecular analysis of microbial guilds. *FEMS Microbiol Ecol.* 2006; 56:236–249.
53. Bremer C, Braker G, Matthies D, Reuter A, Engels C, Conrad R. Impact of plant functional group, plant species, and sampling time on the composition of nirK-type denitrifier communities in soil. *Appl Environ Microb.* 2007; 73:6876–6884.
54. Weinert N, Meincke R, Gottwald C, Radl V, Dong X, Schloter M, Berg G, Smalla K. Effects of genetically modified potatoes with increased zeaxanthin content on the abundance and diversity of *rhizobacteria* with in vitro antagonistic activity do not exceed natural variability among cultivars. *Plant Soil.* 2010; 326:437–452.

55. Bonfante P, Anca IA. Plants, mycorrhizal fungi, and bacteria: a network of interactions. *Annu Rev Microbiol.* 2009; 63:363–383.
56. Mendes R, Kruijt M, De Bruijn I, Dekkers E, van der Voort M, Schneider JHM, Piceno YM, DeSantis TZ, Andersen GL, Bakker PAHM, Raaijmakers JM. Deciphering the rhizosphere microbiome for disease-suppressive bacteria. *Science.* 2011; 332:1097–1100.
57. Meena RS, Meena VS, Meena SK, Verma JP. The needs of healthy soils for a healthy world. *J Cleaner Prod.* 2015; 102:560–561.

Figures



Figure 1

Sampling plots of wheat root. a The occurrence of wheat root rot in this wheat field in 2015 and 2016. b The soil sample collection of wheat root in this wheat field in 2017. Areas with diseased wheat and healthy wheat were marked by inserting white cards at fixed points. The samples, in April and May, were collected from the areas labeled in April.




Figure 2

Comparative analysis of OTUs and genus levels in each community. a Principal coordinate analysis (PCoA) at the OTU level in healthy and diseased groups at the heading stage and filling stage. Distance algorithm is based on bray_curtis. b Sample hierarchical clustering analysis based on OTU level. c Typing analysis of fungi at the genus level.




Figure 4

Hierarchical clustering analysis at the OTU level between physical and chemical soil properties and the 30 most abundant genera.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Tables.pdf](#)