

Transcriptome analysis reveals ethylene-mediated defense responses to *Fusarium oxysporum* f. sp. *cucumerinum* infection in *Cucumis sativus*

Jingping Dong

Yangzhou University

Yuean Wang

Yangzhou University

Qianqian Xian

Yangzhou University

Xuehao Chen (✉ xhchen@yzu.edu.cn)

Jun Xu

Yangzhou University

Research article

Keywords: Cucumber, *Fusarium* wilt, RNA-seq, Exogenous ethylene

Posted Date: January 23rd, 2020

DOI: <https://doi.org/10.21203/rs.2.21765/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on July 16th, 2020. See the published version at <https://doi.org/10.1186/s12870-020-02537-7>.

Abstract

Background: *Fusarium* wilt, caused by *Fusarium oxysporum* f. sp. *cucumerinum* (Foc), is a severe disease affecting cucumber (*Cucumis sativus* L.) production worldwide, but the molecular mechanisms underlying *Fusarium* wilt resistance in cucumber remain unknown. To gain an improved understanding of the defense mechanisms elicited in response to Foc inoculation, RNA sequencing-based transcriptomic profiling of responses of the *Fusarium* wilt-resistant cucumber line 'Rijiecheng' at 0, 24, 48, 96, and 192 h after Foc inoculation was performed.

Results: We identified 4116 genes that were differentially expressed between 0 h and other time points after inoculation. All ethylene-related and pathogenesis-related genes from among the differentially expressed genes were filtered out. Real-time PCR analysis showed that ethylene-related genes were induced in response to Foc infection. Importantly, after Foc infection and exogenous application of ethephon, a donor of ethylene, these genes were highly expressed. In response to exogenous ethephon treatment in conjunction with Foc inoculation, the infection resistance of cucumber seedlings was enhanced and endogenous ethylene biosynthesis increased dramatically.

Conclusion: Collectively, ethylene signaling pathways play a positive role in regulating the defense response of cucumber to Foc infection. The results provide insight into the cucumber *Fusarium* wilt defense mechanisms and provide valuable information for breeding new cucumber cultivars with enhanced *Fusarium* wilt tolerance.

Background

Cucumber is among the most widely cultivated vegetables in the world. Given the continuous cropping systems used in cucumber production in China, cucumber fusarium wilt (FW) incidence is aggravated year-on-year. FW is a typical soil-borne disease caused by *Fusarium oxysporum* f. sp. *cucumerinum* (Foc) [1], which results in severe losses in global cucumber output [2]. Foc comprises four races, designated races 1, 2, 3, and 4, that are prevalent mainly in the United States, Israel, Japan, and China [3, 4, 5], respectively. Foc can survive in soil and seeds for many years or even decades [6]. Hyphae of this pathogen penetrate cucumber roots, then spread to the vascular tissues and occludes the xylem vessels, and also produces a toxin to kill cells, which leads to wilting of the leaves or even the entire plant, until plant death occurs several days or weeks after infection [7, 8]. FW is extremely difficult to treat and changes in the pathogenicity of Foc have led to the ineffectiveness of certain fungicides [9, 10]. Therefore, identification of disease resistance genes, augmentation of germplasm pools, and breeding FW-resistant cucumber cultivars [11] is an efficient strategy to control cucumber FW.

Increasing research attention is focused on the potential of functional genomics to study the host immune system to enhance disease protection [12]. The defense mechanisms induced in a plant upon exposure to a pathogen, including the network of cross-communicating signaling pathways, may halt pathogen infection. Ethylene (ET) is the principal mediator in plants of these signaling pathways [13, 14].

ET was first identified owing to its effects on fruit maturation, senescence, germination, cell elongation, and flowering [15, 16], and was subsequently determined to function as a modulator of the plant immune signaling network [17]. ET regulates the inducible plant defense responses [18]. Previous molecular studies indicate that in plants many diverse plant hormone-regulated pathways involved in resistance to biotic stresses are triggered through biosynthesis of effector proteins by the pathogen [19]. Certain plant hormones are essential for plant immunity [20]. The predominant symptom elicited by Foc infection of cucumber is wilting, which is caused by disturbance of the water balance [21, 22]. Abscisic acid (ABA) might play a crucial regulatory role in modifying stomatal behavior, which has a major impact on disturbance of water loss in cucumber [23]. Disruption of the transcription factor AtMYC2, which is a positive regulator of ABA signaling [24], induces expression of ethylene-responsive genes [25]. In Arabidopsis, ethylene response 1 (ETR1), which belongs to the ET receptor family, positively influences pathogen-mediated responses [26]. It is also known that ET signaling pathways play a crucial role in plant defense response [27]. When soybean was infected with Soybean mosaic virus in the shade, Zhang et al. [28] observed that ET might perform an important role in regulating the defense response. The genes ETR, EBF1, and ERF1 function as positive regulators in 'Guijiao 9', a resistant banana cultivar, in response to infection by *Fusarium oxysporum* f. sp. *cubense* tropical race 4, which activated ET signaling in the disease resistance response. Liu et al. [29] observed that ET signaling pathways might positively regulate the defense response of *Panax notoginseng* to *Fusarium solani*. However, the function and mechanism of ET in biotic stress response remains unclear and rarely reported, especially in cucumber.

In this research, we conducted a comprehensive and global analysis of the transcriptome during the defense response of cucumber to Foc infection using a high-throughput RNA sequencing (RNA-seq) approach, which is an efficient and powerful method for gene discovery. We aimed to identify differentially expressed genes (DEGs) and enriched pathways that contribute to disease resistance after inoculation of cucumber roots with spore suspensions at different time points. The findings provide insight into ET-related candidate genes and pathways associated with resistance to FW, which will contribute to broader knowledge of the resistance mechanism in cucumber to Foc.

Results

Quality analysis of RNA-seq data from cucumber roots infected by Foc

More than 100.75 Gb clean data were obtained from 15 RNA-seq libraries (Additional file1: Table S1; deposited in the NCBI SRA database under accession no. PRJNA472169). For each sample, the Q30 value was higher than 85.27%, indicating that the sequence data were of high quality. The number of reads mapped to the cucumber Chinese Long reference genome exceeded 29.54 million (70.64% of the total reads).

Identification of DEGs and expression profiles of ET-related genes

Using fold change ≥ 1 and false discovery rate (FDR) < 0.01 as cutoff, 4116 DEGs were identified. Among these genes, 1469 genes were upregulated and 2647 genes were downregulated (Additional file2: Figure

S1). Thirty-two ET-related DEGs were filtered out (Fig. 1). Of the filtered genes, group A consisted of ET-responsive transcription factors including 25 DEGs, and group B comprised 1-aminocyclopropane-1-carboxylate (ACC) oxidase, which catalyzes the oxidative cleavage of ACC to form ET. Group A was divided into several subgroups and the expression profiles of DEGs in each subgroup was similar. Quantitative real-time PCR (qPCR) analysis was used to verify the DEGs by selection of six DEGs that showed a high relative expression level and upregulated expression (Fig. 2). The expression trends were consistent across both qPCR and RNA-seq analysis.

ET affects growth of Foc

To clarify whether ET affected the growth of Foc, we cultured Foc on potato dextrose agar (PDA) medium supplemented with 1000 ppm ethephon. Mycelial growth of the Foc strain on PDA supplemented with sterile water (the control) was superior to that on PDA with ethephon (Fig. 3). The average diameter of mycelial colonies on PDA supplemented with sterile water was 74.33 mm, whereas that on PDA supplemented with ethephon was 43.67 mm. The color of mycelia was darker on PDA supplemented with ethephon than that on the control medium. These results indicated that ET could suppress the growth of Foc in the presence of 1000 ppm ethephon.

ET increases resistance of cucumber seedlings to Foc

To test the possibility that ET may enhance the resistance response in cucumber seedlings, cucumber seedlings at the second-true-leaf stage were inoculated with Foc and exogenous ethephon and grown for three weeks. The disease grade was numbered and disease index was calculated. Four days after inoculation, cotyledons of the Foc-sensitive cultivar 'Superina' cultured with Foc began to turn yellow, whereas the same symptoms of Superina seedlings cultured with Foc and ethephon were observed at 10 days after inoculation. Symptoms in seedlings of the Foc-resistant cultivar 'Rijiecheng' were observed at seven days after inoculation with Foc, and at 15 days after inoculation with Foc and exogenous ethephon. The disease indices of Superina and Rijiecheng under Foc inoculation were higher than those under Foc inoculation in conjunction with exogenous ethephon (Fig. 4C, D). Thus, the resistance of cucumber seedlings inoculated with Foc and ET was superior to that of seedlings inoculated only with Foc (Fig. 4A, B) in both a resistant cultivar (Rijiecheng) and a sensitive cultivar (Superina). These results indicated that ET increases the resistance of cucumber seedlings to Foc.

Expression profiles of ET-related genes in response to exogenous ET and Foc infection

Fourteen genes that showed high relative expression levels were selected to confirm the expression levels in response to exogenous ET using the qPCR method (Fig. 1). *Csa2G010390* and *Csa7G318990* encoded pathogenesis-related (PR) proteins, *Csa4G630010*, *Csa7G049230*, *Csa2G382540*, *Csa1G042290*, *Csa4G641590*, *Csa2G354000*, and *Csa6G318160* encoded ethylene-responsive transcription factors, *Csa7G405830* encoded an ethylene receptor, and *Csa3G135690*, *Csa6G511860*, *Csa3G135740*, and *Csa6G160180* encoded ACC oxidases. Six of the 14 genes were chosen as representative genes in the ET pathway for cucumber resistance to Foc. The expression level of the two PR genes improved significantly

in response to exogenous application of ET both in the Foc-resistant and -sensitive cultivars (Fig. 5). The expression level was higher in the Foc-resistant cultivar than that in the sensitive cultivar. Thus, the PR genes may be modulated by ET and the disease resistance of seedlings maybe improved simultaneously. *Csa3G135690* and *Csa6G160180*, which encoding ACC oxidases, and *Csa7G049230* and *Csa6G318160*, which encode ethylene-responsive transcription factors, were upregulated by exogenous ethephon and Foc infection (Fig. 5). The expression levels of the other genes shown in Additional file3: Figure S2. In short, after inoculation with Foc, these ET signaling pathway genes were highly expressed in response to exogenous ethephon. All the primers were shown in Additional file 4: Table S2.

Validation of endogenous ET of cucumber seedlings with exogenous ethephon and Foc infection

Given that the ET-related genes were upregulated by Foc infection and exogenous application of ethephon, we measured the endogenous ET biosynthesis of cucumber seedlings in response to Foc inoculation and exogenous ethephon (Fig. 6). The endogenous ethylene biosynthesis was higher in Rijeicheng than in Superina at 48 and 96 h in response to Foc infection alone. The endogenous ethylene concentration was about four-times higher at 48 h and twice as high at 96 h in Superina than in Rijeicheng in response to exogenous ethephon and Foc infection. Endogenous ET biosynthesis increased markedly after treatment with exogenous ethephon. These experiments indicated that Foc may induce the biosynthesis of endogenous ET in cucumber seedlings and exogenous ethephon may enhance this response.

Discussion

As a common soil-borne disease, FW presents a serious barrier to continuous cropping of cucumber [30]. There is currently no effective measure to control this disease except for breeding resistant cultivars. Exploration of genes associated with pathogen resistance in plants using diverse approaches is a major research focus. The evolution of resistance genes has been observed in numerous model plants [31, 32, 33]. Expression profiling of FW resistance genes in cucumber has not previously been reported. In the present study, we gained insight into the genes associated with resistance to Foc in cucumber based on RNA-seq analysis of the global transcriptome profile in cucumber at different time points after inoculation. Global data analyses can help in elucidating the mechanism and key genes that control resistance to FW in cucumber.

The whole-genome transcriptome analysis revealed that ET-responsive genes were distinctly induced in response to Foc infection. Thus, ET signaling was involved in the interaction between cucumber and Foc, and genes encoding ET receptors were differentially expressed before and after inoculation. In previous studies, ET signaling pathways under physiological stress or other abiotic stresses is often reported. The present experiments suggest that ET is a crucial factor in cucumber resistance to Foc. The role of ET in immune responses remains unknown for the most part. Similar findings were reported for *Medicago truncatula* infected by *Rhizoctonia solani* [34], which led to upregulation of ET signaling. ET receptors are downstream regulatory factors of the ET-signaling pathway [27], which regulate plant growth,

development, and many types of stress response [35]. In tomato, *Trichoderma harzianum* infection induced the expression of PR proteins, which were markers of the ET-dependent signaling pathways [30]. Similarly, two PR genes were upregulated by ET in the present study. PR genes perform a pioneering role in the diagnosis of plant immunity under pathogen challenge. Expression of genes that encode PR proteins appear to be upregulated rapidly in response to pathogen infection. PR-1 was detected in pathogen-stressed *Nicotiana tabacum* plants [36]. Nevertheless, the function of PR-1 may be assigned to other members of the PR protein family [37]. PR-1 encodes a small antimicrobial protein responsive to plant immune signaling, and plants that show enhanced PR-1 expression are strongly resistant to pathogenic oomycetes [37]. These PR proteins are an important component of plant resistance to pathogens [38]. Ohme-Takagi and Shinishi [39] reported that the GCC box of PR genes promoted the functions of an ET-responsive element [27]. Enhanced disease resistance 1 (EDR1) encodes a CTR1-like kinase [40], which is a negative regulator of ET responses [41, 42]. Tang et al. [43] considered that EDR1 may function in ET signaling to promote cell death.

Plant immune systems may also be activated by exogenous application of relevant molecules to modulate the defense response. In the present study, exogenous ethephon was applied to verify that ET may enhance the resistance of cucumber to *Foc*. Exogenous application of ACC, the precursor of ET, enhanced the resistance of NbALD1-transgenic plants of *Nicotiana benthamiana* to Turnip mosaic virus [44]. Liu et al. [29] first reported that exogenous application of methyl jasmonate increased the resistance of *Panax notoginseng* to *Fusarium solani*, which is an important causal agent of root rot disease. The defense response to rice bacterial blight disease may be enhanced by exogenous application of jasmonic acid [45]. Treatments with exogenous ethylene provide a safe and effective method for control of pear browning [46]. Phytohormones regulate internal as well as external environmental signals [47]. ET biosynthesis may be activated after infection by *Magnaporthe oryzae* in rice, and resistant cultivars accumulated higher amounts of ET than susceptible cultivars [48]. The present study revealed that endogenous ET biosynthesis is significantly elevated with exogenous application of ethephon.

The present transcriptome data provided an improved understanding of gene expression profiles in cucumber upon infection by *Foc*. The data provide a comprehensive overview of the functions and effects of genes involved in the cucumber defense response. ET-related genes were induced in response to *Foc* inoculation; ET-related and PR genes were highly expressed in response to exogenous ethephon application, and the disease resistance of cucumber seedlings was enhanced simultaneously. The present results provide a foundation for further discovery of gene functions in cucumber. Future research should focus on overexpression and knockout of candidate genes that enhance resistance to *Foc* and elucidate the molecular mechanisms of resistance in cucumber.

Conclusions

By means of a global transcriptome analysis, we identified 4116 genes that were differentially expressed between 0 h and 24, 48, 96, and 192 h after inoculation with *Foc*. In response to *Foc* infection and exogenous application of ethephon, ET-related and PR genes were confirmed to be highly expressed using

qPCR analysis. Exogenous ethephon treatment in conjunction with Foc inoculation enhanced the disease resistance of cucumber seedlings and endogenous ET biosynthesis was substantially increased. The present results illustrate that ET signaling pathways play a role in positively regulating the defense response of cucumber to Foc. The findings will be helpful for elucidating the cucumber FW defense mechanisms, and information on the candidate genes will enrich the breeding of new cucumber cultivars with enhanced FW tolerance.

Methods

Fungal culture and plant materials

The Foc fungal strain was isolated from the experimental field of Yangzhou University, Jiangsu Province, China, and propagated on PDA plates at 28 °C for 4 days, then cultured in potato dextrose broth on a shaker at 180 rpm at 28 °C for 3 days. The spore suspension was diluted to 1×10^6 spores per milliliter with sterile distilled water.

Seedlings of cucumber 'Rijiecheng', a moderately Foc-resistant cultivar, were grown in 36-well plates filled with an aseptic organic substrate (contents of total nitrogen, phosphorus, and potassium = 40-60 g/kg, content of humus \geq 350 g/kg, pH = 6.5-7.5) at 25 °C/18 °C day/night temperatures with a 16 h/8 h photoperiod. Seedlings were infected with Foc by irrigation of the roots with a fungal spore suspension (5 mL per seedling) at the second-true-leaf stage. Cucumber roots were sampled at 0, 24, 48, 96, and 192 h after inoculation with three biological replicates.

RNA extraction, cDNA library construction, Illumina sequencing, and analysis of sequence reads

Total RNA was isolated using the TaKaRa MiniBEST Plant RNA Extraction Kit (TaKaRa, China). The RNA concentration was measured using the Qubit RNA Assay Kit with a Qubit[®] 2.0 fluorometer (Life Technologies, USA). One microgram of RNA was used as input for the RNA sample preparations. The mRNA was purified from the total RNA using poly-T oligo-attached magnetic beads. Fragmentation was carried out using divalent cations under an elevated temperature in NEBNext[®] First Strand Synthesis Reaction Buffer (5 \times). The first-strand cDNA was synthesized using the mRNA fragments as templates. To select cDNA fragments of preferentially 200–250 bp in length, the library fragments were purified with the AMPure XP system (Beckman Coulter, USA). Eligible cDNAs were selected for PCR amplification, which was performed with Phusion[®] High-Fidelity DNA polymerase, Universal PCR primers, and Index (X) Primer. The PCR products were purified using the AMPure XP system and library quality was assessed with an Agilent Bioanalyzer 2100 system.

Primary cDNA produced using the Illumina HiSeq 2500 platform by BioMarker Technologies (Beijing, China) were termed raw reads. Clean reads were obtained by removing reads containing the adapter, reads containing poly-N, and low-quality reads. In addition, Q20, Q30, GC-content and sequence duplication values of the clean reads were calculated. The trimmed reads were aligned to the cucumber Chinese Long reference genome v2 [35], which was retrieved from

<ftp://www.icugi.org/pub/genome/cucumber/>. All downstream analyses were based on the clean, high-quality data. Quantification of gene expression levels was estimated as fragments per kilobase of transcript per million fragments mapped (FPKM) [49,50] using Cufflinks (version: 2.1.1).

Identification of DEGs and validation of RNA-seq by qPCR

We divided the data into four groups by comparing the data at 0 h with that at the other sampling time points. The analysis of DEGs for the four groups was performed using the DESeq R package (1.10.1). DESeq provides statistical routines for determining differential expression in digital gene expression data using a model based on the negative binomial distribution. The *P* values were adjusted using the Benjamini–Hochberg approach for controlling the FDR. Genes with an adjusted FDR < 0.01 identified by DESeq and \log_2 FPKM (fold change) ≥ 1 were considered to be differentially expressed.

Total RNA of each condition was isolated using the MiniBEST Plant RNA Extraction Kit (TaKaRa, China), then dissolved with UltraPure™ DNase/RNase-free Distilled Water (Invitrogen, USA). Total RNA was reverse transcribed using the PrimeScript™ RT Reagent Kit with gDNA Eraser (TaKaRa, China). Primer sequences for qPCR were designed using Primer Premier 5. The qPCR analysis was performed using SYBR® Premix Ex Taq™ (TaKaRa, China) in accordance with the manufacturer's instructions. SYBR Green PCR cycling was performed on an Iqtm5 Multicolor qPCR detection system (Bio-Rad, USA) using 20 μ L samples with the following temperature program: 95 °C for 3 min, followed by 39 cycles of 95 °C for 10 s, 60 °C for 20 s, and 72 °C for 20 s, then a melting curve analysis was performed. The relative quantification of gene expression was calculated and normalized to the tubulin alpha chain gene (*Csa4G000580*). Three biological replicates from each condition were used for qPCR.

Evaluation of exogenous ethephon on Foc and cucumber seedlings

To ascertain the effect of ET on Foc growth, exogenous ethephon (a donor of ET) was incorporated in PDA medium. To 18 mL PDA medium was added either 2 mL ethephon (final concentration 1000 ppm) or 2 mL sterile water (as the control). Foc was cultured on the PDA medium at 25 °C for 5 days and the colony diameter was measured three times.

To examine the effect of ET on seedling resistance to Foc, cucumber seedlings were inoculated with Foc (spore concentration 10^6 conidia/mL) using the root irrigation method and half of the seedlings were also sprayed with exogenous ethephon solution (concentration 10 ppm). After three weeks, disease grades were recorded and the disease index was calculated using the following formula:

See Formula 1 in the supplemental files.

Each experimental group included at least 15 individual seedlings and three biological replicates were treated. Mock inoculated seedlings were used as the control.

Measurement of endogenous ET biosynthesis

The two cucumber inbred lines, 'Rijiecheng' and 'Superina', which are resistant and sensitive to Foc, respectively, were used in this experiment. Seedlings at the second-true-leaf stage were inoculated by irrigation with Foc (spore concentration 10^6 conidia/mL) and sprayed with exogenous ethephon solution (concentration 10 ppm). Endogenous ET biosynthesis by the root portion of the seedlings was measured at 48 and 96 h after treatment. The root portion was placed in a sealed 30 mL ampoule for 24 h in the dark. For analysis of ET production, gas samples (1 mL) were collected using a syringe and injected into a gas chromatograph (Agilent 7890A, USA) fitted with a flame ionization and electron capture detectors [51]. Experiments were conducted with three biological replicates.

Declarations

Ethics approval and consent to participate

In this manuscript, I have obtained permission from all the authors and I declare that the material and the manuscript have not and will not be published in whole and in part elsewhere, all authors have been personally involved in the substantive work. I testify to the accuracy of the above on behalf of all the authors.

Consent for publication

No applicable.

Availability of data and materials

All transcriptome data associated with this study have been submitted to the NCBI, and can be found using accession number PRJNA472169.

Competing interests

All authors declare that they have no competing interests.

Funding

This research was supported by the National Natural Science Foundation of China (31902015), Natural Science Foundation of Jiangsu Province (BK20190887), Modern Agricultural Industry Technology System of Jiangsu Province (JATS [2018] 308), Jiangsu Agricultural Science and Technology Innovation Fund (CX(19)3029), and Creation of Major New Agricultural Varieties in Jiangsu Province (PZCZ201720).

Authors' contributions

XC and JX contributed to the conceptualization and design of the experiments. JD and YW performed the experiments and JD wrote the manuscript. QX prepared the cucumber seedlings and performed RNA extraction. All authors read and approved the final manuscript.

Acknowledgements

We thank Robert McKenzie, PhD, from Liwen Bianji, Edanz Group China (www.liwenbianji.cn/ac), for editing the English text of a draft of this manuscript.

Author details

¹ School of horticulture and plant protection, Yangzhou University, Wenhui Road, Yangzhou 225009, Jiangsu Province, China.

² State Key Laboratory of Vegetable Germplasm Innovation, Tianjin 300384, China.

Abbreviations

FW: Fusarium wilt; Foc: *Fusarium oxysporum* f. sp. *cucumerinum*; ET: ethylene; ABA: abscisic acid; ETR1: ethylene response 1; RNA-seq: RNA sequencing; DEGs: differentially expressed genes; FDR: false discovery rate; ACC: 1-aminocyclopropane-1-carboxylate; qPCR: quantitative real-time PCR; PDA: potato dextrose agar; PR: pathogenesis-related

References

1. Korolev N, Katan J, Katan T. Vegetative Compatibility Groups of *Verticillium dahliae* in Israel: Their Distribution and Association with Pathogenicity. *Phytopathology*. 2000;90:529-536.
2. Du NS. Effect of vinegar residue compost amendments on cucumber growth and fusarium wilt. [Environmental Science and Pollution Research](#). 2015;22:199133-19141.
3. Jenkins JR, and Wehner TC. Occurrence of *Fusarium oxysporum* f. sp. *cucumerinum* on greenhouse-grown *Cucumis sativus* seed stocks in North Carolina. *Plant Disease*. 1983;67:1024-1025.
4. Mao AJ, Zhang F, Zhang LR. Analysis on the Inheritance of Resistance to Fusarium Wilt Race 4 and Cucumber Scab and Their Linkage in Cucumber WIS2757. *Scientia Agriculture Sinica*. 2008;41:3382-3388.
5. Netzer D, Niego S, Galun E. A dominant gene conferring resistance to Fusarium wilt in cucumber. *Phytopathology*. 1977;67:525-527.
6. Snyder WC, Hansen HN. The species concept in *Fusarium*. [American Journal of Botany](#). 1940;27:64-67.
7. Harding MW, Marques LLR, Howard RJ, Olson ME. Can filamentous fungi form biofilms? *Trends in Microbiology*. 2009;17:475-480.
8. Li PQ, Pu XM, Feng BZ, Yang QY, Shen HF, Zhang JX, Lin BR. *FocVel1* influences asexual production, filamentous growth, biofilm formation, and virulence *Fusarium oxysporum* f. sp. *cucumerinum*. *Frontiers in Plant Science*. 2015;6:312.

9. Chen LH, Huang XQ, Zhang FG, Zhao DK, Yang XM. Application of *Trichoderma harzianum* SQR-T037 bio-organic fertiliser significantly controls *Fusarium* wilt and affects the microbial communities of continuously cropped soil of cucumber. *Journal of the Science of Food and Agriculture*. 2012;92:2465-2470.
10. Huang X, Shi DZ, Sun FQ, Lu HH, Liu JJ, Wang WX. Efficacy of sludge and manure compost amendments against *Fusarium* wilt of cucumber. *Environmental Science and Pollution Research*. 2018;19:3895-3905.
11. Zhang SS, Raza W, Yang XM, Hu J, Huang QW, Xu YC, Liu XH, Ran W, Shen QR. Control of *Fusarium* wilt disease of cucumber plants with the application of a bioorganic fertilizer. *Biology and Fertility of Soils*. 2008;44:1073-1080
12. Pétriacq P, López A, Luna E. Fruit decay to diseases: can induced resistance and priming help? *Plants*. 2018;7:1-16.
13. Pandey D, Rajendran SRCK, Gaur M, Sajeesh PK, Kumar A. Plant defense signaling and responses against necrotrophic fungal pathogens. *Journal of Plant Growth Regulation*. 2016;35:1159-1174.
14. De Vos M, Van Oosten VR, Van Poecke RMP, Van Pelt JA, Pozo MJ, Mueller MJ, Buchala AJ, Métraux JP, Van Loon LC, Dicke M, Pieterse CMJ. Signal signature and transcriptome changes of *Arabidopsis* during pathogen and insect attack. *Molecular Plant Microbe Interaction*. 2005;18:923-937.
15. Bleeker, A.B., Kende, H. Ethylene: a gaseous signal molecule in plants. *Annual Review of Cell and Developmental*. 2000;1:1-18.
16. Xu K, Xu X, Fukao T, Canlas P, Maghirang-Rodriguez R, Heuer S, Heuer S, Ismail AM, Bailey-Serres J, Ronald PC, Mackill DJ. Sub1A is an ethylene-response-factor-like gene that confers submergence tolerance to rice. *Nature*. 2006;442:705-708.
17. Van Loon LC, Geraats BPJ, Linthorst HJM. Ethylene as a modulator of disease resistance in plants. *Trends in Plant Science*. 2006;11:184-191.
18. Denancé N, Sánchez-Vallet A, Gofner D, Molina A. Disease resistance or growth: the role of plant hormones in balancing immune responses and fitness costs. *Frontiers in Plant Science*. 2013;4:155.
19. Shores M, Harman GE, Mastouri F. Induced Systemic Resistance and Plant Responses to Fungal Biocontrol Agents. *Annual Review of Phytopathology*. 2010;48:21-43.
20. Pieterse CMJ, Leon-Reyes A, Van der Ent S, Van Wees SCM. Networking by small-molecule hormones in plant immunity. *Nature Chemical Biology*. 2005;5:308-316.
21. Wang M, Ling N, Dong X, Zhu YY, Shen QR, Guo SW. Thermographic visualization of leaf response in cucumber plants infected with the soil-borne pathogen *Fusarium oxysporum* f. sp. *cucumerinum*. *Plant Physiology and Biochemistry*. 2012;61:153-161.
22. Wang M, Sun YM, Sun GM, Liu XK, Zhai LC, Shen QR. Water balance altered in cucumber plants infected with *Fusarium oxysporum* f. sp. *cucumerinum*. *Scientific Reports*. 2015;5:7722.
23. Rodrigues O, Reshetnyak G, Grondin A, Saijo Y, Leonhardt N, Maurel C, Verdoucq L. Aquaporins facilitate hydrogen peroxide entry into guard cells to mediate ABA and pathogen-triggered stomatal

- closure. Proceedings of the National Academy of Sciences of the United States of America. 2017;114:9200-9205.
24. Abe H, Urao T, Ito T, Seki M, Shinozaki K, Yamaguchi-Shinozaki K. Arabidopsis AtMYC2 (bHLH) and AtMYB2 (MYB) function as transcriptional activators in abscisic acid signaling. *The Plant Cell*. 2003;15:63-78.
 25. Anderson JP, Badruzsaufari E, Schenk PM, Manners JM, Desmond OJ, Ehlert C, Maclean DJ, Ebert PR, Kazan K. Antagonistic interaction between abscisic acid and jasmonate-ethylene signaling pathways modulates defense gene expression and disease resistance in Arabidopsis. *Plant Cell*. 2004;16:3460-3479.
 26. Dong JP, Qi XH, Xu Q, Chen XH. Identification of *Fusarium oxysporum* f. sp. *cucumerinum* and Resistance Analysis in Cucumber. *Molecular Plant Breeding*. 2017;15:3648-3653.
 27. Wang FF, Cui XK, Sun Y, Dong CH. Ethylene signaling and regulation in plant growth and stress responses. *Plant Cell Reports*. 2013;32:1099-1109.
 28. Zhang L, Shang J, Wang WM, Du JB, Li K, Wu XL, Yu L, Liu CY, Khaskheli MI, Yang WY. Comparison of Transcriptome Differences in Soybean Response to Soybean Mosaic Virus under Normal Light and in the Shade. *Viruses*. 2019;11:1-15.
 29. Liu DQ, Zhao Q, Cui XM, Chen R, Li X, Qiu BB, Ge F. A transcriptome analysis uncovers *Panax notoginseng* resistance to *Fusarium solani* induced by methyl jasmonate. *Genes & Genomics*. 2019;41:1383-1396.
 30. Wang YY, Wu FZ, Liu SW. Allelopathic effects of root exudates from wheat, oat and soybean on seed germination and growth of cucumber. *Allelopathy Journal*. 2009;24:103-112.
 31. Leonetti P, Zonno MC, Molinari S, Altomare C. Induction of SA-signaling pathway and ethylene biosynthesis in *Trichoderma harzianum*-treated tomato plants after infection of the root-knot nematode *Meloidogyne incognita*. *Plant Cell Reports*. 2017;36:621-631.
 32. Wu J, Kim SG, Kang KY, Kim JG, Park SR, Gupta R, Kim YH, Wang YM, Kim ST. Overexpression of a Pathogenesis-Related Protein 10 Enhances Biotic and Abiotic Stress Tolerance in Rice Plant *Pathology Journal*. 2016;32:552-562.
 33. Finkina EI, Melnikova DN, Bogdanov IV, Ovchinnikova TV. Plant Pathogenesis-Related Proteins PR-10 and PR-14 as Components of Innate Immunity System and Ubiquitous Allergens. *Current Medicinal Chemistry*. 2017;24:1772-1787.
 34. Brotman Y, Normantovich M, Goldenberg Z, Zvirin Z, Kovalski I, Stovbun N, Doniger T, Bolger AM, Troadec C, Bendahmane A, Cohen R, Katzir N, Pitrat M, Dogimont C, Perl-Treves R. Dual Resistance of Melon to *Fusarium oxysporum* Races 0 and 2 and to Papaya ring-spot virus is Controlled by a Pair of Head-to-Head-Oriented NB-LRR Genes of Unusual Architecture. *Molecular Plant*. 2013;6:235-238.
 35. Chen T, Liu J, Lei G, Liu YF, Li ZG, Tao JJ, Hao YJ, Cao YR, Lin Q, Zhang WK, Ma B, Chen SY, Zhang JS. Effects of tobacco ethylene receptor mutations on receptor kinase activity, plant growth and stress responses. *Plant Cell Physiology*. 2009;50:1636-1650.

36. Gianinazzi S, Martin, C & Vallée, JC. Hypersensibilité aux virus, température et protéines solubles chez le *Nicotiana Xanthi* n.c: Apparition de nouvelles macromolécules lors de la répression de la synthèse virale. *Acta Crystallographica*. 1970;270:2383-2386.
37. Gamir J, Darwiche R, Van't Hof P, Choudhary V, Stumpe M, Sxhneiter R, Mauch F. The sterol-binding activity of PATHOGENESIS-RELATED PROTEIN 1 reveals the mode of action of an antimicrobial protein. *Plant Journal*. 2017;89:502-509.
38. Jones JDG, Vance RE, Dangl JL. Intracellular innate immune surveillance devices in plants and animals. *Science*. 2016;354:1117-1126..
39. Ohme-Takagi M, Shinshi H. Ethylene-inducible DNA binding proteins that interact with an ethylene-responsive element. *Plant Cell*. 1995;7:173-182.
40. Frye CA, Tang D, Innes RW. Negative regulation of defense responses in plants by a conserved MAPKK kinase. *Proceedings of the National Academy Sciences*. 2001;98:373-378
41. Cao H, Glazebrook J, Clarke JD, Volko S, Dong XN. The Arabidopsis NPR1 gene that controls systemic acquired resistance encodes a novel protein containing ankyrin repeats. *Cell*. 1997;88:57-63.
42. Kieber JJ, Rothenberg M, Roman G, Feldmann KA, Ecker JR. CTR1, a negative regulator of the ethylene response pathway in Arabidopsis, encodes a member of the Raf family of protein kinases. *Cell*. 1993;72:427-441.
43. Tang D, Christiansen KM, Innes RW. Regulation of Plant Disease Resistance, Stress Responses, Cell Death and Ethylene Signaling in Arabidopsis by the EDR1 Protein Kinase. *Plant Physiology*. 2005;138:1018-1026.
44. Wang S, Han KL, Peng JJ, Zhao JP, Jiang LL, Lu YW, Zheng HY, Lin L, Chen JP, Yan F. *NbALD1* mediates resistance to turnip mosaic virus by regulating the accumulation of salicylic acid and the ethylene pathway in *Nicotiana benthamiana*. *Molecular Plant Pathology*. 2019;20:990–1004.
45. Taniguchi S, Hosokawa-Shinonaga Y, Tamaoki D, Yamada S, Akimitsu K, Gomi K. Jasmonate induction of the monoterpene linalool confers resistance to rice bacterial blight and its biosynthesis is regulated by JAZ protein in rice. *Plant Cell & Environment*. 2014;37:451–461.
46. Ma YR, Yang MN, Wang JJ, Jiang CZ, Wang QG. Application of Exogenous Ethylene Inhibits Postharvest Peel Browning of 'Huangguan' Pear. *Frontiers in Plant Science*. 2016;7:2029.
47. Kazan K. Diverse roles of jasmonates and ethylene in abiotic stress tolerance. *Trends in Plant Science*. 2015;20:219-229.
48. Yang C, Li W, Cao JD, Meng FW, Yu YQ, Huang JK, Jiang L, Liu MX, Zhang ZG, Chen XW, Miyamoto K, Yamane H, Zhang JS, Chen SY, Liu J. Activation of ethylene signaling pathways enhances disease resistance by regulating ROS and phytoalexin production in rice. *The Plant Journal*. 2017;89:338-353.
49. Florea L, Song L, Salzberg SL. Thousands of exon skipping events differentiate among splicing patterns in sixteen human tissues. *F1000Res*. 2013;2:188.

50. Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, Baren MJ, Salzberg SL, Wold BJ, Pachter L. Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nature Biotechnology*. 2010;28:511-515.
51. Giné-Bordonaba J, Echeverria G, Ubach D, Aguiló-Aguayo I, López ML, Larrigaudière C. Biochemical and physiological changes during fruit development and ripening of two sweet cherry varieties with different levels of cracking tolerance. *Plant Physiol Biochem*. 2017;111,216-225.

Figures




Figure 1

Hierarchical clustering of 32 ethylene-related differentially expressed genes induced by *Fusarium oxysporum* f. sp. *cucumerinum* inoculation. Fold-difference was designated as fragments per kilobase of transcript per million fragments mapped. The maximum value was '1' for each gene. The darker the shade of red, the higher the expression level. 0 h, 24 h, 48 h, 96 h, and 192 h indicate the sampling time points after inoculation.




Figure 2

Changes in relative expression level of six differentially expressed genes (DEGs) from among 32 ethylene-related DEGs analyzed by qPCR and RNA-seq. The left vertical axis and bars indicate the relative expression level determined by qPCR, the right vertical axis and solid blue line indicate the FPKM value determined by RNA-seq. 0 h, 24 h, 48 h, 96 h, and 192 h indicate time points after *Fusarium oxysporum* f. sp. *cucumerinum* (*Foc*) inoculation. R, 'Rijiecheng', a *Foc*-resistant cucumber line, inoculated with *Foc*; CK, 'Rijiecheng' inoculated with sterile water. At least three biological replicates were performed in each experiment group. * P < 0.05, ** P < 0.01.




Figure 3

In vitro phenotypic differences between the *Fusarium oxysporum* f. sp. *cucumerinum* strain cultured on PDA medium supplemented with sterile water (A) and 1000 ppm ethephon (B).



Figure 4

(A, B) Phenotypic differences of cucumber seedlings treated with mock solutions, inoculated with a *Fusarium oxysporum* f. sp. *cucumerinum* strain (Foc), and inoculated with Foc and exogenous ethephon (Foc+ET). (C, D) Disease index for the phenotypes of the cucumber seedlings. R, 'Rijiecheng' (Foc-resistant line); S, 'Superina', (Foc-sensitive line).




Figure 5

Expression profiles of six candidate ethylene-related genes in cucumber at 0 h, 48 h, and 96 h after inoculation with *Fusarium oxysporum* f. sp. *cucumerinum* (Foc) and spray treatment with exogenous ethephon (ET). R, 'Rijiecheng' (Foc-resistant line); S, 'Superina' (Foc-sensitive line).




Figure 6

Endogenous ethylene production by the root of cucumber seedlings at 0, 48, and 96 h after inoculation with *Fusarium oxysporum* f. sp. *cucumerinum* (Foc) and spray treatment with exogenous ethephon. R, 'Rijiecheng' (Foc-resistant line); S, 'Superina' (Foc-sensitive line).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementaryinformation.docx](#)
- [formula1.PNG](#)