

Impact of Cotadutide Drug on Obese Individuals with Type 2 Diabetes Mellitus: A Systematic Review and Meta-analysis

Mahmoud M Ali

Al-Azhar University

Ahmed Hafez

Menoufia University

Mahmoud Shaban Abdelgalil

Ain-shams University

Mohammed Tarek Hasan

Al-Azhar University

Mohammed Magdy El-Ghannam

German University in Cairo

Osama M Ghogar

Al-Azhar University

Asmaa Ahmed Elrashedy

Kafr El-Shaikh University

Mohamed Abd-ElGawad (✉ mohammed.mahmod87@gmail.com)

Fayoum University

Research Article

Keywords: Type 2 diabetes mellitus, Cotadutide, Glucagon-like peptide 1, Weight loss

Posted Date: January 27th, 2022

DOI: <https://doi.org/10.21203/rs.3.rs-1236048/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: The food and drug administration approved many drugs to treat diabetes mellitus, but those drugs do not have a noticeable effect on weight management. Recently, glucagon-like peptide 1 agonist known as Cotadutide was estimated to serve as a potent drug in treating type 2 diabetes by reducing blood glucose levels and body weight indices. This study aimed to explore the safety and efficacy of Cotadutide as a treatment for type 2 diabetes individuals.

Methods: A comprehensive literature search was done on different databases, including PubMed, Scopus, Web of Science, and Cochrane Library to capture all relevant articles using an established search strategy. The inclusion criteria were randomized controlled trials that assessed the safety and efficacy of Cotadutide versus placebo or any anti-diabetes drugs in obese and overweight people with type 2 diabetes mellitus. Data analysis was accomplished using Revman 5.4 software.

Results: We found 663 relevant articles. From which nine studies were included and subjected to qualitative analysis and eight for quantitative analysis. The pooled effect showed that Cotadutide was better than placebo in reducing body weight (Mean difference (MD) = 3.31, $p > 0.00001$), glycated hemoglobin (HbA_{1c}) (MD = -0.68, $p > 0.00001$), glucose area under the plasma concentration curve (AUC [0-4h]) (MD = 30.15, $p > 0.00001$), and fasting plasma glucose over time (mg/dl) (MD = 31.31, $p > 0.00001$).

Conclusion: Cotadutide is safe and effective in reducing blood glucose levels and body weight in individuals with type 2 diabetes.

Study protocol registration: The study protocol was registered on PROSPERO (CRD: CRD42021257670).

Background

Type 2 diabetes mellitus is one of the most common endocrine disorders worldwide, according to the International Diabetes Federation (IDF) its prevalence has surged rapidly to include more than 400 million individuals over the past three decades (1). Type 2 diabetes mellitus is a long-term disease characterized by chronic insulin resistance and hyperglycemia that increases over time, resulting in increasing of insulin resistance leading to weight gain (2,3). Therefore, reducing body weight will prevent more insulin resistance and better control of the body weight condition.

Many medications with different mechanisms are available to control type 2 diabetes mellitus as (a) metformin which acts through various trajectories to inhibit gluconeogenesis and reduce the level of lipopolysaccharide, (b) insulin secretagogues, (c) alpha-glucosidase inhibitors, (d) dipeptidyl peptidase 4 inhibitors, and (e) sodium-glucose co-transporter-2 inhibitor (4). However, none of them is significant in reducing body weight at doses approved for blood glucose reduction. Therefore, weight loss remains an unmet medical need for these people (5).

Glucagon-like peptide-1 (GLP-1) receptor agonist's therapy known as Cotadutide seems to be effective in glycemic control and weight loss. The impact of GLP-1 drugs varies depending on the pharmacokinetic profile (6). Lorenz M et al 2013 showed that short-acting GLP-1 receptor agonists (lixisenatide) at a dose of 20 µg daily lowers postprandial hyperglycemia excursions in individuals with type 2 diabetes mellitus, probably caused by the continuous slowing of stomach emptying (7). In the same way, J van Can et al 2014 investigated the effects of long-acting GLP-1 receptor agonists (liraglutide) on gastric emptying and the result indicated that liraglutide at 3 mg significantly delays the gastric emptying (8). In addition, Daniel R et al reported in 2020 that lixisenatide reduced the gastric emptying rate more than liraglutide (9).

Based on the above-mentioned data, GLP-1 receptor agonists are useful in treating individuals with type 2 diabetes and obesity by controlling hyperglycemia and delaying stomach emptying. They also help people lose weight by reducing the appetite and increasing energy expenditure by optimizing metabolic reactions such as amino acid catabolism, and fatty acid oxidation (10).

Many studies have investigated the effect of Cotadutide (GLP-1 receptor agonist) on type 2 diabetes mellitus. In this study, we aim to summarize, review, and analyze those studies to understand the safety and efficacy profiles of this new medication in controlling type 2 diabetes mellitus and its effect on weight reduction.

Methods

Study design and registration:

This meta-analysis was conducted following the Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA) guideline and Cochrane Handbook of Systematic Reviews of Intervention (11,12). The study protocol was registered on PROSPERO (CRD: CRD42021257670).

Literature search:

PubMed, Cochrane Central Register of Controlled Trials (CENTRAL), Scopus, and Web of Science were searched for articles conducted from January 1979 to June 2021. The search strategy used can be found in **supplementary file 1**.

Eligibility criteria and studies selection:

The inclusion criteria included Randomized Controlled Trials (RCT) evaluating the efficacy and safety of the drug Cotadutide on men or women aged 18 to 65 years with controlled type 2 diabetes and a BMI between 22 kg/m² and 40 kg/m².

We excluded studies with unreliable or insufficient data for extraction. Reviews, book chapters, thesis, editorial, letters, conference papers, and non-English studies. Animal or In vitro studies, cohort, case-control, non-clinical studies, literature reviews, and meta-analysis were excluded.

Two independent authors screened the articles retrieved from the four electronic databases by title, abstract, and full text on an excel sheet for eligibility. Another independent author resolved any disagreements between the other two authors.

Quality assessment:

Cochrane risk-of-bias tool for randomized trials (RoB 2) was applied to assess the quality of the selected RCTs. The Rob2 tool consists of six domains: randomization process, deviations from the intended interventions, missing outcome data, measurement of the outcome, selection of the reported result, and other biases. The evaluators' responses were categorized as yes, probably yes, probably no, no, and no information. Following that, all disputes were discussed and resolved.

Data extraction and study outcomes:

Two independent reviewers extracted data in a pre-defined excel sheet. The excel sheet items were categorized as a summary of the included trials' key features, characteristics of the participants, and Cotadutide safety and efficacy outcomes. Any disagreements were solved by a discussion between the reviewers.

Outcome definition:

Treatment efficacy was assessed by frequency of positive Anti-drug antibodies to Cotadutide, Percent Change from Baseline in Body Weight, Change from Baseline in Glycated Hemoglobin (HbA_{1c}), Mean Percentage Change from Baseline in Glucose Area Under the Plasma Concentration Curve (AUC [0-4h]) as Measured by (MMTT). The safety outcomes included Treatment-Emergent Adverse Events (TEAEs), and Treatment-Emergent Serious Adverse Events (TESAEs).

Data synthesis and assessment of heterogeneity:

We performed all statistical analyses using Revman software Version 5.4.1. The present meta-analysis estimated the pooled risk ratio (RR) for dichotomous data, mean difference (MD) for continuous data with 95% confidence intervals (CI). The significance point was set at P-value less than 0.05.

We assessed the heterogeneity using the I-square and p-value. The analysis was considered heterogeneous if it had a p-value less than 0.05 or an I-square less than 50%. A random-effect model was applied if heterogeneity was detected and a leave one out test was performed to determine which study was causing the heterogeneity (13).

Results

Data collection and study selection:

Our search retrieved 655 records from PubMed, Scopus, Web of Science, and Cochrane library. There were 75 duplicates. After title and abstract screening, we eliminated 557 records. Afterward, we screened 23 full texts for eligibility. Finally, nine records were included in our study: four published clinical trials and five registered protocols from Clinicaltrials.gov, and only eight studies were included in the meta-analysis (Figure. 1).

The total sample size for this meta-analysis was 1258 participants. 262 persons received a placebo and 886 participants received Cotadutide. There were no concomitant treatment modalities except in two studies. In NCT03235050, participants in all study groups received metformin tablets and a separate group was treated with liraglutide to compare it with Cotadutide and placebo. Moreover, during the

treatment period of the study NCT03444584, participants were on metformin and dapagliflozin as well. **Table 1** elucidates the full summary of the included studies. The baseline characteristics of the participants are illustrated in **Table 2**.

Quality assessment results:

The risk of bias summary is illustrated in figures 2 and 3. Regarding the Randomization process bias, all the studies were of low risk in terms of the randomization process except for NCT03550378, which was judged as some concerns because there was inadequate information about the allocation concealment, randomization, and baseline balance.

Regarding the intended interventions bias, most of the included trials had a low risk of bias in terms of deviations from the intended interventions except for NCT03645421 and NCT03745937, which were judged as some concerns. This is because there was no information about the statistical analysis used to estimate the effect of assignments in both of them despite blinding the personnel.

Regarding the missing outcome data bias, most of the included trials had a low risk of bias in terms of the missing outcome data due to applying the intention to treat analysis. We judged NCT03645421 and P.D. Ambery et al. as high risk of bias because the authors applied as-treated analysis (14).

Regarding the measurement outcome bias, we judged the risk of bias in the measurement of the outcome as low risk of bias in most of the studies due to blinding of all outcome assessors and using appropriate methods in measuring the outcomes. We judged NCT03645421 and NCT03550378 as some concerns due to the lack of information about blinding the outcome assessor.

For the selection of the reported results bias, the risk of bias due to the selection of the reported results ranged between low and some concerns. We judged all the registered protocols as some concerns because there is no published data yet to compare it with the protocols. The published studies (6,14–16) were of low risk as all outcomes mentioned in the results were present in the protocols.

For other sourced of bias, we judged almost all the studies as high risk in terms of other potential sources of bias as most of them are registered protocols without any published papers yet. Parker et al (15) stated the lack of statistical power to draw inferences between cohorts and the absence of validated questionnaires as a limitation in their study and so we judged it as having a high risk of bias. Accordingly, Ambery et al (6) had a relatively small population size which we considered as high-risk potential. Only (14,16) showed no other potential sources of bias.

Efficacy endpoints:

Percentage decrease in body weight: The pooled effect estimate of five studies favored Cotadutide 300 mcg over placebo (MD = 3.31, $p < 0.00001$). Pooled data were homogenous under a fixed effect model ($p = 0.80$, $I^2 = 0\%$); Figure. 4.

Decrease in glycated hemoglobin (HbA_{1c}): The pooled effect estimate of five studies showed that Cotadutide is significantly better than placebo (MD = 0.68, 95% CI [0.58, 0.79], $p < 0.00001$). Pooled data were homogenous under a fixed effect model ($p = 0.05$, $I^2 = 55\%$); Figure. 5.

Percentage decrease in glucose area under the plasma concentration curve (AUC [0-4h]): The pooled effect estimate of six studies favored Cotadutide 300 mcg over placebo (MD = 30.15, $p < 0.00001$). Pooled data were heterogeneous under a random effect model and the heterogeneity was best resolved by leaving out NCT03596177 ($p = 0.08$, $I^2 = 48\%$); Figure. 6.

Decrease from baseline in fasting plasma glucose over time (mg/dl): The pooled effect estimate of four studies favored Cotadutide over the placebo (MD = 31.31, $p < 0.00001$). Pooled data were heterogeneous under a random effect model and the heterogeneity was best resolved by leaving out NCT03645421 ($p < 0.17$, $I^2 = 40\%$); Figure. 7.

Anti-drug antibodies (ADA): Nahra et al reported a statistically significant increase in the number of participants with ADA in the Cotadutide group over the placebo (155 out of 256 in the Cotadutide group and three out of 256 in the placebo group); NCT03444584, NCT03550378, and NCT03596177 reported non-significant results on the number of participants having ADA. NCT03444584 reported 2 out of 24 and 1 out of 24 in the Cotadutide group and the placebo group, respectively. In NCT03550378, two out of 21 participants in the Cotadutide group had ADA in comparison to the 20 persons on placebo in which none of them developed ADA. In NCT03596177, three out of 14 and zero out of seven participants experienced ADA in the Cotadutide group and the placebo group, respectively.

Safety endpoints:

Treatment-emergent adverse events (TEAEs): The pooled effect estimate of six studies showed a statistically significant increased risk of TEAEs in the Cotadutide group compared to placebo (RR = 1.40, $p = 0.0007$). Pooled data were homogenous under a fixed-effect model ($p = 0.23$, $I^2 = 26\%$); Figure. 8.

Treatment-emergent serious adverse events (TESAEs): We didn't do a meta-analysis for this outcome because none of the participants suffered any TESAEs in three out of six studies in both the Cotadutide and the placebo group. Only three studies reported some participants having TESAEs and they are relatively very low. In terms of TESAEs, NCT03550378 reported two out of 21 persons in the Cotadutide group and two out of 20 in the placebo group. On the other hand, NCT03596177 reported two persons out of 18 and zero out of seven in the Cotadutide group and the placebo group, respectively. In the MAD portion of the study by P. Amberly et al, they reported one out of seven participants having TESAEs in the Cotadutide 200 mcg group and none out of 19 participants in the placebo group (6).

Discussion

This meta-analysis on 1258 participants with type 2 diabetes revealed that efficacy outcomes, including body weight, fasting blood glucose, HbA_{1c} , and AUC [0-4h], were significantly better in people receiving Cotadutide treatment than placebo. The number of participants with positive ADA to Cotadutide was high but without a significant difference compared to placebo. Furthermore, no significant difference was observed between the Cotadutide group and placebo in TESAEs. Hence, Cotadutide is safe and effective as a hypoglycemic drug in people with controlled type 2 diabetes.

In ten years, more than half of individuals with type 2 diabetes mellitus switch from oral monotherapy (usually Metformin) to insulin therapy to control their blood glucose levels (17). Multiple combination therapies are routinely used before insulin is initiated. Insulin use causes weight gain, which can exceed 6 kg 20 in the first year after starting insulin medication (18). The overall gain in weight can cause an increase in insulin resistance which is associated with high blood pressure, dyslipidemia, and a high risk of cardiovascular mortalities and morbidities such as non-fatal myocardial infarction or stroke, both before and after diagnosis of diabetes (19,20). Pre-clinical findings further suggest that the balance of activities at GLP-1 receptors and glucagon receptors was appropriate for both weight reduction and glycemic management (21). These activities are supposed to be balanced by stimulating insulin release mediated by glucose, delayed gastric emptying, and enhanced oxidation of fatty acids (22,23). GLP-1, including Cotadutide (MEDI0382) and glucagon receptor dual agonists, may have central impacts on appetite as glucagon receptor agonist has been found in animal and human studies to increase energy expenditure (24).

Cotadutide (5-300 μ g) corrected the glucose levels to the normal range in phase one of the first human trial which was conducted on healthy volunteers, with a pharmacokinetic profile that included once-daily treatment (6). Similar to these previous findings (6,14), in phase 2a, Cotadutide (100-300 μ g) significantly lowered blood glucose levels and body weight indices in overweight or obese Japanese people with type 2 diabetes throughout a 48-day treatment period compared to placebo. Parker et al. found a substantial decline in glucose AUC (0-4h) by -21.52% with up titrated Cotadutide (50-300 μ g) in comparison to $+6.32\%$ with placebo. Similarly, a decline in body weight was reported by -3.41% versus -0.08% for Cotadutide versus placebo, respectively (15). Nevertheless, in different study (25), with a lower BMI (26.3-28.8 kg/m²) than Parker et al (31.5 kg/m²) (15), blood glucose and weight reduction with Cotadutide 300 μ g remained significant compared to placebo at -37.86% versus $+2.45\%$ and -3.34% versus -0.82% , respectively.

Cotadutide therapy reduced body weight in a dose-dependent approach, and the highest reductions occurred at 300 μ g. Moreover, Cotadutide improved fasting plasma glucose, fructosamine, HbA_{1c} , percentage of time in hyperglycemia, insulin secretion, and resistance. After 6 weeks of Cotadutide medication, significant decreases in HbA_{1c} were found, with efficacy remaining constant (25).

Cotadutide has also been known to significantly reduce hepatic glycogen and steatosis, as well as having a beneficial effect on hepatic inflammation and fibrosis markers (25,26). The decrease in hepatic glycogen contrasts with what would be expected from a GLP-1 monogonist, which would cause glycogen accumulation and exhibit glucagon receptor interaction (27). Furthermore, the degree of liver fat loss with Cotadutide (39% reduction) was comparable to that shown in a small study of women three months following bariatric surgery (42% reduction) (28). This decrease in liver fat found with Cotadutide was larger than would be expected from weight loss alone—for example, in individuals with documented non-alcoholic fatty liver disease, a 5% decrease in BMI results in a 25.5 percent relative decline in liver fat (29).

MEDI0382 had a linear pharmacokinetic profile in the first human study on healthy volunteers (phase 1), and no participants tested positive for ADA (14). In a previous study, participants were given Cotadutide for a year and had a significant ADA incidence. Only 16% of participants acquired ADAs over a titer of 80, at which point the influence on pharmacokinetics was around two times higher than the population average (16) ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT04019561) identifier NCT04019561).

In the Harmony Outcomes study, albiglutide outperformed placebo in terms of serious adverse cardiovascular problems in people with type 2 diabetes and cardiovascular morbidities with a hazard ratio of 0.78, which implies that GLP-1 agonists can improve cardiovascular outcomes according to these data (30). Due to GLP-1 and glucagon receptor agonism on the heart and vascular system, an increase in heart rate was expected. The rise in heart rate by 6.8 beats per minute observed with Cotadutide was not significantly greater than that seen with the GLP-1 receptor agonist liraglutide which increased by 6 to 9 beats from baseline. Furthermore, the drop in blood pressure was comparable to that seen with GLP-1 receptor agonists (22,31).

Cotadutide plasma concentrations increased in agreement with the anticipated dose titration at all dose levels, with no TEAEs linked to immunogenicity observed (6). In this study, Cotadutide had a higher rate of gastrointestinal co-morbidities such as nausea and vomiting compared to placebo. This outcome is also seen with the GLP-1 receptor mono-agonists (32,33). In addition, Cotadutide's safety profile was equivalent to that of previous global trials (6,14), with a greater incidence of gastrointestinal adverse events.

To lower the gastrointestinal adverse events associated with Cotadutide 300 µg, dose escalation was required upon which a phase 2 study in obese type 2 diabetes participants reported that Cotadutide was effective and well-tolerated with starting doses of 50 µg for 7 days, then gradual dose escalation up to 300 µg (15,34). Despite causing more gastrointestinal upset than placebo, escalated dosages of Cotadutide of up to 300 µg which were given once daily were generally tolerated because the symptoms were mild or moderate in severity (14).

This study comprehensively evaluated the efficacy and safety of Cotadutide for people with type 2 diabetes. Nine RCTs were included in the study, resulting in a valuable evidence level. The included trials varied from low to high quality. The majority of the identified heterogeneity was resolved. Our analysis also has certain limitations, including the small sample size and the small number of included studies. Publication bias could not be detected due to the small number of included studies. The short follow-up period and lack of placebo were the major drawbacks of the study. Most of the included studies were protocols with published results, not articles. Cotadutide medication should also be evaluated for its effects on stomach emptying, energy intake, and energy expenditure in larger studies.

Conclusion

Over a short dosage period, Cotadutide provided considerable metabolic benefits to overweight and obese participants with type 2 diabetes. Cotadutide's safety and pharmacokinetics allow once-daily administration of dosages less than 150 µg, which can be followed by dose escalation. Cotadutide's promising impacts on glycemic control, body weight, and liver fat suggest that it might be a helpful agent for type 2 diabetes individuals with longer-term treatment.

Abbreviations

Mean difference (MD)

Glycated hemoglobin (HbA_{1c})

International Diabetes Federation (IDF)

Glucagon-like peptide-1 (GLP-1)

Preferred Reporting Items for Systematic Reviews and Meta-Analysis (PRISMA)

Cochrane Central Register of Controlled Trials (CENTRAL)

Randomized Controlled Trials (RCT)

Cochrane risk-of-bias tool for randomized trials (RoB 2)

Treatment-Emergent Adverse Events (TEAEs)

Risk Ratio (RR)

Confidence Intervals (CI)

Percentage decrease in glucose area under the plasma concentration curve (AUC [0-4h])

Anti-drug antibodies (ADA)

Treatment-emergent adverse events (TEAEs)

Treatment-emergent serious adverse events (TESAEs)

Declarations

Data Availability:

All data generated or analyzed during this study are included in this published article or in the data repositories listed in References.

Conflicts of interest: Dr. Abd-ElGawad has nothing to disclose. All the authors also declare no conflict of interest.

Funding: not applicable

Ethics approval: not applicable

Consent to participate: not applicable

Consent for publication: not applicable

Availability of data and material: All data generated or analyzed during this study are included in this published article or in the data repositories listed in References.

Code availability: not applicable

Authors contributions:

Mahmoud M Ali leads the team, performed the search strategy and data collection step, solved any conflict in the screening phase, performed the meta-analysis part, and solved any conflict in the quality assessment part, took part in the data extraction phase.

Ahmed Hafez took part in the screening process, data extraction, and meta-analysis in addition to writing the results section and edited the whole manuscript.

Mahmoud Shaban took part in the screening process, data extraction, quality assessment, and drafting the tables.

Mohammed Tarek Hasan took part in data extraction, and meta-analysis.

Mohammed Magdy El-Ghannam took part in quality assessment and writing the introduction section.

Osama M Ghogar took part in study selection and writing the methods section.

Asmaa Ahmed Elrashedy wrote the discussion section and edited the manuscript.

Mohamed Abd-ElGawad supervised the authors in all steps and performed peer-review

All authors reviewed the final manuscript.

References

1. Malone JI, Hansen BC. Does obesity cause type 2 diabetes mellitus (T2DM)? Or is it the opposite? *Pediatr Diabetes*. 2019;20(1).
2. Stumvoll M, Goldstein BJ, van Haeften TW. Type 2 diabetes: principles of pathogenesis and therapy. *Lancet*. 2005;365(9467).
3. Petersen MC, Vatner DF, Shulman GI. Regulation of hepatic glucose metabolism in health and disease. *Nat Rev Endocrinol*. 2017;13(10).
4. Marín-Peñalver JJ, Martín-Timón I, Sevillano-Collantes C, Del Cañizo-Gómez FJ. Update on the treatment of type 2 diabetes mellitus. *World J Diabetes*. 2016; 7 (17): 354–95.
5. Baretic M, Troskot R. How to fight obesity with antidiabetic drugs: targeting gut or kidney? *Minerva Endocrinol*. 2014;40(1):71–83.
6. Ambery P, Parker VE, Stumvoll M, Posch MG, Heise T, Plum-Moerschel L, et al. MEDI0382, a GLP-1 and glucagon receptor dual agonist, in obese or overweight patients with type 2 diabetes: a randomised, controlled, double-blind, ascending dose and phase 2a study. *Lancet* [Internet]. 2018 Jun;391(10140):2607–18. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0140673618307268>

7. Lorenz M, Pfeiffer C, Steinstraer A, Becker RHA, Rutten H, Ruus P, et al. Effects of lixisenatide once daily on gastric emptying in type 2 diabetes – Relationship to postprandial glycemia. *Regul Pept.* 2013;185.
8. van Can J, Sloth B, Jensen CB, Flint A, Blaak EE, Saris WHM. Effects of the once-daily GLP-1 analog liraglutide on gastric emptying, glycemic parameters, appetite and energy metabolism in obese, non-diabetic adults. *Int J Obes.* 2014;38(6).
9. Quast DR, Schenker N, Menge BA, Nauck MA, Kapitza C, Meier JJ. Effects of Lixisenatide Versus Liraglutide (Short- and Long-Acting GLP-1 Receptor Agonists) on Esophageal and Gastric Function in Patients With Type 2 Diabetes. *Diabetes Care.* 2020;43(9).
10. Jiang G, Zhang BB. Glucagon and regulation of glucose metabolism. *Am J Physiol Metab.* 2003;284(4):E671–8.
11. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al. Cochrane handbook for systematic reviews of interventions [Internet]. Higgins JPT, Thomas J, Chandler J, Cumpston M, Li T, Page MJ, et al., editors. *Cochrane Handbook for Systematic Reviews of Interventions.* Wiley; 2019 [cited 2020 Feb 18]. 1–694 p. Available from: <https://onlinelibrary.wiley.com/doi/book/10.1002/9781119536604>
12. Moher D, Liberati A, Tetzlaff J, Altman DG, Altman D, Antes G, et al. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement [Internet]. Vol. 6, *PLoS Medicine.* Public Library of Science; 2009 [cited 2020 Feb 18]. p. e1000097. Available from: <https://dx.plos.org/10.1371/journal.pmed.1000097>
13. Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *Bmj.* 2003;327(7414):557–60.
14. Ambery PD, Klammt S, Posch MG, Petrone M, Pu W, Rondinone C, et al. MEDI0382, a GLP-1/glucagon receptor dual agonist, meets safety and tolerability endpoints in a single-dose, healthy-subject, randomized, Phase 1 study. *Br J Clin Pharmacol* [Internet]. 2018 Oct;84(10):2325–35. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/bcp.13688>
15. Parker VER, Robertson D, Wang T, Hornigold DC, Petrone M, Cooper AT, et al. Efficacy, Safety, and Mechanistic Insights of Cotadutide, a Dual Receptor Glucagon-Like Peptide-1 and Glucagon Agonist. *J Clin Endocrinol Metab* [Internet]. 2020 Mar 1;105(3):803–20. Available from: <https://academic.oup.com/jcem/article/105/3/803/5586886>
16. Nahra R, Wang T, Gadde KM, Oscarsson J, Stumvoll M, Jermutus L, et al. Effects of Cotadutide on Metabolic and Hepatic Parameters in Adults With Overweight or Obesity and Type 2 Diabetes: A 54-Week Randomized Phase 2b Study. *Diabetes Care* [Internet]. 2021 Jun;44(6):1433–42. Available from: <http://care.diabetesjournals.org/lookup/doi/10.2337/dc20-2151>
17. Turner RC. Glycemic Control With Diet, Sulfonylurea, Metformin, or Insulin in Patients With Type 2 Diabetes Mellitus<SUBTITLE>Progressive Requirement for Multiple Therapies (UKPDS 49)</SUBTITLE>. *JAMA* [Internet]. 1999 Jun 2;281(21):2005. Available from: <http://jama.jamanetwork.com/article.aspx?doi=10.1001/jama.281.21.2005>
18. Douek IF, Allen SE, Ewings P, Gale EAM, Bingley PJ. Continuing metformin when starting insulin in patients with Type 2 diabetes: a double-blind randomized placebo-controlled trial. *Diabet Med* [Internet]. 2005 May;22(5):634–40. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/j.1464-5491.2005.01475.x>
19. Wilcox R, Kupfer S, Erdmann E. Effects of pioglitazone on major adverse cardiovascular events in high-risk patients with type 2 diabetes: Results from PROspective pioglitAzone Clinical Trial In macro Vascular Events (PROactive 10). *Am Heart J.* 2008;155(4).
20. Purnell JQ, Hokanson JE, Marcovina SM, Steffes MW, Cleary PA, Brunzell JD. Effect of Excessive Weight Gain With Intensive Therapy of Type 1 Diabetes on Lipid Levels and Blood Pressure. *JAMA.* 1998;280(2).
21. Hernandez AF, Green JB, Janmohamed S, D’Agostino RB, Granger CB, Jones NP, et al. Albiglutide and cardiovascular outcomes in patients with type 2 diabetes and cardiovascular disease (Harmony Outcomes): a double-blind, randomised placebo-controlled trial. *Lancet* [Internet]. 2018 Oct;392(10157):1519–29. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S014067361832261X>
22. Meier JJ, Gethmann A, Gotze O, Gallwitz B, Holst JJ, Schmidt WE, et al. Glucagon-like peptide 1 abolishes the postprandial rise in triglyceride concentrations and lowers levels of non-esterified fatty acids in humans. *Diabetologia* [Internet]. 2006 Mar 31;49(3):452–8. Available from: <http://link.springer.com/10.1007/s00125-005-0126-y>
23. Lynch AM, Pathak N, Flatt YE, Gault VA, O’Harte FPM, Irwin N, et al. Comparison of stability, cellular, glucose-lowering and appetite suppressing effects of oxyntomodulin analogues modified at the N-terminus. *Eur J Pharmacol* [Internet]. 2014 Nov;743:69–78. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0014299914006621>
24. Habegger KM, Stemmer K, Cheng C, Muller TD, Heppner KM, Ottaway N, et al. Fibroblast Growth Factor 21 Mediates Specific Glucagon Actions. *Diabetes* [Internet]. 2013 May 1;62(5):1453–63. Available from: <http://diabetes.diabetesjournals.org/cgi/doi/10.2337/db12-1116>
25. Asano M, Sekikawa A, Kim H, Gasser RA, Robertson D, Petrone M, et al. Pharmacokinetics, safety, tolerability and efficacy of cotadutide, a glucagon-like peptide-1 and glucagon receptor dual agonist, in phase <scp>1 and 2</scp> trials in overweight or obese participants of <scp>A</scp> sian descent with or without type 2 dia. *Diabetes, Obes Metab* [Internet]. 2021 Aug 19;23(8):1859–67. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/dom.14412>

26. Guzman CB, Zhang XM, Liu R, Regev A, Shankar S, Garhyan P, et al. Treatment with LY2409021, a glucagon receptor antagonist, increases liver fat in patients with type 2 diabetes. *Diabetes, Obes Metab* [Internet]. 2017 Nov;19(11):1521–8. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/dom.12958>
27. Boland ML, Laker RC, Mather K, Nawrocki A, Oldham S, Boland BB, et al. Resolution of NASH and hepatic fibrosis by the GLP-1R and GCGR dual-agonist cotadutide via modulating mitochondrial function and lipogenesis. *Nat Metab* [Internet]. 2020 May 21;2(5):413–31. Available from: <http://www.nature.com/articles/s42255-020-0209-6>
28. Heath ML, Kow L, Slavotinek JP, Valentine R, Toouli J, Thompson CH. Abdominal adiposity and liver fat content 3 and 12 months after gastric banding surgery. *Metabolism* [Internet]. 2009 Jun;58(6):753–8. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0026049509000092>
29. Patel NS, Doycheva I, Peterson MR, Hooker J, Kisselva T, Schnabl B, et al. Effect of Weight Loss on Magnetic Resonance Imaging Estimation of Liver Fat and Volume in Patients With Nonalcoholic Steatohepatitis. *Clin Gastroenterol Hepatol* [Internet]. 2015 Mar;13(3):561-568.e1. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1542356514013160>
30. Green JB, Hernandez AF, D'Agostino RB, Granger CB, Janmohamed S, Jones NP, et al. Harmony Outcomes: A randomized, double-blind, placebo-controlled trial of the effect of albiglutide on major cardiovascular events in patients with type 2 diabetes mellitus—Rationale, design, and baseline characteristics. *Am Heart J* [Internet]. 2018 Sep;203:30–8. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S000287031830187X>
31. Kumarathurai P, Anholm C, Fabricius-Bjerre A, Nielsen OW, Kristiansen O, Madsbad S, et al. Effects of the glucagon-like peptide-1 receptor agonist liraglutide on 24-h ambulatory blood pressure in patients with type 2 diabetes and stable coronary artery disease. *J Hypertens* [Internet]. 2017 May;35(5):1070–8. Available from: <https://journals.lww.com/00004872-201705000-00024>
32. Davies MJ, Bergenstal R, Bode B, Kushner RF, Lewin A, Skjøth TV, et al. Efficacy of Liraglutide for Weight Loss Among Patients With Type 2 Diabetes. *JAMA* [Internet]. 2015 Aug 18;314(7):687. Available from: <http://jama.jamanetwork.com/article.aspx?doi=10.1001/jama.2015.9676>
33. Nauck MA, Petrie JR, Sesti G, Mannucci E, Courrèges J-P, Lindegaard ML, et al. A Phase 2, Randomized, Dose-Finding Study of the Novel Once-Weekly Human GLP-1 Analog, Semaglutide, Compared With Placebo and Open-Label Liraglutide in Patients With Type 2 Diabetes. *Diabetes Care* [Internet]. 2015 Sep 10;dc150165. Available from:
34. ROBERTSON D, PARKER VE, AMBERY P, PETRONE M, WANG T, HEISE T, et al. 988-P: MEDI0382, an Oxyntomodulin-Like Peptide with Targeted GLP-1/Glucagon Receptor Activity, Promotes a Dose-Dependent Increase in Gastric Emptying Time. *Diabetes* [Internet]. 2019 Jun 4;68(Supplement 1):988-P. Available from: <http://diabetes.diabetesjournals.org/lookup/doi/10.2337/db19-988-P>

Tables

Table 1. Summary of included studies key features

Study ID	Title	Study design, country, and timing	Criteria	Sample size	MEDI0382 treatment regimen	Control group	study duration
P. Ambery et al. (2018)	MEDI0382, a GLP-1 and glucagon receptor dual agonist, in obese or overweight patients with type 2 diabetes: a randomized, controlled, double-blind, ascending dose and phase 2a study	RCT Germany, Dec 9, 2015, and Feb 24, 2017	<ul style="list-style-type: none"> Men or women aged 18–65 years with controlled T2DM HbA1c levels of 6.5–8.5% at screening BMI between 27 kg/m² and 40 kg/m². Individuals who received metformin monotherapy of 500 mg or more within three months before screening or received an adjunct to metformin were eligible after a 4-week washout period. 	112 patients (67 for MEDI0382, and 45 for placebo)	MEDI0382 100 mcg N=6 MEDI0382 150 mcg N=6 MEDI0382 200 mcg N=7 MEDI0382 200 mcg N=25 MEDI0382 300 mcg N=12 MEDI0382 300 mcg N=12	N= 45 Participants received placebo matched to MEDI0382 dose	3 years
P. D. Ambery (2018)	MEDI0382, a GLP-1/glucagon receptor dual agonist, meets safety and tolerability endpoints in a single-dose, healthy-subject, randomized, Phase 1 study	RCT Germany, 26 February 2015 and 12 August 2015,	<ul style="list-style-type: none"> Healthy volunteers, male or female aged 18–45 years, are chosen approximately to a population of T2DM patients minimizing the chance for concomitant medical conditions. BMI between ≥ 22 kg.m⁻² and ≤ 30 kg.m⁻², Bodyweight ≥ 70 kg. Individuals were required to have venous access suitable for multiple cannulations. 	48 patients (36 for MEDI0382, and 12 for placebo)	MEDI0382 5 µg N=6 MEDI0382 10 µg N=6 MEDI0382 30 µg N=6 MEDI0382 100 µg N=6 MEDI0382 300 µg N=6 MEDI0382 150 µg N=6	N= 12 Participants received placebo matched to MEDI0382 dose	3 years
Rajaa Nahra (2021)	A Study to Evaluate the Efficacy and Safety of MEDI0382 in the Treatment of Overweight and Obese Subjects with Type 2 Diabetes	RCT, Bulgaria, Canada, Czechia, Germany, Mexico, Russian Federation, Slovakia, United States, 2 August 2017 and	<ul style="list-style-type: none"> Male and female subjects aged ≥ 18 years with T2DM BMI ≥ 25 kg/m² HbA1c range of 7.0% to 10.5% Individuals Treated with metformin ≥ 1500 mg/day or more for at least two months before screening or using adjuvant 	834 patients (612 for MEDI0382 ,110 for Liraglutide and 112 for placebo)	MEDI0382 100 mcg N= 100 MEDI0382 200 mcg N= 256 MEDI0382 300 mcg N= 256	N= 112 Participants received placebo / Metformin tablets, total daily dose of ≥ 1500 mg (unless only tolerated at a lower dose)	2 Years

		14 June 2019	medication for up to 2 weeks in the two months before screening is acceptable.				
NCT03745937	A Study to Evaluate the Safety and Tolerability of MEDI0382 in Overweight and Obese Subjects With Type 2 Diabetes Mellitus	RCT, Germany	<ul style="list-style-type: none"> Participants aged 18 to 74 years with T2DM BMI between 27 and 35 kg/m² HbA1c range of 6.5% to 8.5% Individuals Treated with metformin monotherapy where no significant dose change (increase or decrease \geq 500 mg/day) has occurred in the three months before screening. 	20 patients (18 for MEDI0382 and 2 for placebo)	Participants received SC dose of MEDI0382 up-titrated weekly once daily up to 8 weeks during the up-titration period and thereafter once daily in 3-week TEP.	N=2 Participants received placebo matched to MEDI0382 dose	4.5 months
NCT03645421	Safety and Tolerability Study of MEDI0382 in Japanese Pre-obese or Obese Subjects With Type 2 Diabetes	RCT, Japan	<ul style="list-style-type: none"> Individuals diagnosed with T2DM. HbA1c range of 7.0% to 10.5% Individuals with drug naïve at Visit 1. BMI within the range of 24 - 40 kg/m² 	60 patients (54 for MEDI0382 and 16 for placebo)	MEDI0382 100 mcg N= 15 MEDI0382 200 mcg N= 15 MEDI0382 300 mcg N= 15	N=16 Participants received placebo matched to MEDI0382 dose	5 months
NCT03596177	A Study to Evaluate the Effect of MEDI0382 on Energy Balance in Overweight and Obese Subjects With Type 2 Diabetes Mellitus	RCT, United Kingdom	<ul style="list-style-type: none"> Participants aged \geq 30 and \leq 75 years with T2DM Body Mass Index $>$ 28 and \leq 40 kg/m² HbA1c \leq 8.0% Individuals treated with metformin, with or without another adjuvant drug (increase or decrease $>$ 50%) occurred three months before screening. For the participant on dual therapy, a 4-week washout of the non-metformin therapy will be performed before Visit. 	28 patients (19 for MEDI0382 and 9 for placebo)	MEDI0382 titrated up to 300 μ g N= 19	N= 9 Participants received placebo matched to MEDI0382 dose	1 year and 1 month
NCT03444584	Study of MEDI0382 in Combination With Dapagliflozin and Metformin in Overweight/Obese	RCT, Germany, and Hungary	<ul style="list-style-type: none"> Male and female participants aged \geq 18 years with T2DM. BMI between 25 kg/m² and 40 	49 patients (25 for MEDI0382 and 24 for placebo)	MEDI0382 titrated up to 300 μ g N=25	N=24 Participants received placebo matched to	7 months

	Subject With Type 2 Diabetes		<ul style="list-style-type: none"> kg/m². HA1c ranges between 7.0% and 10.0%. Individuals treated with metformin monotherapy (MTD > 1 g) at least eight weeks or treated with stable, oral doses of dapagliflozin 10 mg and metformin (MTD > 1 g) for at least three months before screening. For the participant on dual therapy in addition to metformin, 28 days washout period will be performed before the initial screening. 		MEDI0382 dose		
Parker (2020)	Efficacy, safety, and mechanistic insights of cotadutide a dual receptor glucagon-like peptide-1 and glucagon agonist	RCT, Germany	<ul style="list-style-type: none"> Patients aged ≥18 years had T2DM taking metformin monotherapy, HbA1c ranged from 6.5–8.5% BMI of 27–40 kg/m² Individuals receiving metformin monotherapy were eligible if no significant dose changes (increase or decrease ≥500 mg/day) occurred in the three months before screening. Patients receiving adjuncts to metformin were allowed after a 4-week washout period. 	65 patients (46 for MEDI0382 and 19 for placebo)	MEDI0382 titrated up to 300 µg N=46	N=19 Participants received placebo matched to MEDI0382 dose	6 months
NCT03550378	A Study to Look at the Effect MEDI0382 Has on Blood Sugar in People with Type 2 Diabetes and Kidney Problems and Also to Check That MEDI0382 is Well Tolerated	RCT, Germany, United Kingdom	<ul style="list-style-type: none"> Patients aged ≥ 18 and < 85 years at the screening with T2DM managed with any insulin or oral therapy combination where no significant dose changes of oral therapy of more than 50% have occurred in the three months before screening. Body mass index (BMI) between 25 and 45 kg/m² Haemoglobin A1c (HbA1c) range of 6.5 % to 10.5% eGFR ≥ 30 and < 60 mL/min/1.73 m². 	41 patients (21 for MEDI0382 and 20 for placebo)	MEDI0382 titrated up to 300 µg N= 21	N=20 Participants received placebo matched to MEDI0382 dose	7 months

- Approximately 16 participants (40%) are required to have a screening eGFR ≥ 30 and < 45 mL/min/1.73 m², and at least 16 participants (40%) are required to have screening eGFR ≥ 45 and < 60 mL/min/1.73 m².

Table 1. Summary of the included studies. RCT: Randomized controlled trial. N: number. RAI: radioactive iodine therapy. MBq: megabecquerel. mCi: millicurie.

Table 2. Baseline characteristics of enrolled patients in each included study. Data are expressed as mean and standard deviation (SD) or frequency and percentage.

Study ID	Groups	Number of patients	Age mean \pm SD	Males (%)	Body mass index (BMI) (kg/m ²)	Weight (kg)	Race				
							White	Black	Asian	American Indian or Alaska Native	
P. Ambery et al 2018 (MAD study)	MEDI0382 300 mcg	11	54.8 \pm 6.8	7 (64%)	33.2 \pm 4.2	99.7 \pm 16.7	11 (100%)	0	0	0	
	placebo	19	57.7 \pm 6	12 (63%)	31.2 \pm 3.1	88.5 \pm 11.4	19 (100%)	0	0	0	
P. D. Ambery 2018	MEDI0382 300 mcg	6	28.3 \pm 5.9	6 (100%)	27.4 \pm 2.8	-	6 (100%)	0	0	0	
	placebo	12	32.8 \pm 9.1	12 (100%)	25.7 \pm 2.4	-	11 (91.7%)	0	0	0	
Rajaa Nahra 2021	MEDI0382 300 mcg	256	56.3 \pm 10.2	127 (94.6%)	-	-	252 (98.4%)	3 (1.2%)	1 (0.4%)	0	
	placebo	112	57.3 \pm 9.5	57 (50.9%)	-	-	107 (95.5%)	0	1 (0.9%)	3 (2.7%)	
NCT03645421	MEDI0382 300 mcg	15	57.5 \pm 9.2	13 (86.7%)	-	-	0	0	15 (100%)	0	
	placebo	16	60 \pm 8.6	13 (81%)	-	-	0	0	16 (100%)	0	
NCT03596177	MEDI0382 300 mcg	19	59.5 \pm 8.4	18 (94.7%)	-	-	18 (94.7%)	0	0	0	
	placebo	9	65.2 \pm 7.2	7 (77.8%)	-	-	8 (88.9%)	1 (11.1%)	0	0	
NCT03444584	MEDI0382 300 mcg	25	61 \pm 8.2	15 (60%)	-	-	25 (100%)	0	0	0	
	placebo	24	58.4 \pm 10	12 (50%)	-	-	24 (100%)	0	0	0	
Parker 2020	Cohort 1	MEDI0382 300 mcg	26	58.7 \pm 8.5	19 (73%)	31.5 \pm 3.5	95.6 \pm 17.2	25 (96%)	0	0	0
		placebo	13	-	9 (69%)	31.6 \pm 3.8	93.8 \pm 21	13 (100%)	0	0	0
	Cohort 2	MEDI0382 300 mcg	20	61.9 \pm 6	10 (50%)	31.1 \pm 3.5	92.4 \pm 8.7	20 (100%)	0	0	0
		placebo	6	-	5 (83%)	31.2 \pm 3.5	93.7 \pm 9.6	6 (100%)	0	0	0
NCT03550378	MEDI0382 300 mcg	21	71.1 \pm 7.4	12 (57.1%)	-	-	20 (95.2%)	0	0	0	
	placebo	20	70.9 \pm 4.7	9 (45%)	-	-	20 (100%)	0	0	0	

Figures




Figure 1

Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA).

Figure 2

Risk of bias graph for randomized controlled trials using Excel tool to implement Rob2.




Figure 3

Risk of bias summary for randomized controlled trials using Excel tool to implement Rob2.

Figure 4

Percentage decrease in body weight plot.

Figure 5

Decrease from baseline in Glycated Haemoglobin (HbA_{1c}) plot.

Figure 6

percentage decrease in glucose area under the plasma concentration curve (AUC [0-4h]) plot.

Figure 7

Change from Baseline in Fasting Plasma Glucose over Time (mg/dl) plot.

Figure 8

Treatment-Emergent Adverse Events (TEAEs) plot.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AbdElGawadSupplementaryfile1.docx](#)